

**NON-TRIVIAL DEFORMATION OF
AN ENTIRE FUNCTION $abz + e^{bz} + c$**

Shunsuke Morosawa* and Masahiko Taniguchi

Abstract. In this note, we consider the dynamics of the family of transcendental entire functions $f(z; a, b, c) = abz + e^{bz} + c$. In particular, while studying the case when they have a Baker domains, we investigate the area of Julia sets and the Hausdorff convergence of Julia sets.

Consider the family

$$f(z; a, b, c) = abz + e^{bz} + c.$$

These $f(z; a, b, c)$ with $a, b \neq 0$ are structurally infinite [7], for they have infinitely many critical values. For the sake of simplicity, we assume that a, b and c are real.

Theorem 1. *If $ab \geq 1$, $a > 0$, $c \leq -1$ and*

$$a \log a - a + c < 0,$$

then $f = f(z; a, b, c)$ has a Baker domain $B = B_{a,b,c}$, which is completely invariant.

In particular, the Julia set is the boundary of B .

Proof. We set $D_0 = \{z | \operatorname{Re} z < 0\}$. Elementary calculation shows that D_0 is mapped by f to a proper subdomain in D_0 , and hence D_0 is invariant. The Fatou component B containing D_0 is a Baker domain, for if $z \in D_0$, then

$$\operatorname{Re} f(z) < ab \operatorname{Re} z.$$

Received June 13, 2000.

Communicated by P. Y. Wu.

2001 *Mathematics Subject Classification*: 30D05, 37F50.

Key words and phrases: Transcendental entire function, Baker domain, Julia set, Hausdorff convergence.

*Partially supported by Grant-in-Aid for Scientific Research 09640199.

Also the line $L_k = \{z = x + iy \mid x \geq 0, y = (2k+1)\pi/b\}$ for $k \in \mathbb{Z}$ is mapped by f into D_0 , since the assumptions imply that

$$\operatorname{Re}(abz + e^{bz} + c) = abx - e^{bx} + c < 0$$

for every $z \in L_k$.

Since every preimage of D_0 by f contains either D_0 or some L_k , it intersects with D_0 , which shows that B is completely invariant. ■

The set D_0 in the proof is called a *fundamental set* of the Baker domain B (see [1]).

The function $f = f(z; a, b, c)$ as above has a completely invariant Baker domain B containing the left-half plane and the rays L_k . Since, the set of all the critical points

$$\left\{ z_m = \frac{\log a + (2m+1)\pi i}{b} \right\} \subset D_0 \cup \{L_k\},$$

the orbits of every critical point have the imaginary parts tending to $-\infty$. Hence, a theorem of Hinkkanen, Krauskopf and Kriete [2] (essentially due to Bergweiler [3]) gives the following

Theorem 2. *For such an $f(z; a, b, c)$ as in Theorem 1,*

$$F(f) = B.$$

Proof. First, the boundary of the Baker domain B has a positive Euclidean distance, say $\sigma > 0$, from the post-critical set C^+ , consisting of the forward orbits of all critical points of f .

On the other hand, since B contains all L_k , every other Fatou component U has a bounded distance, say η , from the boundary of B . Then the theorem of Hinkkanen, Krauskopf and Kriete gives the assertion.

But for the sake of convenience, we include a proof, following their preprint. Suppose that there were a Fatou component U , which should be either a wandering domain or a Baker domain disjoint from B . Fix a point $\zeta \in U$, and set $\zeta_n = f^n(\zeta)$. Then the disk D_n with center ζ_n and radius

$$r_n = \sigma + d(\zeta_n, B)$$

satisfies $D_n \cap C^+ = \emptyset$. Then there is a univalent branch ϕ_n of $(f^n)^{-1}$ which sends ζ_n to ζ , and a point $\xi_n \in J(f)$ such that

$$|\zeta_n - \xi_n| = d(\zeta_n, B)$$

for every n . Since functions

$$g_n(z) = \phi_n(r_n z + \zeta_n)$$

on $\{z \mid |z| < 1\}$ map 0 to ζ and the images are disjoint from C^+ , $\{g_n\}$ is a normal family. So assume that g_n converges to g uniformly on compact sets. Then the preimages w_n of $\xi'_n = \phi_n(\xi_n)$ by g_n satisfy

$$1 > \frac{\eta}{\sigma + \eta} > |w_n| > d$$

with some $d > 0$ by Schwarz lemma, since $\xi'_n \in J(f)$ cannot tend to ζ . Thus g is nonconstant and hence a univalent map. The Carathéodory convergence theorem implies that, for an $r < 1$ sufficiently close to 1, $\{f^n\}$ on $D = g(\{z \mid |z| < r\})$ converges uniformly, and hence $D \subset F(f)$. But this contradicts the fact that $\xi'_n \in J(f)$ are contained in D for all but finitely many n . ■

Remark. If every point z in a component U as in the above proof has such orbits whose real parts tend to $+\infty$, then the standard argument due to Eremenko and Lyubich gives the same contradiction.

We further note the following facts.

Theorem 3. *The Julia set of such an $f(z; a, b, c)$ as in Theorem 1 has vanishing area.*

Proof. We can find an $\epsilon > 0$ such that

$$(L_k)_\epsilon = \{w \in \mathbb{C} \mid d(w, L_k) \leq \epsilon\} \subset B$$

for every k . The assertion follows from McMullen-Stallard theorem in [6]. ■

Proposition 4. *Set $a = e^{i\alpha}$ and $b = i\beta$. Then we have*

$$f(z; a, b, c) = \int_0^z 2be^{(\beta t + \alpha)i/2} \frac{\beta t - \alpha + \pi}{2} \prod_{k=1}^{\infty} \left(1 - \frac{(\beta t - \alpha + \pi)^2}{4k^2\pi^2}\right) dt + d.$$

Proof. Since

$$f'(z) = 2be^{(\beta z + \alpha)i/2} \cos \frac{\beta z - \alpha}{2},$$

by using the famous equation

$$\sin \pi z = \pi z \prod_{k=1}^{\infty} \left(1 - \frac{z^2}{k^2}\right),$$

we have

$$f'(z) = 2be^{(\beta z + \alpha)i/2} \frac{\beta z - \alpha + \pi}{2} \prod_{k=1}^{\infty} \left(1 - \frac{(\beta z - \alpha + \pi)^2}{4k^2\pi^2} \right).$$

Hence we obtain the desired equation. \blacksquare

Finally we consider the convergence of Julia sets for two cases.

Theorem 5. Fix $b > 0$ and $c < -3$ satisfying $1 - bc + \log b > 0$ and let $ab < 1$. Then the sequence of the Julia sets of $f_a(z) = f(z; a, b, c)$ converges to the Julia set of $f(z) = z + e^{bz} + c$ in the sense of the Hausdorff convergence as $ab \nearrow 1$.

Proof. The assumption $1 - bc + \log b > 0$ implies $F(f)$ consists of only one Baker domain, which can be seen by the same argument as in the proof of Theorem 1.

In the case $ab < 1$, there exists an attracting fixed point on the negative real axis, say, $-r_a$ ($r_a > 0$). It is easy to see that r_a tends to infinity monotonically as $ab \nearrow 1$. We set

$$D_a = \{z \mid |z + r_a| < r_a\}.$$

For $z \in D_a$, we have

$$|f_a(z) + r_a| < abr_a + 2 < r_a + c + 3 < r_a.$$

This means that D_a is a collapsing disk for the attracting fixed point $-r_a$. The sequence $\{D_a\}$ is monotonically increasing and converges to $D_0 = \{z \mid \operatorname{Re} z < 0\}$ in the sense of the Carathéodory convergence. To see the sequence of the basins B_a of the attracting fixed points $-r_a$ to $F(f)$ in the sense of the Carathéodory convergence, we need to show only that, for an arbitrary compact set $E \subset F(f)$, there exists an $A > 0$ such that $E \subset B_a$ for all $a > A$ with $ab < 1$.

There exists an n such that $f^n(E) \subset D_0$, for D_0 is a fundamental set. Since $\{f_a\}$ converges to f uniformly on compact sets as $ab \nearrow 1$ (see [5]), we have $f^n(E) \subset D_a$, and hence E is contained in B_a for all $a > A$ for some A with $ab < 1$. Therefore, the sequence of the complements $\{B_a^c\}$ converges to the complement $F(f)^c = J(f)$ in the sense of the Hausdorff convergence. Since $J(f)$ has no interior points, we see that $\{J(f_a)\}$ converges to $J(f)$ in the sense of the Hausdorff convergence. \blacksquare

The Fatou set of $z \mapsto z + e^z$ contains infinitely many Baker domains, each of which is contained in the strip

$$S_m = \{z \mid |\operatorname{Im} z - (2m - 1)\pi| < \pi\}$$

for some $m \in \mathbb{Z} \setminus \{0\}$ (see [4]). The orbits of every point in the Baker domain have the real parts tending to $-\infty$.

Theorem 6. *If $f_c(z) = z + e^z + c$ with $c < 0$, then $F(f_c)$ consists of only one Baker Domain. Furthermore, $\{J(f_c)\}$ converges to $J(z + e^z)$ as $c \nearrow 0$ in the sense of the Hausdorff convergence.*

Proof. We set $f(z) = z + e^z$. Let B_m be the Baker domain in S_m . If $z = x + i\{(2m - 1)\pi \pm (\alpha + \pi/2)\}$ with $0 \leq \alpha < \pi/2$, then we have

$$\operatorname{Re} f(z) = x + e^x \sin \alpha > x.$$

If $z = x + i\{(2m - 1)\pi \pm \alpha\}$ with $0 \leq \alpha < \pi/2$ and $x < 0$, then we have

$$\operatorname{Im} f(z) = (2m - 1)\pi \pm \alpha \mp e^x \sin \alpha$$

and thus

$$(2m - 1)\pi - \alpha < \operatorname{Im} f(z) < (2m - 1)\pi + \alpha.$$

Hence, for $z \in B_m$, there exists an N such that

$$|\operatorname{Im} f^n(z) - (2m - 1)\pi| < \frac{\pi}{2}$$

for all $n > N$. Setting

$$K_m = \left\{ z \mid |\operatorname{Im} z - (2m - 1)\pi| < \frac{\pi}{2}, \operatorname{Re} z < 0 \right\},$$

we see that K_m is a fundamental set of B_m .

We consider only the case $c > -1$. Then $D_0 = \{z \mid \operatorname{Re} z < \log(-c)\}$ is a fundamental set of a Baker domain of f_c . Since $f_c(z) = f(z) + c$,

$$D_0 \cup \left(\bigcup_{m \in \mathbb{Z} \setminus \{0\}} K_m \right)$$

is also a fundamental set. By the argument similar to that in the proof of Theorem 1, we see that $F(f_c)$ consists of only one Baker domain. Further by the argument similar to that in the proof of Theorem 5, we see that $\{J(f_c)\}$ converges to $J(f)$ as $c \nearrow 0$ in the sense of the Hausdorff convergence. ■

REFERENCES

1. C. Cowen, Iteration and the solution of functional equations for functions analytic in the unit disk, *Trans. Amer. Math. Soc.* **265** (1981), 69-95.

2. A. Hinkkanen, B. Krauskopf and H. Kriete, Growing a Baker domain from attracting islands I: The dynamics of the limit function, preprint.
3. W. Bergweiler, Invariant domains and singularities, *Math. Proc. Cambridge Philos. Soc.* **117** (1995), 525-532.
4. S. Morosawa, Y. Nishimura, M. Taniguchi and T. Ueda, *Holomorphic Dynamics*, Cambridge Univ. Press, 1999.
5. S. Morosawa, The Carathéodory convergence of Fatou components of polynomials to Baker domains or wandering domains, Proceedings of the Second Congress ISAAC, to appear.
6. G. M. Stallard, Entire functions with Julia sets of zero measure, *Math Proc. Cambridge Philos. Soc.* **108** (1990), 551-557.
7. M. Taniguchi, Maskit surgery and structural finiteness of entire functions, to appear.

Shunsuke Morosawa
Department of Mathematics and Information Science,
Faculty of Science, Kochi University, Kochi 780-8520, Japan

Masahiko Taniguchi
Department of Mathematics, Graduate School of Science,
Kyoto University, Kyoto 606-8502, Japan