

## BAYESIAN ANALYSIS FOR REVERSIBLE MARKOV CHAINS

BY PERSI DIACONIS AND SILKE W. W. ROLLES

*Stanford University and Eindhoven University of Technology*

We introduce a natural conjugate prior for the transition matrix of a reversible Markov chain. This allows estimation and testing. The prior arises from random walk with reinforcement in the same way the Dirichlet prior arises from Pólya's urn. We give closed form normalizing constants, a simple method of simulation from the posterior and a characterization along the lines of W. E. Johnson's characterization of the Dirichlet prior.

**1. Introduction.** Modeling with Markov chains is an important part of time series analysis, genomics and many other applications. *Reversible* Markov chains are a mainstay of computational statistics through the Gibbs sampler, Metropolis algorithm and their many variants. Reversible chains are widely used natural models in physics and chemistry where reversibility (often called detailed balance) is a stochastic analog of the time reversibility of Newtonian mechanics.

This paper develops tools for a Bayesian analysis of the transition probabilities, stationary distribution and future prediction of a reversible Markov chain. We observe  $X_0 = v_0, X_1 = v_1, \dots, X_n = v_n$  from a reversible Markov chain with a finite state space  $V$ . Neither the stationary distribution  $\nu(v)$  nor the transition kernel  $k(v, v')$  is assumed known. Reversibility entails  $\nu(v)k(v, v') = \nu(v')k(v', v)$  for all  $v, v' \in V$ . We also assume we know which transitions are possible [for which  $v, v' \in V$  is  $k(v, v') > 0$ ].

In Section 2 we introduce a family of natural conjugate priors. These are defined via closed form densities and by a generalization of Pólya's urn to random walk with reinforcement on a graph. The density gives normalizing constants needed for testing independence versus reversibility or reversibility versus a full Markovian specification. The random walk gives a simple method of simulating from the posterior (Section 4.5).

Properties of the prior are developed in Section 4. The family is closed under sampling (Proposition 4.1). Mixtures of our conjugates are shown to be dense (Proposition 4.5). A characterization of the priors via predictive properties of the posterior is given (Section 4.2).

A practical example is given in Section 5. Several simple hypotheses are tested for a data set arising from the DNA of the human HLA-B gene. Section 5 also contains remarks about statistical analysis for reversible chains.

---

Received April 2004; revised March 2005.

*AMS 2000 subject classifications.* Primary 62M02; secondary 62C10.

*Key words and phrases.* Bayesian analysis, reversible Markov chains, conjugate priors, hypothesis testing.

**2. A class of prior distributions.** We observe  $X_0 = v_0, X_1 = v_1, \dots, X_n = v_n$  from a reversible Markov chain with a finite state space  $V$  and unknown transition kernel  $k(\cdot, \cdot)$ .

Let  $G = (V, E)$  be the finite graph with vertex set  $V$  and edge set  $E$  defined as follows:  $e = \{v, v'\} \in E$  (i.e., there is an edge between  $v$  and  $v'$ ) if and only if  $k(v, v') > 0$ . We assume that  $k(v, v') > 0$  iff  $k(v', v) > 0$ . In particular, all edges of  $G$  are undirected and an edge is denoted by the set of its endpoints. For some vertices  $v$ , we may have  $k(v, v) > 0$ . Define the simplex

$$(1) \quad \Delta := \left\{ x = (x_e)_{e \in E} \in (0, 1]^E : \sum_{e \in E} x_e = 1 \right\}.$$

REMARK 2.1. The distribution of a reversible Markov chain can be described by putting on the edge between  $v$  and  $v'$  the weight  $x_{\{v, v'\}} := v(v)k(v, v') = v(v')k(v', v)$ . If the weights are normalized so that  $\sum_{e \in E} x_e = 1$ , this is a unique way to describe the distribution of the Markov chain. A transition from  $v$  to  $v'$  is made with probability proportional to the weight  $x_{\{v, v'\}}$ .

Denote by  $Q_{v_0, x}$  the distribution of the Markov chain induced by the weights  $x = (x_e)_{e \in E} \in \Delta$  which starts with probability 1 in  $v_0$ . Using this notation, our assumption says that the observed data comes from a distribution in the class

$$(2) \quad \mathcal{Q} := \{ Q_{v_0, x} : v_0 \in V, x \in \Delta \}.$$

2.1. *A minimal sufficient statistic.* If the endpoints of an edge  $e$  agree, we call  $e$  a loop. Let

$$(3) \quad E_{\text{loop}} := \{ e \in E : e \text{ is a loop} \}.$$

For an edge  $e$ , denote the set of its endpoints by  $\bar{e}$ . For  $x = (x_e)_{e \in E} \in (0, \infty)^E$  and a vertex  $v$ , define  $x_v$  to be the sum of all components  $x_e$  with  $e$  incident to  $v$ ,

$$(4) \quad x_v := \sum_{\{e : v \in \bar{e}\}} x_e.$$

In sums such as this the sum is over edges including loops.

Let  $\pi := (\pi_0, \pi_1, \dots, \pi_n)$  be an admissible path in  $G$ . Define

$$(5) \quad k_v(\pi) := |\{ i \in \{1, 2, \dots, n\} : (v, \pi_i) = (\pi_{i-1}, \pi_i) \}| \quad \text{for } v \in V,$$

$$(6) \quad k_e(\pi) := \begin{cases} |\{ i \in \{1, 2, \dots, n\} : \{\pi_{i-1}, \pi_i\} = e \}|, & \text{for } e \in E \setminus E_{\text{loop}}, \\ 2 \cdot |\{ i \in \{1, 2, \dots, n\} : \{\pi_{i-1}, \pi_i\} = e \}|, & \text{for } e \in E_{\text{loop}}. \end{cases}$$

That is,  $k_v(\pi)$  equals the number of times the path  $\pi$  leaves vertex  $v$ ; for an edge  $e$  which is not a loop,  $k_e(\pi)$  is the number of traversals of  $e$  by  $\pi$ , and for a loop  $e$ ,  $k_e(\pi)$  is twice the number of traversals of  $e$ . Recall that the edges are undirected;

hence,  $k_e(\pi)$  counts the traversals of  $e$  in *both* directions. Set

$$(7) \quad Z_n := (X_0, X_1, \dots, X_n).$$

PROPOSITION 2.2. *The vector of transition counts  $(k_e(Z_n))_{e \in E}$  is a minimal sufficient statistic for the model  $\mathcal{Q}_{v_0} := \{Q_{v_0,x} : x \in \Delta\}$ .*

PROOF. Let  $\pi$  be an admissible path in  $G$ . In order to prove that  $(k_e(Z_n))_{e \in E}$  is a sufficient statistic, we need to show that

$$(8) \quad Q_{v_0,x}(Z_n = \pi | (k_e(Z_n))_{e \in E})$$

does not depend on  $x$ . If  $\pi$  does not start in  $v_0$ , (8) equals zero. Otherwise, we have

$$(9) \quad Q_{v_0,x}(Z_n = \pi) = \frac{\prod_{e \in E \setminus E_{\text{loop}}} x_e^{k_e(\pi)} \prod_{e \in E_{\text{loop}}} x_e^{k_e(\pi)/2}}{\prod_{v \in V} x_v^{k_v(\pi)}}.$$

It is not hard to see that  $k_v(\pi)$  can be expressed in terms of the  $k_e(\pi)$  and the first observation  $v_0$ . Hence, the  $Q_{v_0,x}$ -probability of  $\pi$  depends only on  $k_e(\pi)$ ,  $e \in E$ , and  $v_0$ . Thus, (8) equals one divided by the number of admissible paths  $\pi'$  with starting point  $v_0$  and  $k_e(\pi') = k_e(\pi)$  for all  $e \in E$ , which is independent of  $x$ .

Suppose  $K := (k_e)_{e \in E}$  is not minimal. Then there exists a sufficient statistic  $K'$  which needs less information than  $K$ . Consequently, there exist two admissible paths  $\pi$  and  $\pi'$  starting in  $v_0$  such that  $K(\pi) \neq K(\pi')$  and  $K'(\pi) = K'(\pi')$ . Then

$$(10) \quad \begin{aligned} & \frac{Q_{v_0,x}(Z_n = \pi | K'(Z_n) = K'(\pi))}{Q_{v_0,x}(Z_n = \pi' | K'(Z_n) = K'(\pi'))} \\ &= \frac{Q_{v_0,x}(Z_n = \pi)}{Q_{v_0,x}(Z_n = \pi')} \\ &= \prod_{e \in E \setminus E_{\text{loop}}} x_e^{k_e(\pi) - k_e(\pi')} \prod_{e \in E_{\text{loop}}} x_e^{(k_e(\pi) - k_e(\pi'))/2} \prod_{v \in V} x_v^{k_v(\pi') - k_v(\pi)}. \end{aligned}$$

Since by assumption  $(k_e(\pi))_{e \in E} \neq (k_e(\pi'))_{e \in E}$ , the last quantity depends on  $x$ . This contradicts the fact that  $K'$  is a sufficient statistic.  $\square$

2.2. *Definition of the prior densities.* Our aim is to define a class of prior distributions in terms of measures on  $\Delta$ . We prepare the definition with some notation. We illustrate the definitions by considering a three state process with states  $\{1, 2, 3\}$ , all transitions possible, but no holding. This leads to the graph in Figure 1.

Denote the cardinality of a set  $S$  by  $|S|$ . Recall the definition (3) of the set  $E_{\text{loop}}$ . Set

$$(11) \quad l := |V| + |E_{\text{loop}}| \quad \text{and} \quad m := |E|.$$

For the three state example,  $m = l = 3$ .




FIG. 1. The triangle.

REMARK 2.3. There is a simple way to delineate a *generating set of cycles* of  $G$ . We call a maximal subgraph of  $G$  which contains all loops but no cycle a *spanning tree* of  $G$ . Choose a spanning tree  $T$ . Each edge  $e \in E \setminus E_{\text{loop}}$  which is not in  $T$  forms a cycle  $c_e$  when added to  $T$ . (By definition, a loop is never a cycle and never contained in a cycle.) There are  $m - l + 1$  such cycles and we enumerate them arbitrarily:  $c_1, \dots, c_{m-l+1}$ . This set of cycles forms an additive basis for the homology  $H_1$  and also serves for our purposes.

For the three state example, we may choose  $T$  to have edges  $\{1, 2\}$  and  $\{1, 3\}$ . Then there is one cycle  $c_1$  oriented (say)  $1 \rightarrow 2 \rightarrow 3 \rightarrow 1$ . In Section 3.4 we show how such a basis of cycles can be obtained for the complete graph.

In general, the first Betti number  $\beta_1$  is the dimension of  $H_1$ . For the complete graph,  $\beta_1(K_n) = \binom{n-1}{2}$ . Further details can be found in [8], Section 1.16.

DEFINITION 2.4. Orient the cycles  $c_1, \dots, c_{m-l+1}$  and all edges  $e \in E$  in an arbitrary way. For every  $x \in \Delta$ , define a matrix  $A(x) = (A_{i,j}(x))_{1 \leq i, j \leq m-l+1}$  by

$$(12) \quad A_{i,i}(x) = \sum_{e \in c_i} \frac{1}{x_e}, \quad A_{i,j}(x) = \sum_{e \in c_i \cap c_j} \pm \frac{1}{x_e} \quad \text{for } i \neq j,$$

where the signs in the last sum are chosen to be  $+1$  or  $-1$  depending on whether the edge  $e$  has in  $c_i$  and  $c_j$  the same orientation or not.

In the three state example, the matrix  $A(x)$  is  $1 \times 1$  with entry  $x_{\{1,2\}}^{-1} + x_{\{2,3\}}^{-1} + x_{\{1,3\}}^{-1}$ .

Recall the definition (4) of  $x_v$ . Similarly, define  $a_v$  for  $a := (a_e)_{e \in E} \in (0, \infty)^E$ . The main definition of this section (the conjugate prior) follows.

DEFINITION 2.5. For all  $v_0 \in V$  and  $a := (a_e)_{e \in E} \in (0, \infty)^E$ , define

$$(13) \quad \phi_{v_0,a}(x) := Z_{v_0,a}^{-1} \frac{\prod_{e \in E \setminus E_{\text{loop}}} x_e^{a_e - 1/2} \prod_{e \in E_{\text{loop}}} x_e^{(a_e/2) - 1}}{x_{v_0}^{a_{v_0}/2} \prod_{v \in V \setminus \{v_0\}} x_v^{(a_v + 1)/2}} \sqrt{\det(A(x))}$$

for  $x := (x_e)_{e \in E} \in \Delta$  with

$$(14) \quad Z_{v_0, a} := \frac{\prod_{e \in E} \Gamma(a_e)}{\Gamma(a_{v_0}/2) \prod_{v \in V \setminus \{v_0\}} \Gamma((a_v + 1)/2) \prod_{e \in E_{\text{loop}}} \Gamma((a_e + 1)/2)} \times \frac{(m-1)! \pi^{(l-1)/2}}{2^{1-l + \sum_{e \in E} a_e}}.$$

For the three state example with parameters  $x_{\{1,2\}} = x$ ,  $x_{\{2,3\}} = y$ ,  $x_{\{1,3\}} = z$ ,  $a_{\{1,2\}} = a$ ,  $a_{\{2,3\}} = b$ ,  $a_{\{1,3\}} = c$ , and  $v_0 = 1$ , the prior density  $\phi$  is

$$(15) \quad Z^{-1} \frac{x^{a-1/2} y^{b-1/2} z^{c-1/2}}{(x+z)^{(a+c)/2} (x+y)^{(a+b+1)/2} (y+z)^{(b+c+1)/2}} \sqrt{\frac{1}{x} + \frac{1}{y} + \frac{1}{z}},$$

with the normalizing constant  $Z$  given by

$$(16) \quad \frac{\Gamma(a)\Gamma(b)\Gamma(c)}{\Gamma((a+c)/2)\Gamma((a+b+1)/2)\Gamma((b+c+1)/2)} \cdot \frac{2\pi}{2^{a+b+c-2}}.$$

A derivation of the formula for the density in this special case can be found, for example, in [12]. The density for the triangle with loops is given in (31).

The following proposition shows that the definition of  $\phi_{v_0, a}$  is independent of the choice of cycles  $c_i$  used in the definition of  $A(x)$ .

PROPOSITION 2.6. *For the matrix  $A$  of Definition 2.4, with  $\mathcal{T}$  the set of spanning trees of  $G$ ,*

$$(17) \quad \det A(x) = \sum_{T \in \mathcal{T}} \prod_{e \notin E(T)} \frac{1}{x_e}.$$

PROOF. This identity is proved for graphs without loops in [14], page 145, Theorem 3'. By definition,  $A(x)$  does not depend on  $x_e$ ,  $e \in E_{\text{loop}}$ . Furthermore, since every spanning tree contains all loops, the right-hand side of (17) does not depend on  $x_e$ ,  $e \in E_{\text{loop}}$  either. In particular, both sides of (17) are the same for  $G$  and the graph obtained from  $G$  by removing all loops; hence, they are equal.  $\square$

The prior density  $\phi_{v_0, a}$  arises in a natural extension of Pólya's urn. We treat this topic next.

2.3. *Random walk with reinforcement.* Let  $\sigma$  denote the Lebesgue measure on  $\Delta$ , normalized such that  $\sigma(\Delta) = 1$ . The measures  $\phi_{v_0, a} d\sigma$  on  $\Delta$  arise in the study of edge-reinforced random walk, as was observed by Coppersmith and Diaconis; see [3]. Let us explain this connection:

DEFINITION 2.7. All edges of  $G$  are given a strictly positive weight; at time 0 edge  $e$  has weight  $a_e > 0$ . An edge-reinforced random walk on  $G$  with starting point  $v_0$  is defined as follows: The process starts at  $v_0$  at time 0. In each step, the random walker traverses an edge with probability proportional to its weight. Each time an edge  $e \in E \setminus E_{\text{loop}}$  is traversed, its weight is increased by 1. Each time a loop  $e \in E_{\text{loop}}$  is traversed, its weight is increased by 2.

Denote the set of nonnegative integers by  $\mathbb{N}_0$ . Let  $\Omega$  be the set of all  $(v_i)_{i \in \mathbb{N}_0} \in V^{\mathbb{N}_0}$  such that  $\{v_i, v_{i+1}\} \in E$  for all  $i \in \mathbb{N}_0$ . Let  $X_n : V^{\mathbb{N}_0} \rightarrow V$  denote the projection onto the  $n$ th coordinate. Recall that  $Z_n = (X_0, X_1, \dots, X_n)$ . Denote by  $P_{v_0, a}$  the distribution on  $\Omega$  of an edge-reinforced random walk with starting point  $v_0$  and initial edge weights  $a = (a_e)_{e \in E}$ .

REMARK 2.8. Let  $\alpha_e(Z_n) := k_e(Z_n)/n$  be the proportion of traversals of edge  $e$  up to time  $n$ . For a finite graph without loops, it was observed by Coppersmith and Diaconis that  $\alpha(Z_n) := (\alpha_e(Z_n))_{e \in E}$  converges almost surely to a random variable with distribution  $\phi_{v_0, a} d\sigma$ ; see [3] and also [13]. In particular,  $\phi_{v_0, a} d\sigma$  is a probability measure on  $\Delta$ . This fact is not at all obvious from the definition of  $\phi_{v_0, a}$ .

It turns out that an edge-reinforced random walk on  $G$  is a mixture of reversible Markov chains, where the mixing measure described as a measure on edge weights  $(x_e)_{e \in E}$  is given by  $\phi_{v_0, a} d\sigma$ . This is made precise by the following theorem.

THEOREM 2.9. Let  $(X_n)_{n \in \mathbb{N}_0}$  be an edge-reinforced random walk with initial weights  $a = (a_e)_{e \in E}$  starting at  $v_0$ , and let  $Z_n = (X_0, X_1, \dots, X_n)$ . For any admissible path  $\pi = (v_0, \dots, v_n)$ , the following holds:

$$(18) \quad P_{v_0, a}(Z_n = \pi) = \int_{\Delta} \prod_{i=1}^n \frac{x_{\{v_{i-1}, v_i\}}}{x_{v_{i-1}}} \phi_{v_0, a}(x) d\sigma(x);$$

here  $x := (x_e)_{e \in E}$ . Hence, if  $\mathbb{Q}_{v_0, a}$  is the mixture of Markov chains where the mixing measure, described as a measure on edge weights  $(x_e)_{e \in E}$ , is given by  $\phi_{v_0, a} d\sigma$ , then

$$(19) \quad P_{v_0, a} = \mathbb{Q}_{v_0, a}.$$

PROOF. If  $G$  has no loops, then the claim is true by Theorem 3.1 of [16].

Let  $G$  be a graph with loops. Define a graph  $G' := (V', E')$  as follows: Replace every loop of  $G$  by an edge of degree 1 incident to the same vertex (see Figure 2). More precisely, for all  $e \in E_{\text{loop}}$ , let  $v(e)$  be the vertex  $e$  is incident to and let  $v'(e)$  be an additional vertex, different from all the others. Then, set  $g(e) := \{v(e), v'(e)\}$  and

$$(20) \quad V' := V \cup \{v'(e) : e \in E_{\text{loop}}\},$$

$$(21) \quad E' := [E \setminus E_{\text{loop}}] \cup \{g(e) : e \in E_{\text{loop}}\}.$$




FIG. 2. Transformation of loops.

The graph  $G'$  has no loops and the claim of the theorem is true for  $G'$ .

Let  $P'_{v_0,b}$  be the distribution of a reinforced random walk on  $G'$  starting at  $v_0$  with initial weights  $b = (b_{e'})_{e' \in E'}$  defined by

$$(22) \quad b_{e'} := \begin{cases} a_{e'}, & \text{if } e' \in E \setminus E_{\text{loop}}, \\ a_e, & \text{if } e' = g(e) \text{ for some } e \in E_{\text{loop}}. \end{cases}$$

Any finite admissible path  $\pi = (\pi_0 = v_0, \pi_1, \dots, \pi_n)$  in  $G$  can be mapped to an admissible path  $\pi' = (\pi'_0 = v_0, \pi'_1, \dots, \pi'_n)$  in  $G'$  by mapping every traversal of a loop  $e \in E_{\text{loop}}$  in  $\pi$  to a traversal of  $(v(e), v'(e), v(e))$  in  $\pi'$  [i.e., a traversal of the edge  $g(e)$  back and forth in  $\pi'$ ]. The probability that the reinforced random walk on  $G$  traverses  $\pi$  agrees with the probability that the reinforced random walk on  $G'$  traverses  $\pi'$ . [Note that for  $G$  and  $G'$  the following is true: Between any two successive visits to  $v(e)$ , the sum of the weights of all edges incident to  $v(e)$  increases by 2.] Since the claim of the theorem is true for  $G'$ , it follows that

$$(23) \quad \begin{aligned} P_{v_0,a}(Z_n = \pi) &= P'_{v_0,b}(Z_{n'} = \pi') \\ &= \int_{\Delta} \prod_{i=1}^{n'} \frac{x_{\{\pi'_{i-1}, \pi'_i\}}}{x_{\pi'_{i-1}}} \phi'_{v_0,b}(x) d\sigma(x), \end{aligned}$$

where  $\phi'_{v_0,b}$  denotes the density corresponding to  $G'$ , starting point  $v_0$  and initial weights  $b$ . We claim that the right-hand side of (23) equals

$$(24) \quad \int_{\Delta} \prod_{i=1}^n \frac{x_{\{\pi_{i-1}, \pi_i\}}}{x_{\pi_{i-1}}} \phi_{v_0,a}(x) d\sigma(x).$$

Note that a traversal of  $e \in E_{\text{loop}}$  contributes  $x_e/x_{v(e)}$  to the integrand in (24), whereas a traversal of  $(v(e), v'(e), v(e))$  contributes  $x_{g(e)}/x_{v(e)}$  to the integrand in (23). Furthermore,  $e \in E_{\text{loop}}$  contributes

$$(25) \quad \frac{\Gamma((a_e + 1)/2)}{\Gamma(a_e)} \cdot 2^{a_e} \cdot (x_e)^{(a_e/2)-1}$$

to  $\phi_{v_0,a}$ , whereas the contribution of the edge  $g(e)$  and the vertex  $v'(e)$  to the density  $\phi'_{v_0,b}$  equals

$$\begin{aligned}
 (26) \quad & \frac{\Gamma((a_{v'(e)} + 1)/2)}{\Gamma(a_e)} \cdot 2^{a_e} \cdot \frac{(x_{g(e)})^{a_e-1/2}}{(x_{v'(e)})^{(a_{v'(e)}+1)/2}} \\
 &= \frac{\Gamma((a_e + 1)/2)}{\Gamma(a_e)} \cdot 2^{a_e} \cdot \frac{(x_{g(e)})^{a_e-1/2}}{(x_{g(e)})^{(a_e+1)/2}} \\
 &= \frac{\Gamma((a_e + 1)/2)}{\Gamma(a_e)} \cdot 2^{a_e} \cdot (x_{g(e)})^{(a_e/2)-1}.
 \end{aligned}$$

Finally,  $|V| + |E_{\text{loop}}| = |V'|$  and  $|E| = |E'|$ . Consequently, the expression in (24) agrees with the right-hand side of (23) and the claim follows.  $\square$

**3. The prior density for special graphs.** In this section we write down the densities  $\phi_{v_0,a}$  for some special graphs.

3.1. *The line graph (Birth and death chains).* Consider the line graph with vertex set  $V = \{i : 0 \leq i \leq n\}$  and edge set  $E = \{\{i, i + 1\} : 0 \leq i \leq n - 1\}$ ; see Figure 3. Given  $a = (a_{\{i-1,i\}})_{1 \leq i \leq n}$ , let  $b_i := a_{\{i-1,i\}}$ . The variables in the simplex  $\Delta$  are denoted  $z_i := x_{\{i-1,i\}}$ .

Recall that the density of the beta distribution with parameters  $b_1, b_2 > 0$  is given by

$$(27) \quad \beta[b_1, b_2](p) := \frac{\Gamma(b_1 + b_2)}{\Gamma(b_1)\Gamma(b_2)} p^{b_1-1} (1 - p)^{b_2-1} \quad (0 < p < 1).$$

Set

$$(28) \quad p_i = \frac{z_i}{z_i + z_{i+1}}, \quad 1 \leq i \leq n - 1,$$

and  $p := (p_i)_{1 \leq i \leq n-1}$ . Clearly,  $p_i$  is the probability that the Markov chain with edge weights  $z_i$  makes a transition to  $i - 1$  given it is at  $i$ . If we make the change of variables (28) in the density  $\phi_{v_0,a}$ , then we obtain the transformed density  $\tilde{\phi}_{v_0,a}(p)$




FIG. 3. *The line graph.*



given by

$$\tilde{\phi}_{0,a}(p) = \begin{cases} \prod_{i=1}^{n-1} \beta\left[\frac{b_i+1}{2}, \frac{b_{i+1}}{2}\right](p_i), & \text{if } v_0 = 0, \\ \left[ \prod_{i=1}^{v_0-1} \beta\left[\frac{b_i}{2}, \frac{b_{i+1}+1}{2}\right](p_i) \right] \beta\left[\frac{b_{v_0}}{2}, \frac{b_{v_0}}{2}\right](p_{v_0}) \\ \quad \times \left[ \prod_{i=v_0+1}^{n-1} \beta\left[\frac{b_i+1}{2}, \frac{b_{i+1}}{2}\right](p_i) \right], & \text{if } v_0 \in \{1, 2, \dots, n-1\}, \\ \prod_{i=1}^{n-1} \beta\left[\frac{b_i}{2}, \frac{b_{i+1}+1}{2}\right](p_i), & \text{if } v_0 = n; \end{cases}$$

here the empty product is defined to be 1.

With the change of variables (28), the conjugate prior can be described as a product of independent beta variables with carefully linked parameters. If loops are allowed, the edge weights are independent Dirichlet by a similar argument (see Section 3.2). The next example contains a generalization.

3.2. *Trees with loops.* Recall that the density of the Dirichlet distribution with parameters  $b_i > 0$ ,  $1 \leq i \leq d$ , is given by

$$(29) \quad \begin{aligned} & D[b_i; 1 \leq i \leq d](p_i; 1 \leq i \leq d) \\ & := \frac{\Gamma(\sum_{i=1}^d b_i)}{\prod_{i=1}^d \Gamma(b_i)} \prod_{i=1}^d p_i^{b_i-1} \quad \left( p_i > 0, \sum_{i=1}^d p_i = 1 \right). \end{aligned}$$

Let  $T = (V, E)$  be a tree. Suppose that there is a loop attached to every vertex, that is,  $\{v\} \in E$  for all  $v \in V$ . Let  $v_0 \in V$ . For every  $v \in V \setminus \{v_0\}$ , there exists a unique shortest path from  $v_0$  to  $v$ . Let  $e(v)$  be the unique edge incident to  $v$  which is traversed by the shortest path from  $v_0$  to  $v$ . Let  $E_v := \{e \in E : v \in \bar{e}\}$  be the set of all edges incident to  $v$ . Set

$$(30) \quad p_e := \frac{x_e}{x_v} \quad \text{for } v \in V, e \in E_v,$$

$p := (p_e)_{e \in E}$ , and  $\vec{p}_v := (p_e)_{e \in E_v}$ . If we make the change of variables (30) in the density  $\phi_{v_0,a}$ , the transformed density  $\tilde{\phi}_{v_0,a}(p)$  is given by

$$D\left[\frac{a_e}{2}, e \in E_{v_0}\right](\vec{p}_{v_0}) \prod_{v \in V \setminus \{v_0\}} D\left[\frac{a_{e(v)}+1}{2}, \frac{a_e}{2}, e \in E_v \setminus \{e(v)\}\right](\vec{p}_v).$$

Thus again, in the reparametrization (30), the conjugate prior is seen as a product of independent random variables. This is not true in the following example.

The fact that the density  $\phi_{v_0,a}$  for a tree has this particular form was first observed by Pemantle [15].




FIG. 4. The triangle with loops.

3.3. *The triangle.* Consider the triangle with loops attached to all vertices. Let the vertex set be  $V = \{1, 2, 3\}$  and the edge set  $E = \{\{1\}, \{2\}, \{3\}, \{1, 2\}, \{1, 3\}, \{2, 3\}\}$  (see Figure 4). Let  $b_i$  be the initial weight of the loop at vertex  $i$  and let  $c_i$  be the initial weight of the edge opposite of vertex  $i$ . Similarly, let  $y_i := x_{\{i\}}$  and let  $z_1 := x_{\{2,3\}}$ ,  $z_2 := x_{\{1,3\}}$ ,  $z_3 := x_{\{1,2\}}$ .

The density  $\phi_{1,a}(y_1, y_2, y_3, z_1, z_2, z_3)$  for  $a = (b_1, b_2, b_3, c_1, c_2, c_3)$  is given by

$$(31) \quad Z_{1,a}^{-1} \cdot y_1^{(b_1/2)-1} y_2^{(b_2/2)-1} y_3^{(b_3/2)-1} z_1^{c_1-1} z_2^{c_2-1} z_3^{c_3-1} \sqrt{z_1 z_2 + z_1 z_3 + z_2 z_3} \\ \times ((y_1 + z_2 + z_3)^{(b_1+c_2+c_3)/2} (y_2 + z_1 + z_3)^{(b_2+c_1+c_3+1)/2} \\ \times (y_3 + z_1 + z_2)^{(b_3+c_1+c_2+1)/2})^{-1},$$

with

$$(32) \quad Z_{1,a} = \Gamma(c_1)\Gamma(c_2)\Gamma(c_3)\Gamma(b_1/2)\Gamma(b_2/2)\Gamma(b_3/2) \\ \times \left( \Gamma\left(\frac{b_1 + c_2 + c_3}{2}\right) \Gamma\left(\frac{b_2 + c_1 + c_3 + 1}{2}\right) \right. \\ \left. \times \Gamma\left(\frac{b_3 + c_1 + c_2 + 1}{2}\right) \right)^{-1} \cdot \frac{480\pi}{2^{c_1+c_2+c_3}}.$$

To calculate  $Z_{1,a}$  from (14), use the identity

$$(33) \quad \frac{\Gamma(b_i)}{2^{b_i} \Gamma((b_i + 1)/2)} = \frac{\Gamma(b_i/2)}{2\sqrt{\pi}} \quad (i = 1, 2, 3).$$

3.4. *The complete graph.* Perhaps the most important example is where all transitions are possible. This involves the complete graph  $K_n$  on  $n$  vertices with loops attached to all vertices. Let  $V = \{1, 2, 3, \dots, n\}$ . Let  $T_n$  be the spanning tree with edges  $\{1, i\}$  and loops  $\{i\}$ ,  $1 \leq i \leq n$ . This spanning tree induces the basis of cycles given by all triangles  $(1, i, j)$ ,  $2 \leq i < j \leq n$ . Figure 5 shows  $K_3, K_4$  and  $K_5$  together with  $T_3, T_4$  and  $T_5$ .




FIG. 5. The complete graphs  $K_3, K_4, K_5$  with loops and a spanning tree.

We remark that a different basis of cycles is given by  $(i, i + 1, j)$  for  $1 \leq i < j + 1 \leq n$ . This may be proved by induction using the Mayer–Vietoris decomposition theorem based on  $K_{n-1}$  and a point.

Let  $a = (a_{\{i,j\}})_{1 \leq i, j \leq n}$  be given. For  $K_n$ , set  $b_i := a_{\{i\}}$ ,  $a_i = \sum_{j=1}^n a_{\{i,j\}}$  and  $b := \sum_{1 \leq i, j \leq n} a_{\{i,j\}}$ . The variables of the simplex are  $x = (x_{\{i,j\}})_{1 \leq i, j \leq n}$ . Abbreviating  $y_i := x_{\{i\}}$  and  $x_i = \sum_{j=1}^n x_{\{i,j\}}$ , the density  $\phi_{1,a}$  is given by

$$(34) \quad \phi_{1,a}(x) = Z_{1,a}^{-1} \cdot \frac{\prod_{1 \leq i < j \leq n} x_{\{i,j\}}^{a_{\{i,j\}} - 1/2} \prod_{i=1}^n y_i^{(b_i/2) - 1}}{x_1^{a_1/2} \prod_{i=2}^n x_i^{(a_i+1)/2}} \sqrt{\det(A_n(x))},$$

with  $A_n(x)$  defined in (12) and

$$(35) \quad Z_{1,a} = \frac{\prod_{1 \leq i, j \leq n} \Gamma(a_{\{i,j\}})}{\Gamma(a_1/2) \prod_{i=2}^n \Gamma((a_i + 1)/2) \prod_{i=1}^n \Gamma((b_i + 1)/2)} \times \frac{((n(n + 1)/2) - 1)! \pi^{n-1/2}}{2^{1-2n+b}}.$$

**4. Properties of the family of priors.** For  $v_0 \in V$  and  $a = (a_e)_{e \in E} \in (0, \infty)^E$ , abbreviate

$$(36) \quad \mathbb{P}_{v_0,a} := \phi_{v_0,a} d\sigma;$$

that is,  $\mathbb{P}_{v_0,a}$  is the measure on  $\Delta$  with density  $\phi_{v_0,a}$ . Recall that  $\mathbb{Q}_{v_0,a}$  denotes the mixture of Markov chains where the mixing measure, described as a measure on edge weights  $(x_e)_{e \in E}$ , is given by  $\mathbb{P}_{v_0,a}$ . In this section we study properties of the set of prior distributions

$$(37) \quad \mathcal{D} := \{\mathbb{P}_{v_0,a} : v_0 \in V, a = (a_e)_{e \in E} \in (0, \infty)^E\}.$$

4.1. *Closure under sampling.* Recall the definition (6) of  $k_e(\pi)$  and recall that  $Z_n = (X_0, \dots, X_n)$ .

PROPOSITION 4.1. *Under the prior distribution  $\mathbb{P}_{v_0,a}$  with observations  $X_0 = v_0, X_1 = v_1, \dots, X_n = v_n$ , the posterior is given by  $\mathbb{P}_{v_n,(a_e+k_e(Z_n))_{e \in E}}$ . In particular, the family  $\mathcal{D}$  is closed under sampling.*

PROOF. Suppose the prior distribution is  $\mathbb{P}_{v_0,a}$  and we are given  $n + 1$  observations  $\pi = (\pi_0, \pi_1, \dots, \pi_n)$  sampled from  $\mathbb{Q}_{v_0,a}$ . Then  $\pi_0 = v_0$ . We claim that the posterior is given by  $\mathbb{P}_{\pi_n,(a_e+k_e(\pi))_{e \in E}}$ . By Theorem 2.9,  $\mathbb{Q}_{v_0,a} = P_{v_0,a}$ . The  $P_{v_0,a}$ -distribution of  $\{X_{n+k}\}_{k \geq 0}$  given  $Z_n = \pi$  is the distribution of an edge-reinforced random walk starting at the vertex  $\pi_n$  with initial values  $a_e + k_e(\pi)$ . Using the identity (19) again, it follows that the  $P_{v_0,a}$ -distribution of  $\{X_{n+k}\}_{k \geq 0}$  given  $Z_n = \pi$  equals  $\mathbb{Q}_{\pi_n,(a_e+k_e(\pi))_{e \in E}}$ . Thus, the posterior equals  $\mathbb{P}_{\pi_n,(a_e+k_e(\pi))_{e \in E}}$ , which is an element of  $\mathcal{D}$ .  $\square$

4.2. *Uniqueness.* In this section we give a characterization of our priors along the lines of W. E. Johnson’s characterization of the Dirichlet prior. See [18] for history and [19] for a version for nonreversible chains. The closely related topic of de Finetti’s theorem for Markov chains is developed by Freedman [7] and Diaconis and Freedman [4]. See also [6].

DEFINITION 4.2. Two finite admissible paths  $\pi$  and  $\pi'$  are called *equivalent* if they have the same starting point and satisfy  $k_e(\pi) = k_e(\pi')$  for all  $e \in E$ . We define  $P$  to be *partially exchangeable* if  $P(Z_n = \pi) = P(Z_n = \pi')$  for any equivalent paths  $\pi$  and  $\pi'$  of length  $n$ .

For  $n \in \mathbb{N}_0$  and  $v \in V$ , define

$$(38) \quad k_n(v) := |\{i \in \{0, 1, \dots, n\} : X_i = v\}|.$$

It seems natural to take a class  $\mathcal{P}$  of distributions for  $(X_n)_{n \in \mathbb{N}_0}$  with the following properties:

- P1. For all  $P \in \mathcal{P}$ , there exists  $v_0 \in V$  such that  $P(X_0 = v_0) = 1$ .
- P2. For all  $P \in \mathcal{P}$ ,  $v_0$  as in P1, and any admissible path  $\pi$  of length  $n \geq 1$  starting at  $v_0$ , we have  $P(Z_n = \pi) > 0$ .
- P3. Every  $P \in \mathcal{P}$  is partially exchangeable.
- P4. For all  $P \in \mathcal{P}$ ,  $v \in V$  and  $e \in E$ , there exists a function  $f_{P,v,e}$  taking values in  $[0, 1]$  such that, for all  $n \geq 0$ ,

$$P(X_{n+1} = v | Z_n) = f_{P,X_n,\{X_n,v\}}(k_n(X_n), k_{\{X_n,v\}}(Z_n)).$$

The condition P4 says that, given  $X_0, X_1, \dots, X_n$ , the probability that  $X_{n+1} = v$  depends only on the following quantities: the observation  $X_n$ , the number of times  $X_n$  has been observed so far, the edge  $\{X_n, v\}$  and the number of times transitions between  $X_n$  and  $v$  (and between  $v$  and  $X_n$ ) have been observed so far.

We make the following assumptions on the graph  $G$ :

- G1. For all  $v \in V$ ,  $\text{degree}(v) \neq 2$ .
- G2. The graph  $G$  is 2-edge-connected, that is, removing an edge does not make  $G$  disconnected.

For example, a triangle with loops or the complete graph  $K_n$ ,  $n \geq 4$ , with or without loops, satisfies G1 and G2, while a path fails both G1 and G2.

Recall that  $Q_{v_0,x}$  is the distribution of the reversible Markov chain starting in  $v_0$ , making a transition from  $v$  to  $v'$  with probability proportional to  $x_{\{v,v'\}}$  whenever  $\{v, v'\} \in E$ .

**THEOREM 4.3.** *Suppose the graph  $G$  satisfies G1 and G2.*

- (a) *The set  $\mathcal{M} := \{Q_{v_0,a} : v_0 \in V, a = (a_e)_{e \in E} \in (0, \infty)^E\}$  satisfies P1–P4.*
- (b) *On the other hand, if P1–P4 are satisfied for a set  $\mathcal{P}$  of probability distributions, then for all  $P \in \mathcal{P}$  there exist  $v_0 \in V$  and  $a \in (0, \infty)^E$  such that either*

$$\begin{aligned}
 &P(X_{n+1} = v | Z_n, k_n(X_n) \geq 3) \\
 &= Q_{v_0,a}(X_{n+1} = v | Z_n, k_n(X_n) \geq 3) \quad \forall n \geq 0 \quad \text{or} \\
 (39) \quad &P(X_{n+1} = v | Z_n, k_n(X_n) \geq 3) \\
 &= Q_{v_0,a}(X_{n+1} = v | Z_n, k_n(X_n) \geq 3) \quad \forall n \geq 0.
 \end{aligned}$$

The second part of the theorem states that either  $P$  and  $Q_{v_0,a}$  or  $P$  and  $Q_{v_0,a}$  essentially agree; only the conditional probabilities to leave from a state which has been visited at most twice could be different.

**PROOF OF THEOREM 4.3.** It is straightforward to check that  $\mathcal{M}$  has the properties P1–P4. For the converse, let  $P \in \mathcal{P}$ . If  $G$  has no loops, then Theorem 1.2 of Rolles [16] implies that there exist  $v_0 \in V$  and  $a \in (0, \infty)^E$  such that either (39) holds or  $P(X_{n+1} = v | Z_n, k_n(X_n) \geq 3) = P_{v_0,a}(X_{n+1} = v | Z_n, k_n(X_n) \geq 3)$  for all  $n$ . In this case, the claim follows from (19).

If  $G$  has loops, consider the graph  $G'$  defined in the proof of Theorem 2.9 and the induced process  $X' := (X'_n)_{n \in \mathbb{N}_0}$  on  $G'$  with reflection at the vertices  $v'(e)$ ,  $e \in E_{\text{loop}}$ . The process  $X'$  satisfies P1–P4. Hence, the claim holds for  $X'$  and, consequently, for  $(X_n)_{n \in \mathbb{N}_0}$ .  $\square$

**REMARK 4.4.** The preceding theorem holds under the assumption that the graph  $G$  is 2-edge-connected (G2). If  $G$  is not 2-edge-connected, a similar statement can be proved for a different class of priors: One replaces the class  $\mathcal{D}$  by the

mixing measures of a so-called *modified edge-reinforced random walk*; for the definition of this process, see Definition 2.1 of [16]. A uniqueness statement similar to Theorem 4.3 follows from Theorem 2.1 of [16].

4.3. *The priors are dense.* As shown by Dalal and Hall [2] and Diaconis and Ylvisaker [5] for classical exponential families, mixtures of conjugate priors are dense in the space of all priors. This holds for reversible Markov chains.

PROPOSITION 4.5. *The set of convex combinations of priors in  $\mathcal{D}$  is weak-star dense in the set of all prior distributions on reversible Markov chains on  $G$ .*

PROOF. For an infinite admissible path  $\pi = (\pi_0, \pi_1, \pi_2, \dots)$  in  $G$ , define  $\alpha(\pi) := (\alpha_e(\pi))_{e \in E}$  by  $\alpha_e(\pi) := \lim_{n \rightarrow \infty} k_e(\pi_0, \pi_1, \dots, \pi_n)/n$  to be the limiting fraction of crossings of the edge  $e$  by the path  $\pi$ . Let  $Z_\infty := (X_0, X_1, X_2, \dots)$ . Note that  $\alpha(Z_\infty)$  is defined  $\mathbb{Q}_{v_0, a}$ -a.s. Define  $\tau_n$  to be the  $n$ th return time to  $v_0$ . Since  $G$  is finite,  $\tau_n < \infty$   $\mathbb{Q}_{v_0, a}$ -a.s. for all  $n \in \mathbb{N}$  and all  $a \in (0, \infty)^E$ .

Let  $f : \Delta \rightarrow \mathbb{R}$  be bounded and continuous. Denote the expectation with respect to  $\mathbb{Q}_{v_0, a}$  by  $\mathbb{E}_{v_0, a}$ . Since  $X_{\tau_n} = v_0$ , Theorem 2.9 implies that

$$(40) \quad \mathbb{E}_{v_0, (a_e + k_e(Z_{\tau_n}))_{e \in E}} [f(\alpha(Z_\infty))] = \mathbb{E}_{v_0, a} [f(\alpha(Z_\infty)) | Z_{\tau_n}] := M_n.$$

Clearly,  $(M_n)_{n \geq 0}$  is a bounded martingale. Hence, by the martingale convergence theorem,

$$(41) \quad \lim_{n \rightarrow \infty} \mathbb{E}_{v_0, (a_e + k_e(Z_{\tau_n}))_{e \in E}} [f(\alpha(Z_\infty))] = \mathbb{E}_{v_0, a} [f(\alpha(Z_\infty)) | Z_\infty] = f(\alpha(Z_\infty)) \\ = \int f d\delta_{\alpha(Z_\infty)}$$

$\mathbb{Q}_{v_0, a}$ -a.s.; here  $\delta_b$  denotes the point mass in  $b$ . Since  $\Delta$  is compact, there is a countable dense subset of the set of bounded continuous functions on  $\Delta$ . Hence, the above shows that, for  $\mathbb{Q}_{v_0, a}$ -almost all  $Z_\infty$ ,

$$(42) \quad \mathbb{Q}_{v_0, (a_e + k_e(Z_{\tau_n}))_{e \in E}} (\alpha(Z_\infty) \in \cdot) \Rightarrow \delta_{\alpha(Z_\infty)}(\cdot) \quad \text{weakly as } n \rightarrow \infty.$$

The  $\mathbb{Q}_{v_0, a}$ -distribution of  $\alpha(Z_\infty)$  equals  $\mathbb{P}_{v_0, a}$ . Thus,

$$(43) \quad \mathbb{P}_{v_0, (a_e + k_e(Z_{\tau_n}))_{e \in E}} (\cdot) \Rightarrow \delta_{\alpha(Z_\infty)}(\cdot) \quad \text{weakly as } n \rightarrow \infty$$

for  $\mathbb{Q}_{v_0, a}$ -almost all  $Z_\infty$ . Recall that the  $\mathbb{Q}_{v_0, a}$ -distribution of  $\alpha(Z_\infty)$  (viz.,  $\mathbb{P}_{v_0, a}$ ) is absolutely continuous with respect to Lebesgue measure on  $\Delta$  with the density  $\phi_{v_0, a}$  which is strictly positive in the interior of  $\Delta$ . Hence, for Lebesgue-almost all  $a \in \Delta$ , there is a sequence  $a_n \in \Delta$  such that  $\mathbb{P}_{v_0, a_n} \Rightarrow \delta_a$  weakly. By the Krein–Milman theorem, convex combinations of point masses are weak-star dense in the set of all measures on  $\Delta$ . Using a standard argument, it follows that the set of convex combinations of the distributions  $\mathbb{P}_{v_0, a}$  is dense in the set of all probability measures on  $\Delta$ . This completes the proof of the proposition.  $\square$

4.4. *Computing some moments.* For any edge  $e_0 \in E$ , we can calculate the probability that the mixture of Markov chains with mixing measure  $\phi_{v_0,a} d\sigma$  traverses  $e_0$  back and forth starting at an endpoint of  $e_0$ . This gives a closed form for certain moments of the prior  $\mathbb{P}_{v_0,a}$ .

PROPOSITION 4.6. *For  $e_0 \in E \setminus E_{\text{loop}}$  with endpoints  $v$  and  $v'$ , we have*

$$(44) \quad \int_{\Delta} \frac{(x_{e_0})^2}{x_v x_{v'}} \phi_{v_0,a}(x) d\sigma(x) = \begin{cases} \frac{a_{e_0}(a_{e_0} + 1)}{(a_v + 1)(a_{v'} + 1)}, & \text{if } v_0 \notin \{v, v'\}, \\ \frac{a_{e_0}(a_{e_0} + 1)}{a_v(a_{v'} + 1)}, & \text{if } v = v_0. \end{cases}$$

For a loop  $e_0 \in E_{\text{loop}}$  incident to  $v$ , we have

$$(45) \quad \int_{\Delta} \frac{x_{e_0}}{x_v} \phi_{v_0,a}(x) d\sigma(x) = \begin{cases} \frac{a_{e_0}}{a_v + 1}, & \text{if } v \neq v_0, \\ \frac{a_{e_0}}{a_v}, & \text{if } v = v_0. \end{cases}$$

PROOF. *Case  $e_0 \in E \setminus E_{\text{loop}}$ .* Suppose  $v_0$  is an endpoint of  $e_0$ , say,  $v = v_0$ . Then

$$(46) \quad \int_{\Delta} \frac{(x_{e_0})^2}{x_v x_{v'}} \phi_{v_0,a}(x) d\sigma(x) = \mathbb{Q}_{v_0,a}(X_0 = v_0, X_1 = v', X_2 = v_0);$$

this is the probability that the mixture of Markov chains traverses the edge  $e_0$  back and forth starting at  $v_0$ . By (19)  $\mathbb{Q}_{v_0,a} = P_{v_0,a}$ . Hence, (46) equals the probability that an edge-reinforced random walk traverses  $e_0$  back and forth, namely,

$$(47) \quad P_{v_0,a}(X_0 = v_0, X_1 = v', X_2 = v_0) = \frac{a_{e_0}(a_{e_0} + 1)}{a_v(a_{v'} + 1)}.$$

Here we used the fact that the sum of the weights of all edges incident to  $v'$  equals  $a_{v'} + 1$  after  $e_0$  has been traversed once. This proves the claim in the case  $v_0 \in \bar{e}_0$ .

Suppose  $v_0 \notin \bar{e}_0$ . Define  $b := (b_e)_{e \in E}$  by  $b_{e_0} := a_{e_0} + 2$  and  $b_e := a_e$  for  $e \in E \setminus \{e_0\}$ . Then, using the definition of  $\phi_{v_0,a}$ , we obtain

$$(48) \quad \frac{(x_{e_0})^2}{x_v x_{v'}} \phi_{v_0,a}(x) = \frac{Z_{v_0,b}}{Z_{v_0,a}} \phi_{v_0,b}(x) \quad \text{for all } x \in \Delta.$$

Using the definition of the normalizing constants  $Z_{v_0,a}$  and  $Z_{v_0,b}$  and the identity  $\Gamma(z + 1) = z\Gamma(z)$ , it follows that

$$(49) \quad \frac{Z_{v_0,b}}{Z_{v_0,a}} = \frac{\Gamma((a_v + 1)/2)\Gamma((a_{v'} + 1)/2)\Gamma(a_{e_0} + 2)}{4\Gamma((a_v + 3)/2)\Gamma((a_{v'} + 3)/2)\Gamma(a_{e_0})} = \frac{a_{e_0}(a_{e_0} + 1)}{(a_v + 1)(a_{v'} + 1)}.$$

Since  $\int_{\Delta} \phi_{v_0,b}(x) d\sigma(x) = 1$ , the claim follows by integrating both sides of (48) over  $\Delta$ .

Case  $e_0 \in E_{\text{loop}}$ . The proof follows the same line as in the case  $e_0 \notin E_{\text{loop}}$ . Let  $e_0 = \{v\}$  be incident to  $v$ . We prove only the case  $v \neq v_0$ . Defining  $b$  as above, (48) is valid with

$$\begin{aligned}
 \frac{Z_{v_0,b}}{Z_{v_0,a}} &= \frac{\Gamma((a_v + 1)/2)\Gamma((a_{e_0} + 1)/2)\Gamma(a_{e_0} + 2)}{4\Gamma((a_v + 3)/2)\Gamma((a_{e_0} + 3)/2)\Gamma(a_{e_0})} \\
 (50) \qquad &= \frac{a_{e_0}(a_{e_0} + 1)}{(a_v + 1)(a_{e_0} + 1)} = \frac{a_{e_0}}{a_v + 1};
 \end{aligned}$$

here we used again the identity  $\Gamma(z + 1) = z\Gamma(z)$ . The claim follows.  $\square$

Recall the definitions (5) and (6) of  $k_v(\pi)$  and  $k_e(\pi)$  for a finite admissible path  $\pi$  in  $G$ . Abbreviate  $k_v := k_v(\pi)$ ,  $k_e := k_e(\pi)$ . For  $x = (x_e)_{e \in E} \in \Delta$ , denote by  $Q_x(\pi)$  the probability that the reversible Markov chain with transition probabilities induced by the weights  $(x_e)_{e \in E}$  on the edges traverses the path  $\pi$ . Note that if  $\pi$  is a closed path, that is, if the starting point and endpoint of  $\pi$  agree, then  $Q_x(\pi)$  is independent of the starting point of  $\pi$ . An argument as in the proof of Proposition 4.6 yields the following:

PROPOSITION 4.7. *For any finite admissible path  $\pi$  starting at  $v_0$ , we have*

$$\begin{aligned}
 (51) \qquad &\int_{\Delta} Q_x(\pi)\phi_{v_0,a}(x) d\sigma(x) \\
 &= \frac{[\prod_{e \in E \setminus E_{\text{loop}}} \prod_{i=0}^{k_e-1} (a_e + i)][\prod_{e \in E_{\text{loop}}} \prod_{i=0}^{k_e/2-1} (a_e + 2i)]}{\prod_{i=0}^{k_{v_0}-1} (a_{v_0} + 2i) \prod_{v \in V \setminus \{v_0\}} \prod_{i=0}^{k_v-1} (a_v + 1 + 2i)}.
 \end{aligned}$$

For any finite admissible path  $\pi$  with the same starting point and endpoint which avoids  $v_0$ , we have

$$\begin{aligned}
 (52) \qquad &\int_{\Delta} Q_x(\pi)\phi_{v_0,a}(x) d\sigma(x) \\
 &= \frac{[\prod_{e \in E \setminus E_{\text{loop}}} \prod_{i=0}^{k_e-1} (a_e + i)][\prod_{e \in E_{\text{loop}}} \prod_{i=0}^{k_e/2-1} (a_e + 2i)]}{\prod_{v \in V} \prod_{i=0}^{k_v-1} (a_v + 1 + 2i)}.
 \end{aligned}$$

Here the empty product is defined to be 1.

If  $\pi$  is a closed path, we call  $Q_x(\pi)$  a cycle probability. The transition probabilities of a Markov chain with finite state space  $V$  that visits every state with probability 1 are completely determined by all its cycle probabilities (see, e.g., [7], Corollary on page 116).



4.5. *Simulating from the posterior.* In this subsection we show how the posterior distribution of the unknown stationary distribution for the underlying Markov chain can be simulated using reinforced random walks.

Suppose our posterior distribution is  $\mathbb{P}_{v_0,a} = \phi_{v_0,a} d\sigma$ . Let  $X^{(i)} := (X_n^{(i)})_{n \geq 0}$ ,  $i \geq 1$ , be independent reinforced random walks with the same initial edge weights  $a = (a_e)_{e \in E}$ . Let  $Z_n^{(i)} := (X_0^{(i)}, X_1^{(i)}, \dots, X_n^{(i)})$  and recall that  $k_e(Z_n^{(i)})$  equals the number of traversals of edge  $e$  by the process  $X^{(i)}$  up to time  $n$ .

PROPOSITION 4.8. *For any interval  $I \subseteq \mathbb{R}$  and all  $e \in E$ , we have*

$$(53) \quad \lim_{n \rightarrow \infty} \lim_{m \rightarrow \infty} \frac{1}{m} \left| \left\{ i \leq m : \frac{k_e(Z_n^{(i)})}{n} \in I \right\} \right| = \mathbb{P}_{v_0,a}(x_e \in I) \quad \text{a.s.}$$

PROOF. For every  $n$ , the random variables  $k_e(Z_n^{(i)})/n$ ,  $i \geq 1$ , are i.i.d. Hence, by the Glivenko–Cantelli theorem, a.s. for all  $x \in \mathbb{R}$ ,

$$(54) \quad \begin{aligned} \lim_{m \rightarrow \infty} \frac{1}{m} \left| \left\{ i \leq m : \frac{k_e(Z_n^{(i)})}{n} \leq x \right\} \right| &= P_{v_0,a} \left( \frac{k_e(Z_n)}{n} \leq x \right) \\ &= Q_{v_0,a} \left( \frac{k_e(Z_n)}{n} \leq x \right). \end{aligned}$$

For the last equality we used (19). Since  $Q_{v_0,a}$  is a mixture of Markov chains,  $k_e(Z_n)/n$  converges to the normalized weight of the edge  $e$   $Q_{v_0,a}$ -a.s. and, hence, weakly. Since the limiting distribution is continuous,

$$(55) \quad \lim_{n \rightarrow \infty} Q_{v_0,a} \left( \frac{k_e(Z_n)}{n} \leq x \right) = \mathbb{P}_{v_0,a}(x_e \leq x),$$

and the claim follows.  $\square$

PROPOSITION 4.9. *For all  $e \in E$ ,*

$$(56) \quad \lim_{n \rightarrow \infty} \int \frac{k_e(Z_n)}{n} dP_{v_0,a} = \int x_e d\mathbb{P}_{v_0,a}.$$

PROOF. By (19),  $P_{v_0,a} = Q_{v_0,a}$ . Since the proportion  $k_n(e)/n$  converges  $Q_{v_0,a}$ -a.s. to the normalized weight of the edge  $e$ , the claim follows from the dominated convergence theorem.  $\square$

REMARK 4.10. The Markov chain with distribution induced by the edge weights  $(x_e)_{e \in E} \in \Delta$  has the stationary distribution  $\nu(v) = \frac{x_v}{2} = \frac{1}{2} \sum_{e \in E_v} x_e$ . Thus, Propositions 4.8 and 4.9 allow simulation of the  $\mathbb{P}_{v_0,a}$ -distribution and the mean of  $\nu(v)$ .

**5. Applications.** Reversibility can serve as a natural intermediate between independence and fully nonparametric Markovian dependence. On  $|V|$  states,

TABLE 1  
*Degrees of freedom for independent, reversible and full Markov specification*

$ V $	3	4	5	10	20	50	100	1000
Independent $ V  - 1$	2	3	4	9	19	49	99	999
Reversible $ V ( V  - 1)/2 - 1$	2	5	9	44	189	1224	4949	499499
Full Markov $ V ( V  - 1)$	6	12	20	90	380	2450	9900	999000

with no restrictions the number of free parameters is  $|V| - 1$  with independence,  $|V|(|V| - 1)$  for full Markov and  $\frac{|V|(|V|-1)}{2} - 1$  for reversibility. As Table 1 indicates, these numbers vary widely for  $|V|$  large.

In this section we illustrate the use of our priors for testing a variety of simple hypotheses. Table 2 shows a genetic data set from the DNA sequence of the humane HLA-B gene. This gene plays a central role in the immune system. The data displayed in Table 2 is downloaded from the webpage of the National Center for Biotechnology Information ([www.ncbi.nlm.nih.gov/genome/guide/human/](http://www.ncbi.nlm.nih.gov/genome/guide/human/)).

In Example A, we test i.i.d.  $\frac{1}{4}$  versus i.i.d. for the DNA-data. In Example B, we test i.i.d. versus reversible. In Example C, we test reversible versus full Markov. In Example D, we compare i.i.d. with full Markov.

Let  $n_a, n_c, n_g$  and  $n_t$  denote the number of occurrences of  $a, c, g$  and  $t$ , respectively, in the data displayed in Table 2. Then

$$(57) \quad n_a = 621, \quad n_c = 974, \quad n_g = 1064, \quad n_t = 711.$$

EXAMPLE A. A Bayes test of  $H_0$ : i.i.d. ( $\frac{1}{4}$ ) versus  $H_1$ : i.i.d.(unknown).  
 A “standard” test can be based on the Bayes factor

$$\frac{P(data|H_0)}{P(data|H_1)}.$$

See [9] for an extensive discussion. For  $H_1$ , we use a Dirichlet(1, 1, 1, 1) prior. This yields

$$\begin{aligned} P(data|H_0) &= \left(\frac{1}{4}\right)^{3370} \approx 1.142429015368253 \cdot 10^{-2029}, \\ P(data|H_1) &= \frac{\Gamma(4)\Gamma(n_a + 1)\Gamma(n_c + 1)\Gamma(n_g + 1)\Gamma(n_t + 1)}{\Gamma(n_a + n_c + n_g + n_t + 4)} \\ &= \frac{\Gamma(4)\Gamma(622)\Gamma(975)\Gamma(1065)\Gamma(712)}{\Gamma(3374)} \\ &\approx 1.140417804695619 \cdot 10^{-1999}, \\ \frac{P(data|H_0)}{P(data|H_1)} &\approx 1.00176 \cdot 10^{-30}. \end{aligned}$$

TABLE 2  
*The humane HLA-B gene. Part of the DNA sequence of length 3370*

1	tggtgtagga	gaagagggat	caggacgaag	tcccaggtcc	cggacggggc	tctcagggtc
61	tcaggctccg	agggcccgct	ctgcaatggg	gaggcgcagc	ggttggggatt	ccccactccc
121	ctgagtttca	cttcttctcc	caacttgtgt	cgggtccttc	ttccaggata	ctcgtgacgc
181	gtcccactt	cccactccca	ttgggtattg	gatatactaga	gaagccaatc	agcgtcgccg
241	cggccccagt	tctaaagtcc	ccacgcacc	acccggactc	agagtctct	cagacgcga
301	gatgctggtc	atggcgcccc	gaaccgtcct	cctgtgtctc	tcggcgcccc	tggccctgac
361	cgagacctgg	gccggtgagt	gcggtcggg	agggaaatgg	cctctgccg	gaggagcgag
421	gggaccgcag	gcgggggcgc	aggacctgag	gagccgcgcc	gggaggagg	tcggggcggt
481	ctcagccct	cctcaccccc	aggctcccac	tcctagaggt	attctacac	ctcctgttcc
541	cggcccggcc	gcggggagcc	ccgtctcatc	tcagtgggct	acgtggacga	caccagttc
601	gtgaggttcg	acagcgacgc	cgcgagtccg	agagaggagc	cgcggggcgc	gtggatagag
661	caggcggggc	gagagatttc	ggaccggaac	acacagatct	acaaggccca	ggcacagact
721	gaccgagaga	gcctcgggaa	cctgcgcgcc	tactacaacc	agagcgagcc	cggtgagtga
781	ccccggcccc	ggcgcgaggt	cacgactccc	catccccac	gtacggcccc	ggtcgcccc
841	agtctccggg	tcagagatcc	gcctccctga	gcccgcggga	ccgcccaga	ccctcgaccg
901	gcgagagccc	caggcgcggt	tacccggttt	cattttcagt	tgaggccaaa	atcccccgcg
961	gttggtcggg	gcgggggcgc	gctcggggga	ctgggctgac	cgcggggcgc	gggccagggt
1021	ctcacaccct	agggagatgc	tacggtcgcg	acgtggggcc	gacggggcgc	ctcctcgctg
1081	ggcatgacca	gtacgcctac	gacggcaagg	attacatcgc	cctgaacgag	gacctgcgct
1141	cttgaccgcg	cgcggacacg	gcggtcaga	tcaccacgag	caagtgggag	gcgcccctgt
1201	agggcgagca	gagagagacc	tacctggagg	gagagtgcgt	ggagtggctc	gcgacatacc
1261	tggagaacgg	gaaggacaag	ctggagcgcg	ctggtaccag	gggcagtggt	gagccttccc
1321	catctcctat	aggtcgcccg	ggatggcctc	ccacgagaag	aggaggaaa	tgggatcagc
1381	gtcaggaatg	gcctccctgc	tgaatggaga	atggcatgag	tttctcagag	ttctctctga
1441	gggccccctc	ttctctctag	acaattaagg	aatgacgtct	ctgaggaat	ggaggggaa
1501	acagtcctca	gaatactgat	caggggctcc	ctttgacccc	tgacgacgc	ttgggaaccg
1561	tgacttttcc	tcagagcctg	tgttctctgc	ctcacactca	ggtgttttgg	ggctctgatt
1621	ccagcacttc	tgagtcaact	tacctccact	cagatcagga	gcagaagtcc	ctgttccccg
1681	ctcagagact	cgaactttcc	aatgaaatag	agattatccc	aggtgcctgc	gtccaggctg
1741	gtgtctgggt	tctgtgcccc	ttccccacc	caggtgtcct	gtccattctc	aggtcgttca
1801	catgggtggt	cctagggtgt	cccataaaag	atgcaaagcg	cctgaatttt	ctgactcttc
1861	ccatcagacc	ccccaaagc	acacgtgacc	caccacccca	tctctgacca	tgaggccacc
1921	ctgaggtgct	ggccctcggg	tttctaccct	gcgagatca	cactgacctg	gcagcgggat
1981	ggcagggacc	aaactcagga	cactgagctt	gtggagacca	gaccagcagg	agatagaacc
2041	ttccagaagt	gggcagctgt	ggtggtgctc	tctggagaag	agcagagata	cacatgccat
2101	gtacagcatg	gaagccctgc	gaagccctgc	accctgagat	gggtaagga	gggggatgag
2161	gggtcatatc	tcttctcagg	gaaagcagga	gcccttcacg	agggtcaggg	cccctcatct
2221	tccccctctt	tcccagagcc	gtcttcccag	ttcacccgtc	ccatcgttgg	catgttctgt
2281	ggctcggctg	tcctagcagt	tgtgtctatc	ggagctgttg	tcgctgtctg	gatgtgtagg
2341	aggaagagtt	caggtagggg	aggggtgagg	ggtggggtct	gggttttctt	gtcccactgg
2401	gggtttcaag	cccaggtag	aagtgttccc	tgctcatta	ctgggaagca	gcatgcacac
2461	agggcttaac	gcccctcggg	accctgtgtg	ccagcactta	ctgttttgg	cagcacatgt
2521	gacaatgaag	gatggatgta	tcacctgat	ggttgtgggt	ttggggctct	gattccagca
2581	ttcatgagtc	aggggaaggt	ccctgctaag	gacagacctt	aggagggcag	ttggtccagg
2641	accacactt	gctttctctg	tgtttctctg	tctgcccctg	ggtctgtagt	catacttctg
2701	gaaattcctt	ttgggtccaa	gactaggagg	ttcctctaag	atctcatggc	cctgcttctt
2761	cccagtgccc	tcacaggaca	ttttcttccc	acaggtggaa	aaggagggag	ctactctcag
2821	gctgcgtgta	agtgggtggg	gtgggagtgt	ggaggagctc	accacccca	taattctctc
2881	tgtcccacgt	ctcctcgggg	ctctgaccag	gtcctgtttt	tgttctactc	caggcagcga
2941	cagtgcccag	ggctctgatg	tgtctctcac	agcttgaaaa	ggtgagattc	tggggctcta
3001	gagtgggtgg	ggtggcgggg	ctgggggtgg	gtggggcaga	gggaaaaggc	ctgggtaatg
3061	gggattcttt	gattggtgatg	tttcgcgtgt	gtgggtggct	ggttagagta	tcactcgctta
3121	ccatgactaa	ccagaatttg	ttcatgactg	ttgtttctgt	tgacctgaga	cagctgctct
3181	gtgagggact	gagatgcagg	atctcttca	gcctcccctt	tgtgacttca	agagcctctg
3241	gcatctcttt	ctgcaaaggc	acctgaatgt	gtctgcgtcc	ctgttagcat	aatgtgagga
3301	ggtggagaga	cagcccacc	ttgtgtccac	tgtgaccctt	gttcgcagtc	tgacctgtgt
3361	ttcctcccca					

Thus,  $H_0$  is strongly rejected. This is not surprising since the observed numbers of  $a$ ,  $c$ ,  $g$ ,  $t$  are  $n_a = 621$ ,  $n_c = 974$ ,  $n_g = 1064$ ,  $n_t = 711$ , respectively.

TABLE 3  
Occurrences  $N_{ij}$  of the string  $ij$  for  $i, j \in \{a, c, g, t\}$

	<i>a</i>	<i>c</i>	<i>g</i>	<i>t</i>
<i>a</i>	91	160	261	108
<i>c</i>	213	351	161	249
<i>g</i>	251	224	388	201
<i>t</i>	66	239	254	152

EXAMPLE B. A Bayes test of  $H_0$ : i.i.d.(unknown) versus  $H_1$ : reversible.

Here we use a Dirichlet(1, 1, 1, 1) prior for the null hypothesis and the prior based on the complete graph  $K_4$  with loops (see Figure 5) and all edge weights equal to 1. The probability  $P(data|H_0)$  is calculated in Example A. In order to calculate  $P(data|H_1)$ , we first determine the transition counts  $k_e$  for our data (see Table 4) and also  $k_v = n_v - \delta_a(v)$ :

$$(58) \quad k_a = 620, \quad k_c = 974, \quad k_g = 1064, \quad k_t = 711.$$

We abbreviate  $E' = \{\{a, c\}, \{a, g\}, \{a, t\}, \{c, g\}, \{c, t\}, \{g, t\}\}$ . By the first part of Proposition 4.7,

$$(59) \quad \begin{aligned} P(data|H_1) &= \frac{\prod_{e \in E'} \prod_{i=0}^{k_e-1} (1+i) \prod_{j \in \{a,c,g,t\}} \prod_{i=0}^{k_{(j)}/2-1} (1+2i)}{\prod_{i=0}^{k_t-1} (4+2i) \prod_{j \in \{a,c,g\}} \prod_{i=0}^{k_j-1} (5+2i)} \\ &= (373)!(512)!(174)!(385)!(488)!(455)! \\ &\quad \times \prod_{i=0}^{90} (1+2i) \prod_{i=0}^{350} (1+2i) \prod_{i=0}^{387} (1+2i) \prod_{i=0}^{151} (1+2i) \\ &\quad \times \left( \prod_{i=0}^{710} (4+2i) \prod_{i=0}^{619} (5+2i) \prod_{i=0}^{973} (5+2i) \prod_{i=0}^{1063} (5+2i) \right)^{-1} \\ &\approx 2.166939224648291 \cdot 10^{-1961}. \end{aligned}$$

TABLE 4  
The undirected transition counts  $k_{\{i,j\}}$ ,  $i, j \in \{a, c, g, t\}$

	<i>a</i>	<i>c</i>	<i>g</i>	<i>t</i>
<i>a</i>	182	373	512	174
<i>c</i>	373	702	385	488
<i>g</i>	512	385	776	455
<i>t</i>	174	488	455	304

So the Bayes factor is

$$\frac{P(\text{data}|H_0)}{P(\text{data}|H_1)} \approx 5.2628 \cdot 10^{-39}$$

and the null hypothesis is strongly rejected.

EXAMPLE C. A Bayes test of  $H_0$ : reversible versus  $H_1$ : full Markov.

Here we use our conjugate prior on reversible chains with all constants chosen as one. We use product Dirichlet measure for the rows in the full Markov case. This yields

$$\begin{aligned} P(\text{data}|H_1) &= \prod_{i \in \{a,c,g,t\}} \Gamma(4) \frac{\prod_{j \in \{a,c,g,t\}} \Gamma(N_{ij} + 1)}{\Gamma(k_i + 4)} \\ &= \Gamma(4)^4 \cdot \frac{\Gamma(92)\Gamma(161)\Gamma(262)\Gamma(109)}{\Gamma(624)} \\ &\quad \times \frac{\Gamma(214)\Gamma(352)\Gamma(162)\Gamma(250)}{\Gamma(978)} \\ &\quad \times \frac{\Gamma(252)\Gamma(225)\Gamma(389)\Gamma(202)}{\Gamma(1068)} \cdot \frac{\Gamma(67)\Gamma(240)\Gamma(255)\Gamma(153)}{\Gamma(715)} \\ &\approx 4.16382063735625 \cdot 10^{-1956}. \end{aligned}$$

The probability  $P(\text{data}|H_0)$  was calculated in Example B. Hence,

$$\frac{P(\text{data}|H_0)}{P(\text{data}|H_1)} \approx 5.20421 \cdot 10^{-6}.$$

We see that a straightforward Bayes test rejects reversibility.

EXAMPLE D. A Bayes test of  $H_0$ : i.i.d.(unknown) versus  $H_1$ : full Markov.

Using the Bayes factors computed above, the null hypothesis is strongly rejected:

$$\frac{P(\text{data}|H_0)}{P(\text{data}|H_1)} \approx 2.73887 \cdot 10^{-44}.$$

Of course, an i.i.d. process is a reversible Markov chain.

In using the Dirichlet prior for testing uniformity with multinomial data and for testing independence in contingency tables, I. J. Good found the symmetric Dirichlet prior with density proportional to  $\prod_{i=1}^d x_i^{c-1}$  an important tool. Good's many insights into these testing problems may be accessed through his book [9] and the survey article [10].

We have used the analog of the symmetric Dirichlet for the reversible Markov chain context with all edge weights  $a_e$  equal to a constant  $c$  say. As  $c$  tends to

infinity, this prior tends to a point mass supported on the simple random walk on the graph. As  $c$  tends to zero, this prior tends to an improper prior which gives the maximum likelihood as its posterior.

Good also worked with  $c$ -mixtures of symmetric Dirichlet priors. We suspect that parallel, useful things can be done in our case as well.

We have not found *any* literature about statistical analysis of reversible Markov chains with unknown transitions and append two data analytic remarks here. First, under reversibility, the count  $N_{vv'}$  of  $v$  to  $v'$  transitions has the same expectation as the count  $N_{v'v}$  of  $v'$  to  $v$  transitions, namely,  $v(v)k(v, v')$ . This suggests looking at ratios  $N_{vv'}/N_{v'v}$  or differences  $N_{vv'} - N_{v'v}$ . For example, from Table 3,  $N_{ac}/N_{ca} = 160/213$ ,  $N_{ag}/N_{ga} = 261/251$ ,  $N_{at}/N_{ta} = 108/66$ ,  $N_{cg}/N_{gc} = 161/224$ ,  $N_{ct}/N_{tc} = 249/239$ ,  $N_{gt}/N_{tg} = 201/254$ ; most of these are way off.

In large samples, these counts have limiting normal distributions by results of Höglund [11]. A second data analytic tool would be to estimate the stationary distribution [perhaps by the method of moments estimator  $\hat{v}(v) = \frac{1}{n}|\{i \leq n : X_i = v\}|$ ] and also estimate the transition matrix, and then compare  $\hat{v}(v)\hat{k}(v, v')$  with  $\hat{v}(v')\hat{k}(v', v)$ .

An interesting problem not tackled here is finding natural priors on the set of reversible Markov chains *with a fixed stationary distribution*. For definiteness, consider the uniform stationary distribution. Then the problem is to put a prior on  $\mathcal{S}(n)$ , the symmetric doubly stochastic  $n \times n$  matrices. We make two remarks. First, determining the Euclidean volume of  $\mathcal{S}(n)$  is a long-standing open problem; see [1] for recent results. Second,  $\mathcal{S}(n)$  is a compact, convex subset of  $\mathbb{R}^{n^2}$ . Its extreme points are well known to be the *symmetrized* permutation matrices (see [17]). Thus, if  $\pi$  is a permutation matrix on  $n$  letters with  $e(\pi)$  the usual  $n \times n$  permutation matrix, let  $\tilde{e}(\pi) = \frac{1}{2}[e(\pi) + e(\pi^{-1})]$ . The extreme points of  $\mathcal{S}(n)$  are  $(\tilde{e}(\pi))$  as  $\pi$  ranges over permutations in  $S_n$ . We may put a prior on  $\mathcal{S}(n)$  by taking a random convex combination of the  $\tilde{e}(\pi)$ . Alas,  $\mathcal{S}(n)$  is *not* a simplex, so symmetric weights on the extreme points may not lead to symmetric measures on  $\mathcal{S}(n)$ .

**Acknowledgment.** We would like to thank Franz Merkl for some interesting discussions.

## REFERENCES

- [1] CHAN, C. S., ROBBINS, D. P. and YUEN, D. S. (2000). On the volume of a certain polytope. *Experiment. Math.* **9** 91–99. [MR1758803](#)
- [2] DALAL, S. R. and HALL, W. J. (1983). Approximating priors by mixtures of natural conjugate priors. *J. Roy. Statist. Soc. Ser. B* **45** 278–286. [MR0721753](#)
- [3] DIACONIS, P. (1988). Recent progress on de Finetti's notions of exchangeability. In *Bayesian Statistics 3* (J.-M. Bernardo, M. H. DeGroot, D. V. Lindley and A. F. M. Smith, eds.) 111–125. Oxford Univ. Press, New York. [MR1008047](#)

- [4] DIACONIS, P. and FREEDMAN, D. (1980). de Finetti's theorem for Markov chains. *Ann. Probab.* **8** 115–130. [MR0556418](#)
- [5] DIACONIS, P. and YLVISAKER, D. (1985). Quantifying prior opinion (with discussion). In *Bayesian Statistics 2* (J.-M. Bernardo, M. H. DeGroot, D. V. Lindley and A. F. M. Smith, eds.) 133–156. North-Holland, Amsterdam. [MR0862488](#)
- [6] FORTINI, S., LADELLI, L., PETRIS, G. and REGAZZINI, E. (2002). On mixtures of distributions of Markov chains. *Stochastic Process. Appl.* **100** 147–165. [MR1919611](#)
- [7] FREEDMAN, D. A. (1962). Mixtures of Markov processes. *Ann. Math. Statist.* **33** 114–118. [MR0137156](#)
- [8] GIBLIN, P. J. (1981). *Graphs, Surfaces and Homology. An Introduction to Algebraic Topology*, 2nd ed. Chapman and Hall, London. [MR0643363](#)
- [9] GOOD, I. J. (1968). *The Estimation of Probabilities. An Essay on Modern Bayesian Methods*. MIT Press, Cambridge, MA. [MR0245135](#)
- [10] GOOD, I. J. and CROOK, J. F. (1987). The robustness and sensitivity of the mixed-Dirichlet Bayesian test for “independence” in contingency tables. *Ann. Statist.* **15** 670–693. [MR0888433](#)
- [11] HÖGLUND, T. (1974). Central limit theorems and statistical inference for finite Markov chains. *Z. Wahrsch. Verw. Gebiete* **29** 123–151. [MR0373201](#)
- [12] KEANE, M. S. (1990). Solution to problem 288. *Statist. Neerlandica* **44** 95–100.
- [13] KEANE, M. S. and ROLLES, S. W. W. (2000). Edge-reinforced random walk on finite graphs. In *Infinite Dimensional Stochastic Analysis* (Ph. Clément, F. den Hollander, J. van Neerven and B. de Pagter, eds.) 217–234. R. Neth. Acad. Arts Sci., Amsterdam. [MR1832379](#)
- [14] MAURER, S. B. (1976). Matrix generalizations of some theorems on trees, cycles and cocycles in graphs. *SIAM J. Appl. Math.* **30** 143–148. [MR0392635](#)
- [15] PEMANTLE, R. (1988). Phase transition in reinforced random walk and RWRE on trees. *Ann. Probab.* **16** 1229–1241. [MR0942765](#)
- [16] ROLLES, S. W. W. (2003). How edge-reinforced random walk arises naturally. *Probab. Theory Related Fields* **126** 243–260. [MR1990056](#)
- [17] STANLEY, R. P. (1978). Generating functions. In *Studies in Combinatorics* (G.-C. Rota, ed.) 100–141. Math. Assoc. Amer., Washington. [MR0513004](#)
- [18] ZABELL, S. L. (1982). W. E. Johnson's “sufficientness” postulate. *Ann. Statist.* **10** 1090–1099. [MR0673645](#)
- [19] ZABELL, S. L. (1995). Characterizing Markov exchangeable sequences. *J. Theoret. Probab.* **8** 175–178. [MR1308676](#)

DEPARTMENT OF STATISTICS  
 STANFORD UNIVERSITY  
 STANFORD, CALIFORNIA 94305-4065  
 USA  
 E-MAIL: [diaconis@math.stanford.edu](mailto:diaconis@math.stanford.edu)

EINDHOVEN UNIVERSITY OF TECHNOLOGY  
 DEPARTMENT OF MATHEMATICS  
 AND COMPUTER SCIENCE  
 P.O. BOX 513  
 5600 MB EINDHOVEN  
 THE NETHERLANDS  
 E-MAIL: [srolles@win.tue.nl](mailto:srolles@win.tue.nl)