

Statistical inference versus mean field limit for Hawkes processes

Sylvain Delattre

*Laboratoire de Probabilités et Modèles Aléatoires, UMR 7599, Université Paris Diderot,
Case courrier 7012, avenue de France, 75205 Paris Cedex 13, France*

e-mail: sylvain.delattre@univ-paris-diderot.fr

and

Nicolas Fournier

*Laboratoire de Probabilités et Modèles Aléatoires, UMR 7599, Université Pierre-et-Marie
Curie, Case 188, 4 place Jussieu, F-75252 Paris Cedex 5, France*

e-mail: nicolas.fournier@upmc.fr

Abstract: We consider a population of N individuals, of which we observe the number of *actions* until time t . For each couple of individuals (i, j) , j may or not *influence* i , which we model by i.i.d. Bernoulli(p)-random variables, for some unknown parameter $p \in (0, 1]$. Each individual acts *autonomously* at some unknown rate $\mu > 0$ and acts *by mimetism* at some rate proportional to the sum of some function φ of the ages of the actions of the individuals which influence him. The function φ is unknown but assumed, roughly, to be decreasing and with fast decay. The goal of this paper is to estimate p , which is the main characteristic of the *graph of interactions*, in the asymptotic $N \rightarrow \infty, t \rightarrow \infty$. The main issue is that the mean field limit (as $N \rightarrow \infty$) of this model is unidentifiable, in that it only depends on the parameters μ and $p\varphi$. Fortunately, this mean field limit is not valid for large times. We distinguish the subcritical case, where, roughly, the mean number m_t of actions per individual increases linearly and the supercritical case, where m_t increases exponentially. Although the nuisance parameter φ is non-parametric, we are able, in both cases, to estimate p without estimating φ in a nonparametric way, with a precision of order $N^{-1/2} + N^{1/2}m_t^{-1}$, up to some arbitrarily small loss. We explain, using a Gaussian toy model, the reason why this rate of convergence might be (almost) optimal.

MSC 2010 subject classifications: 62M09, 60J75, 60K35.

Keywords and phrases: Multivariate Hawkes processes, point processes, statistical inference, interaction graph, stochastic interacting particle systems, propagation of chaos, mean field limit.

Received September 2015.

1. Introduction and main results

1.1. Setting

We consider some unknown parameters $p \in (0, 1]$, $\mu > 0$ and $\varphi : [0, \infty) \mapsto [0, \infty)$. For $N \geq 1$, we consider an i.i.d. family $(\pi^i(dt, dz))_{i=1, \dots, N}$ of Poisson measures

on $[0, \infty) \times [0, \infty)$ with intensity measure $dt dz$, independent of an i.i.d. family $(\theta_{ij})_{i,j=1,\dots,N}$ of Bernoulli(p)-distributed random variables. We also consider the system of equations, for $i = 1, \dots, N$,

$$Z_t^{i,N} = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq \lambda_s^{i,N}\}} \pi^i(ds, dz),$$

$$\text{where } \lambda_t^{i,N} = \mu + \frac{1}{N} \sum_{j=1}^N \theta_{ij} \int_0^{t-} \varphi(t-s) dZ_s^{j,N}. \quad (1)$$

Here and in the whole paper, \int_0^t means $\int_{[0,t]}$ and \int_0^{t-} means $\int_{[0,t)}$. The solution $((Z_t^{i,N})_{t \geq 0})_{i=1,\dots,N}$ is a family of N counting processes (that is, a.s. integer-valued, càdlàg and non-decreasing). The following well-posedness result is more or less well-known, see e.g. Brémaud-Massoulié [9] and [13] (we will apply directly the latter reference).

Proposition 1. *Assume that φ is locally integrable and fix $N \geq 1$. The system (1.1) has a unique càdlàg $(\mathcal{F}_t)_{t \geq 0}$ -adapted solution $((Z_t^{i,N})_{t \geq 0})_{i=1,\dots,N}$ such that $\sum_{i=1}^N \mathbb{E}[Z_t^{i,N}] < \infty$ for all $t \geq 0$, where $\mathcal{F}_t = \sigma(\pi^i(A) : A \in \mathcal{B}([0, t] \times [0, \infty)), i = 1, \dots, N) \vee \sigma(\theta_{ij} : i, j = 1, \dots, N)$.*

Let us provide a brief heuristic description of this process. We have N individuals and $Z_t^{i,N}$ stands for the number of actions of the i -th individual until t . We say that j influences i if and only if $\theta_{ij} = 1$ (with possibly $i = j$). Each individual i acts, at time t , with rate $\lambda_t^{i,N}$. In other words, each individual has an autonomous rate of action μ as well as a subordinate rate of action $N^{-1} \sum_{j=1}^N \theta_{ij} \int_0^t \varphi(t-s) dZ_s^{j,N}$, which depends on the number of actions of the individuals that influence him, with a weight N^{-1} and taking into account the age of these actions through φ . If for example $\varphi = a \mathbf{1}_{[0,K]}$, then the subordinate rate of action of i is simply a/N times the total number of actions, during $[t-K, t]$, of all the individuals that influence him.

As is well-known, a phase-transition occurs for such a model, see Hawkes-Oakes [18] (or [13] for such considerations on large networks): setting $\Lambda = \int_0^\infty \varphi(t) dt$,

- in the subcritical case where $\Lambda p < 1$, we will see that $Z_t^{1,N}$ increases linearly with time, at least on the event where the family $(\theta_{ij})_{i,j=1,\dots,N}$ behaves reasonably;

- in the supercritical case where $\Lambda p > 1$, we will see that $Z_t^{1,N}$ increases exponentially fast with time, at least on the event where the family $(\theta_{ij})_{i,j=1,\dots,N}$ behaves reasonably.

The limit theorems, and thus the statistical inference, completely differ in both cases, so that the present paper contains essentially two independent parts.

We will not study the critical case where $\Lambda p = 1$ because it is a very particular case. However, it would be very interesting to understand what happens near the critical case. Our results say nothing about this problem.

1.2. Assumptions

Recalling that $\Lambda = \int_0^\infty \varphi(s) ds$, we will work under one of the two following conditions: either for some $q \geq 1$,

$$\mu \in (0, \infty), \quad \Lambda p \in (0, 1) \quad \text{and} \quad \int_0^\infty s^q \varphi(s) ds < \infty \quad (H(q))$$

or

$$\mu \in (0, \infty), \quad \Lambda p \in (1, \infty) \quad \text{and} \quad \int_0^t |d\varphi(s)| \text{ increases at most polynomially.} \quad (A)$$

In many applications, φ is smooth and has a fast decay, so that, except in the critical case, either $H(q)$ is satisfied for all $q \geq 1$ or A is satisfied.

1.3. References and fields of application

Hawkes processes have been introduced by Hawkes [17] and Oakes-Hawkes [18] have found a noticeable representation of such processes in terms of Galton-Watson trees. Since then, there has been a huge literature on Hawkes processes, see e.g. Daley and Vere-Jones [12] for an introduction, Massoulié [24], Brémaud-Massoulié [9] and [13] for stability results, Brémaud-Nappo-Torrisi [10], Zhu [35, 36] and [3] for limit theorems, etc. Hawkes processes are used in various fields of applications:

- earthquake replicas in seismology, see Helmstetter-Sornette [19], Kagan [23], Ogata [26], Bacry-Muzy [5],
- spike trains for brain activity in neuroscience, see Grün et al. [15], Okatan et al. [27], Pillow et al. [28], Reynaud et al. [31, 32],
- genome analysis, see Reynaud-Schbath [30],
- various fields of mathematical finance, see Aït-Sahalia et al. [1], Bauwens-Hautsch [6], Hewlett [20], Bacry et al. [2], Bacry-Muzy [4, 5],
- social networks interactions, see Blundell et al. [8] and Zhou et al. [34].

Concerning the statistical inference for Hawkes processes, only the case of fixed finite dimension N has been studied, to our knowledge, in the asymptotic $t \rightarrow \infty$ (for possibly more general shapes of interaction). Some parametric and nonparametric estimation procedures for μ and φ have been proposed, with or without rigorous proofs. Let us mention Ogata [25], Bacry-Muzy [5], [2], the various recent results of Hansen et al. [16] and Reynaud et al. [30, 31, 32], as well as the Bayesian study of Rasmussen [29].

1.4. Goals and motivation

In many applications, the number of individuals is very large (think of neurons, financial agents or of social networks). Then we need some estimators in the asymptotic where N and t tend simultaneously to infinity. This problem seems to be completely open.

We assume that we observe $(Z_s^{i,N})_{i=1,\dots,N,s\in[0,t]}$ (or $(Z_s^{i,N})_{i=1,\dots,N,s\in[0,2t]}$ for convenience), that is all the actions of the individuals on some (large) time interval.

In our point of view, we only observe the activity of the individuals, we do not know the *graph of interactions*. A very similar problem was studied in [32], although in fixed finite dimension N . Our goal is to estimate p , which can be seen as the main characteristic of the graph of interactions, since it represents the proportion of open edges. We consider μ and φ as nuisance parameters, although this is debatable. In the supercritical case, we will be able to estimate p without estimating μ nor φ . In the subcritical case, we will be able to recover p estimating only μ and the integral Λ of φ . In any case, we will *not* need to provide a nonparametric estimation of φ , and we believe it is a very good point: it would require regularity assumptions and would complicate a lot the study.

The main goal of this paper is to provide the basic tools for the statistical estimation of Hawkes processes when both the graph size and the observation time increase. Of course, this is only a toy model and we have no precise idea of real world applications, although we can think e.g. of neurons spiking: they are clearly numerous (so N is large), we can only observe their activities (each time they spike), and we would like to have an idea of the graph of interactions. See again [32] for a more convincing biological background. Think also of financial agents: they are also numerous, we can observe their actions (each time they buy or sell a product), and we would like to recover the interaction graph.

1.5. Mean field limit

We quickly describe the expected *chaotic* behavior of $((Z_t^{i,N})_{t\geq 0})_{i=1,\dots,N}$ as $N \rightarrow \infty$. We refer to Sznitman [33] for an introduction to *propagation of chaos*. Extending the method of [13, Theorem 8], it is not hard to check, assuming that $\int_0^\infty \varphi^2(s)ds < \infty$, that for each given $k \geq 1$ and $T > 0$, the sample $((Z_t^{i,N})_{t\in[0,T]})_{i=1,\dots,k}$ goes in law, as $N \rightarrow \infty$, to a family $((Y_t^i)_{t\in[0,T]})_{i=1,\dots,k}$ of i.i.d. inhomogeneous Poisson processes with intensity $(\lambda_t)_{t\geq 0}$, unique locally bounded nonnegative solution to $\lambda_t = \mu + \int_0^t p\varphi(t-s)\lambda_s ds$.

On the one hand, approximate independence is of course a good point for statistical inference. On the other hand, the mean-field limit (i.e. the $(Y_t^i)_{t\geq 0}$'s) depends on p and φ only through $(\lambda_t)_{t\geq 0}$ and thus through $p\varphi$, which is a negative point: the mean-field limit is *unidentifiable*. The situation is however not hopeless because roughly, the mean-field limit *does not* hold true for the whole sample $(Z_t^{i,N})_{i=1,\dots,N}$ and is less and less true as time becomes larger and larger.

1.6. Main result in the subcritical case

For $N \geq 1$ and for $((Z_t^{i,N})_{t\geq 0})_{i=1,\dots,N}$ the solution to (1.1), we introduce $\bar{Z}_t^N = N^{-1} \sum_{i=1}^N Z_t^{i,N}$. We mention in the following remark, that we will prove later,

that the number of actions per individual increases linearly in the subcritical case.

Remark 2. Assume $H(1)$. Then for all $\varepsilon > 0$,

$$\lim_{(N,t) \rightarrow (\infty, \infty)} \Pr \left(\left| \frac{\bar{Z}_t^N}{t} - \frac{\mu}{1 - \Lambda p} \right| \geq \varepsilon \right) = 0.$$

We next introduce

$$\mathcal{E}_t^N = \frac{\bar{Z}_{2t}^N - \bar{Z}_t^N}{t}, \quad \mathcal{V}_t^N = \sum_{i=1}^N \left(\frac{Z_{2t}^{i,N} - Z_t^{i,N}}{t} - \mathcal{E}_t^N \right)^2 - \frac{N}{t} \mathcal{E}_t^N,$$

$$\mathcal{W}_{\Delta,t}^N = 2\mathcal{Z}_{2\Delta,t}^N - \mathcal{Z}_{\Delta,t}^N, \quad \text{where} \quad \mathcal{Z}_{\Delta,t}^N = \frac{N}{t} \sum_{k=t/\Delta+1}^{2t/\Delta} \left(\bar{Z}_{k\Delta}^N - \bar{Z}_{(k-1)\Delta}^N - \Delta \mathcal{E}_t^N \right)^2.$$

In the last expression, $\Delta \in (0, t)$ is required to be such that $t/(2\Delta) \in \mathbb{N}^*$.

Theorem 3. We assume $H(q)$ for some $q > 3$. For $t \geq 1$, we put $\Delta_t = t/(2\lfloor t^{1-4/(q+1)} \rfloor)$: it holds that $t/(2\Delta_t) \in \mathbb{N}^*$ and that $\Delta_t \sim t^{4/(q+1)}/2$ as $t \rightarrow \infty$. There is a constant C depending only on p, μ, φ and q such that for all $\varepsilon \in (0, 1)$, all $N \geq 1$, all $t \geq 1$,

$$\begin{aligned} \Pr \left(\left| \mathcal{E}_t^N - \frac{\mu}{1 - \Lambda p} \right| \geq \varepsilon \right) &\leq \frac{C}{\varepsilon} \left(\frac{1}{N} + \frac{1}{\sqrt{Nt}} + \frac{1}{t^q} \right), \\ \Pr \left(\left| \mathcal{V}_t^N - \frac{\mu^2 \Lambda^2 p(1-p)}{(1-\Lambda p)^2} \right| \geq \varepsilon \right) &\leq \frac{C}{\varepsilon} \left(\frac{\sqrt{N}}{t} + \frac{1}{\sqrt{N}} \right), \\ \Pr \left(\left| \mathcal{W}_{\Delta_t,t}^N - \frac{\mu}{(1-\Lambda p)^3} \right| \geq \varepsilon \right) &\leq \frac{C}{\varepsilon} \left(\frac{1}{N} + \frac{N}{t^2} + \frac{1}{\sqrt{t^{1-4/(q+1)}}} \right). \end{aligned}$$

We will easily deduce the following corollary.

Corollary 4. We assume $H(q)$ for some $q > 3$. For $t \geq 1$, we put $\Delta_t = t/(2\lfloor t^{1-4/(q+1)} \rfloor)$. There is a constant C depending only on p, μ, φ and q such that for all $\varepsilon \in (0, 1)$, all $N \geq 1$, all $t \geq 1$,

$$\begin{aligned} \Pr \left(\left\| \Psi \left(\mathcal{E}_t^N, \mathcal{V}_t^N, \mathcal{W}_{\Delta_t,t}^N \right) - (\mu, \Lambda, p) \right\| \geq \varepsilon \right) &\leq \frac{C}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{t^{1-4/(q+1)}}} \right) \\ &\leq \frac{2C}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{t^{1-4/(q+1)}} \right), \end{aligned}$$

where $\Psi = \mathbf{1}_D \Phi$ with $D = \{(u, v, w) \in \mathbb{R}^3 : w > u > 0 \text{ and } v \geq 0\}$ and $\Phi : D \mapsto \mathbb{R}^3$ defined by

$$\begin{aligned} \Phi_1(u, v, w) &= u \sqrt{\frac{u}{w}}, \quad \Phi_2(u, v, w) = \frac{v + [u - \Phi_1(u, v, w)]^2}{u[u - \Phi_1(u, v, w)]}, \\ \Phi_3(u, v, w) &= \frac{1 - u^{-1} \Phi_1(u, v, w)}{\Phi_2(u, v, w)}. \end{aligned}$$

We did not optimize the dependence in q : in many applications, $H(q)$ holds for all $q \geq 1$.

1.7. Main result in the supercritical case

For $N \geq 1$ and for $((Z_t^{i,N})_{t \geq 0})_{i=1, \dots, N}$ the solution to (1.1), we set $\bar{Z}_t^N = N^{-1} \sum_{i=1}^N Z_t^{i,N}$. We will check later the following remark, which states that the mean number of actions per individual increases exponentially in the supercritical case.

Remark 5. Assume A and consider $\alpha_0 > 0$ defined by $p \int_0^\infty e^{-\alpha_0 t} \varphi(t) dt = 1$. Then

$$\text{for all } \eta > 0, \quad \lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} \Pr \left(\bar{Z}_t^N \in [e^{(\alpha_0 - \eta)t}, e^{(\alpha_0 + \eta)t}] \right) = 1.$$

We next introduce

$$\mathcal{U}_t^N = \left[\sum_{i=1}^N \left(\frac{Z_t^{i,N} - \bar{Z}_t^N}{\bar{Z}_t^N} \right)^2 - \frac{N}{\bar{Z}_t^N} \right] \mathbf{1}_{\{\bar{Z}_t^N > 0\}} \quad \text{and} \quad \mathcal{P}_t^N = \frac{1}{\mathcal{U}_t^N + 1} \mathbf{1}_{\{\mathcal{U}_t^N \geq 0\}}.$$

Theorem 6. Assume A and consider $\alpha_0 > 0$ defined in Remark 5. For all $\eta > 0$, there is a constant $C_\eta > 0$ (depending only on p, μ, φ, η) such that for all $N \geq 1$, all $t \geq 1$, all $\varepsilon \in (0, 1)$,

$$\Pr \left(|\mathcal{P}_t^N - p| \geq \varepsilon \right) \leq \frac{C_\eta e^{\eta t}}{\varepsilon} \left(\frac{\sqrt{N}}{e^{\alpha_0 t}} + \frac{1}{\sqrt{N}} \right).$$

1.8. Detecting subcriticality and supercriticality

In practise, we may of course not know if we are in the subcritical or supercritical case.

Proposition 7. (i) Under $H(1)$, there are some constants $0 < c < C$ depending only on p, μ, φ such that for all $N \geq 1$, all $t \geq 1$, $\Pr(\log(\bar{Z}_t^N) \geq (\log t)^2) \leq C(e^{-cN} + t^{1/2} e^{-(\log t)^2})$.

(ii) Under A, for all $\eta > 0$, there is a constant C_η depending only on p, μ, φ, η such that for all $N \geq 1$, all $t \geq 1$, $\Pr(\log(\bar{Z}_t^N) \leq (\log t)^2) \leq C_\eta e^{\eta t} (N^{-1/2} + e^{-\alpha_0 t})$.

It is then not hard to check that, with the notation of Corollary 4 and Theorem 6, under $H(q)$ (for some $q > 3$) or A, the estimator

$$\hat{p}_t^N = \mathbf{1}_{\{\log(\bar{Z}_t^N) < (\log t)^2\}} \Psi_3(\mathcal{E}_{t/2}^N, \mathcal{V}_{t/2}^N, \mathcal{W}_{\Delta_{t/2}, t/2}^N) + \mathbf{1}_{\{\log(\bar{Z}_t^N) > (\log t)^2\}} \mathcal{P}_t^N,$$

which is based on the observation of $(Z_s^{i,N})_{s \in [0, t], i=1, \dots, N}$, converges in probability to p , with the same speed of convergence as in Corollary 4 (under $H(q)$ for some $q > 3$) or as in Theorem 6 (under A).

1.9. About optimality

In Subection 2.3, we will see on a toy model that there is no real hope to find an estimator of p with a better precision than $N^{-1/2} + N^{1/2} m_t^{-1}$, where m_t is something like the mean number of jumps per individual during $[0, t]$.

Consequently, we believe that the precision we found in Corollary 4 is almost optimal, since then $m_t \simeq t$ by Remark 2 and since we reach the precision $N^{-1/2} + N^{1/2}t^{\alpha-1}$ for any $\alpha > 0$ (if φ has a fast decay), so that the loss is arbitrarily small. Similarly, the precision found in Theorem 6 is rather satisfying, since then $m_t \simeq e^{\alpha_0 t}$ by Remark 5 and since we reach the precision $e^{\eta t}(N^{-1/2} + N^{1/2}e^{-\alpha_0 t})$ for any $\eta > 0$, so that the loss is, here also, arbitrarily small.

The main default of the present paper is that the constants in Corollary 4 and in Theorem 6 strongly depend on the parameters μ, Λ, p . They also depend on q in the subcritical case. In particular, it would be quite delicate to understand how they behave when approaching, from below or from above, the critical case.

1.10. About the modeling

There are two main limitations in our setting.

Assuming that the θ_{ij} 's are i.i.d. is of course a strong assumption. What we really need is that the family $(\theta_{ij})_{i,j=1,\dots,N}$ satisfies similar properties as those shown in Subsection 4.1 (in the subcritical case) and in Subsection 5.1 (in the supercritical case). This clearly requires that the family $(\theta_{ij})_{i,j=1,\dots,N}$ is not too far from being i.i.d., and it does *not* suffice that $\lim_{N \rightarrow \infty} N^{-2} \sum_{i,j=1}^N \theta_{ij} = p$. However, we believe that all the conclusions of the present paper are still true if one assumes that $(\theta_{ij})_{1 \leq i \leq j \leq N}$ is i.i.d. and that $\theta_{ji} = \theta_{ij}$ for all $1 \leq i < j \leq N$, which might be the case in some applications where the interactions are symmetric. A rigorous proof would require some work but should not be too hard. We will study this problem numerically at the end of the paper.

Assuming that we observe all the population is also rather stringent. It would be interesting to study what happens if one observes only $(Z_s^{i,N})_{i=1,\dots,K,s \in [0,t]}$, for some K large but smaller than N . It is not difficult to guess how to adapt the estimators to such a situation (see Section 7 for precise formulae). The theoretical analysis would require a careful and tedious study. Again, we will discuss this numerically.

1.11. Notation

We denote by \Pr_θ the conditional probability knowing $(\theta_{ij})_{i,j=1,\dots,N}$. We introduce $\mathbb{E}_\theta, \mathbb{V}\text{ar}_\theta$ and Cov_θ accordingly.

For two functions $f, g : [0, \infty) \mapsto \mathbb{R}$, we introduce (if it exists) $(f \star g)(t) = \int_0^t f(t-s)g(s)ds$. The functions $\varphi^{\star n}$ will play an important role in the paper. Observe that, since $\int_0^\infty \varphi(s)ds = \Lambda$, $\int_0^\infty \varphi^{\star n}(s)ds = \Lambda^n$. We adopt the conventions $\varphi^{\star 0}(s)ds = \delta_0(ds)$ and $\varphi^{\star 0}(t-s)ds = \delta_t(ds)$. We also adopt the convention that $\varphi^{\star n}(s) = 0$ for $s < 0$.

All the finite constants used in the upperbounds are denoted by C , the positive constants used in the lowerbounds are denoted by c and their values change from line to line. They are allowed to depend only on μ, p and φ (and on q under $H(q)$), but never on N nor on t . Any other dependence will be indicated

in subscript. For example, C_η is a finite constant depending only on μ , p , φ and η (and on q under $H(q)$).

1.12. Plan of the paper

In the next section, we try to give the main reasons why our estimators should be convergent, which should help the reader to understand the strategies of the proofs. We also briefly and formally introduce a Gaussian toy model in Section 2.3 to show that the rates of convergence we obtain are not far from being the best we can hope for. In Section 3, we prove Proposition 1 (strong existence and uniqueness of the process) and check a few more or less explicit formulae concerning $(Z_t^{i,N})_{i=1,\dots,N,t\geq 0}$ of constant use. Section 4 is devoted to the proof of Theorem 3 and Corollary 4 (main results in the subcritical case). Theorem 6 (main result in the supercritical case) is proved in Section 5. We check Proposition 7 in Section 6. Finally, we illustrate numerically the results of the paper and some possible extensions in the last section.

2. Heuristics

This section is completely informal and the symbol \simeq means nothing precise. For example, “ $Z_t^{i,N} \simeq \mathbb{E}_\theta[Z_t^{i,N}]$ for t large” should be understood as “we hope that $Z_t^{i,N}/\mathbb{E}_\theta[Z_t^{i,N}]$ tends to 1 as $t \rightarrow \infty$ in probability or in another sense.”

2.1. The subcritical case

We assume that $\Lambda p \in [0, 1)$ and try to explain roughly the asymptotics of $(Z_t^{i,N})_{i=1,\dots,N,t\geq 0}$ and where the three estimators \mathcal{E}_t^N , \mathcal{V}_t^N and $\mathcal{W}_{\Delta,t}^N$ come from. We introduce the matrices $A_N(i, j) = N^{-1}\theta_{ij}$ and $Q_N = (I - \Lambda A_N)^{-1}$, which exists with high probability because $\Lambda p < 1$. We also set $\ell_N(i) = \sum_{j=1}^N Q_N(i, j)$ and $c_N(i) = \sum_{j=1}^N Q_N(j, i)$.

Fixing N and knowing $(\theta_{ij})_{i,j=1,\dots,N}$, we expect that $Z_t^{i,N} \simeq \mathbb{E}_\theta[Z_t^{i,N}]$ for t large by a law of large numbers. Next, it is not hard to check that $\mathbb{E}_\theta[Z_t^{i,N}] = \mu t + N^{-1} \sum_{j=1}^N \theta_{ij} \int_0^t \varphi(t-s) \mathbb{E}_\theta[Z_s^{j,N}] ds$. Assume that $\gamma_N(i) = \lim_{t \rightarrow \infty} t^{-1} \mathbb{E}_\theta[Z_t^{i,N}]$ exists for each $i = 1, \dots, N$. Then, using that $\int_0^t \varphi(t-s) ds \simeq \Lambda t$ for t large, we find that the vector γ_N must solve $\gamma_N = \mu \mathbf{1}_N + \Lambda A_N \gamma_N$, where $\mathbf{1}_N$ is the N -dimensional vector with all coordinates equal to 1. This implies that $\gamma_N = \mu(I - \Lambda A_N)^{-1} \mathbf{1}_N = \mu \ell_N$. We thus expect that $Z_t^{i,N} \simeq \mathbb{E}_\theta[Z_t^{i,N}] \simeq \mu \ell_N(i) t$.

Based on this and setting $\bar{\ell}_N = N^{-1} \sum_{i=1}^N \ell_N(i)$, we expect that $\bar{Z}_t^N \simeq \mu \bar{\ell}_N t$ for large values of t , whence $\bar{\mathcal{E}}_t^N := t^{-1} \bar{Z}_t^N \simeq \mu \bar{\ell}_N$.

Knowing $(\theta_{ij})_{i,j=1,\dots,N}$, $Z_t^{1,N}$ should resemble, roughly, a Poisson process, so that it should approximately hold true that $\text{Var}_\theta(Z_t^{1,N}) \simeq \mathbb{E}_\theta[Z_t^{1,N}]$. Consequently, $N^{-1} \sum_{i=1}^N (Z_t^{i,N} - \bar{Z}_t^N)^2$ should resemble $\text{Var}(Z_t^{1,N}) = \text{Var}(\mathbb{E}_\theta[Z_t^{1,N}]) +$

$\mathbb{E}[\text{Var}_\theta(Z_t^{1,N})] \simeq \text{Var}(\mathbb{E}_\theta[Z_t^{1,N}] + \mathbb{E}[Z_t^{1,N}])$, which itself should be approximately equal to $N^{-1} \sum_{i=1}^N (\mathbb{E}_\theta[Z_t^{i,N}] - \mathbb{E}_\theta[\bar{Z}_t^N])^2 + \bar{Z}_t^N \simeq N^{-1} \mu^2 t^2 \sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2 + \bar{Z}_t^N$. All in all, we expect that $\tilde{\mathcal{V}}_t^N := t^{-2} [\sum_{i=1}^N (Z_t^{i,N} - \bar{Z}_t^N)^2 - N \bar{Z}_t^N] \simeq \mu^2 \sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2$ for t large.

The temporal empirical variance $\Delta t^{-1} \sum_{k=1}^{t/\Delta} [\bar{Z}_{k\Delta}^N - \bar{Z}_{(k-1)\Delta}^N - \Delta t^{-1} \bar{Z}_t^N]^2$ should resemble $\text{Var}_\theta[\bar{Z}_\Delta^N]$ if $1 \ll \Delta \ll t$. Thus $\tilde{\mathcal{W}}_{\Delta,t}^N := N t^{-1} \sum_{k=1}^{t/\Delta} [\bar{Z}_{k\Delta}^N - \bar{Z}_{(k-1)\Delta}^N - \Delta t^{-1} \bar{Z}_t^N]^2 \simeq N \Delta^{-1} \text{Var}_\theta[\bar{Z}_\Delta^N]$. Introducing the martingales $M_t^{i,N} = Z_t^{i,N} - C_t^{i,N}$ (where $C_t^{i,N}$ is the compensator of $Z_t^{i,N}$), the centered processes $U_t^{i,N} = Z_t^{i,N} - \mathbb{E}_\theta[Z_t^{i,N}]$, and the N -dimensional vectors \mathbf{U}_t^N and \mathbf{M}_t^N with coordinates $U_t^{i,N}$ and $M_t^{i,N}$, we will see in Section 3 that $\mathbf{U}_t^N = \mathbf{M}_t^N + A_N \int_0^t \varphi(t-s) \mathbf{U}_s^N ds$, so that for large times, $\mathbf{U}_t^N \simeq \mathbf{M}_t^N + \Lambda A_N \mathbf{U}_t^N$ and thus $\mathbf{U}_t^N \simeq Q_N \mathbf{M}_t^N$. Consequently, we hope that $\bar{U}_t^N \simeq \bar{Q}_N \bar{\mathbf{M}}_t^N$, where $\bar{U}_t^N = N^{-1} \sum_{i=1}^N U_t^{i,N}$ and $\bar{Q}_N \bar{\mathbf{M}}_t^N = N^{-1} \sum_{i=1}^N (Q_N \mathbf{M}_t^N)_i$. A little study shows that the martingales $M_t^{j,N}$ are orthogonal and that $[M_t^{j,N}, M_t^{j,N}] = Z_t^{j,N} \simeq \mu \ell_N(j)t$, so that $\text{Var}_\theta(\bar{Q}_N \bar{\mathbf{M}}_t^N) \simeq \mu t N^{-2} \sum_{j=1}^N (\sum_{i=1}^N Q_N(i,j))^2 \ell_N(j)$, which is nothing but $\mu t N^{-2} \sum_{j=1}^N (c_N(j))^2 \ell_N(j)$. Finally, $\text{Var}_\theta[\bar{Z}_t^N] = \text{Var}_\theta[\bar{U}_t^N]$ thus resembles $\mu t N^{-2} \sum_{j=1}^N (c_N(j))^2 \ell_N(j)$ and we hope that $\tilde{\mathcal{W}}_{\Delta,t}^N \simeq N \Delta^{-1} \text{Var}_\theta[\bar{Z}_\Delta^N] \simeq \mu N^{-1} \sum_{j=1}^N (c_N(j))^2 \ell_N(j)$ if $1 \ll \Delta \ll t$.

We need to find the limits as $N \rightarrow \infty$ of $\bar{\ell}_N$, $\sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2$ and $N^{-1} \sum_{i=1}^N \ell_N(i) (c_N(i))^2$. It is not easy to make rigorous, but it holds true that $\ell_N(i) \simeq 1 + \Lambda(1 - \Lambda p)^{-1} L_N(i)$, where $L_N(i) = \sum_{j=1}^N A_N(i,j)$. This comes from $\sum_{j=1}^N A_N^2(i,j) = \sum_{j=1}^N A_N(i,j) \sum_{k=1}^N A_N(j,k) \simeq p \sum_{j=1}^N A_N(i,j) = p L_N(i)$, $\sum_{j=1}^N A_N^3(i,j) \simeq p^2 L_N(i)$ for similar reasons, etc. It is very rough, but it will imply that $\ell_N(i) = \sum_{n \geq 0} \Lambda^n \sum_{j=1}^N A_N^n(i,j) \simeq 1 + \sum_{n \geq 1} \Lambda^n p^{n-1} L_N(i) = 1 + \Lambda(1 - \Lambda p)^{-1} L_N(i)$. Once this is seen (as well as a similar fact for the columns), we get convinced, $N L_N$ being a vector of N i.i.d. Binomial(N, p)-distributed random variables, that $\bar{\ell}_N \simeq 1/(1 - \Lambda p)$, that $\sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2 \simeq \Lambda^2 p(1 - p)/(1 - \Lambda p)^2$ and that $N^{-1} \sum_{i=1}^N \ell_N(i) (c_N(i))^2 \simeq 1/(1 - \Lambda p)^3$.

At the end, it should be more or less true that, for t, Δ and N large enough and in a suitable regime, $\tilde{\mathcal{E}}_t^N \simeq \mu/(1 - \Lambda p)$, $\tilde{\mathcal{V}}_t^N \simeq \mu^2 \Lambda^2 p(1 - p)/(1 - \Lambda p)^2$, and $\tilde{\mathcal{W}}_{\Delta,t}^N \simeq \mu/(1 - \Lambda p)^3$. Of course, all this is completely informal and many points have to be clarified.

Observe that concerning $\tilde{\mathcal{V}}_t^N$, we use that $Z_t^{1,N}$ resembles a Poisson process, while concerning $\tilde{\mathcal{W}}_{\Delta,t}^N$, we use that \bar{Z}_t^N does not resemble a Poisson process.

The three estimators $\mathcal{E}_t^N, \mathcal{V}_t^N, \mathcal{W}_{\Delta,t}^N$ we study in the paper resemble much $\tilde{\mathcal{E}}_t^N, \tilde{\mathcal{V}}_t^N, \tilde{\mathcal{W}}_{\Delta,t}^N$ and should converge to the same limits. Let us explain why we have modified the expressions. We started this subsection by the observation that $\mathbb{E}_\theta[Z_t^{i,N}] \simeq \mu \ell_N(i)t$, on which the construction of the estimators relies. A detailed study shows that, under $H(q)$, $\mathbb{E}_\theta[Z_t^{i,N}] = \mu \ell_N(i)t + \chi_i^N \pm t^{1-q}$, for some finite random variable χ_i^N . As a consequence, $t^{-1} \mathbb{E}_\theta[Z_{2t}^{i,N} - Z_t^{i,N}]$ con-

verges to $\mu\ell_N(i)$ considerably faster (with an error in t^{-q}) than $t^{-1}\mathbb{E}_\theta[Z_t^{i,N}]$ (for which the error is of order t^{-1}). This explains our modifications and why these modifications are crucial.

Let us conclude this subsection with a technical issue. If $\Lambda > 1$ (which is not forbidden even in the subcritical case), there is a positive probability that an anomalously high proportion of the θ_{ij} 's equal 1, so that $I - \Lambda A_N$ is not invertible and our multivariate Hawkes process is supercritical (on this event with small probability). We will thus work on an event Ω_N^1 on which such problems do not occur and show that this event has a high probability.

2.2. The supercritical case

We now assume that $\Lambda p > 1$ and explain the asymptotics of $(Z_t^{i,N})_{i=1,\dots,N,t \geq 0}$ and where the estimator \mathcal{U}_t^N comes from. We introduce $A_N(i, j) = N^{-1}\theta_{ij}$.

Fixing N and knowing $(\theta_{ij})_{i,j=1,\dots,N}$, we expect that $Z_t^{i,N} \simeq H_N \mathbb{E}_\theta[Z_t^{i,N}]$, for some random $H_N > 0$ not depending on i (and with H_N almost constant for N large). This is typically a supercritical phenomenon, that can already be observed on Galton-Watson processes. Fortunately, we will not really need to check it nor to study H_N , essentially because we will use the ratios $Z_t^{i,N}/\bar{Z}_t^N$, which makes disappear H_N .

Next, we believe that $\mathbb{E}_\theta[Z_t^{i,N}] \simeq \gamma_N(i)e^{\alpha_N t}$ for t large, for some vector γ_N with positive entries and some exponent $\alpha_N > 0$. Inserting this into $\mathbb{E}_\theta[Z_t^{i,N}] = \mu t + N^{-1} \sum_{j=1}^N \theta_{ij} \int_0^t \varphi(t-s) \mathbb{E}_\theta[Z_s^{j,N}] ds$, we find $\gamma_N = A_N \gamma_N \int_0^\infty e^{-\alpha_N s} \varphi(s) ds$. The vector γ_N being positive, it is necessarily a Perron-Frobenius eigenvector of A_N , so that $\rho_N = (\int_0^\infty e^{-\alpha_N s} \varphi(s) ds)^{-1}$ is its Perron-Frobenius eigenvalue (i.e. its spectral radius). We now consider the normalized Perron-Frobenius eigenvector V_N such that $\sum_{i=1}^N (V_N(i))^2 = N$ and conclude that $Z_t^{i,N} \simeq K_N V_N(i) e^{\alpha_N t}$ for all $i = 1, \dots, N$, where $K_N = [N^{-1} \sum_{i=1}^N (\gamma_N(i))^2]^{1/2} H_N$.

As in the subcritical case, the empirical variance $N^{-1} \sum_{i=1}^N (Z_t^{i,N} - \bar{Z}_t^N)^2$ resembles $N^{-1} \sum_{i=1}^N (\mathbb{E}_\theta[Z_t^{i,N}] - \mathbb{E}_\theta[\bar{Z}_t^N])^2 + \bar{Z}_t^N \simeq N^{-1} K_N^2 e^{2\alpha_N t} \sum_{i=1}^N (V_N(i) - \bar{V}_N)^2 + \bar{Z}_t^N$. We also guess that $\bar{Z}_t^N \simeq K_N \bar{V}_N e^{\alpha_N t}$, where $\bar{V}_N = N^{-1} \sum_{i=1}^N V_N(i)$. Hence we expect, that for t large, $\mathcal{U}_t^N = (\bar{Z}_t^N)^{-2} [\sum_{i=1}^N (Z_t^{i,N} - \bar{Z}_t^N)^2 - N \bar{Z}_t^N] \simeq (\bar{V}_N)^{-2} \sum_{i=1}^N (V_N(i) - \bar{V}_N)^2$.

We now search for the limit of $(\bar{V}_N)^{-2} \sum_{i=1}^N (V_N(i) - \bar{V}_N)^2$ as $N \rightarrow \infty$. Roughly, $A_N^2(i, j) \simeq p^2/N$, whence, starting from $A_N^2 V_N = \rho_N^2 V_N$, we see that $\rho_N^2 V_N \simeq p^2 \bar{V}_N \mathbf{1}_N$, where $\mathbf{1}_N$ is the N -dimensional vector with all coordinates equal to 1. Consequently, $V_N = (A_N V_N)/\rho_N \simeq \kappa_N A_N \mathbf{1}_N$, where $\kappa_N = (p^2/\rho_N^3) \bar{V}_N$. In other words, V_N is almost colinear to $L_N := A_N \mathbf{1}_N$, and $N L_N$ is a vector of N i.i.d. Binomial(N, p)-distributed random variables. It is thus reasonable to expect that $(\bar{V}_N)^{-2} \sum_{i=1}^N (V_N(i) - \bar{V}_N)^2 \simeq (\bar{L}_N)^{-2} \sum_{i=1}^N (L_N(i) - \bar{L}_N)^2 \simeq p^{-2} p(1-p) = 1/p - 1$.

All in all, we hope that for N and t large and in a suitable regime, $\mathcal{U}_t^N \simeq 1/p - 1$.

Let us mention that $\alpha_N \simeq \alpha_0$ (recall Remark 5) because $\int_0^\infty e^{-\alpha_N s} \varphi(s) ds = 1/\rho_N$, because $\int_0^\infty e^{-\alpha_0 s} \varphi(s) ds = 1/p$ and because $\rho_N \simeq p$. This last assertion follows from the fact that $A_N^2(i, j) \simeq p^2/N$, so that the largest eigenvalue of A_N^2 should resemble p^2 , whence that of A_N should resemble p .

Of course, all this is not clear and has to be made rigorous. Let us mention that we will use a quantified version of the Perron-Frobenius of G. Birkhoff [7]. As we will see, the projection onto the eigenvector V_N will be very fast (almost immediate for N very large).

As in the subcritical case, we will have to work on an event Ω_N^2 , of high probability, on which the θ_{ij} 's behave reasonably. For example, to apply the Perron-Frobenius theorem, we have to be sure that the matrix A_N is irreducible, which is not a.s. true.

2.3. About optimality: A related toy model

Consider $\alpha_0 \geq 0$ and two unknown parameters $\Gamma > 0$ and $p \in (0, 1]$. For $N \geq 1$, consider an i.i.d. family $(\theta_{ij})_{i,j=1,\dots,N}$ of Bernoulli(p)-distributed random variables, put $\lambda_t^{i,N} = N^{-1}\Gamma e^{\alpha_0 t} \sum_{j=1}^N \theta_{ij}$ and, conditionally on $(\theta_{ij})_{i,j=1,\dots,N}$, a family $(Z_t^{1,N})_{t \geq 0}, \dots, (Z_t^{N,N})_{t \geq 0}$ of independent inhomogeneous Poisson processes with intensities $(\lambda_t^{1,N})_{t \geq 0}, \dots, (\lambda_t^{N,N})_{t \geq 0}$. We observe $(Z_s^{i,N})_{s \in [0,t], i=1,\dots,N}$ and we want to estimate p in the asymptotic $(N, t) \rightarrow (\infty, \infty)$.

This problem can be seen as a strongly simplified version of the one studied in the present paper, with $\alpha_0 = 0$ in the subcritical case and $\alpha_0 > 0$ in the supercritical case. Roughly, the mean number of jumps per individual resembles $m_t = \int_0^t e^{\alpha_0 s} ds$, which is of order t when $\alpha_0 = 0$ and $e^{\alpha_0 t}$ else.

There is classically no loss of information, since α_0 is known, if we only observe $(Z_t^{i,N})_{i=1,\dots,N}$: after a (deterministic and known) change of time, the processes $(Z_t^{i,N})_{i=1,\dots,N}$ become *homogeneous* Poisson processes with unknown parameters (conditionally on $(\theta_{ij})_{i,j=1,\dots,N}$), and the conditional law of a Poisson process on $[0, t]$ knowing its value at time t does not depend on its parameter.

We next proceed to a Gaussian approximation: we have $\lambda_t^{i,N} \simeq \Gamma e^{\alpha_0 t} [p + \sqrt{N^{-1}p(1-p)}]G_i$ and $Z_t^{i,N} \simeq \int_0^t \lambda_s^{i,N} ds + \sqrt{\int_0^t \lambda_s^{i,N} ds} H_i$, for two independent i.i.d. families $(G_i)_{i=1,\dots,N}, (H_i)_{i=1,\dots,N}$ of $\mathcal{N}(0, 1)$ -distributed random variables. Using finally that $(m_t)^{-1} N^{-1/2} \ll (m_t)^{-1}$ in our asymptotic, we conclude that $(m_t)^{-1} Z_t^{i,N} \simeq \Gamma p + \Gamma \sqrt{N^{-1}p(1-p)} G_i + \sqrt{(m_t)^{-1} \Gamma p} H_i$, of which the law is $\mathcal{N}(\Gamma p, N^{-1} \Gamma^2 p(1-p) + (m_t)^{-1} \Gamma p)$.

Our toy problem is thus the following: estimate p when observing a N -sample $(X_t^{i,N})_{i=1,\dots,N}$ of the $\mathcal{N}(\Gamma p, N^{-1} \Gamma^2 p(1-p) + (m_t)^{-1} \Gamma p)$ -distribution. We assume that Γp is known, which can only make easier the estimation of p . As is well-known the statistic $S_t^N = N^{-1} \sum_{i=1}^N (X_t^{i,N} - \Gamma p)^2$ is then *sufficient* and is the best estimator (in all the usual senses), for $N \geq 1$ and $t \geq 1$ fixed, of $N^{-1} \Gamma^2 p(1-p) + (m_t)^{-1} \Gamma p$, so that $T_t^N = N(\Gamma p)^{-2} (S_t^N - m_t^{-1} \Gamma p)$ is more or less the best estimator of $(1/p - 1)$. But $\text{Var } S_t^N = 2N^{-1} (N^{-1} \Gamma^2 p(1-p) + (m_t)^{-1} \Gamma p)^2$, whence $\text{Var } T_t^N = 2(\Gamma p)^{-4} (N^{-1/2} \Gamma^2 p(1-p) + N^{1/2} (m_t)^{-1} \Gamma p)^2$. It is thus not

possible to estimate $(1/p - 1)$ with a better precision than $N^{-1/2} + N^{1/2}(m_t)^{-1}$. This of course implies that we cannot estimate p with a better precision than $N^{-1/2} + N^{1/2}(m_t)^{-1}$.

3. Well-posedness and explicit formulae

We first give the

Proof of Proposition 1. Conditionally on $(\theta_{ij})_{i,j=1,\dots,N}$, we can apply directly [13, Theorem 6], of which the assumption is satisfied here, see [13, Remark 5-(i)]: conditionally on $(\theta_{ij})_{i,j=1,\dots,N}$, there is a unique solution $(Z_t^{i,N})_{t \geq 0, i=1,\dots,N}$ to (1.1) such that $\sum_{i=1}^N \mathbb{E}_\theta[Z_t^{i,N}] < \infty$ for all $t \geq 0$. Since now $(\theta_{ij})_{i,j=1,\dots,N}$ can only take a finite number of values, we immediately deduce that indeed $\sum_{i=1}^N \mathbb{E}[Z_t^{i,N}] < \infty$ for all $t \geq 0$. \square

We carry on with a classical lemma. Recall that $\varphi^{\star 0}(t - s)ds = \delta_t(ds)$ by convention.

Lemma 8. *Consider $d \geq 1$, $A \in \mathcal{M}_{d \times d}(\mathbb{R})$, $m, g : [0, \infty) \mapsto \mathbb{R}^d$ locally bounded and assume that $\varphi : [0, \infty) \mapsto [0, \infty)$ is locally integrable. If $m_t = g_t + \int_0^t \varphi(t - s)Am_s ds$ for all $t \geq 0$, then $m_t = \sum_{n \geq 0} \int_0^t \varphi^{\star n}(t - s)A^n g_s ds$.*

Proof. The equation $m_t = g_t + \int_0^t \varphi(t - s)Am_s ds$ with unknown m has at most one locally bounded solution. Indeed, consider two such solutions m, \tilde{m} , observe that $u = |m - \tilde{m}|$ satisfies $u_t \leq |A| \int_0^t \varphi(t - s)u_s ds$, and conclude that $u = 0$ by the generalized Gronwall lemma, see e.g. [13, Lemma 23-(i)]. We thus just have to prove that $m_t := \sum_{n \geq 0} \int_0^t \varphi^{\star n}(t - s)A^n g_s ds$ is locally bounded and solves $m = g + A\varphi \star m$. We introduce $k_t^n = |A|^n \int_0^t \varphi^{\star n}(s) ds$, which is locally bounded because φ is locally integrable and which satisfies $k_t^{n+1} \leq |A| \int_0^t k_s^n \varphi(t - s) ds$. We use [13, Lemma 23-(ii)] to conclude that $\sum_{n \geq 0} k_t^n$ is locally bounded. Consequently, $|m_t| \leq \sup_{[0,t]} |g_s| \times \sum_{n \geq 0} k_t^n$ is locally bounded. Finally, we write $m = g + \sum_{n \geq 1} A^n \varphi^{\star n} \star g = g + A\varphi \star \sum_{n \geq 0} A^n \varphi^{\star n} \star g = g + A\varphi \star m$ as desired. \square

We next introduce a few processes.

Notation 9. *Assume only that φ is locally integrable, fix $N \geq 1$ and consider the solution $(Z_t^{i,N})_{t \geq 0, i=1,\dots,N}$ to (1.1). For each $i = 1, \dots, N$, we introduce the martingale (recall that $\lambda^{i,N}$ was defined in (1.1))*

$$M_t^{i,N} = \int_0^t \int_0^\infty \mathbf{1}_{\{z \leq \lambda_s^{i,N}\}} \tilde{\pi}^i(ds, dz),$$

where $\tilde{\pi}^i(ds, dz) = \pi^i(ds, dz) - dsdz$ is the compensated Poisson measure associated to π^i . We also introduce $M_t^{i,N,*} = \sup_{[0,t]} |M_s^{i,N}|$, as well as the (conditionally) centered process

$$U_t^{i,N} = Z_t^{i,N} - \mathbb{E}_\theta[Z_t^{i,N}].$$

For each $t \geq 0$, we denote by \mathbf{Z}_t^N (resp. $\mathbf{M}_t^N, \mathbf{M}_t^{N,*}, \mathbf{U}_t^N$) the N -dimensional vector with coordinates $Z_t^{i,N}$ (resp. $M_t^{i,N}, M_t^{i,N,*}, U_t^{i,N}$). We also set $\bar{Z}_t^N = N^{-1} \sum_{i=1}^N Z_t^{i,N}$, $\bar{M}_t^N = N^{-1} \sum_{i=1}^N M_t^{i,N}$ and $\bar{U}_t^N = N^{-1} \sum_{i=1}^N U_t^{i,N}$.

We refer to Jacod-Shiryaev [22, Chapter 1, Section 4e] for definitions and properties of pure jump martingales and of their quadratic variations.

Remark 10. Since the Poisson measures π^i are independent, the martingales $M^{i,N}$ are orthogonal. More precisely, we have $[M^{i,N}, M^{j,N}]_t = 0$ if $i \neq j$, while $[M^{i,N}, M^{i,N}]_t = Z_t^{i,N}$ (because $Z_t^{i,N}$ counts the jumps of $M^{i,N}$, which are all of size 1). Consequently, $\mathbb{E}_\theta[M_s^{i,N} M_t^{j,N}] = \mathbf{1}_{\{i=j\}} \mathbb{E}_\theta[Z_s^{i,N}]$.

We now give some more or less explicit formulas. We denote by $\mathbf{1}_N$ the N -dimensional vector with all entries equal to 1 and we set $A_N(i, j) = N^{-1} \theta_{ij}$ for $i, j = 1, \dots, N$.

Lemma 11. Assume only that φ is locally integrable. We have (recall that $\varphi^{*0}(t - s)ds = \delta_t(ds)$):

$$\mathbf{Z}_t^N = \mathbf{M}_t^N + \mu \mathbf{1}_N t + \int_0^t \varphi(t - s) A_N \mathbf{Z}_s^N ds, \tag{2}$$

$$\mathbb{E}_\theta[\mathbf{Z}_t^N] = \mu \sum_{n \geq 0} \left[\int_0^t s \varphi^{*n}(t - s) ds \right] A_N^n \mathbf{1}_N, \tag{3}$$

$$\mathbf{U}_t^N = \sum_{n \geq 0} \int_0^t \varphi^{*n}(t - s) A_N^n \mathbf{M}_s^N ds. \tag{4}$$

Proof. The first expression is not difficult: starting from (1.1),

$$Z_t^{i,N} = M_t^{i,N} + \int_0^t \lambda_s^{i,N} ds = M_t^{i,N} + \mu t + \sum_{j=1}^N A_N(i, j) \int_0^t \int_0^s \varphi(s - u) dZ_u^{j,N} ds.$$

Using [13, Lemma 22], we see that $\int_0^t \int_0^s \varphi(s - u) dZ_u^{j,N} ds = \int_0^t \varphi(t - s) Z_s^{j,N} ds$, whence indeed,

$$Z_t^{i,N} = M_t^{i,N} + \mu t + \int_0^t \varphi(t - s) \sum_{j=1}^N A_N(i, j) Z_s^{j,N} ds,$$

which is nothing but (2). Taking conditional expectations in (2), we find that $\mathbb{E}_\theta[\mathbf{Z}_t^N] = \mu \mathbf{1}_N t + \int_0^t \varphi(t - s) A_N \mathbb{E}_\theta[\mathbf{Z}_s^N] ds$ and thus also $\mathbf{U}_t^N = \mathbf{M}_t^N + \int_0^t \varphi(t - s) A_N \mathbf{U}_s^N ds$. Since now φ is (a.s.) locally integrable, since $\mu \mathbf{1}_N t$ and \mathbf{M}_t^N are (a.s.) locally bounded, as well as $\mathbb{E}_\theta[\mathbf{Z}_t^N]$ and \mathbf{U}_t^N , (3) and (4) directly follow from Lemma 8. \square

4. The subcritical case

Here we consider the subcritical case. We first study the large N -asymptotic of the matrix $Q_N = (I - \Lambda A_N)^{-1}$, which plays a central role in the rest of the

section. In Subsection 4.2, we finely study the behavior of φ^{*n} . In Subsection 4.3, we handle a few computations to be used several times later. Subsections 4.4, 4.5 and 4.6 are devoted to the studies of the three estimators \mathcal{E}_t^N , \mathcal{V}_t^N and $\mathcal{W}_{\Delta,t}^N$. We conclude the proofs of Theorem 3 and Corollary 4 in Subsection 4.7.

4.1. Study of a random matrix

We use the following standard notation: for $x = (x_1, \dots, x_N) \in \mathbb{R}^N$ and $r \in [1, \infty)$, we set $\|x\|_r = (\sum_{i=1}^N |x_i|^r)^{1/r}$ and $\|x\|_\infty = \max_{i=1, \dots, N} |x_i|$. For $r \in [1, \infty]$, we denote by $\|\cdot\|_r$ the operator norm on $\mathcal{M}_{N \times N}(\mathbb{R})$ associated to $\|\cdot\|_r$. We recall that

$$\|M\|_1 = \sup_{j=1, \dots, N} \sum_{i=1}^N |M_{ij}|, \quad \|M\|_\infty = \sup_{i=1, \dots, N} \sum_{j=1}^N |M_{ij}|$$

and that for all $r \in (1, \infty)$,

$$\|M\|_r \leq \|M\|_1^{1/r} \|M\|_\infty^{1-1/r}. \quad (5)$$

Notation 12. We assume that $\Lambda p < 1$. For each $N \geq 1$, we introduce the $N \times N$ random matrix A_N defined by $A_N(i, j) = N^{-1} \theta_{ij}$, as well as the event

$$\Omega_N^1 = \left\{ \Lambda \|A_N\|_r \leq a \text{ for all } r \in [1, \infty] \right\}, \text{ where } a = \frac{1 + \Lambda p}{2} \in (\Lambda p, 1). \quad (6)$$

On Ω_N^1 , the $N \times N$ matrix $Q_N = \sum_{n \geq 0} \Lambda^n A_N^n = (I - \Lambda A_N)^{-1}$ is well-defined and we introduce, for each $i = 1, \dots, N$, $\ell_N(i) = \sum_{j=1}^N Q_N(i, j)$, $c_N(i) = \sum_{j=1}^N Q_N(j, i)$, as well as $\bar{\ell}_N = N^{-1} \sum_{i=1}^N \ell_N(i)$ and $\bar{c}_N = N^{-1} \sum_{i=1}^N c_N(i)$. We of course have $\bar{\ell}_N = \bar{c}_N$.

Let us remark once for all that, with $C = 1/(1 - a) < \infty$,

$$\Omega_N^1 \subset \left\{ \|Q_N\|_r \leq C \text{ for all } r \in [1, \infty] \right\} \subset \left\{ \sup_{1 \leq i \leq N} \max\{\ell_N(i), c_N(i)\} \leq C \right\}, \quad (7)$$

$$\Omega_N^1 \subset \left\{ \mathbf{1}_{\{i=j\}} \leq Q_N(i, j) \leq \mathbf{1}_{\{i=j\}} + \Lambda C N^{-1} \text{ for all } i, j = 1, \dots, N \right\}. \quad (8)$$

Indeed, (7) is straightforward since $Q_N = \sum_{n \geq 0} \Lambda^n A_N^n$. To check (8), we first observe that $Q_N(i, j) \geq \Lambda^0 A_N^0(i, j) = \mathbf{1}_{\{i=j\}}$. Next, we use that $A_N(i, j) \leq N^{-1}$ while, if $n \geq 2$, $A_N^n(i, j) = \sum_{k=1}^N A_N(i, k) A_N^{n-1}(k, j) \leq N^{-1} \sum_{k=1}^N A_N^{n-1}(k, j) \leq N^{-1} \|A_N^{n-1}\|_1 \leq N^{-1} \|A_N\|_1^{n-1}$. Thus $A_N^n(i, j) \leq N^{-1} \|A_N\|_1^{n-1}$ for all $n \geq 1$. Hence on Ω_N^1 , it holds that $Q_N(i, j) \leq \mathbf{1}_{\{i=j\}} + N^{-1} \sum_{n \geq 1} \Lambda^n \|A_N\|_1^{n-1} \leq \mathbf{1}_{\{i=j\}} + N^{-1} \Lambda / (1 - a)$ as desired.

Lemma 13. Assume that $\Lambda p < 1$. It holds that $\Pr(\Omega_N^1) \geq 1 - C \exp(-cN)$.

Proof. By (5), it suffices to prove that $\Pr(\Lambda \|A_N\|_1 > a) \leq C \exp(-cN)$ and $\Pr(\Lambda \|A_N\|_\infty > a) \leq C \exp(-cN)$. Since $\|A_N\|_\infty = \|A_N^t\|_1$ and since A_N^t (the transpose of A_N) has the same law as A_N , it actually suffices to verify the first inequality. First, $N \|A_N\|_1 = \max\{X_1^N, \dots, X_N^N\}$, where $X_i^N = \sum_{j=1}^N \theta_{ij}$ is Binomial(N, p)-distributed for each i . Consequently, $\Pr(\Lambda \|A_N\|_1 > a) \leq N \Pr(X_1^N \geq Na/\Lambda) \leq N \Pr(|X_1^N - Np| \geq N(a/\Lambda - p))$. Since $a/\Lambda > p$, we can use the Hoeffding inequality [21] to obtain $\Pr(\Lambda \|A_N\|_1 > a) \leq 2N \exp(-2N(a/\Lambda - p)^2) \leq C \exp(-N(a/\Lambda - p)^2)$ as desired. \square

The next result is much harder but crucial.

Proposition 14. *Assume that $\Lambda p < 1$. It holds that*

$$\begin{aligned} \mathbb{E} \left[\mathbf{1}_{\Omega_N^1} \left| \bar{\ell}_N - \frac{1}{1 - \Lambda p} \right|^2 \right] &\leq \frac{C}{N^2}, \\ \mathbb{E} \left[\mathbf{1}_{\Omega_N^1} \left| \frac{1}{N} \sum_{i=1}^N \ell_N(i) (c_N(i))^2 - \frac{1}{(1 - \Lambda p)^3} \right|^2 \right] &\leq \frac{C}{N^2}, \\ \mathbb{E} \left[\mathbf{1}_{\Omega_N^1} \left| \sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2 - \frac{\Lambda^2 p(1 - p)}{(1 - \Lambda p)^2} \right| \right] &\leq \frac{C}{\sqrt{N}}. \end{aligned}$$

Proof. Recall that $\mathbf{1}_N$ is the N -dimensional vector of which all the coordinates equal 1. Let ℓ_N (resp. c_N) be the vector with coordinates $\ell_N(1), \dots, \ell_N(N)$ (resp. $c_N(1), \dots, c_N(N)$). We also introduce, for all $i = 1, \dots, N$, $L_N(i) = \sum_{j=1}^N A_N(i, j)$ and $C_N(i) = \sum_{j=1}^N A_N(j, i)$, as well as the corresponding vectors L_N and C_N . Let us observe that, with obvious notation, $\bar{\ell}_N = \bar{c}_N$ and $\bar{L}_N = \bar{C}_N$. Finally, we introduce the vectors

$$x_N = \ell_N - \bar{\ell}_N \mathbf{1}_N, \quad y_N = c_N - \bar{c}_N \mathbf{1}_N, \quad X_N = L_N - \bar{L}_N \mathbf{1}_N, \quad Y_N = C_N - \bar{C}_N \mathbf{1}_N.$$

We recall that $a = (1 + \Lambda p)/2 \in (0, 1)$ and we introduce $b = (2 + \Lambda p)/3 \in (a, 1)$.

Step 1. We introduce the event

$$\mathcal{A}_N = \left\{ \|L_N - p \mathbf{1}_N\|_2 + \|C_N - p \mathbf{1}_N\|_2 \leq N^{1/4} \right\} \subset \left\{ \|X_N\|_2 + \|Y_N\|_2 \leq N^{1/4} \right\}.$$

The inclusion comes from the fact that a.s., $\|X_N\|_2 = \|L_N - \bar{L}_N \mathbf{1}_N\|_2 \leq \|L_N - x \mathbf{1}_N\|_2$ for any $x \in \mathbb{R}$. Since $N L_N = (Z_1^N, \dots, Z_N^N)$ with Z_i^N i.i.d. and Binomial(N, p)-distributed, it is very classical that for any $\alpha > 0$, $\mathbb{E}[\|L_N - p \mathbf{1}_N\|_2^\alpha] \leq C_\alpha$ (uniformly in N), we have similarly $\mathbb{E}[\|C_N - p \mathbf{1}_N\|_2^\alpha] \leq C_\alpha$, so that

$$\Pr(\mathcal{A}_N) \geq 1 - C_\alpha N^{-\alpha/4}.$$

Step 2. We now check the following points: (i) $\mathbb{E}[\|\bar{L}_N - p\|^2] \leq CN^{-2}$, (ii) $\mathbb{E}[\|X_N\|_2^4] \leq C$, (iii) $\mathbb{E}[(\|X_N\|_2^2 - p(1 - p))^2] \leq CN^{-1}$ and (iv) $\mathbb{E}[\|A_N X_N\|_2^2] \leq CN^{-1}$.

Point (i) is clear, because $\bar{L}_N = N^{-2} \sum_{i,j=1}^N \theta_{ij}$ is nothing but the empirical mean of N^2 independent Bernoulli(p)-random variables. Points (ii) and (iii)

are very classical, since $N\|X_N\|_2^2$ is the empirical variance of N independent Binomial(N, p)-random variables. We now prove (iv):

$$\begin{aligned}\mathbb{E}[\|A_N X_N\|_2^2] &= \sum_{i=1}^N \mathbb{E} \left[\left(\sum_{j=1}^N \frac{\theta_{ij}}{N} (L_N(j) - \bar{L}_N) \right)^2 \right] \\ &= \frac{1}{N} \mathbb{E} \left[\left(\sum_{j=1}^N \theta_{1j} (L_N(j) - \bar{L}_N) \right)^2 \right]\end{aligned}$$

by symmetry. We now write $\mathbb{E}[\|A_N X_N\|_2^2] \leq 4N^{-1}(I_N + J_N + K_N)$, where

$$\begin{aligned}I_N &= \mathbb{E} \left[(\bar{L}_N - p)^2 \left(\sum_{j=1}^N \theta_{1j} \right)^2 \right], \quad J_N = \mathbb{E} \left[\left(\theta_{11} (L_N(1) - p) \right)^2 \right], \\ K_N &= \mathbb{E} \left[\left(\sum_{j=2}^N \theta_{1j} (L_N(j) - p) \right)^2 \right].\end{aligned}$$

First, $I_N \leq N^2 \mathbb{E}[(\bar{L}_N - p)^2] \leq C$ by (i). Next, it is obvious that $J_N \leq 1$ (because $\theta_{11} \in \{0, 1\}$ and $L_N(1) \in [0, 1]$). Finally, the random variables $\theta_{1j}(L_N(j) - p)$ being i.i.d. and centered (for $j = 2, \dots, N$), we may write

$$K_N = (N-1) \mathbb{E} \left[\left(\theta_{12} (L_N(2) - p) \right)^2 \right] \leq (N-1) \mathbb{E} \left[(L_N(2) - p)^2 \right] \leq C,$$

since $NL_N(2)$ follows a Binomial(N, p)-distribution. This completes the step.

Step 3. We next prove that (i) $x_N = \Lambda A_N x_N - \Lambda r_N \mathbf{1}_N + \Lambda \bar{\ell}_N X_N$ on Ω_N^1 , where $r_N = N^{-2} \sum_{i,j=1}^N (\theta_{ij} - p)x_N(j)$ and that (ii) $|r_N| \leq N^{-3/4} \|x_N\|_2$ on $\Omega_N^1 \cap \mathcal{A}_N$.

We start from $\ell_N = Q_N \mathbf{1}_N = (I - \Lambda A_N)^{-1} \mathbf{1}_N$, whence $\ell_N = \mathbf{1}_N + \Lambda A_N \ell_N$. Since $\bar{\ell}_N = N^{-1}(\ell_N, \mathbf{1}_N)$, we see that $\bar{\ell}_N = 1 + \Lambda N^{-1}(A_N \ell_N, \mathbf{1}_N)$ (here (\cdot, \cdot) is the usual scalar product on \mathbb{R}^N) and thus

$$\begin{aligned}x_N &= \Lambda A_N \ell_N - \Lambda N^{-1}(A_N \ell_N, \mathbf{1}_N) \mathbf{1}_N \\ &= \Lambda A_N x_N - \Lambda N^{-1}(A_N x_N, \mathbf{1}_N) \mathbf{1}_N + \bar{\ell}_N \Lambda A_N \mathbf{1}_N - \bar{\ell}_N \Lambda N^{-1}(A_N \mathbf{1}_N, \mathbf{1}_N) \mathbf{1}_N.\end{aligned}$$

It only remains to verify that $N^{-1}(A_N x_N, \mathbf{1}_N) = r_N$, which follows from the facts that $N^{-1}(A_N x_N, \mathbf{1}_N) = N^{-2} \sum_{i,j=1}^N \theta_{ij} x_N(j)$, that $\sum_{j=1}^N x_N(j) = 0$; and that $A_N \mathbf{1}_N - N^{-1}(A_N \mathbf{1}_N, \mathbf{1}_N) \mathbf{1}_N = X_N$, which is clear since $A_N \mathbf{1}_N = L_N$.

To verify (ii), we observe that $r_N = N^{-1} \sum_{j=1}^N (C_N(j) - p)x_N(j)$, whence, by the Cauchy-Schwarz inequality, $|r_N| \leq N^{-1} \|x_N\|_2 \|C_N - p \mathbf{1}_N\|_2 \leq N^{-3/4} \|x_N\|_2$ on $\Omega_N^1 \cap \mathcal{A}_N$.

Step 4. Let N_0 be the smallest integer such that $a + \Lambda N_0^{-1/4} \leq b$. We check that for all $N \geq N_0$,

$$\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \|x_N\|_2 \leq C \|X_N\|_2.$$

Using Step 3 and that $\|\mathbf{1}_N\|_2 = N^{1/2}$, we write $\|x_N\|_2 \leq \Lambda \|A_N\|_2 \|x_N\|_2 + \Lambda N^{-1/4} \|x_N\|_2 + \Lambda |\bar{\ell}_N| \|X_N\|_2$. But on Ω_N^1 , $\Lambda \|A_N\|_2 \leq a$ and $|\bar{\ell}_N| \leq C$,

see (6) and (7). Hence, for $N \geq N_0$, on $\Omega_N^1 \cap \mathcal{A}_N$, we have $\|x_N\|_2 \leq (a + \Lambda N^{-1/4})\|x_N\|_2 + C\|X_N\|_2 \leq b\|x_N\|_2 + C\|X_N\|_2$. Since $b < 1$, the conclusion follows.

Step 5. We now prove that for $N \geq N_0$,

$$\mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| \bar{\ell}_N - \frac{1}{1 - \Lambda p} \right|^2\right] \leq \frac{C}{N^2}.$$

Using Step 3, we know that on $\Omega_N^1 \cap \mathcal{A}_N$, $\ell_N = \mathbf{1}_N + \Lambda A_N \ell_N$, whence

$$\bar{\ell}_N = 1 + \frac{\Lambda}{N} \sum_{i,j=1}^N A_N(i,j) \ell_N(j) = 1 + \frac{\Lambda}{N} \sum_{j=1}^N C_N(j) \ell_N(j) = 1 + \Lambda p \bar{\ell}_N + S_N,$$

where $S_N = \Lambda N^{-1} \sum_{j=1}^N (C_N(j) - p) \ell_N(j)$. Consequently, $\bar{\ell}_N = (1 - \Lambda p)^{-1}(1 + S_N)$, and we only have to prove that $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} S_N^2] \leq CN^{-2}$. To this end, we write $S_N = \Lambda N^{-1}(a_N + b_N)$, where $a_N = \sum_{j=1}^N (C_N(j) - p)x_N(j)$ and $b_N = \bar{\ell}_N \sum_{j=1}^N (C_N(j) - p)$. First, since $|\bar{\ell}_N| \leq C$ on Ω_N^1 by (7), we can write $\mathbb{E}[\mathbf{1}_{\Omega_N^1} b_N^2] \leq C \mathbb{E}[(\sum_{j=1}^N (C_N(j) - p))^2] = CN^2 \mathbb{E}[(\bar{C}_N - p)^2] \leq C$, the last inequality coming from Step 2-(i) since $\bar{C}_N = \bar{L}_N$. Next, we use the Cauchy-Schwarz inequality: $a_N^2 \leq \|C_N - p\mathbf{1}_N\|_2 \|x_N\|_2 \leq C \|C_N - p\mathbf{1}_N\|_2 \|X_N\|_2$ on $\Omega_N^1 \cap \mathcal{A}_N$ by Step 4. Consequently, $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} a_N^2] \leq C \mathbb{E}[\|X_N\|_2^2]^{1/2} \mathbb{E}[\|C_N - p\mathbf{1}_N\|_2^2]^{1/2}$. But $\mathbb{E}[\|X_N\|_2^2] \leq C$ by Step 2-(ii) and we have seen at the end of Step 1 that $\mathbb{E}[\|C_N - p\mathbf{1}_N\|_2^2] \leq C$.

Step 6. Here we verify that, still for $N \geq N_0$,

$$\mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| \frac{1}{N} \sum_{i=1}^N \ell_N(i) (c_N(i))^2 - \frac{1}{(1 - \Lambda p)^3} \right|^2\right] \leq \frac{C}{N^2}.$$

We write, using that $\bar{c}_N = \bar{\ell}_N$,

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N \ell_N(i) (c_N(i))^2 &= \frac{1}{N} \sum_{i=1}^N \ell_N(i) (c_N(i) - \bar{c}_N)^2 + (\bar{\ell}_N)^3 \\ &\quad + \frac{2}{N} \bar{\ell}_N \sum_{i=1}^N \ell_N(i) (c_N(i) - \bar{c}_N). \end{aligned}$$

First, since $|\bar{\ell}_N| \leq C$ on Ω_N^1 , we have $|(\bar{\ell}_N)^3 - (1 - \Lambda p)^{-3}| \leq C|\bar{\ell}_N - (1 - \Lambda p)^{-1}|$, whence $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} |(\bar{\ell}_N)^3 - (1 - \Lambda p)^{-3}|^2] \leq CN^{-2}$ by Step 5. It thus suffices to verify that $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} ((a'_N)^2 + (b'_N)^2)] \leq C$, where $a'_N = \sum_{i=1}^N \ell_N(i) (c_N(i) - \bar{c}_N)^2$ and $b'_N = \sum_{i=1}^N \ell_N(i) (c_N(i) - \bar{c}_N)$.

First, $b'_N = \sum_{i=1}^N \ell_N(i) y_N(i) = \sum_{i=1}^N x_N(i) y_N(i)$ because $\sum_{i=1}^N y_N(i) = 0$. Hence $|b'_N| \leq \|x_N\|_2 \|y_N\|_2$. But on $\Omega_N^1 \cap \mathcal{A}_N$, we know from Step 4 that $\|x_N\|_2 \leq C\|X_N\|_2$, and it obviously also holds true that $\|y_N\|_2 \leq C\|Y_N\|_2$. We thus conclude that $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} (b'_N)^2] \leq C \mathbb{E}[\|X_N\|_2^4]^{1/2} \mathbb{E}[\|Y_N\|_2^4]^{1/2} = \mathbb{E}[\|X_N\|_2^4]$ by

symmetry. Using finally Step 2-(ii), we deduce that indeed, $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} (b'_N)^2] \leq C$. Next, since $|\ell_N(i)| \leq C$ on Ω_N^1 by (7), we can write $|a'_N| \leq C \|c_N - \bar{c}_N \mathbf{1}_N\|_2^2 = C \|y_N\|_2^2$. We conclude as previously that $\mathbb{E}[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} (a'_N)^2] \leq C$.

Step 7. The goal of this step is to establish that, for all $N \geq N_0$,

$$\mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| \|x_N\|_2^2 - \frac{\Lambda^2 p(1-p)}{(1-\Lambda p)^2} \right|\right] \leq \frac{C}{\sqrt{N}}.$$

Starting from Step 3, we write

$$x_N - \Lambda \bar{\ell}_N X_N = \Lambda A_N x_N - \Lambda r_N \mathbf{1}_N = \Lambda A_N (x_N - \Lambda \bar{\ell}_N X_N) + \Lambda^2 \bar{\ell}_N A_N X_N - \Lambda r_N \mathbf{1}_N.$$

Thus

$$\begin{aligned} \|x_N - \Lambda \bar{\ell}_N X_N\|_2 &\leq \Lambda \|A_N\|_2 \|x_N - \Lambda \bar{\ell}_N X_N\|_2 + \Lambda^2 |\bar{\ell}_N| \|A_N X_N\|_2 \\ &\quad + \Lambda N^{-1/2} \|C_N - p \mathbf{1}_N\|_2 \|x_N\|_2, \end{aligned}$$

where we used that $\|\mathbf{1}_N\|_2 = N^{1/2}$ and that $|r_N| \leq N^{-1} \|C_N - p \mathbf{1}_N\|_2 \|x_N\|_2$ on $\Omega_N^1 \cap \mathcal{A}_N$, as checked at the end of Step 3. Using now that $\Lambda \|A_N\|_2 \leq a < 1$ and $|\bar{\ell}_N| \leq C$ on Ω_N^1 and that $\|x_N\|_2 \leq C \|X_N\|_2$ on $\Omega_N^1 \cap \mathcal{A}_N$ by Step 4, we conclude that, still on $\Omega_N^1 \cap \mathcal{A}_N$,

$$\|x_N - \Lambda \bar{\ell}_N X_N\|_2^2 \leq C (\|A_N X_N\|_2^2 + C N^{-1} \|C_N - p \mathbf{1}_N\|_2^2 \|X_N\|_2^2).$$

Since now $\mathbb{E}[\|A_N X_N\|_2^2] \leq C N^{-1}$ by Step 2-(iv), since $\mathbb{E}[\|X_N\|_2^4] \leq C$ by Step 2-(ii) and since $\mathbb{E}[\|C_N - p \mathbf{1}_N\|_2^4] \leq C$ (see the end of Step 1), we deduce that

$$\mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \|x_N - \Lambda \bar{\ell}_N X_N\|_2^2\right] \leq \frac{C}{N}.$$

Next, we observe that $\left| \|x_N\|_2^2 - (\Lambda \bar{\ell}_N)^2 \|X_N\|_2^2 \right| \leq \|x_N - \Lambda \bar{\ell}_N X_N\|_2 (\|x_N\|_2 + \Lambda |\bar{\ell}_N| \|X_N\|_2) \leq C \|x_N - \Lambda \bar{\ell}_N X_N\|_2 \|X_N\|_2$ on $\Omega_N^1 \cap \mathcal{A}_N$ by Step 4 and since $\bar{\ell}_N$ is bounded on Ω_N^1 . Hence

$$\mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| \|x_N\|_2^2 - (\Lambda \bar{\ell}_N)^2 \|X_N\|_2^2 \right|\right] \leq \frac{C}{\sqrt{N}} \mathbb{E}[\|X_N\|_2^2]^{1/2} \leq \frac{C}{\sqrt{N}}$$

by Step 2-(ii). To complete the step, it only remains to verify that

$$d_N = \mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| (\bar{\ell}_N)^2 \|X_N\|_2^2 - p(1-p)(1-\Lambda p)^{-2} \right|\right] \leq \frac{C}{\sqrt{N}}.$$

We naturally write $d_N \leq a''_N + b''_N$, where

$$\begin{aligned} a''_N &= \mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| (\bar{\ell}_N)^2 - (1-\Lambda p)^{-2} \right| \|X_N\|_2^2 \right], \\ b''_N &= (1-\Lambda p)^{-2} \mathbb{E}\left[\mathbf{1}_{\Omega_N^1 \cap \mathcal{A}_N} \left| \|X_N\|_2^2 - p(1-p) \right|\right]. \end{aligned}$$

Step 2-(iii) directly implies that $b''_N \leq CN^{-1/2}$. Using that $\bar{\ell}_N$ is bounded on Ω^1_N , we deduce that $|(\bar{\ell}_N)^2 - (1 - \Lambda p)^{-2}| \leq C|\bar{\ell}_N - (1 - \Lambda p)^{-1}|$. Thus

$$a''_N \leq C\mathbb{E}\left[\mathbf{1}_{\Omega^1_N \cap \mathcal{A}_N} \left| \bar{\ell}_N - (1 - \Lambda p)^{-1} \right|^2\right]^{1/2} \mathbb{E}[\|X_N\|_2^4]^{1/2}.$$

Step 2-(ii) and Step 5 imply that $a''_N \leq CN^{-1} \leq CN^{-1/2}$ as desired.

Step 8. It remains to conclude. It clearly suffices to treat the case where $N \geq N_0$, because $\ell_N(i)$ and $c_N(i)$ are uniformly bounded on Ω^1_N by (7), so that the inequalities of the statement are trivial when $N \leq N_0$ (if the constant C is large enough). Since $\bar{\ell}_N$ is (uniformly) bounded on Ω^1_N , we have

$$\mathbb{E}\left[\mathbf{1}_{\Omega^1_N} \left| \bar{\ell}_N - \frac{1}{1 - \Lambda p} \right|^2\right] \leq \mathbb{E}\left[\mathbf{1}_{\Omega^1_N \cap \mathcal{A}_N} \left| \bar{\ell}_N - \frac{1}{1 - \Lambda p} \right|^2\right] + C\Pr((\mathcal{A}_N)^c).$$

The first term is bounded by CN^{-2} (by Step 5), as well as the second one (use the last inequality of Step 1 with $\alpha = 8$).

Similarly, using Step 6 and that $\ell_N(i)$ and $c_N(i)$ are (uniformly) bounded on Ω^1_N , we see that

$$\mathbb{E}\left[\mathbf{1}_{\Omega^1_N} \left| \frac{1}{N} \sum_{i=1}^N \ell_N(i)(c_N(i))^2 - \frac{1}{(1 - \Lambda p)^3} \right|^2\right] \leq \frac{C}{N^2} + C\Pr((\mathcal{A}_N)^c) \leq \frac{C}{N^2}.$$

Finally, observe that $\sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2 = \|x_N\|_2^2$ is bounded by CN on Ω^1_N , so that by Step 7,

$$\mathbb{E}\left[\mathbf{1}_{\Omega^1_N} \left| \sum_{i=1}^N (\ell_N(i) - \bar{\ell}_N)^2 - \frac{\Lambda^2 p(1 - p)}{(1 - \Lambda p)^2} \right|\right] \leq \frac{C}{\sqrt{N}} + CN\Pr((\mathcal{A}_N)^c) \leq \frac{C}{\sqrt{N}}.$$

We used the last inequality of Step 1 with $\alpha = 6$. □

4.2. Preliminary analytic estimates

In view of (3) and (4), it will be necessary for our purpose to study very precisely the behavior of φ^{*n} , which we now do. The following statements may seem rather tedious, but they are exactly the ones we need. Recall that $\varphi^{*0}(t - s)ds = \delta_t(ds)$ and that $\varphi^{*n}(s) = 0$ for $s < 0$ by convention.

Lemma 15. *Recall that $\varphi : [0, \infty) \mapsto [0, \infty)$ and that $\Lambda = \int_0^\infty \varphi(s)ds$. Assume that there is $q \geq 1$ such that $\int_0^\infty s^q \varphi(s)ds < \infty$ and set $\kappa = \Lambda^{-1} \int_0^\infty s \varphi(s)ds$.*

(i) *For $n \geq 0$ and $t \geq 0$, we have $\int_0^t s \varphi^{*n}(t - s)ds = \Lambda^n t - n\Lambda^n \kappa + \varepsilon_n(t)$, where*

$$0 \leq \varepsilon_n(t) \leq Cn^q \Lambda^n t^{1-q} \quad \text{and} \quad \varepsilon_n(t) \leq n\Lambda^n \kappa.$$

(ii) *For $n \geq 0$, for $0 \leq t \leq z$ and $s \in [0, z]$, we set $\beta_n(t, z, s) = \varphi^{*n}(z - s) - \varphi^{*n}(t - s)$. Then $\int_0^z |\beta_n(t, z, s)|ds \leq 2\Lambda^n$ and for all $0 \leq \Delta \leq t$ and all $z \in [t, t + \Delta]$,*

$$\left| \int_0^z \beta_n(t, z, s)ds \right| \leq Cn^q \Lambda^n t^{-q},$$

$$\int_0^{t-\Delta} |\beta_n(t, z, s)| ds + \left| \int_{t-\Delta}^z \beta_n(t, z, s) ds \right| \leq Cn^q \Lambda^n \Delta^{-q}.$$

(iii) For $m, n \geq 0$, for $0 \leq t \leq z$, we put

$$\gamma_{m,n}(t, z) = \int_0^z \int_0^z (s \wedge u) \beta_m(t, z, s) \beta_n(t, z, u) duds.$$

It holds that $0 \leq \gamma_{m,n}(t, t+\Delta) \leq \Lambda^{m+n} \Delta$, for all $t \geq 0$, all $\Delta \geq 0$. Furthermore, there is a family $\kappa_{m,n}$ satisfying $0 \leq \kappa_{m,n} \leq (m+n)\kappa$ such that, for all $0 \leq \Delta \leq t$,

$$\gamma_{m,n}(t, t+\Delta) = \Delta \Lambda^{m+n} - \kappa_{m,n} \Lambda^{m+n} + \varepsilon_{m,n}(t, t+\Delta),$$

with $|\varepsilon_{m,n}(t, t+\Delta)| \leq C(m+n)^q \Lambda^{m+n} t \Delta^{-q}$.

Proof. We introduce some i.i.d. random variables X_1, X_2, \dots with density $\Lambda^{-1}\varphi$ and set $S_0 = 0$ as well as $S_n = X_1 + \dots + X_n$ for all $n \geq 1$. We observe that, by the Minkowski inequality, $\mathbb{E}[S_n^q] \leq n^q \mathbb{E}[X_1^q] \leq Cn^q$, since $\mathbb{E}[X_1^q] = \Lambda^{-1} \int_0^\infty s^q \varphi(s) ds < \infty$ by assumption.

To check (i), we use that S_n has for density $\Lambda^{-n}\varphi^{*n}$, so that we can write

$$\int_0^t s \varphi^{*n}(t-s) ds = \int_0^t (t-s) \varphi^{*n}(s) ds = \Lambda^n \mathbb{E}[(t-S_n)_+] = \Lambda^n t - \Lambda^n \mathbb{E}[S_n] + \varepsilon_n(t),$$

where $\varepsilon_n(t) = \Lambda^n \mathbb{E}[(S_n - t) \mathbf{1}_{\{S_n \geq t\}}]$. We clearly have that $\mathbb{E}[S_n] = n\kappa$, that $\varepsilon_n(t) \geq 0$ and that $\varepsilon_n(t) \leq \Lambda^n \mathbb{E}[S_n] = n\Lambda^n \kappa$. Finally, $\varepsilon_n(t) \leq \Lambda^n \mathbb{E}[S_n \mathbf{1}_{\{S_n \geq t\}}] \leq \Lambda^n t^{1-q} \mathbb{E}[S_n^q] \leq Cn^q \Lambda^n t^{1-q}$.

To check (ii), we observe that $\int_0^z |\beta_n(t, z, s)| ds \leq 2\Lambda^n$ is obvious because $\int_0^\infty \varphi^{*n}(s) ds = \Lambda^n$ and that, since $\mathbb{E}[S_n^q] \leq Cn^q$,

$$\int_r^\infty \varphi^{*n}(u) du = \Lambda^n \Pr(S_n \geq r) \leq Cn^q \Lambda^n r^{-q}.$$

We write $\int_0^z \beta_n(t, z, s) ds = \int_0^z \varphi^{*n}(z-s) ds - \int_0^t \varphi^{*n}(t-s) ds = \int_t^z \varphi^{*n}(u) du$, which implies that $|\int_0^z \beta_n(t, z, s) ds| \leq \int_t^\infty \varphi^{*n}(u) du \leq Cn^q \Lambda^n t^{-q}$. Next, we see that $\int_0^{t-\Delta} |\beta_n(t, z, s)| ds \leq \int_0^{t-\Delta} \varphi^{*n}(z-u) du + \int_0^{t-\Delta} \varphi^{*n}(t-u) du$, which is bounded by $2 \int_\Delta^\infty \varphi^{*n}(u) du \leq Cn^q \Lambda^n \Delta^{-q}$. Finally, using the two previous bounds, we find $|\int_{t-\Delta}^z \beta_n(t, z, s) ds| \leq |\int_0^z \beta_n(t, z, s) ds| + |\int_0^{t-\Delta} \beta_n(t, z, s) ds| \leq Cn^q \Lambda^n t^{-q} + Cn^q \Lambda^n \Delta^{-q} \leq Cn^q \Lambda^n \Delta^{-q}$ because $\Delta \in [0, t]$ by assumption.

We finally prove (iii) and thus consider $0 \leq \Delta \leq t$ and $m, n \geq 0$. We start from

$$\begin{aligned} \gamma_{m,n}(t, t+\Delta) &= \int_0^{t+\Delta} \int_0^{t+\Delta} (s \wedge u) \left[\varphi^{*m}(t+\Delta-s) \varphi^{*n}(t+\Delta-u) \right. \\ &\quad + \varphi^{*m}(t-s) \varphi^{*n}(t-u) - \varphi^{*m}(t+\Delta-s) \varphi^{*n}(t-u) \\ &\quad \left. - \varphi^{*m}(t-s) \varphi^{*n}(t+\Delta-u) \right] duds. \end{aligned}$$

Using another (independent) i.i.d. family Y_1, Y_2, \dots of random variables with density $\Lambda^{-1}\varphi$ and setting $T_m = Y_1 + \dots + Y_m$ (or $T_m = 0$ if $m = 0$), we may write

$$\begin{aligned} & \gamma_{m,n}(t, t + \Delta) \\ &= \Lambda^{m+n} \mathbb{E} \left[(t + \Delta - T_m)_+ \wedge (t + \Delta - S_n)_+ + (t - T_m)_+ \wedge (t - S_n)_+ \right. \\ & \quad \left. - (t + \Delta - T_m)_+ \wedge (t - S_n)_+ - (t - T_m)_+ \wedge (t + \Delta - S_n)_+ \right]. \end{aligned}$$

This precisely rewrites $\gamma_{m,n}(t, t + \Delta) = \Lambda^{m+n} \mathbb{E}[(t + \Delta - T_m \vee S_n)_+ - (t - T_m \wedge S_n)_+]$, which implies that $0 \leq \gamma_{m,n}(t, t + \Delta) \leq \Lambda^{m+n} \Delta$. We next introduce

$$\delta_{m,n}(t, t + \Delta) = \Lambda^{m+n} \mathbb{E}[(t + \Delta - T_m \vee S_n) - (t - T_m \wedge S_n)],$$

which equals $\delta_{m,n}(t, t + \Delta) = \Lambda^{m+n}(\Delta - \kappa_{m,n})$, where $\kappa_{m,n} = \mathbb{E}[|T_m - S_n|]$ satisfies $0 \leq \kappa_{m,n} \leq \kappa(m + n)$. Thus $\gamma_{m,n}(t, t + \Delta) = \Lambda^{m+n}(\Delta - \kappa_{m,n}) + \varepsilon_{m,n}(t, t + \Delta)$, where $\varepsilon_{m,n}(t, t + \Delta) = \gamma_{m,n}(t, t + \Delta) - \delta_{m,n}(t, t + \Delta)$. Finally, it is clear that, since $0 \leq \Delta \leq t$,

$$\begin{aligned} |\varepsilon_{m,n}(t, t + \Delta)| &\leq \Lambda^{m+n}(t + \Delta) \Pr(T_m \vee S_n \geq t + \Delta \text{ or } T_m \wedge S_n \geq t \\ & \quad \text{or } |T_m - S_n| \geq \Delta) \\ &\leq 2\Lambda^{m+n}t \Pr(T_m \geq \Delta \text{ or } S_n \geq \Delta). \end{aligned}$$

This is, as usual, bounded by $C\Lambda^{m+n}t(m^q + n^q)\Delta^{-q}$. □

4.3. Preliminary stochastic analysis

We handle once for all a number of useful computations concerning the processes introduced in Notation 9.

Lemma 16. *We assume $H(q)$ for some $q \geq 1$. Recall that Ω_N^1 and ℓ_N were defined in Notation 12 and that all the processes below have been introduced in Notation 9.*

(i) *For any $r \in [1, \infty]$, for all $t \geq 0$,*

$$\mathbf{1}_{\Omega_N^1} \|\mathbb{E}_\theta[\mathbf{Z}_t^N]\|_r \leq Ct \|\mathbf{1}_N\|_r.$$

(ii) *For any $r \in [1, \infty]$, for all $t \geq s \geq 0$,*

$$\mathbf{1}_{\Omega_N^1} \left\| \mathbb{E}_\theta \left[\mathbf{Z}_t^N - \mathbf{Z}_s^N \right] - \mu(t - s)\ell_N \right\|_r \leq C(1 \wedge s^{1-q}) \|\mathbf{1}_N\|_r.$$

(iii) *For all $t \geq s + 1 \geq 1$,*

$$\begin{aligned} & \mathbf{1}_{\Omega_N^1} \sup_{i=1, \dots, N} \mathbb{E}_\theta \left[(Z_t^{i,N} - Z_s^{i,N})^2 + \sup_{[s,t]} |M_r^{i,N} - M_s^{i,N}|^4 \right] + \mathbf{1}_{\Omega_N^1} \mathbb{E}_\theta \left[(\bar{Z}_t^N - \bar{Z}_s^N)^2 \right] \\ & \leq C(t - s)^2. \end{aligned}$$

Proof. Recall (3), which asserts that $\mathbb{E}_\theta[\mathbf{Z}_t^N] = \mu \sum_{n \geq 0} [\int_0^t s \varphi^{*n}(t-s) ds] A_N^n \mathbf{1}_N$. Since $\int_0^t s \varphi^{*n}(t-s) ds \leq t \Lambda^n$, we get $\|\mathbb{E}_\theta[\mathbf{Z}_t^N]\|_r \leq \mu t \sum_{n \geq 0} \Lambda^n \|A_N\|_r^n \|\mathbf{1}_N\|_r$. This is clearly bounded, on Ω_N^1 , by $Ct \|\mathbf{1}_N\|_r$, which proves (i).

Using next Lemma 15-(i), $\mathbb{E}_\theta[\mathbf{Z}_t^N] = \mu \sum_{n \geq 0} [\Lambda^n t - n \Lambda^n \kappa + \varepsilon_n(t)] A_N^n \mathbf{1}_N$, where $0 \leq \varepsilon_n(t) \leq C n^q \Lambda^n (t^{1-q} \wedge 1)$. Hence

$$\mathbb{E}_\theta[\mathbf{Z}_t^N] - \mathbb{E}_\theta[\mathbf{Z}_s^N] = \mu(t-s) \sum_{n \geq 0} \Lambda^n A_N^n \mathbf{1}_N + \mu \sum_{n \geq 0} [\varepsilon_n(t) - \varepsilon_n(s)] A_N^n \mathbf{1}_N.$$

But $\sum_{n \geq 0} \Lambda^n A_N^n \mathbf{1}_N = Q_N \mathbf{1}_N = \ell_N$ on Ω_N^1 . Thus, still on Ω_N^1 , since $s \leq t$ and $q \geq 1$,

$$\begin{aligned} \left\| \mathbb{E}_\theta[\mathbf{Z}_t^N - \mathbf{Z}_s^N] - \mu(t-s)\ell_N \right\|_r &\leq C(1 \wedge s^{1-q}) \sum_{n \geq 0} n^q \Lambda^n \|A_N\|_r^n \|\mathbf{1}_N\|_r \\ &\leq C(1 \wedge s^{1-q}) \|\mathbf{1}_N\|_r. \end{aligned}$$

Since $[M^{i,N}, M^{i,N}]_t = Z_t^{i,N}$ by Remark 10, the Doob inequality implies that $\mathbb{E}_\theta[\sup_{[s,t]} |M_r^{i,N} - M_s^{i,N}|^4] \leq C \mathbb{E}_\theta[(Z_t^{i,N} - Z_s^{i,N})^2]$. Also, the Cauchy-Schwarz inequality tells us that $\mathbb{E}_\theta[(\bar{Z}_t^N - \bar{Z}_s^N)^2] \leq N^{-1} \sum_{i=1}^N \mathbb{E}_\theta[(Z_t^{i,N} - Z_s^{i,N})^2] \leq \sup_{i=1, \dots, N} \mathbb{E}_\theta[(Z_t^{i,N} - Z_s^{i,N})^2]$.

Hence we just have to prove that $\sup_{i=1, \dots, N} \mathbb{E}_\theta[(Z_t^{i,N} - Z_s^{i,N})^2] \leq C(t-s)^2$. Recalling that $Z_t^{i,N} = U_t^{i,N} + \mathbb{E}_\theta[Z_t^{i,N}]$, we have to show that, on Ω_N^1 , (a) $(\mathbb{E}_\theta[Z_t^{i,N}] - \mathbb{E}_\theta[Z_s^{i,N}])^2 \leq C(t-s)^2$ and (b) $\mathbb{E}_\theta[(U_t^{i,N} - U_s^{i,N})^2] \leq C(t-s)^2$.

To prove (a), we use (ii) with $r = \infty$ and find that, on Ω_N^1 , $\mathbb{E}_\theta[Z_t^{i,N}] - \mathbb{E}_\theta[Z_s^{i,N}] \leq \mu(t-s) \|\ell_N\|_\infty + C \|\mathbf{1}_N\|_\infty \leq C(t-s)$, since ℓ_N is bounded on Ω_N^1 and since $t-s \geq 1$ by assumption.

To prove (b), we use (4) to write

$$U_t^{i,N} - U_s^{i,N} = \sum_{n \geq 0} \int_0^t \beta_n(s, t, r) \sum_{j=1}^N A_N^n(i, j) M_r^{j,N} dr,$$

where we have set $\beta_n(s, t, r) = \varphi^{*n}(t-r) - \varphi^{*n}(s-r)$ as in Lemma 15. We deduce that

$$\begin{aligned} \mathbb{E}[(U_t^{i,N} - U_s^{i,N})^2] &= \sum_{m, n \geq 0} \int_0^t \int_0^t \beta_m(s, t, u) \beta_n(s, t, v) \\ &\quad \sum_{j, k=1}^N A_N^m(i, j) A_N^n(i, k) \mathbb{E}_\theta[M_u^{j,N} M_v^{k,N}] dv du. \end{aligned}$$

By Remark 10, $\mathbb{E}_\theta[M_u^{j,N} M_v^{k,N}] = \mathbf{1}_{\{j=k\}} \mathbb{E}_\theta[Z_{u \wedge v}^{j,N}]$. Using now (ii) with $s = 0$ and $r = \infty$, we see that $x_t^{j,N} := \mathbb{E}_\theta[Z_t^{j,N}] - \mu t \ell_N(j)$ satisfies $\sup_{t \geq 0, j=1, \dots, N} |x_t^{j,N}| \leq C$ on Ω_N^1 . We thus write $\mathbb{E}_\theta[(U_t^{i,N} - U_s^{i,N})^2] = I + J$, where

$$I = \mu \sum_{m, n \geq 0} \int_0^t \int_0^t \beta_m(s, t, u) \beta_n(s, t, v) \sum_{j=1}^N A_N^m(i, j) A_N^n(i, j) (u \wedge v) \ell_N(j) du dv,$$

$$J = \sum_{m,n \geq 0} \int_0^t \int_0^t \beta_m(s, t, u) \beta_n(s, t, v) \sum_{j=1}^N A_N^m(i, j) A_N^n(i, j) x_{u \wedge v}^{j, N} du dv.$$

First, using that $x_t^{j, N}$ is bounded on Ω_N^1 and that $\int_0^t |\beta_m(s, t, u)| du \leq 2\Lambda^m$, we find $|J| \leq C \sum_{m,n \geq 0} \Lambda^{m+n} \sum_{j=1}^N A_N^m(i, j) A_N^n(i, j) = C \sum_{j=1}^N (Q_N(i, j))^2$ on Ω_N^1 , whence $|J| \leq C \sum_{j=1}^N (\mathbf{1}_{\{i=j\}} + N^{-1})^2$ by (8). We conclude that $|J| \leq C \leq C(t - s)^2$. Next, we realize that, with the notation of Lemma 15-(iii),

$$I = \mu \sum_{m,n \geq 0} \gamma_{m,n}(s, t) \sum_{j=1}^N A_N^m(i, j) A_N^n(i, j) \ell_N(j).$$

But we know that $0 \leq \gamma_{m,n}(s, t) \leq \Lambda^{m+n}(t - s)$, so that

$$I \leq \mu(t - s) \sum_{j=1}^N (Q_N(i, j))^2 \ell_N(j) \leq C(t - s),$$

since ℓ_N is bounded on Ω_N^1 and since, as already seen, $\sum_{j=1}^N (Q_N(i, j))^2$ is also bounded on Ω_N^1 . We conclude that $\mathbb{E}_\theta[(U_t^{i, N} - U_r^{i, N})^2] \leq C(t - s) \leq C(t - s)^2$ on Ω_N^1 , as desired. \square

4.4. First estimator

We recall that $\mathcal{E}_t^N = (\bar{Z}_{2t}^N - \bar{Z}_t^N)/t$, that the matrices A_N and Q_N and the event Ω_N^1 were defined in Notation 12, as well as $\ell_N(i) = \sum_{j=1}^N Q_N(i, j)$ and $\bar{\ell}_N = N^{-1} \sum_{i=1}^N \ell_N(i)$. The goal of this subsection is to establish the following estimate.

Proposition 17. *Assume $H(q)$ for some $q \geq 1$. Then for $t \geq 1$,*

$$\mathbf{1}_{\Omega_N^1} \mathbb{E}_\theta \left[\left| \mathcal{E}_t^N - \mu \bar{\ell}_N \right|^2 \right] \leq C \left(\frac{1}{t^{2q}} + \frac{1}{Nt} \right).$$

We start with the following lemma (recall that \bar{U}^N was defined in Notation 9).

Lemma 18. *Assume $H(q)$ for some $q \geq 1$. Then on Ω_N^1 , for $t \geq 1$,*

$$\left| \mathbb{E}_\theta[\mathcal{E}_t^N] - \mu \bar{\ell}_N \right| \leq Ct^{-q} \quad \text{and} \quad \mathbb{E}_\theta[|\bar{U}_t^N|^2] \leq CtN^{-1}.$$

Proof. Applying Lemma 16-(ii) with $r = 1$, we immediately find, on Ω_N^1 ,

$$\left| \mathbb{E}_\theta[\mathcal{E}_t^N] - \mu \bar{\ell}_N \right| \leq N^{-1} \left\| \mathbb{E}_\theta \left[\frac{\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N}{t} \right] - \mu \ell_N \right\|_1 \leq CN^{-1}t^{-q} \|\mathbf{1}_N\|_1 = Ct^{-q}.$$

Next, $\bar{U}_t^N = N^{-1} \sum_{n \geq 0} \int_0^t \varphi^{*n}(t - s) \sum_{i,j=1}^N A_N^n(i, j) M_s^{j, N} ds$ by (4). Hence

$$\mathbb{E}_\theta[|\bar{U}_t^N|^2]^{1/2} \leq N^{-1} \sum_{n \geq 0} \int_0^t \varphi^{*n}(t-s) \mathbb{E}_\theta \left[\left(\sum_{i,j=1}^N A_N^n(i,j) M_s^{j,N} \right)^2 \right]^{1/2} ds$$

by the Minkowski inequality. But recalling Remark 10, i.e. $\mathbb{E}_\theta[M_s^{j,N} M_s^{l,N}] = \mathbf{1}_{\{j=l\}} \mathbb{E}_\theta[Z_s^{j,N}]$,

$$\begin{aligned} \mathbb{E}_\theta \left[\left(\sum_{i,j=1}^N A_N^n(i,j) M_s^{j,N} \right)^2 \right] &= \sum_{j=1}^N \left(\sum_{i=1}^N A_N^n(i,j) \right)^2 \mathbb{E}_\theta[Z_s^{j,N}] \\ &\leq \|A_N\|_1^{2n} \sum_{j=1}^N \mathbb{E}_\theta[Z_s^{j,N}]. \end{aligned}$$

We know from Lemma 16-(i) with $r = 1$ that $\sum_{j=1}^N \mathbb{E}_\theta[Z_s^{j,N}] \leq CNs$ on Ω_N^1 . Hence, still on Ω_N^1 ,

$$\mathbb{E}_\theta[|\bar{U}_t^N|^2]^{1/2} \leq \frac{C}{N} \sum_{n \geq 0} \|A_N\|_1^n \int_0^t \sqrt{Ns} \varphi^{*n}(t-s) ds \leq \frac{Ct^{1/2}}{N^{1/2}} \sum_{n \geq 0} \Lambda^n \|A_N\|_1^n,$$

which is smaller than $Ct^{1/2}N^{-1/2}$ as desired. □

We can now give the

Proof of Proposition 17. It suffices to write

$$\mathbb{E}_\theta \left[\left| \mathcal{E}_t^N - \mu \bar{\ell}_N \right|^2 \right] \leq 2 \mathbb{E}_\theta \left[\left| \mathcal{E}_t^N - \mathbb{E}_\theta[\mathcal{E}_t^N] \right|^2 \right] + 2 \left| \mathbb{E}_\theta[\mathcal{E}_t^N] - \mu \bar{\ell}_N \right|^2$$

and to observe that $|\mathcal{E}_t^N - \mathbb{E}_\theta[\mathcal{E}_t^N]| = |\bar{U}_{2t}^N - \bar{U}_t^N|/t \leq |\bar{U}_{2t}^N|/t + |\bar{U}_t^N|/t$, whence finally

$$\mathbb{E}_\theta \left[\left| \mathcal{E}_t^N - \mu \bar{\ell}_N \right|^2 \right] \leq \frac{4}{t^2} (\mathbb{E}_\theta[|\bar{U}_{2t}^N|^2] + \mathbb{E}_\theta[|\bar{U}_t^N|^2]) + 2 \left| \mathbb{E}_\theta[\mathcal{E}_t^N] - \mu \bar{\ell}_N \right|^2.$$

Then the proposition immediately follows from Lemma 18. □

4.5. Second estimator

We recall that $\mathcal{V}_t^N = \sum_{i=1}^N [(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mathcal{E}_t^N]^2 - N\mathcal{E}_t^N/t$ where $\mathcal{E}_t^N = (\bar{Z}_{2t}^N - \bar{Z}_t^N)/t$, that the matrices A_N and Q_N and the event Ω_N^1 were defined in Notation 12, as well as $\ell_N(i) = \sum_{j=1}^N Q_N(i,j)$ and $\bar{\ell}_N = N^{-1} \sum_{i=1}^N \ell_N(i)$. We also introduce $\mathcal{V}_\infty^N = \mu^2 \sum_{i=1}^N [\ell_N(i) - \bar{\ell}_N]^2$.

Proposition 19. Assume $H(q)$ for some $q \geq 1$. Then for $t \geq 1$, a.s.,

$$\mathbf{1}_{\Omega_N^1} \mathbb{E}_\theta \left[\left| \mathcal{V}_t^N - \mathcal{V}_\infty^N \right| \right] \leq C \left(1 + \sum_{i=1}^N [\ell_N(i) - \bar{\ell}_N]^2 \right)^{1/2} \left(\frac{N}{t^q} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{t}} \right).$$

Observe that the term $\sum_{i=1}^N [\ell_N(i) - \bar{\ell}_N]^2$ will not cause any problem, since its expectation (restricted to Ω_N^1) is uniformly bounded, see Proposition 14.

We write $|\mathcal{V}_t^N - \mathcal{V}_\infty^N| \leq \Delta_t^{N,1} + \Delta_t^{N,2} + \Delta_t^{N,3}$, where

$$\begin{aligned} \Delta_t^{N,1} &= \left| \sum_{i=1}^N [(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mathcal{E}_t^N]^2 - \sum_{i=1}^N [(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mu\bar{\ell}_N]^2 \right|, \\ \Delta_t^{N,2} &= \left| \sum_{i=1}^N [(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mu\ell_N(i)]^2 - N\mathcal{E}_t^N/t \right|, \\ \Delta_t^{N,3} &= 2 \left| \sum_{i=1}^N [(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mu\ell_N(i)][\mu\ell_N(i) - \mu\bar{\ell}_N] \right|. \end{aligned}$$

We next write $\Delta_t^{N,2} \leq \Delta_t^{N,21} + \Delta_t^{N,22} + \Delta_t^{N,23}$, where

$$\begin{aligned} \Delta_t^{N,21} &= \left| \sum_{i=1}^N \left[(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mathbb{E}_\theta[(Z_{2t}^{i,N} - Z_t^{i,N})/t] \right]^2 - N\mathcal{E}_t^N/t \right|, \\ \Delta_t^{N,22} &= \sum_{i=1}^N \left[\mathbb{E}_\theta[(Z_{2t}^{i,N} - Z_t^{i,N})/t] - \mu\ell_N(i) \right]^2, \\ \Delta_t^{N,23} &= 2 \left| \sum_{i=1}^N \left[(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mathbb{E}_\theta[(Z_{2t}^{i,N} - Z_t^{i,N})/t] \right] \right. \\ &\quad \left. \times \left[\mathbb{E}_\theta[(Z_{2t}^{i,N} - Z_t^{i,N})/t] - \mu\ell_N(i) \right] \right|. \end{aligned}$$

We will also need to write, recalling that $U_t^{i,N} = Z_t^{i,N} - \mathbb{E}_\theta[Z_t^{i,N}]$,

$$\Delta_t^{N,21} = \left| \sum_{i=1}^N \left[(U_{2t}^{i,N} - U_t^{i,N})/t \right]^2 - N\mathcal{E}_t^N/t \right| \leq \Delta_t^{N,211} + \Delta_t^{N,212} + \Delta_t^{N,213},$$

where

$$\begin{aligned} \Delta_t^{N,211} &= \left| \sum_{i=1}^N \left\{ \left((U_{2t}^{i,N} - U_t^{i,N})/t \right)^2 - \mathbb{E}_\theta \left[\left((U_{2t}^{i,N} - U_t^{i,N})/t \right)^2 \right] \right\} \right|, \\ \Delta_t^{N,212} &= \left| \sum_{i=1}^N \mathbb{E}_\theta \left[\left((U_{2t}^{i,N} - U_t^{i,N})/t \right)^2 \right] - \mathbb{E}_\theta[N\mathcal{E}_t^N/t] \right|, \\ \Delta_t^{N,213} &= \left| N\mathcal{E}_t^N/t - \mathbb{E}_\theta[N\mathcal{E}_t^N/t] \right|. \end{aligned}$$

Finally, we will use that $\Delta_t^{N,3} \leq \Delta_t^{N,31} + \Delta_t^{N,32}$, where

$$\begin{aligned} \Delta_t^{N,31} &= 2 \left| \sum_{i=1}^N \left[(Z_{2t}^{i,N} - Z_t^{i,N})/t - \mathbb{E}_\theta[(Z_{2t}^{i,N} - Z_t^{i,N})/t] \right] \left[\mu\ell_N(i) - \mu\bar{\ell}_N \right] \right|, \\ \Delta_t^{N,32} &= 2 \left| \sum_{i=1}^N \left[\mathbb{E}_\theta[(Z_{2t}^{i,N} - Z_t^{i,N})/t] - \mu\ell_N(i) \right] \left[\mu\ell_N(i) - \mu\bar{\ell}_N \right] \right|. \end{aligned}$$

To summarize, we have to bound $\Delta_t^{N,1}$, $\Delta_t^{N,211}$, $\Delta_t^{N,212}$, $\Delta_t^{N,213}$, $\Delta_t^{N,22}$, $\Delta_t^{N,23}$, $\Delta_t^{N,31}$ and $\Delta_t^{N,32}$. Only the term $\Delta_t^{N,211}$ is really difficult.

In the following lemma, we treat the easy terms. We do not try to be optimal when not useful: for example in (iv) below, some sharper estimate could probably be obtained with more work, but since we already have a term in $N^{1/2}t^{-1}$ (see Lemma 24), this would be useless. We also recall that we do not really try to optimize the dependence in q : it is likely that t^{-q} could be replaced by t^{-2q} here and there.

Lemma 20. *Assume $H(q)$ for some $q \geq 1$. Then a.s. on Ω_N^1 , for $t \geq 1$,*

- (i) $\mathbb{E}_\theta[\Delta_t^{N,1}] \leq C(Nt^{-2q} + t^{-1})$,
- (ii) $\mathbb{E}_\theta[\Delta_t^{N,22}] \leq CNt^{-2q}$,
- (iii) $\mathbb{E}_\theta[\Delta_t^{N,23}] \leq CNt^{-q}$,
- (iv) $\mathbb{E}_\theta[\Delta_t^{N,213}] \leq CN^{1/2}t^{-3/2}$,
- (v) $\mathbb{E}_\theta[\Delta_t^{N,32}] \leq CNt^{-q}$.

Proof. We work on Ω_N^1 during the whole proof.

Using that $\mathcal{E}_t^N = N^{-1} \sum_{i=1}^N [(Z_{2t}^{i,N} - Z_t^{i,N})/t]$, one easily checks that $\Delta_t^{N,1} = N|\mathcal{E}_t^N - \mu\bar{\ell}_N|^2$. Thus point (i) follows from Proposition 17.

Next, we observe that $\Delta_t^{N,22} = \|\mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t] - \mu\ell_N\|_2^2$. Applying Lemma 16-(ii) with $r = 2$, we conclude that indeed, $\Delta_t^{N,22} \leq Ct^{-2q}\|\mathbf{1}_N\|_2^2 = CNt^{-2q}$.

We write

$$\Delta_t^{N,23} \leq 2\left\|(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t - \mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t]\right\|_1 \left\|\mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t] - \mu\ell_N\right\|_\infty.$$

Applying Lemma 16-(ii) with $r = \infty$, we deduce that $\|\mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t] - \mu\ell_N\|_\infty \leq Ct^{-q}$. Lemma 16-(i) with $r = 1$ gives us that $\mathbb{E}_\theta[\|(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t - \mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t]\|_1] \leq 2t^{-1}\|\mathbb{E}_\theta[\mathbf{Z}_{2t}^N + \mathbf{Z}_t^N]\|_1 \leq CN$. We thus find that indeed, $\mathbb{E}_\theta[\Delta_t^{N,23}] \leq CNt^{-q}$.

Since $\Delta_t^{N,213} = (N/t)|\mathcal{E}_t^N - \mathbb{E}_\theta[\mathcal{E}_t^N]| = Nt^{-2}|\bar{U}_{2t}^N - \bar{U}_t^N| \leq Nt^{-2}(|\bar{U}_{2t}^N| + |\bar{U}_t^N|)$, we deduce from Lemma 18 that $\mathbb{E}_\theta[\Delta_t^{N,213}] \leq CNt^{-2}\sqrt{t/N} = CN^{1/2}t^{-3/2}$.

Finally, starting from $\Delta_t^{N,32} \leq 2\mu\|\mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t] - \mu\ell_N\|_\infty\|\ell_N - \bar{\ell}_N\mathbf{1}_N\|_1$ and using that, as already seen when studying $\Delta_t^{N,23}$, $\|\mathbb{E}_\theta[(\mathbf{Z}_{2t}^N - \mathbf{Z}_t^N)/t] - \mu\ell_N\|_\infty \leq Ct^{-q}$, we conclude that $\Delta_t^{N,32} \leq Ct^{-q}\|\ell_N - \bar{\ell}_N\mathbf{1}_N\|_1 \leq CNt^{-q}$, since ℓ_N is bounded (see (7)) on Ω_N^1 . \square

Next, we treat the term $\Delta_t^{N,212}$.

Lemma 21. *Assume $H(q)$ for some $q \geq 1$. Then a.s. on Ω_N^1 , for $t \geq 1$, $\mathbb{E}_\theta[\Delta_t^{N,212}] \leq Ct^{-1}$.*

Proof. We work on Ω_N^1 . Recalling that $N\mathcal{E}_t^N = t^{-1} \sum_{i=1}^N (Z_{2t}^{i,N} - Z_t^{i,N})$, we may write $\mathbb{E}_\theta[\Delta_t^{N,212}] \leq t^{-2} \sum_{i=1}^N a_i$, where $a_i = |\mathbb{E}_\theta[(U_{2t}^{i,N} - U_t^{i,N})^2 - (Z_{2t}^{i,N} - Z_t^{i,N})]|$. We infer from (4) that $U_t^{i,N} = M_t^{i,N} + \sum_{n \geq 1} \int_0^t \varphi^{*n}(t-s) \sum_{j=1}^N A_N^n(i, j) M_s^{j,N} ds$,

so that $U_{2t}^{i,N} - U_t^{i,N} = M_{2t}^{i,N} - M_t^{i,N} + R_t^{i,N}$, where

$$R_t^{i,N} = \sum_{n \geq 1} \int_0^{2t} \beta_n(t, 2t, s) \sum_{j=1}^N A_N^n(i, j) M_s^{j,N} ds.$$

We have set $\beta_n(t, 2t, s) = \varphi^{*n}(2t - s) - \varphi^{*n}(t - s)$ as in Lemma 15 and the only thing we will use is that $\int_0^{2t} |\beta_n(t, 2t, s)| ds \leq 2\Lambda^n$. Recalling that $M^{i,N}$ is a martingale with quadratic variation $[M^{i,N}, M^{i,N}]_t = Z_t^{i,N}$, see Remark 10, we deduce that $\mathbb{E}_\theta[(M_{2t}^{i,N} - M_t^{i,N})^2] = \mathbb{E}_\theta[Z_{2t}^{i,N} - Z_t^{i,N}]$. Hence

$$a_i = \mathbb{E}_\theta[(R_t^{i,N})^2] + 2\mathbb{E}_\theta[(M_{2t}^{i,N} - M_t^{i,N})R_t^{i,N}] = b_i + d_i,$$

the last equality standing for a definition. We first write

$$b_i = \sum_{m,n \geq 1} \int_0^{2t} \int_0^{2t} \beta_m(t, 2t, s) \beta_n(t, 2t, u) \sum_{j,k=1}^N A_N^m(i, j) A_N^n(i, k) \mathbb{E}_\theta[M_s^{j,N} M_u^{k,N}] duds.$$

But we know that $\mathbb{E}_\theta[M_s^{j,N} M_u^{k,N}] = \mathbf{1}_{\{j=k\}} \mathbb{E}_\theta[Z_{s \wedge u}^{j,N}]$ by Remark 10 and that $\mathbb{E}_\theta[Z_{s \wedge u}^{j,N}] \leq Ct$ on Ω_N^1 by Lemma 16-(i) (with $r = \infty$). Hence

$$b_i \leq Ct \sum_{m,n \geq 1} \Lambda^{m+n} \sum_{j=1}^N A_N^m(i, j) A_N^n(i, j) = Ct \sum_{j=1}^N \left(\sum_{n \geq 1} \Lambda^n A_N^n(i, j) \right)^2.$$

But $\sum_{n \geq 1} \Lambda^n A_N^n(i, j) = Q_N(i, j) - \mathbf{1}_{\{i=j\}} \leq CN^{-1}$ on Ω_N^1 by (8), so that $b_i \leq CtN^{-1}$.

Next, we start from

$$d_i = 2 \sum_{n \geq 1} \int_0^{2t} \beta_n(t, 2t, s) \sum_{j=1}^N A_N^n(i, j) \mathbb{E}_\theta[(M_{2t}^{i,N} - M_t^{i,N}) M_s^{j,N}] ds.$$

As previously, we see that $\mathbb{E}_\theta[(M_{2t}^{i,N} - M_t^{i,N}) M_s^{j,N}] = 0$ if $i \neq j$ and that $\mathbb{E}_\theta[(M_{2t}^{i,N} - M_t^{i,N}) M_s^{i,N}] = \mathbb{E}_\theta[Z_{2t \wedge s}^{i,N} - Z_{t \wedge s}^{i,N}] \leq Ct$ on Ω_N^1 (by Lemma 16-(i)), whence

$$d_i \leq Ct \sum_{n \geq 1} \Lambda^n A_N^n(i, i) = Ct(Q_N(i, i) - 1) \leq CtN^{-1}$$

on Ω_N^1 by (8) again. Finally, $a_i \leq CtN^{-1}$, so that $\mathbb{E}_\theta[\Delta_t^{N,212}] \leq t^{-2} \sum_{i=1}^N a_i \leq Ct^{-1}$ on Ω_N^1 . \square

We next compute some covariances in the following tedious lemma.

Lemma 22. *Assume $H(q)$ for some $q \geq 1$. Then a.s., on Ω_N^1 , for all $t \geq 1$, all $k, l, a, b \in \{1, \dots, N\}$, all $r, s, u, v \in [0, t]$,*

- (i) $|\text{Cov}_\theta(Z_r^{k,N}, Z_s^{l,N})| = |\text{Cov}_\theta(U_r^{k,N}, U_s^{l,N})| \leq Ct(N^{-1} + \mathbf{1}_{\{k=l\}}),$
(ii) $|\text{Cov}_\theta(Z_r^{k,N}, M_s^{l,N})| = |\text{Cov}_\theta(U_r^{k,N}, M_s^{l,N})| \leq Ct(N^{-1} + \mathbf{1}_{\{k=l\}}),$
(iii) $|\text{Cov}_\theta(Z_r^{k,N}, \int_0^s M_{\tau-}^{l,N} dM_\tau^{l,N})| = |\text{Cov}_\theta(U_r^{k,N}, \int_0^s M_{\tau-}^{l,N} dM_\tau^{l,N})|$ is smaller than $Ct^{3/2}(N^{-1} + \mathbf{1}_{\{k=l\}}),$
(iv) $|\mathbb{E}_\theta[M_r^{k,N} M_s^{l,N} M_u^{l,N}]| \leq CN^{-1}t$ if $\#\{k, l\} = 2,$
(v) $|\text{Cov}_\theta(M_r^{k,N} M_s^{l,N}, M_u^{a,N} M_v^{b,N})| = 0$ if $\#\{k, l, a, b\} = 4,$
(vi) $|\text{Cov}_\theta(M_r^{k,N} M_s^{k,N}, M_u^{a,N} M_v^{b,N})| \leq CN^{-2}t$ if $\#\{k, a, b\} = 3,$
(vii) $|\text{Cov}_\theta(M_r^{k,N} M_s^{k,N}, M_u^{a,N} M_v^{a,N})| \leq CN^{-1}t^{3/2}$ if $\#\{k, a\} = 2,$
(viii) $|\text{Cov}_\theta(M_r^{k,N} M_s^{l,N}, M_u^{a,N} M_v^{b,N})| \leq Ct^2$ without condition.

Proof. We work on Ω_N^1 and start with point (i). First, it is clear, since $U_t^{k,N} = Z_t^{k,N} - \mathbb{E}_\theta[Z_t^{k,N}]$, that $\text{Cov}_\theta(Z_r^{k,N}, Z_s^{l,N}) = \text{Cov}_\theta(U_r^{k,N}, U_s^{l,N})$. Then we infer from (4) that

$$\begin{aligned} \text{Cov}_\theta(U_r^{k,N}, U_s^{l,N}) &= \sum_{m,n \geq 0} \int_0^r \int_0^s \varphi^{*m}(r-x) \varphi^{*n}(s-y) \\ &\quad \sum_{i,j=1}^N A_N^m(k,i) A_N^n(l,j) \text{Cov}_\theta(M_x^{i,N}, M_y^{j,N}) dy dx. \end{aligned}$$

But we know (see Remark 10) that $\text{Cov}_\theta(M_x^{i,N}, M_y^{j,N}) = \mathbf{1}_{\{i=j\}} \mathbb{E}_\theta[Z_{x \wedge y}^{i,N}] \leq C \mathbf{1}_{\{i=j\}} t$ by Lemma 16-(i) (with $r = \infty$). Thus

$$\begin{aligned} |\text{Cov}_\theta(U_r^{k,N}, U_s^{l,N})| &\leq Ct \sum_{m,n \geq 0} \Lambda^{m+n} \sum_{i=1}^N A_N^m(k,i) A_N^n(l,i) \\ &= Ct \sum_{i=1}^N Q_N(k,i) Q_N(l,i). \end{aligned}$$

Recalling (8), $\sum_{i=1}^N Q_N(k,i) Q_N(l,i) \leq C \sum_{i=1}^N (N^{-1} + \mathbf{1}_{\{k=i\}})(N^{-1} + \mathbf{1}_{\{l=i\}}) \leq C(N^{-1} + \mathbf{1}_{\{k=l\}})$. Point (i) is checked.

For point (ii), we again have $\text{Cov}_\theta(Z_r^{k,N}, M_s^{l,N}) = \text{Cov}_\theta(U_r^{k,N}, M_s^{l,N})$ and, using again (4),

$$\text{Cov}_\theta(U_r^{k,N}, M_s^{l,N}) = \sum_{n \geq 0} \int_0^r \varphi^{*n}(r-x) \sum_{i=1}^N A_N^n(k,i) \text{Cov}_\theta(M_x^{i,N}, M_s^{l,N}) dx.$$

Since $|\text{Cov}_\theta(M_x^{i,N}, M_s^{l,N})| \leq C \mathbf{1}_{\{i=l\}} t$ as in (i), we conclude that

$$|\text{Cov}_\theta(U_r^{k,N}, M_s^{l,N})| \leq Ct \sum_{n \geq 0} \Lambda^n A_N^n(k,l) = Ct Q_N(k,l) \leq Ct(N^{-1} + \mathbf{1}_{\{k=l\}}).$$

Point (iii) is checked similarly as point (ii): we just have to verify that $|\text{Cov}_\theta(M_x^{i,N}, \int_0^s M_{\tau-}^{l,N} dM_\tau^{l,N})| \leq C \mathbf{1}_{\{i=l\}} t^{3/2}$. This is obvious if $i \neq l$ because

the martingales $M^{i,N}$ and $\int_0^\cdot M_{\tau-}^{l,N} dM_\tau^{l,N}$ are orthogonal, and relies on the fact, if $i = l$, that

$$\begin{aligned} \left| \text{Cov}_\theta \left(M_x^{i,N}, \int_0^s M_{\tau-}^{i,N} dM_\tau^{i,N} \right) \right| &\leq \mathbb{E}_\theta[|M_x^{i,N}|^2]^{1/2} \mathbb{E}_\theta \left[\left| \int_0^s M_{\tau-}^{i,N} dM_\tau^{i,N} \right|^2 \right]^{1/2} \\ &\leq Ct^{3/2}. \end{aligned}$$

The last inequality uses that $\mathbb{E}_\theta[|M_x^{i,N}|^2] = \mathbb{E}_\theta[Z_x^{i,N}] \leq Ct$ by Remark 10 and Lemma 16-(i) and that $\mathbb{E}_\theta[|\int_0^s M_{\tau-}^{i,N} dM_\tau^{i,N}|^2] \leq Ct^2$. Indeed, we have $[\int_0^\cdot M_{\tau-}^{i,N} dM_\tau^{i,N}, \int_0^\cdot M_{\tau-}^{i,N} dM_\tau^{i,N}]_s = \int_0^s (M_{\tau-}^{i,N})^2 dZ_\tau^{i,N} \leq (M_s^{i,N,*})^2 Z_s^{i,N}$, whence

$$\mathbb{E}_\theta \left[\left| \int_0^s M_{\tau-}^{i,N} dM_\tau^{i,N} \right|^2 \right] \leq \mathbb{E}_\theta[(M_s^{i,N,*})^2 Z_s^{i,N}] \leq \mathbb{E}_\theta[(M_s^{i,N,*})^4]^{1/2} \mathbb{E}_\theta[(Z_s^{i,N})^2]^{1/2},$$

which is bounded by Ct^2 by Lemma 16-(iii).

For point (iv), we assume e.g. that $r \leq s$ and first note that

$$\mathbb{E}_\theta[M_r^{k,N} M_s^{k,N} M_u^{l,N}] = \mathbb{E}_\theta[M_r^{k,N} \mathbb{E}_\theta[M_s^{k,N} M_u^{l,N} | \mathcal{F}_r]] = \mathbb{E}_\theta[(M_r^{k,N})^2 M_{u \wedge r}^{l,N}]$$

because the martingales $M^{k,N}$ and $M^{l,N}$ are orthogonal. Since $[M^{k,N}, M^{k,N}]_r = Z_r^{k,N}$, we have $(M_r^{k,N})^2 = 2 \int_0^r M_{\tau-}^{k,N} dM_\tau^{k,N} + Z_r^{k,N}$. Since $\int_0^\cdot M_{\tau-}^{k,N} dM_\tau^{k,N}$ and $M^{l,N}$ are orthogonal, we conclude that $\mathbb{E}_\theta[(M_r^{k,N})^2 M_{u \wedge r}^{l,N}] = \mathbb{E}_\theta[Z_r^{k,N} M_{u \wedge r}^{l,N}] = \text{Cov}_\theta(Z_r^{k,N}, M_{u \wedge r}^{l,N})$. Since $k \neq l$, we conclude using point (ii).

Point (v) is obvious, since when k, l, a, b are pairwise different, the martingales $M^{k,N}, M^{l,N}, M^{a,N}$ and $M^{b,N}$ are orthogonal.

Point (vi) is harder. Recall that $\#\{k, a, b\} = 3$, so that clearly,

$$\text{Cov}_\theta(M_r^{k,N} M_s^{k,N}, M_u^{a,N} M_v^{b,N}) = \mathbb{E}_\theta[M_r^{k,N} M_s^{k,N} M_u^{a,N} M_v^{b,N}].$$

We assume e.g. that $r \leq s$ and we observe that

$$\begin{aligned} \mathbb{E}_\theta[M_r^{k,N} M_s^{k,N} M_u^{a,N} M_v^{b,N}] &= \mathbb{E}_\theta[M_r^{k,N} \mathbb{E}_\theta[M_s^{k,N} M_u^{a,N} M_v^{b,N} | \mathcal{F}_r]] \\ &= \mathbb{E}_\theta[(M_r^{k,N})^2 M_{u \wedge r}^{a,N} M_{v \wedge r}^{b,N}] \end{aligned}$$

because $M^{k,N}, M^{a,N}$ and $M^{b,N}$ are orthogonal. We thus have to prove that for all $r, u, v \in [0, t]$ with $u, v \leq r$, $|\mathbb{E}_\theta[(M_r^{k,N})^2 M_u^{a,N} M_v^{b,N}]| \leq CN^{-2}t$. We write $(M_r^{k,N})^2 = 2 \int_0^r M_{\tau-}^{k,N} dM_\tau^{k,N} + Z_r^{k,N}$ as in the proof of (iv). The three martingales $\int_0^\cdot M_{\tau-}^{k,N} dM_\tau^{k,N}, M^{a,N}$ and $M^{b,N}$ being orthogonal, we find that $\mathbb{E}_\theta[(M_r^{k,N})^2 M_u^{a,N} M_v^{b,N}] = \mathbb{E}_\theta[Z_r^{k,N} M_u^{a,N} M_v^{b,N}] = \mathbb{E}_\theta[U_r^{k,N} M_u^{a,N} M_v^{b,N}]$. We now write, starting again from (4),

$$\mathbb{E}_\theta[U_r^{k,N} M_u^{a,N} M_v^{b,N}] = \sum_{n \geq 0} \int_0^r \varphi^{*n}(r-x) \sum_{j=1}^N A_N^n(k, j) \mathbb{E}_\theta[M_x^{j,N} M_u^{a,N} M_v^{b,N}] dx.$$

But $|\mathbb{E}_\theta[M_x^{j,N} M_u^{a,N} M_v^{b,N}]|$ is zero if $j \notin \{a, b\}$ because the martingales $M^{j,N}, M^{a,N}$ and $M^{b,N}$ are orthogonal, and is bounded by $CN^{-1}t$ else by point (iv).

As a consequence,

$$\begin{aligned} |\mathbb{E}_\theta[U_r^{k,N} M_u^{a,N} M_v^{b,N}]| &\leq CN^{-1}t \sum_{n \geq 0} \Lambda^n (A_N^n(k, a) + A_N^n(k, b)) \\ &= CN^{-1}t(Q_N(k, a) + Q_N(k, b)). \end{aligned}$$

Since $k \neq a$ and $k \neq b$, this is bounded by $CN^{-2}t$ by (8).

For (vii), we assume e.g. that $r \leq s$ and $u \leq v$ and we recall that $k \neq a$. We have

$$\begin{aligned} &\text{Cov}_\theta (M_r^{k,N} M_s^{k,N}, M_u^{a,N} M_v^{a,N}) \\ &= \text{Cov}_\theta ((M_r^{k,N})^2, (M_u^{a,N})^2) + \text{Cov}_\theta (M_r^{k,N} (M_s^{k,N} - M_r^{k,N}), (M_u^{a,N})^2) \\ &\quad + \text{Cov}_\theta ((M_r^{k,N})^2, M_u^{a,N} (M_v^{a,N} - M_u^{a,N})) \\ &\quad + \text{Cov}_\theta (M_r^{k,N} (M_s^{k,N} - M_r^{k,N}), M_u^{a,N} (M_v^{a,N} - M_u^{a,N})) \\ &= I + J + K + L. \end{aligned}$$

First, $L = 0$. Indeed, assuming e.g. that $r \geq u$, we have

$$\begin{aligned} L &= \mathbb{E}_\theta[M_r^{k,N} (M_s^{k,N} - M_r^{k,N}) M_u^{a,N} (M_v^{a,N} - M_r^{a,N} + M_r^{a,N} - M_u^{a,N})] \\ &= \mathbb{E}_\theta[M_r^{k,N} M_u^{a,N} \mathbb{E}_\theta[(M_s^{k,N} - M_r^{k,N})(M_v^{a,N} - M_r^{a,N}) | \mathcal{F}_r]] \\ &\quad + \mathbb{E}_\theta[M_r^{k,N} M_u^{a,N} (M_r^{a,N} - M_u^{a,N}) \mathbb{E}_\theta[M_s^{k,N} - M_r^{k,N} | \mathcal{F}_r]] \end{aligned}$$

and in both terms, the conditional expectation vanishes. Next, we write as usual $(M_r^{k,N})^2 = 2 \int_0^r M_{\tau-}^{k,N} dM_\tau^{k,N} + Z_r^{k,N}$ and $(M_u^{a,N})^2 = 2 \int_0^u M_{\tau-}^{a,N} dM_\tau^{a,N} + Z_u^{a,N}$. By orthogonality of the martingales $\int_0^r M_{\tau-}^{k,N} dM_\tau^{k,N}$ and $\int_0^u M_{\tau-}^{a,N} dM_\tau^{a,N}$, we find

$$\begin{aligned} I &= \text{Cov}_\theta (Z_r^{k,N}, Z_u^{a,N}) + 2\text{Cov}_\theta \left(Z_r^{k,N}, \int_0^u M_{\tau-}^{a,N} dM_\tau^{a,N} \right) \\ &\quad + 2\text{Cov}_\theta \left(\int_0^r M_{\tau-}^{k,N} dM_\tau^{k,N}, Z_u^{a,N} \right). \end{aligned}$$

We deduce from points (i) and (iii), since $k \neq a$, that $|I| \leq C(N^{-1}t + N^{-1}t^{3/2}) \leq CN^{-1}t^{3/2}$. We now treat K . It vanishes if $u \geq r$, because $\mathbb{E}_\theta[M_v^{a,N} - M_u^{a,N} | \mathcal{F}_u] = 0$. We thus assume that $u < r$. We write as usual $(M_r^{k,N})^2 = (M_u^{k,N})^2 + 2 \int_u^r M_{\tau-}^{k,N} dM_\tau^{k,N} + Z_r^{k,N} - Z_u^{k,N}$ and

$$\begin{aligned} K &= \mathbb{E}_\theta[(M_u^{k,N})^2 M_u^{a,N} (M_v^{a,N} - M_u^{a,N})] \\ &\quad + 2\mathbb{E} \left[\left(\int_u^r M_{\tau-}^{k,N} dM_\tau^{k,N} \right) M_u^{a,N} (M_v^{a,N} - M_u^{a,N}) \right] \\ &\quad + \mathbb{E}_\theta[(Z_r^{k,N} - Z_u^{k,N}) M_u^{a,N} (M_v^{a,N} - M_u^{a,N})]. \end{aligned}$$

The first term vanishes (because $\mathbb{E}_\theta[M_v^{a,N} - M_u^{a,N} | \mathcal{F}_u] = 0$), as well as the second one (because $\mathbb{E}_\theta[(\int_u^r M_{\tau-}^{k,N} dM_\tau^{k,N})(M_v^{a,N} - M_u^{a,N}) | \mathcal{F}_u] = 0$ by orthogonality of the involved martingales). Consequently,

$$K = \mathbb{E}_\theta[(Z_r^{k,N} - Z_u^{k,N}) M_u^{a,N} (M_v^{a,N} - M_u^{a,N})]$$

$$= \mathbb{E}_\theta[(U_r^{k,N} - U_u^{k,N})M_u^{a,N}(M_v^{a,N} - M_u^{a,N})].$$

Using (4) and recalling that $\beta_n(u, r, x) = \varphi^{*n}(r - x) - \varphi^{*n}(u - x)$, we find

$$K = \sum_{n \geq 0} \int_0^r \beta_n(u, r, x) \sum_{j=1}^N A_N^n(k, j) \mathbb{E}_\theta[M_x^{j,N} M_u^{a,N} (M_v^{a,N} - M_u^{a,N})] dx.$$

But $|\mathbb{E}_\theta[M_x^{j,N} M_u^{a,N} (M_v^{a,N} - M_u^{a,N})]| \leq CN^{-1}t$ if $a \neq j$ by (iv), while Lemma 16-(iii) tells us that $|\mathbb{E}_\theta[M_x^{j,N} M_u^{a,N} (M_v^{a,N} - M_u^{a,N})]| \leq Ct^{3/2}$ if $a = j$. Thus

$$\begin{aligned} |K| &\leq C \sum_{n \geq 0} \Lambda^n \left[A_N^n(k, a) t^{3/2} + \sum_{j=1}^N A_N^n(k, j) N^{-1} t \right] \\ &\leq C \left[Q_N(k, a) t^{3/2} + N^{-1} \sum_{j=1}^N Q_N(k, j) t \right]. \end{aligned}$$

But $k \neq a$ implies that $Q_N(k, a) \leq CN^{-1}$ by (8), while $N^{-1} \sum_{j=1}^N Q_N(k, j) \leq CN^{-1} \|Q_N\|_\infty \leq CN^{-1}$. As a conclusion, $|K| \leq CN^{-1}(t^{3/2} + t) \leq CN^{-1}t^{3/2}$. Of course, J is treated similarly, and this completes the proof of point (vii).

Point (viii) is obvious: it suffices to use the Hölder inequality to find

$$\begin{aligned} &|\text{Cov}_\theta(M_r^{k,N} M_s^{l,N}, M_u^{a,N} M_v^{b,N})| \\ &\leq \mathbb{E}_\theta[(M_r^{k,N})^4]^{1/4} \mathbb{E}_\theta[(M_s^{l,N})^4]^{1/4} \mathbb{E}_\theta[(M_u^{a,N})^4]^{1/4} \mathbb{E}_\theta[(M_v^{b,N})^4]^{1/4}, \end{aligned}$$

which is bounded by Ct^2 by Lemma 16-(iii). □

We can now easily bound $\Delta_t^{N,31}$.

Lemma 23. Assume $H(q)$ for some $q \geq 1$. Then a.s., on Ω_N^1 , for $t \geq 1$,

$$\mathbb{E}_\theta[(\Delta_t^{N,31})^2] \leq Ct^{-1} \sum_{i=1}^N [\ell_N(i) - \bar{\ell}_N]^2.$$

Proof. We first note that

$$\Delta_t^{N,31} = 2\mu t^{-1} \left| \sum_{i=1}^N [U_{2t}^{i,N} - U_t^{i,N}] [\ell_N(i) - \bar{\ell}_N] \right|.$$

Since $U_{2t}^{i,N} - U_t^{i,N}$ is centered (its conditional expectation \mathbb{E}_θ vanishes),

$$\begin{aligned} \mathbb{E}_\theta[(\Delta_t^{N,31})^2] &= 4\mu^2 t^{-2} \sum_{i,j=1}^N [\ell_N(i) - \bar{\ell}_N] [\ell_N(j) - \bar{\ell}_N] \\ &\quad \times \text{Cov}_\theta(U_{2t}^{i,N} - U_t^{i,N}, U_{2t}^{j,N} - U_t^{j,N}). \end{aligned}$$

Using now Lemma 22-(i), we deduce that $|\text{Cov}_\theta(U_{2t}^{i,N} - U_t^{i,N}, U_{2t}^{j,N} - U_t^{j,N})| \leq Ct(\mathbf{1}_{\{i=j\}} + N^{-1})$ on Ω_N^1 . Using furthermore that $[\ell_N(i) - \bar{\ell}_N][\ell_N(j) - \bar{\ell}_N] \leq [\ell_N(i) - \bar{\ell}_N]^2 + [\ell_N(j) - \bar{\ell}_N]^2$ and a symmetry argument, we conclude that

$$\mathbb{E}_\theta[(\Delta_t^{N,31})^2] \leq Ct^{-1} \sum_{i,j=1}^N \left[\ell_N(i) - \bar{\ell}_N \right]^2 (\mathbf{1}_{\{i=j\}} + N^{-1}) = Ct^{-1} \sum_{i=1}^N \left[\ell_N(i) - \bar{\ell}_N \right]^2,$$

which was our goal. □

We can finally estimate $\Delta_t^{N,211}$.

Lemma 24. *Assume $H(q)$ for some $q \geq 1$. Then a.s., on Ω_N^1 , for $t \geq 1$, $\mathbb{E}_\theta[(\Delta_t^{N,211})^2] \leq CNt^{-2}$.*

Proof. We as usual work on Ω_N^1 . We note that $\mathbb{E}_\theta[(\Delta_t^{N,211})^2] = t^{-4} \sum_{i,j=1}^N a_{ij}$, where

$$a_{ij} = \text{Cov}_\theta((U_{2t}^{i,N} - U_t^{i,N})^2, (U_{2t}^{j,N} - U_t^{j,N})^2).$$

But recalling (4) and setting $\alpha_N(s, t, i, k) = \sum_{n \geq 0} A_N^n(i, k)[\varphi^{*n}(2t - s) - \varphi^{*n}(t - s)]$ for all $0 \leq s \leq 2t$ and $i, k \in \{1, \dots, N\}$,

$$U_{2t}^{i,N} - U_t^{i,N} = \int_0^{2t} \sum_{k=1}^N \alpha_N(s, t, i, k) M_s^{k,N} ds. \tag{9}$$

Concerning α_N , we will only use that, on Ω_N^1 ,

$$\int_0^{2t} |\alpha_N(s, t, i, k)| ds \leq 2 \sum_{n \geq 0} \Lambda^n A_N^n(i, k) = 2Q_N(i, k) \leq C(\mathbf{1}_{\{i=k\}} + N^{-1}), \tag{10}$$

the last inequality coming from (8). A direct computation starting from (9) shows that

$$a_{ij} = \sum_{k,l,a,b=1}^N \int_0^{2t} \int_0^{2t} \int_0^{2t} \int_0^{2t} \alpha_N(r, t, i, k) \alpha_N(s, t, i, l) \alpha_N(u, t, j, a) \alpha_N(v, t, j, b) \text{Cov}_\theta(M_r^{k,N} M_s^{l,N}, M_u^{a,N} M_v^{b,N}) dv du ds dr.$$

Let us denote by $\Gamma_{k,l,a,b}(t) = \sup_{r,s,u,v \in [0,2t]} |\text{Cov}_\theta(M_r^{k,N} M_s^{l,N}, M_u^{a,N} M_v^{b,N})|$. We can write, recalling (10),

$$\sum_{i,j=1}^N a_{ij} \leq C \sum_{i,j,k,l,a,b=1}^N (\mathbf{1}_{\{i=k\}} + N^{-1})(\mathbf{1}_{\{i=l\}} + N^{-1})(\mathbf{1}_{\{j=a\}} + N^{-1})(\mathbf{1}_{\{j=b\}} + N^{-1}) \Gamma_{k,l,a,b}(t).$$

Using some symmetry arguments, we find that $\sum_{i,j=1}^N a_{ij} \leq C[R_1 + \dots + R_6]$, where

$$R_1 = N^{-4} \sum_{i,j,k,l,a,b=1}^N \Gamma_{k,l,a,b}(t) = N^{-2} \sum_{k,l,a,b=1}^N \Gamma_{k,l,a,b}(t),$$

$$\begin{aligned}
 R_2 &= N^{-3} \sum_{i,j,k,l,a,b=1}^N \mathbf{1}_{\{i=k\}} \Gamma_{k,l,a,b}(t) = N^{-2} \sum_{k,l,a,b=1}^N \Gamma_{k,l,a,b}(t), \\
 R_3 &= N^{-2} \sum_{i,j,k,l,a,b=1}^N \mathbf{1}_{\{i=k\}} \mathbf{1}_{\{j=a\}} \Gamma_{k,l,a,b}(t) = N^{-2} \sum_{k,l,a,b=1}^N \Gamma_{k,l,a,b}(t), \\
 R_4 &= N^{-2} \sum_{i,j,k,l,a,b=1}^N \mathbf{1}_{\{i=k\}} \mathbf{1}_{\{i=l\}} \Gamma_{k,l,a,b}(t) = N^{-1} \sum_{k,a,b=1}^N \Gamma_{k,k,a,b}(t), \\
 R_5 &= N^{-1} \sum_{i,j,k,l,a,b=1}^N \mathbf{1}_{\{i=k\}} \mathbf{1}_{\{i=l\}} \mathbf{1}_{\{j=a\}} \Gamma_{k,l,a,b}(t) = N^{-1} \sum_{k,a,b=1}^N \Gamma_{k,k,a,b}(t), \\
 R_6 &= \sum_{i,j,k,l,a,b=1}^N \mathbf{1}_{\{i=k\}} \mathbf{1}_{\{i=l\}} \mathbf{1}_{\{j=a\}} \mathbf{1}_{\{j=b\}} \Gamma_{k,l,a,b}(t) = \sum_{k,a=1}^N \Gamma_{k,k,a,a}(t).
 \end{aligned}$$

Using Lemma 22-(v)-(viii), from which $\Gamma_{k,l,a,b}(t) \leq Ct^2 \mathbf{1}_{\{\#\{k,l,a,b\} < 4\}}$, we deduce that $R_1 = R_2 = R_3 \leq CNt^2$. Next we use Lemma 22-(vi)-(viii), that is $\Gamma_{k,k,a,b}(t) \leq C(\mathbf{1}_{\{\#\{k,a,b\}=3\}}N^{-2}t + \mathbf{1}_{\{\#\{k,a,b\} < 3\}}t^2)$, whence $R_4 = R_5 \leq Ct + CNt^2 \leq CNt^2$. Finally, we use Lemma 22-(vii)-(viii), i.e. $\Gamma_{k,k,a,a}(t) \leq C(\mathbf{1}_{\{\#\{k,a\}=2\}}N^{-1}t^{3/2} + \mathbf{1}_{\{\#\{k,a\}=1\}}t^2)$ and find that $R_6 \leq CNt^{3/2} + CNt^2 \leq CNt^2$. All in all, we have proved that $\sum_{i,j=1}^N a_{ij} \leq CNt^2$, which completes the proof. \square

We can finally give the

Proof of Proposition 19. It suffices to recall that $|\mathcal{V}_t^N - \mathcal{V}_\infty^N| \leq \Delta_t^{N,1} + \Delta_t^{N,211} + \Delta_t^{N,212} + \Delta_t^{N,213} + \Delta_t^{N,22} + \Delta_t^{N,23} + \Delta_t^{N,31} + \Delta_t^{N,32}$ and to use Lemmas 20, 21, 23 and 24: this gives, on Ω_N^1 ,

$$\mathbb{E}_\theta[|\mathcal{V}_t^N - \mathcal{V}_\infty^N|] \leq C \left(\frac{N}{t^{2q}} + \frac{1}{t} + \frac{N}{t^q} + \frac{N^{1/2}}{t^{3/2}} + \left[\sum_{i=1}^N [\ell_N(i) - \bar{\ell}_N]^2 \right]^{1/2} \frac{1}{t^{1/2}} + \frac{N^{1/2}}{t} \right).$$

Recalling that $t \geq 1$, the conclusion immediately follows. \square

4.6. Third estimator

We recall that, for $\Delta > 0$ such that $t/(2\Delta)$ is an integer, we have set $\mathcal{E}_t^N = (\bar{Z}_{2t}^N - \bar{Z}_t^N)/t$, $\mathcal{Z}_{\Delta,t}^N = (N/t) \sum_{a=t/\Delta+1}^{2t/\Delta} [\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \Delta \mathcal{E}_t^N]^2$ and $\mathcal{W}_{\Delta,t}^N = 2\mathcal{Z}_{2\Delta,t}^N - \mathcal{Z}_{\Delta,t}^N$. The matrices A_N and Q_N and the event Ω_N^1 were defined in Notation 12, as well as $\ell_N(i) = \sum_{j=1}^N Q_N(i,j)$ and $c_N(i) = \sum_{j=1}^N Q_N(j,i)$. We finally introduce $\mathcal{W}_{\infty,\infty}^N = \mu N^{-1} \sum_{i=1}^N \ell_N(i)(c_N(i))^2$. The aim of the subsection is to verify the following result.

Proposition 25. *Assume $H(q)$ for some $q \geq 3$. Then a.s., for $t \geq 4$ and $\Delta \in [1, t/4]$ such that $t/(2\Delta)$ is a positive integer,*

$$\mathbf{1}_{\Omega_N^1} \mathbb{E}_\theta \left[\left| \mathcal{W}_{\Delta,t}^N - \mathcal{W}_{\infty,\infty}^N \right| \right] \leq C \left(\sqrt{\frac{\Delta}{t}} + \frac{N}{\Delta^{(q+1)/2}} + \frac{t}{\Delta^{q/2+1}} \right).$$

Recall that we do not try to optimize the dependence in q . We first write

$$|\mathcal{W}_{\Delta,t}^N - \mathcal{W}_{\infty,\infty}^N| \leq D_{\Delta,t}^{N,1} + 2D_{2\Delta,t}^{N,1} + D_{\Delta,t}^{N,2} + 2D_{2\Delta,t}^{N,2} + D_{\Delta,t}^{N,3} + 2D_{2\Delta,t}^{N,3} + D_{\Delta,t}^{N,4},$$

where

$$D_{\Delta,t}^{N,1} = \frac{N}{t} \left| \sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \Delta \mathcal{E}_t^N \right]^2 - \sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \Delta \mu \bar{\ell}_N \right]^2 \right|,$$

$$D_{\Delta,t}^{N,2} = \frac{N}{t} \left| \sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \Delta \mu \bar{\ell}_N \right]^2 - \sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right]^2 \right|,$$

$$D_{\Delta,t}^{N,3} = \frac{N}{t} \left| \sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right]^2 - \mathbb{E}_\theta \left[\sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right]^2 \right] \right|,$$

$$D_{\Delta,t}^{N,4} = \left| \frac{2N}{t} \mathbb{E}_\theta \left[\sum_{a=t/(2\Delta)+1}^{t/\Delta} \left[\bar{Z}_{2a\Delta}^N - \bar{Z}_{2(a-1)\Delta}^N - \mathbb{E}_\theta[\bar{Z}_{2a\Delta}^N - \bar{Z}_{2(a-1)\Delta}^N] \right]^2 \right] - \frac{N}{t} \mathbb{E}_\theta \left[\sum_{a=t/\Delta+1}^{2t/\Delta} \left[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right]^2 \right] - \mathcal{W}_{\infty,\infty}^N \right|.$$

We treat these four terms one by one.

Lemma 26. *Assume $H(q)$ for some $q \geq 1$. Then a.s. on Ω_N^1 , for $1 \leq \Delta \leq t$, $\mathbb{E}_\theta[D_{\Delta,t}^{N,1}] \leq C\Delta[t^{-1} + Nt^{-2q}]$.*

Proof. Using that $(\Delta/t) \sum_{a=t/\Delta+1}^{2t/\Delta} (\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N) = \Delta \mathcal{E}_t^N$, we find that

$$D_{\Delta,t}^{N,1} = \frac{N}{t} \frac{t}{\Delta} (\Delta \mu \bar{\ell}_N - \Delta \mathcal{E}_t^N)^2 = N \Delta (\mu \bar{\ell}_N - \mathcal{E}_t^N)^2,$$

whence, on Ω_N^1 , $\mathbb{E}_\theta[D_{\Delta,t}^{N,1}] \leq CN\Delta(t^{-2q} + (Nt)^{-1}) \leq C\Delta(Nt^{-2q} + t^{-1})$, see Proposition 17. \square

The second term is also easy.

Lemma 27. *Assume $H(q)$ for some $q \geq 1$. Then on Ω_N^1 , for $1 \leq \Delta \leq t$, $\mathbb{E}_\theta[D_{\Delta,t}^{N,2}] \leq CNt^{1-q}$.*

Proof. Using that $|(A - x)^2 - (A - y)^2| \leq |x - y|(|x| + |y| + 2|A|)$,

$$D_{\Delta,t}^{N,2} \leq \frac{N}{t} \sum_{a=t/\Delta+1}^{2t/\Delta} \left| \Delta\mu\bar{\ell}_N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right| \times \left[\Delta\mu\bar{\ell}_N + \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] + 2(Z_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N) \right],$$

whence

$$\mathbb{E}_\theta[D_{\Delta,t}^{N,2}] \leq \frac{N}{t} \sum_{a=t/\Delta+1}^{2t/\Delta} \left| \Delta\mu\bar{\ell}_N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right| \left[\Delta\mu\bar{\ell}_N + 3\mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right].$$

But we deduce from Lemma 16-(ii) with $r = 1$ that, since $(a - 1)\Delta \geq t$,

$$\left| \Delta\mu\bar{\ell}_N - \mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \right| \leq Ct^{1-q},$$

whence also $\mathbb{E}_\theta[\bar{Z}_{a\Delta}^N - \bar{Z}_{(a-1)\Delta}^N] \leq \Delta\mu\bar{\ell}_N + Ct^{1-q} \leq \Delta\mu\bar{\ell}_N + C$. We conclude that

$$\mathbb{E}_\theta[D_{\Delta,t}^{N,2}] \leq C \frac{N}{t} \sum_{a=t/\Delta+1}^{2t/\Delta} t^{1-q} [4\Delta\mu\bar{\ell}_N + C].$$

Since $\bar{\ell}_N$ is bounded on Ω_N^1 and since $\Delta \geq 1 \geq t^{1-q}$, we find $\mathbb{E}_\theta[D_{\Delta,t}^{N,2}] \leq C(N/t)(t/\Delta)t^{1-q}\Delta \leq CNt^{1-q}$. \square

To treat $D_{\Delta,t}^{N,4}$, we need the following lemma.

Lemma 28. *Assume $H(q)$ for some $q \geq 1$. Almost surely on Ω_N^1 , for all $1 \leq \Delta \leq x/2$,*

$$\text{Var}_\theta(\bar{U}_{x+\Delta}^N - \bar{U}_x^N) = \frac{\Delta}{N} \mathcal{W}_{\infty,\infty}^N - \mathcal{X}^N + r_N(x, \Delta),$$

where \mathcal{X}_N is a $\sigma((\theta_{ij})_{i,j=1,\dots,N})$ -measurable finite random variable and where r_N satisfies, for some deterministic constant C , the inequality $|r_N(x, \Delta)| \leq Cx\Delta^{-q}N^{-1}$.

Proof. We set $V_{x,\Delta}^N = \text{Var}_\theta(\bar{U}_{x+\Delta}^N - \bar{U}_x^N)$.

Step 1. Recalling (4) and setting $\beta_n(x, x + \Delta, s) = \varphi^{*n}(x + \Delta - s) - \varphi^{*n}(x - s)$ as in Lemma 15, we get

$$\bar{U}_{x+\Delta}^N - \bar{U}_x^N = \sum_{n \geq 0} \int_0^{x+\Delta} \beta_n(x, x + \Delta, s) N^{-1} \sum_{i,j=1}^N A_N^n(i, j) M_s^{j,N} ds.$$

Hence

$$V_{x,\Delta}^N = \sum_{m,n \geq 0} \int_0^{x+\Delta} \int_0^{x+\Delta} \beta_m(x, x + \Delta, r) \beta_n(x, x + \Delta, s) N^{-2}$$

$$\sum_{i,j,k,l=1}^N A_N^m(i,j)A_N^n(k,l)\text{Cov}_\theta(M_r^{j,N}, M_s^{l,N})drds.$$

Using Remark 10, we find

$$V_{x,\Delta}^N = \sum_{m,n \geq 0} \int_0^{x+\Delta} \int_0^{x+\Delta} \beta_m(x, x + \Delta, r)\beta_n(x, x + \Delta, s) \\ \times N^{-2} \sum_{i,j,k=1}^N A_N^m(i,j)A_N^n(k,j)\mathbb{E}_\theta[Z_{r \wedge s}^{j,N}]drds.$$

Step 2. Here we show that $\mathbb{E}_\theta[Z_s^{j,N}] = \mu \ell_N(j)s - X_j^N + R_j^N(s)$, with, for some constant C , for all $j = 1, \dots, N$,

$$0 \leq X_j^N \leq C \quad \text{and} \quad |R_j^N(s)| \leq C(s^{1-q} \wedge 1).$$

By (3), we have $\mathbb{E}_\theta[Z_s^{j,N}] = \mu \sum_{n \geq 0} (\int_0^s r \varphi^{*n}(s-r)dr) \sum_{l=1}^N A_N^n(j,l)$, whence by Lemma 15-(i),

$$\mathbb{E}_\theta[Z_s^{j,N}] = \mu \sum_{n \geq 0} (\Lambda^n s - n\Lambda^n \kappa + \varepsilon_n(s)) \sum_{l=1}^N A_N^n(j,l) = \mu \ell_N(j)s - X_j^N + R_j^N(s).$$

We used that $\sum_{n \geq 0} \Lambda^n \sum_{l=1}^N A_N^n(j,l) = \sum_{l=1}^N Q_N(j,l) = \ell_N(j)$ and we set $X_j^N = \mu \kappa \sum_{n \geq 0} n\Lambda^n \sum_{l=1}^N A_N^n(j,l)$ and $R_j^N(s) = \mu \sum_{n \geq 0} \varepsilon_n(s) \sum_{l=1}^N A_N^n(j,l)$. We obviously have $0 \leq X_j^N \leq \mu \kappa \sum_{n \geq 0} n\Lambda^n \|A_N\|_\infty^n \leq C$ on Ω_N^1 and, since $\varepsilon_n(s) \leq Cn^q \Lambda^n (s^{1-q} \wedge 1)$ by Lemma 15-(i), still on Ω_N^1 ,

$$|R_j^N(s)| \leq C(s^{1-q} \wedge 1) \sum_{n \geq 0} n^q \Lambda^n \|A_N\|_\infty^n \leq C(s^{1-q} \wedge 1).$$

Step 3. Gathering Steps 1 and 2, we now write $V_{x,\Delta}^N = I - J + K$, where

$$I = \sum_{m,n \geq 0} \int_0^{x+\Delta} \int_0^{x+\Delta} \beta_m(x, x + \Delta, r)\beta_n(x, x + \Delta, s) \\ \times N^{-2} \sum_{i,j,k=1}^N A_N^m(i,j)A_N^n(k,j)\mu \ell_N(j)(r \wedge s)drds, \\ J = \sum_{m,n \geq 0} \int_0^{x+\Delta} \int_0^{x+\Delta} \beta_m(x, x + \Delta, r)\beta_n(x, x + \Delta, s) \\ \times N^{-2} \sum_{i,j,k=1}^N A_N^m(i,j)A_N^n(k,j)X_j^N drds, \\ K = \sum_{m,n \geq 0} \int_0^{x+\Delta} \int_0^{x+\Delta} \beta_m(x, x + \Delta, r)\beta_n(x, x + \Delta, s)$$

$$\times N^{-2} \sum_{i,j,k=1}^N A_N^m(i,j)A_N^n(k,j)R_j^N(r \wedge s)drds.$$

Step 4. Here we check that $|J| \leq Cx^{-2q}N^{-1}$ on Ω_N^1 . Since $|\int_0^{x+\Delta} \beta_m(x, x + \Delta, r)dr| \leq Cn^q\Lambda^n x^{-q}$ by Lemma 15-(ii) and since X_j^N is bounded by Step 2 (and does not depend on time),

$$\begin{aligned} |J| &\leq C \sum_{m,n \geq 0} m^q n^q \Lambda^{m+n} x^{-2q} N^{-2} \sum_{i,j,k=1}^N A_N^m(i,j)A_N^n(k,j) \\ &\leq Cx^{-2q}N^{-1} \sum_{m,n \geq 0} m^q n^q \Lambda^{m+n} \|A_N\|_1^{m+n}. \end{aligned}$$

The conclusion follows, since $\Lambda \|A_N\|_1 \leq a < 1$ on Ω_N^1 .

Step 5. We next check that $|K| \leq Cx\Delta^{-q}N^{-1}$ on Ω_N^1 . Using the bound on R_j^N (see Step 2), we start from

$$\begin{aligned} |K| &\leq C \sum_{m,n \geq 0} \int_0^{x+\Delta} \int_0^{x+\Delta} |\beta_m(x, x + \Delta, r)| |\beta_n(x, x + \Delta, s)| \\ &\quad \times N^{-1} \|A_N\|_1^{m+n} [(r \wedge s)^{1-q} \wedge 1] drds \\ &\leq C(K_1 + K_2), \end{aligned}$$

where, using that $x - \Delta \geq x/2$ (whence $(r \wedge s)^{1-q} \leq Cx^{1-q}$ if $r \wedge s \geq x - \Delta$) and a symmetry argument,

$$\begin{aligned} K_1 &= x^{1-q} \sum_{m,n \geq 0} \int_{x-\Delta}^{x+\Delta} \int_{x-\Delta}^{x+\Delta} |\beta_m(x, x + \Delta, r)| |\beta_n(x, x + \Delta, s)| \\ &\quad \times N^{-1} \|A_N\|_1^{m+n} drds, \\ K_2 &= \sum_{m,n \geq 0} \int_0^{x-\Delta} \int_0^{x+\Delta} |\beta_m(x, x + \Delta, r)| |\beta_n(x, x + \Delta, s)| \\ &\quad N^{-1} \|A_N\|_1^{m+n} drds. \end{aligned}$$

First, on Ω_N^1 ,

$$K_1 \leq Cx^{1-q} \sum_{m,n \geq 0} \Lambda^{m+n} N^{-1} \|A_N\|_1^{m+n} \leq CN^{-1}x^{1-q} \leq Cx\Delta^{-q}N^{-1}$$

since $x \geq \Delta$. Next, using that $\int_0^{x-\Delta} |\beta_m(x, x + \Delta, r)|dr \leq Cm^q\Lambda^m\Delta^{-q}$ by Lemma 15-(ii) and that $\int_0^{x+\Delta} |\beta_n(x, x + \Delta, s)|ds \leq 2\Lambda^n$, still on Ω_N^1 ,

$$K_2 \leq C\Delta^{-q} \sum_{m,n \geq 0} m^q \Lambda^{m+n} N^{-1} \|A_N\|_1^{m+n} \leq C\Delta^{-q}N^{-1} \leq Cx\Delta^{-q}N^{-1},$$

since $x \geq 1$ by assumption.

Step 6. Finally recall that $\gamma_{m,n}(x, x + \Delta) = \int_0^{x+\Delta} \int_0^{x+\Delta} (s \wedge u) \beta_m(x, x + \Delta, s) \beta_n(x, x + \Delta, u) ds du = \Delta \Lambda^{m+n} - \kappa_{m,n} \Lambda^{m+n} + \varepsilon_{m,n}(x, x + \Delta)$ with the notation of Lemma 15-(iii). We thus may write

$$I = \mu \sum_{m,n \geq 0} \gamma_{m,n}(x, x + \Delta) N^{-2} \sum_{i,j,k=1}^N A_N^m(i, j) A_N^n(k, j) \ell_N(j) = I_1 - I_2 + I_3,$$

where

$$\begin{aligned} I_1 &= \mu \Delta \sum_{m,n \geq 0} \Lambda^{m+n} N^{-2} \sum_{i,j,k=1}^N A_N^m(i, j) A_N^n(k, j) \ell_N(j), \\ I_2 &= \mu \sum_{m,n \geq 0} \kappa_{m,n} \Lambda^{m+n} N^{-2} \sum_{i,j,k=1}^N A_N^m(i, j) A_N^n(k, j) \ell_N(j), \\ I_3 &= \mu \sum_{m,n \geq 0} \varepsilon_{m,n}(x, x + \Delta) N^{-2} \sum_{i,j,k=1}^N A_N^m(i, j) A_N^n(k, j) \ell_N(j). \end{aligned}$$

First, we clearly have

$$I_1 = \mu \Delta N^{-2} \sum_{i,j,k=1}^N Q_N(i, j) Q_N(k, j) \ell_N(j) = \mu \Delta N^{-2} \sum_{j=1}^N (c_N(j))^2 \ell_N(j),$$

which equals $\Delta N^{-1} \mathcal{W}_{\infty, \infty}^N$.

We next simply set $\mathcal{X}^N = I_2$, which is clearly $\sigma((\theta_{i,j})_{i,j=1,\dots,N})$ -measurable and well-defined on Ω_N^1 . Finally, since $\varepsilon_{m,n}(x, x + \Delta) \leq C(m+n)^q \Lambda^{m+n} x \Delta^{-q}$ by Lemma 15-(iii), since ℓ_N is bounded on Ω_N^1 and since, as already seen, $\sum_{i,j,k=1}^N A_N^m(i, j) A_N^n(k, j) \leq N \|A_N\|_1^{m+n}$,

$$|I_3| \leq C x \Delta^{-q} N^{-1} \sum_{m,n \geq 0} (m+n)^q \Lambda^{m+n} \|A_N\|_1^{m+n} \leq C x \Delta^{-q} N^{-1}.$$

All this implies that $|I - \Delta N^{-1} \mathcal{W}_{\infty, \infty}^N + \mathcal{X}^N| \leq C x \Delta^{-q} N^{-1}$. Since $V_{x, \Delta}^N = I - J + K$ by Step 3 and since we have seen in Steps 4 and 5 that $|J| \leq C x^{-2q} N^{-1} \leq C x \Delta^{-q} N^{-1}$ and $|K| \leq C x \Delta^{-q} N^{-1}$, we conclude that, on Ω_N^1 , $|V_{x, \Delta}^N - \Delta N^{-1} \mathcal{W}_{\infty, \infty}^N + \mathcal{X}^N| \leq C x \Delta^{-q} N^{-1}$ as desired. \square

We can now study the term $D_{\Delta, t}^{N,4}$.

Lemma 29. *Assume $H(q)$ for some $q \geq 1$. Then a.s. on Ω_N^1 , for $1 \leq \Delta \leq t/4$, $D_{\Delta, t}^{N,4} \leq C t \Delta^{-1-q}$.*

Proof. We clearly have

$$D_{\Delta, t}^{N,4} = \left| \frac{2N}{t} \sum_{a=t/(2\Delta)+1}^{t/\Delta} \text{Var}_\theta(\bar{U}_{2a\Delta}^N - \bar{U}_{2(a-1)\Delta}^N) \right|$$

$$-\frac{N}{t} \sum_{a=t/\Delta+1}^{2t/\Delta} \left| \text{Var}_\theta (\bar{U}_{a\Delta}^N - \bar{U}_{(a-1)\Delta}^N) - \mathcal{W}_{\infty,\infty}^N \right|.$$

Using Lemma 28, (observe that for $a \in \{t/(2\Delta)+1, \dots, t/\Delta\}$, $x = 2(a-1)\Delta \geq t$ satisfies $2\Delta \leq x/2$ and, for $a \in \{t/\Delta + 1, \dots, 2t/\Delta\}$, $x = (a-1)\Delta \geq t$ satisfies $\Delta \leq x/2$), we get

$$D_{\Delta,t}^{N,4} = \left| \frac{2N}{t} \sum_{a=t/(2\Delta)+1}^{t/\Delta} \left[\frac{2\Delta}{N} \mathcal{W}_{\infty,\infty}^N - \mathcal{X}^N + r_N(2(a-1)\Delta, 2\Delta) \right] - \frac{N}{t} \sum_{a=t/\Delta+1}^{2t/\Delta} \left[\frac{\Delta}{N} \mathcal{W}_{\infty,\infty}^N - \mathcal{X}^N + r_N((a-1)\Delta, \Delta) \right] - \mathcal{W}_{\infty,\infty}^N \right|.$$

This rewrites

$$D_{\Delta,t}^{N,4} = \left| \frac{2N}{t} \sum_{a=t/(2\Delta)+1}^{t/\Delta} r_N(2(a-1)\Delta, 2\Delta) - \frac{N}{t} \sum_{a=t/\Delta+1}^{2t/\Delta} r_N((a-1)\Delta, \Delta) \right|.$$

Since $r_N(x, \Delta) \leq Cx\Delta^{-q}N^{-1}$, we find that $D_{\Delta,t}^{N,4} \leq C(N/t)(t/\Delta)(t\Delta^{-q}N^{-1}) = Ct\Delta^{-1-q}$. □

The following tedious lemma will allow us to treat the last term $D_{\Delta,t}^{N,3}$.

Lemma 30. Assume $H(q)$ for some $q \geq 1$. On Ω_N^1 , for all $t, x, \Delta \geq 1$ with $t/2 \leq x - \Delta \leq x + \Delta \leq 2t$,

$$\text{Var}_\theta ((\bar{U}_{x+\Delta}^N - \bar{U}_x^N)^2) \leq C \left(\frac{\Delta^2}{N^2} + \frac{t^2}{N^2\Delta^{4q}} \right)$$

and, if $t/2 \leq y - \Delta \leq y + \Delta \leq x - 2\Delta \leq x + \Delta \leq 2t$,

$$\text{Cov}_\theta ((\bar{U}_{x+\Delta}^N - \bar{U}_x^N)^2, (\bar{U}_{y+\Delta}^N - \bar{U}_y^N)^2) \leq C \left(\frac{t^{1/2}}{N\Delta^{q-1}} + \frac{t^2}{N^2\Delta^{4q}} + \frac{t^{1/2}}{N^2\Delta^{q-3/2}} \right).$$

Proof. We divide the proof in several steps. We work on Ω_N^1 .

Step 1. For $i = 1, \dots, N$ and $z \in [x, x + \Delta]$, we can write, recalling (4) and that $\beta_n(x, z, r) = \varphi^{*n}(z-r) - \varphi^{*n}(x-r)$,

$$U_z^{i,N} - U_x^{i,N} = \sum_{n \geq 0} \int_0^z \beta_n(x, z, r) \sum_{j=1}^N A_N^n(i, j) M_r^{j,N} dr = \Gamma_{x,z}^{i,N} + X_{x,z}^{i,N},$$

where

$$\Gamma_{x,z}^{i,N} = \sum_{n \geq 0} \int_{x-\Delta}^z \beta_n(x, z, r) \sum_{j=1}^N A_N^n(i, j) (M_r^{j,N} - M_{x-\Delta}^{j,N}) dr,$$

$$X_{x,z}^{i,N} = \sum_{n \geq 0} \left(\int_{x-\Delta}^z \beta_n(x, z, r) dr \right) \sum_{j=1}^N A_N^n(i, j) M_{x-\Delta}^{j,N} \\ + \sum_{n \geq 0} \int_0^{x-\Delta} \beta_n(x, z, r) \sum_{j=1}^N A_N^n(i, j) M_r^{j,N} dr,$$

and we set as usual $\bar{\Gamma}_{x,z}^N = N^{-1} \sum_{i=1}^N \Gamma_{x,z}^{i,N}$ and $\bar{X}_{x,z}^N = N^{-1} \sum_{i=1}^N X_{x,z}^{i,N}$.

Step 2. We now show that, on Ω_N^1 , for $z \in [x, x + \Delta]$,

$$\sup_{i=1, \dots, N} \mathbb{E}_\theta[(X_{x,z}^{i,N})^4] \leq Ct^2 \Delta^{-4q} \quad \text{and} \quad \mathbb{E}_\theta[(\bar{X}_{x,z}^N)^4] \leq Ct^2 N^{-2} \Delta^{-4q}.$$

Using that $|\int_{x-\Delta}^z \beta_n(x, z, r) dr| + \int_0^{x-\Delta} |\beta_n(x, z, r)| dr \leq Cn^q \Lambda^n \Delta^{-q}$ by Lemma 15-(ii)

$$|X_{x,z}^{i,N}| \leq C \sum_{n \geq 0} n^q \Lambda^n \Delta^{-q} \sum_{j=1}^N A_N^n(i, j) \sup_{[0, 2t]} |M_r^{j,N}|.$$

But we now from Lemma 16-(iii) that $\sup_{j=1, \dots, N} \mathbb{E}_\theta[\sup_{[0, 2t]} |M_r^{j,N}|^4] \leq Ct^2$. We thus deduce from the Minkowski inequality that, still on Ω_N^1 ,

$$\mathbb{E}_\theta[(X_{x,z}^{i,N})^4]^{1/4} \leq Ct^{1/2} \Delta^{-q} \sum_{n \geq 0} n^q \Lambda^n \sum_{j=1}^N A_N^n(i, j) \\ \leq Ct^{1/2} \Delta^{-q} \sum_{n \geq 0} n^q \Lambda^n \| \|A_N\| \|_\infty^n \leq Ct^{1/2} \Delta^{-q}.$$

We next observe that

$$\bar{X}_{x,z}^N = \sum_{n \geq 0} \left(\int_{x-\Delta}^z \beta_n(x, z, r) dr \right) O_{x-\Delta}^{N,n} + \sum_{n \geq 0} \int_0^{x-\Delta} \beta_n(x, z, r) O_r^{N,n} dr,$$

where the martingale

$$O_r^{N,n} = N^{-1} \sum_{i,j=1}^N A_N^n(i, j) M_r^{j,N}$$

has for quadratic variation $[O^{N,n}, O^{N,n}]_r = N^{-2} \sum_{j=1}^N (\sum_{i=1}^N A_N^n(i, j))^2 Z_r^{j,N} \leq N^{-1} \| \|A_N\| \|_1^{2n} \bar{Z}_r^N$ by Remark 10. By Lemma 16-(iii), we conclude that, on Ω_N^1 ,

$$\mathbb{E}_\theta \left[\sup_{[0, 2t]} (O_r^{N,n})^4 \right] \leq CN^{-2} \| \|A_N\| \|_1^{4n} \mathbb{E}_\theta[(\bar{Z}_{2t}^N)^2] \leq CN^{-2} \| \|A_N\| \|_1^{4n} t^2.$$

Using again that $|\int_{x-\Delta}^z \beta_n(x, z, r) dr| + \int_0^{x-\Delta} |\beta_n(x, z, r)| dr \leq Cn^q \Lambda^n \Delta^{-q}$ by Lemma 15-(ii),

$$|\bar{X}_{x,z}^N| \leq C \sum_{n \geq 0} n^q \Lambda^n \Delta^{-q} \sup_{[0, 2t]} |O_r^{N,n}|.$$

Thus, we infer from the Minkowski inequality that, still on Ω_N^1 ,

$$\mathbb{E}[(\bar{X}_{x,z}^N)^4]^{1/4} \leq C \sum_{n \geq 0} n^q \Lambda^n \Delta^{-q} N^{-1/2} \| \|A_N\| \|_1^n t^{1/2} \leq C \Delta^{-q} N^{-1/2} t^{1/2}.$$

Step 3. We next check that $\mathbb{E}_\theta[(\bar{\Gamma}_{x,z}^N)^4] \leq C\Delta^2 N^{-2}$ for any $z \in [x, x + \Delta]$, on Ω_1^N . Using the same martingale $O^{N,n}$ as in Step 2,

$$\bar{\Gamma}_{x,z}^N = \sum_{n \geq 0} \int_{x-\Delta}^z \beta_n(x, z, r) [O_r^{N,n} - O_{x-\Delta}^{N,n}] dr.$$

Recalling that $[O_r^{N,n}, O_{x-\Delta}^{N,n}]_r = N^{-2} \sum_{j=1}^N (\sum_{i=1}^N A_N^n(i, j))^2 Z_r^{j,N}$ with moreover $\sum_{i=1}^N A_N^n(i, j) \leq \|A_N\|_1^n$,

$$\begin{aligned} \mathbb{E}_\theta \left[\sup_{[x-\Delta, z]} (O_r^{N,n} - O_{x-\Delta}^{N,n})^4 \right] &\leq CN^{-4} \|A_N\|_1^{4n} \mathbb{E}_\theta \left[\left(\sum_{j=1}^N (Z_z^{j,N} - Z_{x-\Delta}^{j,N}) \right)^2 \right] \\ &= CN^{-2} \|A_N\|_1^{4n} \mathbb{E}_\theta \left[\left(\bar{Z}_z^N - \bar{Z}_{x-\Delta}^N \right)^2 \right]. \end{aligned}$$

We conclude from Lemma 16-(iii) that (recall that $z \in [x, x + \Delta]$)

$$\mathbb{E}_\theta \left[\sup_{[x-\Delta, z]} (O_r^{N,n} - O_{x-\Delta}^{N,n})^4 \right] \leq C\Delta^2 N^{-2} \|A_N\|_1^{4n}.$$

Using that $\int_{x-\Delta}^z |\beta_n(x, z, r)| dr \leq 2\Lambda^n$ and the Minkowski inequality,

$$\mathbb{E}[(\bar{\Gamma}_{x,z}^N)^4]^{1/4} \leq C \sum_{n \geq 0} \Lambda^n \Delta^{1/2} N^{-1/2} \|A_N\|_1^n \leq C\Delta^{1/2} N^{-1/2}.$$

Step 4. Recalling Step 1, $(\bar{U}_{x+\Delta}^N - \bar{U}_x^N)^4 = (\bar{\Gamma}_{x,x+\Delta}^N + \bar{X}_{x,x+\Delta}^N)^4 \leq 8(\bar{\Gamma}_{x,x+\Delta}^N)^4 + 8(\bar{X}_{x,x+\Delta}^N)^4$. We deduce from Steps 2 and 3 that $\text{Var}_\theta((\bar{U}_{x+\Delta}^N - \bar{U}_x^N)^2) \leq C(\Delta^2 N^{-2} + t^2 N^{-2} \Delta^{-4q})$.

Step 5. Here we show that

$$\begin{aligned} &\left| \text{Cov}_\theta((\bar{U}_{x+\Delta}^N - \bar{U}_x^N)^2, (\bar{U}_{y+\Delta}^N - \bar{U}_y^N)^2) \right| \\ &\leq \left| \text{Cov}_\theta((\bar{\Gamma}_{x,x+\Delta}^N)^2, (\bar{\Gamma}_{y,y+\Delta}^N)^2) \right| + \frac{C}{N^2} \left(\frac{t^2}{\Delta^{4q}} + \frac{t^{1/2}}{\Delta^{q-3/2}} \right). \end{aligned}$$

It suffices to write $(\bar{U}_{x+\Delta}^N - \bar{U}_x^N)^2 = (\bar{\Gamma}_{x,x+\Delta}^N)^2 + (\bar{X}_{x,x+\Delta}^N)^2 + 2\bar{\Gamma}_{x,x+\Delta}^N \bar{X}_{x,x+\Delta}^N$, the same formula with y instead of x , and to use the bilinearity of the covariance: we have the term $\text{Cov}_\theta((\bar{\Gamma}_{x,x+\Delta}^N)^2, (\bar{\Gamma}_{y,y+\Delta}^N)^2)$, and the other ones are bounded by

$$\begin{aligned} &\mathbb{E}_\theta \left[(\bar{\Gamma}_{x,x+\Delta}^N)^2 (\bar{X}_{y,y+\Delta}^N)^2 + 2(\bar{\Gamma}_{x,x+\Delta}^N)^2 |\bar{\Gamma}_{y,y+\Delta}^N \bar{X}_{y,y+\Delta}^N| + (\bar{X}_{x,x+\Delta}^N)^2 (\bar{\Gamma}_{y,y+\Delta}^N)^2 \right. \\ &\quad + (\bar{X}_{x,x+\Delta}^N)^2 (\bar{X}_{y,y+\Delta}^N)^2 + 2(\bar{X}_{x,x+\Delta}^N)^2 |\bar{\Gamma}_{y,y+\Delta}^N \bar{X}_{y,y+\Delta}^N| \\ &\quad + 2|\bar{\Gamma}_{x,x+\Delta}^N \bar{X}_{x,x+\Delta}^N| (\bar{\Gamma}_{y,y+\Delta}^N)^2 + 2|\bar{\Gamma}_{x,x+\Delta}^N \bar{X}_{x,x+\Delta}^N| (\bar{X}_{y,y+\Delta}^N)^2 \\ &\quad \left. + 4|\bar{\Gamma}_{x,x+\Delta}^N \bar{X}_{x,x+\Delta}^N \bar{\Gamma}_{y,y+\Delta}^N \bar{X}_{y,y+\Delta}^N| \right]. \end{aligned}$$

We bound all these terms, using only the Hölder inequality and recalling that $\mathbb{E}[(\bar{\Gamma}_{x,x+z}^N)^4] \leq C\Delta^2 N^{-2}$ and $\mathbb{E}[(\bar{X}_{x,x+z}^N)^4] \leq Ct^2 N^{-2} \Delta^{-4q}$ and that the same bounds hold with y instead of x . We finally remove a few terms using the inequality $a + a^{3/4}b^{1/4} + a^{1/2}b^{1/2} + a^{1/4}b^{3/4} \leq 4(a + a^{1/4}b^{3/4})$ with $a = t^2 N^{-2} \Delta^{-4q}$ and $b = \Delta^2 N^{-2}$.

Step 6. Recall that $y + \Delta \leq x - 2\Delta$. We check here that for any $r, s \in [x - \Delta, x + \Delta]$, any $u, v \in [y - \Delta, y + \Delta]$, any $i, j, k, l \in \{1, \dots, N\}$,

$$\begin{aligned} & \left| \text{Cov}_\theta \left((M_r^{i,N} - M_{x-\Delta}^{i,N})(M_s^{j,N} - M_{x-\Delta}^{j,N}), (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right) \right| \\ & \leq C \mathbf{1}_{\{i=j\}} t^{1/2} \Delta^{1-q}. \end{aligned}$$

If $i \neq j$, $\mathbb{E}_\theta[(M_r^{i,N} - M_{x-\Delta}^{i,N})(M_s^{j,N} - M_{x-\Delta}^{j,N}) | \mathcal{F}_{x-\Delta}] = 0$ so that the covariance vanishes since $u, v \leq y + \Delta \leq x - \Delta$. We next assume that $i = j$ and w.l.o.g. that $r \leq s$. Conditioning with respect to \mathcal{F}_r , we easily find, since $u, v \leq x - \Delta \leq r$,

$$\begin{aligned} K & := \text{Cov}_\theta \left((M_r^{i,N} - M_{x-\Delta}^{i,N})(M_s^{i,N} - M_{x-\Delta}^{i,N}), (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right) \\ & = \text{Cov}_\theta \left((M_r^{i,N} - M_{x-\Delta}^{i,N})^2, (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right). \end{aligned}$$

We write as usual $(M_r^{i,N} - M_{x-\Delta}^{i,N})^2 = 2 \int_{x-\Delta}^r M_{\tau-}^{i,N} dM_\tau^{i,N} + Z_r^{i,N} - Z_{x-\Delta}^{i,N}$, because $[M^{i,N}, M^{i,N}]_\tau = Z_\tau^{i,N}$ by Remark 10. Since $\mathbb{E}[\int_{x-\Delta}^r M_{\tau-}^{i,N} dM_\tau^{i,N} | \mathcal{F}_{x-\Delta}] = 0$ and since $u, v \leq x - \Delta$, we find that

$$\begin{aligned} K & = \text{Cov}_\theta \left(Z_r^{i,N} - Z_{x-\Delta}^{i,N}, (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right) \\ & = \text{Cov}_\theta \left(U_r^{i,N} - U_{x-\Delta}^{i,N}, (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right) \\ & = \text{Cov}_\theta \left(\Gamma_{x-\Delta,r}^{i,N} + X_{x-\Delta,r}^{i,N}, (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right) \end{aligned}$$

with the notation of Step 1. But $\Gamma_{x-\Delta,r}^{i,N}$ involves only increments of martingales of the form $M_\tau^{j,N} - M_{x-2\Delta}^{j,N}$, of which the conditional expectation knowing $\mathcal{F}_{x-2\Delta}$ vanishes. Since now $u, v \leq y + \Delta \leq x - 2\Delta$, we deduce that

$$K = \text{Cov}_\theta \left(X_{x-\Delta,r}^{i,N}, (M_u^{k,N} - M_{y-\Delta}^{k,N})(M_v^{l,N} - M_{y-\Delta}^{l,N}) \right),$$

whence

$$|K| \leq \mathbb{E}_\theta[(X_{x-\Delta,r}^{i,N})^2]^{1/2} \mathbb{E}[(M_u^{k,N} - M_{y-\Delta}^{k,N})^4]^{1/4} \mathbb{E}[(M_v^{l,N} - M_{y-\Delta}^{l,N})^4]^{1/4}.$$

Using Step 2, Lemma 16-(iii) and that $u - (y - \Delta) \leq 2\Delta$ and $v - (y - \Delta) \leq 2\Delta$, we easily conclude that indeed, $|K| \leq Ct^{1/2} \Delta^{-q}$.

Step 7. We now show, recalling that $y + \Delta \leq x - 2\Delta$, that

$$\left| \text{Cov}_\theta \left((\bar{\Gamma}_{x,x+\Delta}^N)^2, (\bar{\Gamma}_{y,y+\Delta}^N)^2 \right) \right| \leq CN^{-1} t^{1/2} \Delta^{1-q}.$$

$$\begin{aligned} |K_{a,b}| &\leq \left(\text{Var}_\theta \left((\bar{U}_{a\Delta}^N - \bar{U}_{(a-1)\Delta}^N)^2 \right) \text{Var}_\theta \left((\bar{U}_{b\Delta}^N - \bar{U}_{(b-1)\Delta}^N)^2 \right) \right)^{1/2} \\ &\leq C \left(\frac{\Delta^2}{N^2} + \frac{t^2}{N^2 \Delta^{4q}} \right). \end{aligned}$$

We finally used the first estimate of Lemma 30, which is valid since $x = (a-1)\Delta$ satisfies $x \geq t$ and thus $t/2 \leq x - \Delta \leq x + \Delta \leq 2t$ and $x = (b-1)\Delta$ satisfies the same conditions. If now $|a-b| \geq 3$ and w.l.o.g. $a > b$, we use the second estimate of Lemma 30, which is valid since $x = (a-1)\Delta$ and $y = (b-1)\Delta$ satisfy the required conditions (in particular, $y + \Delta \leq x - 2\Delta$). This gives

$$|K_{a,b}| \leq C \left(\frac{t^{1/2}}{N\Delta^{q-1}} + \frac{t^2}{N^2\Delta^{4q}} + \frac{t^{1/2}}{N^2\Delta^{q-3/2}} \right).$$

We end with

$$\begin{aligned} \mathbb{E}_\theta[(D_{\Delta,t}^{N,3})^2] &\leq C \frac{N^2}{t^2} \frac{t}{\Delta} \left(\frac{\Delta^2}{N^2} + \frac{t^2}{N^2\Delta^{4q}} \right) \\ &\quad + C \frac{N^2}{t^2} \frac{t^2}{\Delta^2} \left(\frac{t^{1/2}}{N\Delta^{q-1}} + \frac{t^2}{N^2\Delta^{4q}} + \frac{t^{1/2}}{N^2\Delta^{q-3/2}} \right). \end{aligned}$$

The conclusion follows. \square

We can at last give the

Proof of Proposition 25. Gathering Lemmas 26, 27, 29 and 31, we see that, on Ω_N^1 , if $1 \leq \Delta \leq t/4$,

$$\begin{aligned} \mathbb{E}_\theta[|\mathcal{W}_{\Delta,t}^N - \mathcal{W}_{\infty,\infty}^N|] &\leq \mathbb{E}_\theta[D_{\Delta,t}^{N,1} + 2D_{2\Delta,t}^{N,1} + D_{\Delta,t}^{N,2} + 2D_{2\Delta,t}^{N,2} + D_{\Delta,t}^{N,3} \\ &\quad + 2D_{2\Delta,t}^{N,3} + D_{\Delta,t}^{N,4}] \\ &\leq C \left(\frac{\Delta}{t} + \frac{N\Delta}{t^{2q}} + \frac{N}{t^{q-1}} + \frac{t}{\Delta^{q+1}} \right. \\ &\quad \left. + \sqrt{\frac{\Delta}{t} + \frac{t}{\Delta^{4q+1}} + \frac{Nt^{1/2}}{\Delta^{q+1}} + \frac{t^2}{\Delta^{4q+2}} + \frac{t^{1/2}}{\Delta^{q+1/2}}} \right). \end{aligned}$$

Using that $q \geq 3$ (whence in particular $2q-1 \geq q-1 \geq (q+1)/2$) and that $1 \leq \Delta \leq t$, we easily deduce that $\Delta/t \leq (\Delta/t)^{1/2}$, that $N\Delta t^{-2q} \leq N\Delta^{1-2q} \leq N\Delta^{-(q+1)/2}$, that $Nt^{1-q} \leq N\Delta^{1-q} \leq N\Delta^{-(q+1)/2}$, that $t\Delta^{-q-1} \leq t\Delta^{-q/2-1}$, that $t^{1/2}\Delta^{-2q-1/2} \leq t\Delta^{-2q-1} \leq t\Delta^{-q/2-1}$, that we have $N^{1/2}t^{1/4}\Delta^{-(q+1)/2} \leq N\Delta^{-(q+1)/2} + t^{1/2}\Delta^{-(q+1)/2} \leq N\Delta^{-(q+1)/2} + t\Delta^{-q/2-1}$, as well as $t\Delta^{-2q-1} \leq t\Delta^{-q/2-1}$ and $t^{1/4}\Delta^{-q/2-1/4} \leq t\Delta^{-q/2-1}$. This gives, still on Ω_N^1 ,

$$\mathbb{E}_\theta[|\mathcal{W}_{\Delta,t}^N - \mathcal{W}_{\infty,\infty}^N|] \leq C \left(\sqrt{\frac{\Delta}{t}} + \frac{N}{\Delta^{(q+1)/2}} + \frac{t}{\Delta^{q/2+1}} \right)$$

as desired. \square

4.7. Conclusion

We now have all the weapons to check our main result.

Proof of Theorem 3. Recall that we assume $H(q)$ for some $q > 3$ and that $\Delta_t = t/(2\lfloor t^{1-4/(q+1)} \rfloor) \sim t^{4/(q+1)}/2$ (for t large). We can of course assume that $t \geq 4$ is large enough so that $\Delta_t \in [1, t/4]$, because else the inequalities of the statement are trivial. Using Propositions 14 and 17, we find

$$\begin{aligned} \mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \mathcal{E}_t^N - \frac{\mu}{1-\Lambda p} \right| \right] &\leq \mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \mathcal{E}_t^N - \mu \bar{\ell}_N \right| \right] + \mu \mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \bar{\ell}_N - \frac{1}{1-\Lambda p} \right| \right] \\ &\leq C \left(\frac{1}{N} + \frac{1}{\sqrt{Nt}} + \frac{1}{t^q} \right). \end{aligned}$$

Since now $\Pr((\Omega_N^1)^c) \leq Ce^{-cN}$ by Lemma 13, we conclude that for any $\varepsilon \in (0, 1)$,

$$\Pr\left(\left| \mathcal{E}_t^N - \frac{\mu}{1-\Lambda p} \right| \geq \varepsilon\right) \leq Ce^{-cN} + \frac{C}{\varepsilon} \left(\frac{1}{N} + \frac{1}{\sqrt{Nt}} + \frac{1}{t^q} \right) \leq \frac{C}{\varepsilon} \left(\frac{1}{N} + \frac{1}{\sqrt{Nt}} + \frac{1}{t^q} \right).$$

Similarly, Propositions 14 and 19 imply, since $\mathcal{V}_\infty^N = \mu^2 \sum_{i=1}^N |\ell_N(i) - \bar{\ell}_N|^2$,

$$\begin{aligned} &\mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \mathcal{V}_t^N - \frac{\mu^2 \Lambda^2 p(1-p)}{(1-\Lambda p)^2} \right| \right] \\ &\leq \mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \mathcal{V}_t^N - \mathcal{V}_\infty^N \right| \right] + \mu^2 \mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \sum_{i=1}^N |\ell_N(i) - \bar{\ell}_N|^2 - \frac{\Lambda^2 p(1-p)}{(1-\Lambda p)^2} \right| \right] \\ &\leq \frac{C}{\sqrt{N}} + C \mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left(1 + \sum_{i=1}^N \left[\ell_N(i) - \bar{\ell}_N \right]^2 \right)^{1/2} \right] \left(\frac{N}{t^q} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{t}} \right) \\ &\leq C \left(\frac{1}{\sqrt{N}} + \frac{N}{t^q} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{t}} \right). \end{aligned}$$

The last inequality uses a second time Proposition 14. We conclude, using Lemma 13 as previously, that for any $\varepsilon \in (0, 1)$,

$$\begin{aligned} \Pr\left(\left| \mathcal{V}_t^N - \frac{\mu^2 \Lambda^2 p(1-p)}{(1-\Lambda p)^2} \right| \geq \varepsilon\right) &\leq Ce^{-cN} + \frac{C}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{N}{t^q} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{t}} \right) \\ &\leq \frac{C}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{N}{t^q} + \frac{\sqrt{N}}{t} \right) \end{aligned}$$

because $t^{-1/2} = (N^{1/4}t^{-1/2})N^{-1/4} \leq N^{1/2}t^{-1} + N^{-1/2}$. This implies that

$$\Pr\left(\left| \mathcal{V}_t^N - \frac{\mu^2 \Lambda^2 p(1-p)}{(1-\Lambda p)^2} \right| \geq \varepsilon\right) \leq \frac{C}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{t} \right).$$

Indeed, either $\sqrt{N} > t$ and the inequality is trivial or $\sqrt{N} \leq t$ and then $Nt^{-q} \leq Nt^{-2} \leq N^{1/2}t^{-1}$.

Finally, by Propositions 14 and 25, since $\mathcal{W}_{\infty, \infty}^N = \mu N^{-1} \sum_{i=1}^N \ell_N(i)(c_N(i))^2$,

$$\mathbb{E}\left[\mathbf{1}_{\Omega_N^1} \left| \mathcal{W}_{\Delta_t, t}^N - \frac{\mu}{(1-\Lambda p)^3} \right| \right]$$

$$\begin{aligned} &\leq \mathbb{E} \left[\mathbf{1}_{\Omega_N^1} \left| \mathcal{W}_{\Delta_t, t}^N - \mathcal{W}_{\infty, \infty}^N \right| \right] + \mu \mathbb{E} \left[\mathbf{1}_{\Omega_N^1} \left| N^{-1} \sum_{i=1}^N \ell_N(i) (c_N(i))^2 - \frac{1}{(1 - \Lambda p)^3} \right| \right] \\ &\leq C \left(\frac{1}{N} + \sqrt{\frac{\Delta_t}{t}} + \frac{N}{\Delta_t^{(q+1)/2}} + \frac{t}{\Delta_t^{q/2+1}} \right), \end{aligned}$$

whence as usual by Lemma 13, for $\varepsilon \in (0, 1)$,

$$\begin{aligned} \Pr \left(\left| \mathcal{W}_{\Delta_t, t}^N - \frac{\mu}{(1 - \Lambda p)^3} \right| \geq \varepsilon \right) &\leq C e^{-cN} + \frac{C}{\varepsilon} \left(\frac{1}{N} + \sqrt{\frac{\Delta_t}{t}} + \frac{N}{\Delta_t^{(q+1)/2}} + \frac{t}{\Delta_t^{q/2+1}} \right) \\ &\leq \frac{C}{\varepsilon} \left(\frac{1}{N} + \frac{1}{\sqrt{t^{1-4/(q+1)}}} + \frac{N}{t^2} \right). \end{aligned}$$

We finally used that $\Delta_t \sim t^{4/(q+1)}/2$, which of course implies that $\sqrt{\Delta_t/t} \sim \sqrt{1/(2t^{1-4/(q+1)})}$, that $N/\Delta_t^{(q+1)/2} \sim 2^{(q+1)/2} N t^{-2}$ and also that $t/\Delta_t^{q/2+1} \sim 2^{q/2+1} t^{-(q+3)/(q+1)} \leq 2^{q/2+1}/\sqrt{t^{1-4/(q+1)}}$. \square

Proof of Corollary 4. Recall that we assume $H(q)$ for some $q > 3$. We fix $\mu > 0$, $\Lambda > 0$ and $p \in (0, 1]$ such that $\Lambda p \in (0, 1)$. We define $u = \mu/(1 - \Lambda p)$, $v = \mu^2 \Lambda^2 p(1 - p)/(1 - \Lambda p)^2$ and $w = \mu/(1 - \Lambda p)^3$. It holds that $(u, v, w) \in D$ (which would not be the case if $\Lambda p = 0$) and $\Psi(u, v, w) = (\mu, \Lambda, p)$. Furthermore, Ψ is obviously of class C^∞ on D , it is in particular locally Lipschitz continuous. As a consequence, there is a constant $c \in (0, 1)$ (depending on μ, Λ, p) such that for any $N \geq 1$, any $t \geq 1$, any $\varepsilon \in (0, 1/c)$,

$$\begin{aligned} &\Pr \left(\left\| \Psi(\mathcal{E}_t^N, \mathcal{V}_t^N, \mathcal{W}_{\Delta_t, t}^N) - (\mu, \Lambda, p) \right\| \geq \varepsilon \right) \\ &\leq \Pr \left(|\mathcal{E}_t^N - u| + |\mathcal{V}_t^N - v| + |\mathcal{W}_{\Delta_t, t}^N - w| \geq c\varepsilon \right) \\ &\leq \frac{C}{\varepsilon} \left(\frac{1}{N} + \frac{1}{\sqrt{N}t} + \frac{1}{t^q} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{N}} + \frac{1}{N} + \frac{N}{t^2} + \frac{1}{\sqrt{t^{1-4/(q+1)}}} \right) \end{aligned}$$

by Theorem 3. Using next that $q > 3$, that $t \geq 1$ and $N \geq 1$, and that either $N^{1/2}t^{-1} \geq 1$ (whence the inequality below is trivial) or $N^{1/2}t^{-1} < 1$ (whence $Nt^{-2} \leq N^{1/2}t^{-1}$), we find

$$\Pr \left(\left\| \Psi(\mathcal{E}_t^N, \mathcal{V}_t^N, \mathcal{W}_{\Delta_t, t}^N) - (\mu, \Lambda, p) \right\| \geq \varepsilon \right) \leq \frac{C}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{t} + \frac{1}{\sqrt{t^{1-4/(q+1)}}} \right).$$

But $t^{-(1-4/(q+1))/2} = \lceil N^{1/4}t^{-(1-4/(q+1))/2} \rceil N^{-1/4} \leq N^{-1/2} + N^{1/2}t^{-(1-4/(q+1))}$, which concludes the proof. \square

We finally give the

Proof of Remark 2. Lemma 16-(ii) with $r = 1$ and $s = 0$ tells us that on Ω_N^1 , $|\mathbb{E}_\theta[\bar{Z}_t^N] - \mu \ell_N t| \leq C$. By Lemma 18, we know that $\mathbb{E}_\theta[|\bar{Z}_t^N - \mathbb{E}_\theta[\bar{Z}_t^N]|] =$

$\mathbb{E}_\theta[|\bar{U}_t^N|] \leq C(t/N)^{1/2}$, still on Ω_N^1 and, by Proposition 14, $\mathbb{E}[\mathbf{1}_{\Omega_N^1}|\bar{\ell}_N - 1/(1 - \Lambda p)|] \leq CN^{-1}$. We easily deduce that

$$\mathbb{E}[\mathbf{1}_{\Omega_N^1}|\bar{Z}_t^N - \mu(1 - \Lambda p)^{-1}t|] \leq C(1 + (t/N)^{1/2} + t/N) \leq C(1 + t/N).$$

Since $\Pr(\Omega_N^1) \geq 1 - Ce^{-cN}$ by Lemma 13, we find that for any $\varepsilon > 0$,

$$\Pr\left(\left|\frac{\bar{Z}_t^N}{t} - \frac{\mu}{1 - \Lambda p}\right| \geq \varepsilon\right) \leq Ce^{-cN} + \frac{C}{\varepsilon}\left(\frac{1}{t} + \frac{1}{N}\right).$$

The conclusion follows. □

5. The supercritical case

The goal of this section is to prove Theorem 6. In Subsection 5.1, we study precisely the Perron-Frobenius eigenvalue and eigenvector of the matrix with nonnegative entries $A_N(i, j) = N^{-1}\theta_{ij}$. In Subsection 5.2, we state and prove a few results on some series involving φ^{*n} . A few preliminary stochastic analysis is handled in Subsection 5.3. We finally conclude the proof in Subsection 5.4.

5.1. Perron-Frobenius analysis of the random matrix A_N

We recall that the norms $\|\cdot\|_r$ on \mathbb{R}^N and $|||\cdot|||_r$ on $\mathcal{M}_{N \times N}(\mathbb{R})$ were defined in Subsection 4.1. We denote by (e_1, \dots, e_N) the canonical basis of \mathbb{R}^N and by $\mathbf{1}_N = \sum_{i=1}^N e_i$ the vector with all entries equal to 1.

Notation 32. We consider the matrix $A_N(i, j) = N^{-1}\theta_{ij}$ and the event

$$\Omega_N^2 = \left\{ \frac{1}{N} \sum_{i,j=1}^N A_N(i, j) > \frac{p}{2} \right. \\ \left. \text{and for all } i, j = 1, \dots, N, |NA_N^2(i, j) - p^2| < \frac{p^2}{2N^{3/8}} \right\}.$$

Actually, $3/8$ could be replaced by any other exponent in $[3/8, 1/2)$. We first show that Ω_N^2 has a high probability.

Lemma 33. Assume that $p \in (0, 1]$. It holds that $\Pr(\Omega_N^2) \geq 1 - Ce^{-cN^{1/4}}$.

Proof. We recall the Hoeffding inequality [21] for a Binomial(n, q) random variable X : for all $x \geq 0$, it holds that $\Pr(|X - nq| \geq x) \leq 2\exp(-2x^2/n)$.

Since $N \sum_{i,j=1}^N A_N(i, j) = \sum_{i,j=1}^N \theta_{ij} \sim \text{Binomial}(N^2, p)$, we deduce that $\Pr(N^{-1} \sum_{i,j=1}^N A_N(i, j) \leq p/2) \leq \Pr(|N \sum_{i,j=1}^N A_N(i, j) - N^2p| \geq N^2p/2) \leq 2\exp(-N^2p^2/2)$.

For each $i \neq j$, we write $N^2A_N^2(i, j) = \sum_{k=1}^N \theta_{ik}\theta_{kj} = Z_{ij}^N + \theta_{ii}\theta_{ij} + \theta_{ij}\theta_{jj}$, where Z_{ij}^N follows a Binomial($N-2, p^2$) distribution. We thus have $|N^2A_N^2(i, j) -$

$Z_{ij}^N| \leq 2$. This obviously extends to the case where $i = j$. Hence for any i, j , $|NA_N^2(i, j) - p^2| \geq p^2/(2N^{3/8})$ implies that $|Z_{ij}^N - (N-2)p^2| \geq p^2N^{5/8}/2 - 4$ and thus, if $N \geq (16/p^2)^{8/5}$, that $|Z_{ij}^N - (N-2)p^2| \geq p^2N^{5/8}/4$. By the Hoeffding inequality, $\Pr(|NA_N^2(i, j) - p^2| \geq p^2/(2N^{3/8})) \leq 2 \exp(-p^4N^{5/4}/(8(N-2))) \leq 2 \exp(-p^4N^{1/4}/8)$.

All this shows that $\Pr((\Omega_N^2)^c) \leq 2 \exp(-N^2p^2/2) + 2N^2 \exp(-p^4N^{1/4}/8)$ for all $N \geq (16/p^2)^{8/5}$. The conclusion easily follows: we can find $0 < c < C$ depending only on p such that for all $N \geq 1$, $\Pr((\Omega_N^2)^c) \leq Ce^{-cN^{1/4}}$. \square

Next, we apply the Perron-Frobenius theorem.

Lemma 34. *Assume that $p \in (0, 1]$. On Ω_N^2 , the spectral radius ρ_N of A_N is a simple eigenvalue of A_N and $\rho_N \in [p(1 - 1/(2N^{3/8})), p(1 + 1/(2N^{3/8}))]$. There is a unique eigenvector $V_N \in (\mathbb{R}_+)^N$ of A_N for the eigenvalue ρ_N such that $\|V_N\|_2 = \sqrt{N}$. We also have $V_N(i) > 0$ for all $i = 1, \dots, N$.*

Proof. The matrix A_N has nonnegative entries and is irreducible on Ω_N^2 since A_N^2 has positive entries. We thus infer from the Perron-Frobenius theorem that on Ω_N^2 , ρ_N is a simple eigenvalue of A_N , that there is a unique corresponding eigenvector V_N with nonnegative entries such that $\|V_N\|_2 = \sqrt{N}$ and that $V_N(i) > 0$ for all $i = 1, \dots, N$.

Since $NA_N^2(i, j) \in [p^2(1 - 1/(2N^{3/8})), p^2(1 + 1/(2N^{3/8}))]$ for all i, j on Ω_N^2 we deduce from $\rho_N^2 V_N = A_N^2 V_N$ that $\rho_N^2 \|V_N\|_1 = \sum_{i,j=1}^N A_N^2(i, j) V_N(j) \leq p^2(1 + 1/(2N^{3/8})) \|V_N\|_1$, whence $\rho_N^2 \leq p^2(1 + 1/(2N^{3/8}))$ and thus $\rho_N \leq p(1 + 1/(2N^{3/8}))$. Similarly, we can write $\rho_N^2 \|V_N\|_1 = \sum_{i,j=1}^N A_N^2(i, j) V_N(j) \geq p^2(1 - 1/(2N^{3/8})) \|V_N\|_1$, whence $\rho_N^2 \geq p^2(1 - 1/(2N^{3/8}))$ and thus $\rho_N \geq p(1 - 1/(2N^{3/8}))$. \square

We now gather a number of important facts.

Lemma 35. *Assume that $p \in (0, 1]$. There is $N_0 \geq 1$ (depending only on p) such that for all $N \geq N_0$, on Ω_N^2 , the following properties hold true for all $i, j, k, l = 1, \dots, N$:*

- (i) for all $n \geq 2$, $A_N^n(i, j) \leq (3/2)A_N^n(k, l)$,
- (ii) $V_N(i) \in [1/2, 2]$,
- (iii) for all $n \geq 0$, $\|A_N^n \mathbf{1}_N\|_2 \in [\sqrt{N}\rho_N^n/2, 2\sqrt{N}\rho_N^n]$,
- (iv) for all $n \geq 2$, $A_N^n(i, j) \in [\rho_N^n/(3N), 3\rho_N^n/N]$,
- (v) for all $n \geq 0$, all $r \in [1, \infty]$, $\| \|A_N^n \mathbf{1}_N\|_r^{-1} A_N^n \mathbf{1}_N - \|V_N\|_r^{-1} V_N \|_r \leq 3(2N^{-3/8})^{\lfloor n/2 \rfloor + 1}$,
- (vi) for all $n \geq 0$, all $r \in [1, \infty]$, $\| \|A_N^n e_j\|_r^{-1} A_N^n e_j - \|V_N\|_r^{-1} V_N \|_r \leq 12(2N^{-3/8})^{\lfloor n/2 \rfloor}$,
- (vii) for all $n \geq 1$, $\|A_N^n e_j\|_2 \leq 3\rho_N^n/(p\sqrt{N})$ and for all $n \geq 0$, $\|A_N^n \mathbf{1}_N\|_\infty \leq 3\rho_N^n/p$.

The proof requires a quantitative version of the Perron-Frobenius theorem due to G. Birkhoff [7]. It is based on the use of the Hilbert projective distance.

Notation 36. For $x = (x_i)_{i=1,\dots,N}$ and $y = (y_i)_{i=1,\dots,N}$ in $(0, \infty)^N$, we set

$$d_N(x, y) = \log \left(\frac{\max_{i=1,\dots,N} (x_i/y_i)}{\min_{i=1,\dots,N} (x_i/y_i)} \right).$$

We have $d_N(x, y) = d_N(y, x) = d_N(x, \lambda y)$ for all $\lambda > 0$ and $d_N(x, y) \leq d_N(x, z) + d_N(z, y)$. Finally, $d_N(x, y) = 0$ if and only if x and y are colinear.

The result of Birkhoff quantifies the projection on the Perron-Frobenius vector.

Theorem 37 (Birkhoff [7], Cavazos-Cadena [11]). For any $A \in \mathcal{M}_{N \times N}(\mathbb{R})$ with positive entries and any x and y in $(0, \infty)^N$, we have $d_N(Ax, Ay) \leq k_A d_N(x, y)$, where

$$\Gamma_A = \max_{i,j,k,l=1,\dots,N} \frac{A(i,k)A(j,l)}{A(i,l)A(j,k)} \geq 1 \quad \text{and} \quad k_A = \frac{\sqrt{\Gamma_A} - 1}{\sqrt{\Gamma_A} + 1} \leq \frac{\Gamma_A - 1}{4}.$$

In our context, this gives the following estimates.

Remark 38. Assume that $p \in (0, 1]$. Then on Ω_N^2 , it holds that for all $x, y \in (0, \infty)^N$, we have (i) $d_N(A_N x, A_N y) \leq d_N(x, y)$ and (ii) $d_N(A_N^2 x, A_N^2 y) \leq 2N^{-3/8} d_N(x, y)$.

Proof. On Ω_N^2 , we have

$$A_N^2(i, j) \in [p^2 N^{-1} (1 - 1/(2N^{3/8})), p^2 N^{-1} (1 + 1/(2N^{3/8}))]. \quad (11)$$

This implies that for each $i = 1, \dots, N$, $\sum_{k=1}^N A_N(i, k) > 0$ (because else, $A_N^2(i, j)$ would vanish for all $j = 1, \dots, N$). Thus for $x, y \in (0, \infty)^N$, we have $A_N x, A_N y \in (0, \infty)^N$ so that $d_N(A_N x, A_N y)$ is well-defined. We put $m = \min_i (x_i/y_i)$ and $M = \max_i (x_i/y_i)$. We then have $m(A_N y)_i \leq (A_N x)_i \leq M(A_N y)_i$ for all i , whence $d_N(A_N x, A_N y) \leq \log(M/m) = d_N(x, y)$, which proves (i). For point (ii), it suffices to use Theorem 37, and to note that, by (11),

$$\Gamma_{A_N^2} = \max_{i,j,k,l=1,\dots,N} \frac{A_N^2(i,k)A_N^2(j,l)}{A_N^2(i,l)A_N^2(j,k)} \leq \frac{(1 + 1/(2N^{3/8}))^2}{(1 - 1/(2N^{3/8}))^2} \leq 1 + 8N^{-3/8},$$

whence $k_{A_N^2} \leq (\Gamma_{A_N^2} - 1)/4 \leq 2N^{-3/8}$. \square

We will also use the following easy remark.

Lemma 39. For all $r \in [1, \infty]$ and all $x, y \in (0, \infty)^N$ such that $d_N(x, y) \leq 1$, we have the inequality $|\|x\|_r^{-1} x - \|y\|_r^{-1} y|_r \leq 3d_N(x, y)$.

Proof. We fix $r \in [1, \infty]$ and assume without loss of generality that $\|x\|_r = \|y\|_r = 1$. We set $m = \min_i (x_i/y_i)$ and $M = \max_i (x_i/y_i)$. Since $\|x\|_r = \|y\|_r$, it holds that $m \leq 1 \leq M$. Using that $1 \geq d_N(x, y) = \log(1 + (M - m)/m)$, we deduce that $(M - m)/m \leq e - 1 \leq 2$. Since $\log(1 + u) \geq u/3$ on $[0, 2]$, we conclude that $d_N(x, y) \geq (M - m)/(3m) \geq (M - m)/3$. But for all i , we have $x_i \in [my_i, My_i]$, whence $|x_i - y_i| \leq (M - m)y_i$. Thus $\|x - y\|_r \leq (M - m)\|y\|_r = (M - m) \leq 3d_N(x, y)$. \square

We can now give the

Proof of Lemma 35. We work on Ω_N^2 during the whole proof.

Step 1. We first check that $d_N(\mathbf{1}_N, V_N) \leq 2N^{-3/8}$. We start from $A_N^2 V_N = \rho_N V_N$, so that for all i , $V_N(i) = \rho_N^{-2} \sum_{j=1}^N A_N^2(i, j) V_N(j)$. Using (11) and setting $\kappa_N = p^2 \rho_N^{-2} N^{-1} \sum_{j=1}^N V_N(j)$, we conclude that $V_N(i) \in [\kappa_N(1 - 1/(2N^{3/8}))]$, $\kappa_N(1 + 1/(2N^{3/8}))]$. Consequently, $\max_i V_N(i) / \min_i V_N(i)$ is smaller than $(1 + 1/(2N^{3/8})) / (1 - 1/(2N^{3/8})) \leq 1 + 2N^{-3/8}$. Hence

$$\begin{aligned} d_N(\mathbf{1}_N, V_N) &\leq \log[(1 + 1/(2N^{3/8})) / (1 - 1/(2N^{3/8}))] \\ &\leq \log(1 + 2N^{-3/8}) \leq 2N^{-3/8}. \end{aligned}$$

Step 2. Here we show that for all i , $V_N(i) \in [(1 + 2N^{-3/8})^{-1}, (1 + 2N^{-3/8})]$. This will imply point (ii) (for N large enough so that $2N^{-3/8} \leq 1$). We introduce $m = \min_i V_N(i)$ and $M = \max_i V_N(i)$. We have seen in Step 1 that $M/m \leq 1 + 2N^{-3/8}$. Recalling that $\|V_N\|_2 = \sqrt{N}$ by definition, we deduce that $N = \sum_{i=1}^N (V_N(i))^2 \leq NM^2 \leq N(1 + 2N^{-3/8})^2 m^2$, whence $m \geq (1 + 2N^{-3/8})^{-1}$. Similarly, $N = \sum_{i=1}^N (V_N(i))^2 \geq Nm^2 \geq N(1 + 2N^{-3/8})^{-2} M^2$, whence $M \leq (1 + 2N^{-3/8})$.

Step 3. We verify that for all $n \geq 0$, $d_N(A_N^n \mathbf{1}_N, V_N) \leq (2N^{-3/8})^{\lfloor n/2 \rfloor + 1}$. By Lemma 39, this will imply point (v) for all N large enough so that $2N^{-3/8} \leq 1$. Using that $A_N^n V_N = \rho_N^n V_N$, we find that $d_N(A_N^n \mathbf{1}_N, V_N) = d_N(A_N^n \mathbf{1}_N, A_N^n V_N)$. Hence for all n even, we deduce from Remark 38-(ii) and Step 1 that we have $d_N(A_N^n \mathbf{1}_N, V_N) \leq (2N^{-3/8})^{n/2} d_N(\mathbf{1}_N, V_N) \leq (2N^{-3/8})^{n/2+1}$. When n is odd, we simply use that $d_N(A_N^n \mathbf{1}_N, V_N) = d_N(A_N^n \mathbf{1}_N, A_N V_N) \leq d_N(A_N^{n-1} \mathbf{1}_N, V_N)$ by Remark 38-(i).

Step 4. We now prove (vi). We fix $r \in [1, \infty]$ and $j \in \{1, \dots, N\}$. The result is obvious if $n = 0$ or $n = 1$ because then $\| \|A_N^n e_j\|_r^{-1} A_N^n e_j - \|V_N \mathbf{1}_N\|_r^{-1} V_N \|_r \leq 2 \leq 12(2N^{-3/8})^{\lfloor n/2 \rfloor}$.

By Remark 38-(ii), we see that $d_N(A_N^{2k} e_j, V_N) = d_N(A_N^{2k} e_j, A_N^{2k} V_N) \leq (2N^{-3/8})^{k-1} d_N(A_N^2 e_j, V_N)$ for all $k \geq 1$.

We next write $d_N(A_N^2 e_j, V_N) \leq d_N(A_N^2 e_j, \mathbf{1}_N) + d_N(\mathbf{1}_N, V_N)$. By Step 1, we have $d_N(\mathbf{1}_N, V_N) \leq \log[(1 + N^{-3/8}/2) / (1 - N^{-3/8}/2)]$. Furthermore, we deduce from (11) that $d_N(A_N^2 e_j, \mathbf{1}_N) = \log[\max_i (A_N^2(i, j)) / \min_i (A_N^2(i, j))] \leq \log[(1 + N^{-3/8}/2) / (1 - N^{-3/8}/2)]$. All in all, we find that $d_N(A_N^2 e_j, V_N) \leq \log[(1 + N^{-3/8}/2)^2 / (1 - N^{-3/8}/2)^2] \leq \log(1 + 8N^{-3/2}) \leq 8N^{-3/2}$.

Hence for all $k \geq 1$, $d_N(A_N^{2k} e_j, V_N) \leq 8N^{-3/8} (2N^{-3/8})^{k-1} = 4(2N^{-3/8})^k$. We also have, by Remark 38-(i), $d_N(A_N^{2k+1} e_j, V_N) = d_N(A_N^{2k+1} e_j, A_N V_N) \leq d_N(A_N^{2k} e_j, V_N)$. Thus for all $n \geq 2$, $d_N(A_N^n e_j, V_N) \leq 4(2N^{-3/8})^{\lfloor n/2 \rfloor}$. This implies that indeed, $\| \|A_N^n e_j\|_r^{-1} A_N^n e_j - \|V_N\|_r^{-1} V_N \|_r \leq 12(2N^{-3/8})^{\lfloor n/2 \rfloor}$ by Lemma 39, if N is large enough so that $2N^{-3/8} \leq 1/4$.

Step 5. We check (i). Using Step 2, we see that for all $j = 1, \dots, N$, all $n \geq 2$,

$$d_N(A_N^n e_j, V_N) = \log \left(\frac{\max_i (A_N^n(i, j) / V_N(i))}{\min_i (A_N^n(i, j) / V_N(i))} \right)$$

$$\geq \log \left(\frac{\max_i A_N^n(i, j)}{\min_i A_N^n(i, j)} \times (1 + 2N^{-3/8})^{-2} \right).$$

But for all $n \geq 2$, using Remark 38-(i), $d_N(A_N^n e_j, V_N) = d_N(A_N^n e_j, A_N^{n-2} V_N) \leq d_N(A_N^2 e_j, V_N) \leq \log(1 + 8N^{-3/8})$ as seen in Step 4. We conclude that

$$\frac{\max_i A_N^n(i, j)}{\min_i A_N^n(i, j)} \leq (1 + 2N^{-3/8})^2 (1 + 8N^{-3/8}).$$

Using the same arguments with the transpose matrix A_N^t (which satisfies exactly the same assumptions as A_N on Ω_N^2), we see that for all $i = 1, \dots, N$,

$$\frac{\max_j A_N^n(i, j)}{\min_j A_N^n(i, j)} \leq (1 + 2N^{-3/8})^2 (1 + 8N^{-3/8}).$$

Finally, we conclude that for all $n \geq 2$,

$$\frac{\max_{i,j} A_N^n(i, j)}{\min_{i,j} A_N^n(i, j)} \leq (1 + 2N^{-3/8})^4 (1 + 8N^{-3/8})^2.$$

This is indeed smaller than $3/2$ if N is large enough.

Step 6. We now verify (iii). We write $A_N^n \mathbf{1}_N = \|A_N^n \mathbf{1}_N\|_2 (N^{-1/2} V_N + Z_{N,n})$, where $Z_{N,n} = \|A_N^n \mathbf{1}_N\|_2^{-1} A_N^n \mathbf{1}_N - N^{-1/2} V_N$. We know by (v) (with $r = 2$) that $\|Z_{N,n}\|_2 \leq 3(2N^{-3/8})^{\lfloor n/2 \rfloor + 1}$. We next write, for each $n \geq 0$, $A_N^{n+1} \mathbf{1}_N = \|A_N^n \mathbf{1}_N\|_2 (N^{-1/2} \rho_N V_N + A_N Z_{N,n})$. Since $\|V_N\|_2 = \sqrt{N}$ and $\|A_N\|_2 \leq 1$ (which immediately follows from the fact that $0 \leq A_N(i, j) \leq 1/N$), we conclude that $|\|A_N^{n+1} \mathbf{1}_N\|_2 - \rho_N \|A_N^n \mathbf{1}_N\|_2| \leq 3\|A_N^n \mathbf{1}_N\|_2 (2N^{-3/8})^{\lfloor n/2 \rfloor + 1}$.

We now set $x_n = \|A_N^n \mathbf{1}_N\|_2 / (\sqrt{N} \rho_N^n)$. For all $n \geq 0$, we have

$$|x_{n+1} - x_n| \leq 3x_n (2N^{-3/8})^{\lfloor n/2 \rfloor + 1} / \rho_N \leq 6x_n (2N^{-3/8})^{\lfloor n/2 \rfloor + 1} / p,$$

because $\rho_N \geq p/2$ on Ω_N^2 , see Lemma 34. If now N is large enough so that $6(2N^{-3/8})^{1/2} / p \leq 1/2$, we easily conclude, using that $x_0 = 1$, that, for all $n \geq 1$,

$$x_n \in \left[\prod_{k=1}^n (1 - 6(2N^{-3/8})^{\lfloor k/2 \rfloor + 1} / p), \prod_{k=1}^n (1 + 6(2N^{-3/8})^{\lfloor k/2 \rfloor + 1} / p) \right],$$

which is included in $[1/2, 2]$ if N is large enough (depending only on p). Since $x_0 = 1$, we thus have $x_n \in [1/2, 2]$ for all $n \geq 0$, and thus $\|A_N^n \mathbf{1}_N\|_2 \in [\sqrt{N} \rho_N^n / 2, 2\sqrt{N} \rho_N^n]$ for all $n \geq 0$.

Step 7. Here we prove (iv). We fix $n \geq 2$ and set $m = \min_{i,j} A_N^n(i, j)$ and $M = \max_{i,j} A_N^n(i, j)$. We know from (i) that $M/m \leq 3/2$. Starting from point (iii), we write $\sqrt{N} \rho_N^n / 2 \leq \|A_N^n \mathbf{1}_N\|_2 = (\sum_{i=1}^N (\sum_{j=1}^N A_N^n(i, j))^2)^{1/2} \leq N^{3/2} M \leq 3N^{3/2} m / 2$, whence $m \geq \rho_N^n / (3N)$. By the same way, $2\sqrt{N} \rho_N^n \geq \|A_N^n \mathbf{1}_N\|_2 \geq N^{3/2} m \geq 2N^{3/2} M / 3$, whence $M \leq 3\rho_N^n / N$.

Step 8. It only remains to check (vii). We know from (iv) that for all $n \geq 2$, $A_N^n(i, j) \leq 3\rho_N^n / N \leq 3\rho_N^n / (pN)$. And for $n = 1$, $A_N(i, j) \leq 1/N \leq 3\rho_N / (pN)$

because $\rho_N \geq p/3$ on Ω_N^2 , see Lemma 34. We conclude that for all $n \geq 1$, $A_N^n(i, j) \leq 3\rho_N^n/(pN)$. This immediately implies that for all $n \geq 1$, $\|A_N^n e_j\|_2 = (\sum_{i=1}^N (A_N^n(i, j))^2)^{1/2} \leq 3\rho_N^n/(p\sqrt{N})$ and $\|A_N^n \mathbf{1}_N\|_\infty = \max_i \sum_{j=1}^N A_N^n(i, j) \leq 3\rho_N^n/p$. Finally, for $n = 0$, we of course have $\|A_N^0 \mathbf{1}_N\|_\infty = 1 \leq 3\rho_N^0/p$. \square

Finally, the following tedious result is crucial for our estimation method.

Proposition 40. *We assume that $p \in (0, 1]$ and we introduce, on Ω_N^2 , $\bar{V}_N = N^{-1} \sum_{i=1}^N V_N(i)$ and*

$$\mathcal{U}_\infty^N = \sum_{i=1}^N \left(\frac{V_N(i) - \bar{V}_N}{\bar{V}_N} \right)^2.$$

There is $N_0 \geq 1$ and $C > 0$ (depending only on p) such that for all $N \geq N_0$,

$$\mathbb{E} \left[\mathbf{1}_{\Omega_N^2} \left| \mathcal{U}_\infty^N - \left(\frac{1}{p} - 1 \right) \right| \right] \leq \frac{C}{\sqrt{N}} \quad \text{and} \quad \mathbb{E}[\mathbf{1}_{\Omega_N^2} \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2] \leq C.$$

Proof. We work with N large enough so that we can apply Lemma 35. We introduce the vectors $L_N = A_N \mathbf{1}_N$ and $\mathcal{L}_N = A_N^6 \mathbf{1}_N$, we set $\bar{L}_N = N^{-1} \sum_{i=1}^N L_N(i)$, $\bar{\mathcal{L}}_N = N^{-1} \sum_{i=1}^N \mathcal{L}_N(i)$,

$$H_N = \sum_{i=1}^N \left(\frac{L_N(i) - \bar{L}_N}{\bar{L}_N} \right)^2 \quad \text{and} \quad \mathcal{H}_N = \sum_{i=1}^N \left(\frac{\mathcal{L}_N(i) - \bar{\mathcal{L}}_N}{\bar{\mathcal{L}}_N} \right)^2.$$

We checked in the proof of Proposition 14-Step 2 that (i) $\mathbb{E}[\|\bar{L}_N - p\|^2] \leq CN^{-2}$, (ii) $\mathbb{E}[\|L_N - \bar{L}_N \mathbf{1}_N\|_2^4] \leq C$, (iii) $\mathbb{E}[(\|L_N - \bar{L}_N \mathbf{1}_N\|_2^2 - p(1-p))^2] \leq CN^{-1}$, (iv) $\mathbb{E}[\|A_N L_N - \bar{L}_N L_N\|_2^2] \leq CN^{-1}$.

We also recall that $\bar{L}_N \leq 1$ and $\|A_N\|_2 \leq 1$ (simply because $0 \leq A_N(i, j) \leq 1/N$). Furthermore, on Ω_N^2 , it holds that $\bar{L}_N = N^{-1} \sum_{i,j=1}^N A_N(i, j) \geq p/2$, that $\bar{\mathcal{L}}_N = N^{-1} \sum_{i,j=1}^N A_N^6(i, j) \geq \rho_N^6/3 \geq p^6/192$ (by Lemma 35-(iv) and because $\rho_N \geq p/2$ by Lemma 34) and that $\bar{V}_N \geq 1/2$ (by Lemma 35-(ii)).

Step 1. We show that on Ω_N^2 , $\Delta_N = |\mathcal{U}_\infty^N - \mathcal{H}_N| \leq CN^{-1/2}$. A simple computation shows that

$$\begin{aligned} \Delta_N &= \left| \sum_{i=1}^N \left[\left(\frac{V_N(i)}{\bar{V}_N} \right)^2 - \left(\frac{\mathcal{L}_N(i)}{\bar{\mathcal{L}}_N} \right)^2 \right] \right| \\ &\leq \left(\sum_{i=1}^N \left| \frac{V_N(i)}{\bar{V}_N} - \frac{\mathcal{L}_N(i)}{\bar{\mathcal{L}}_N} \right| \right) \left(\max_i \left(\frac{V_N(i)}{\bar{V}_N} + \frac{\mathcal{L}_N(i)}{\bar{\mathcal{L}}_N} \right) \right) = S_N T_N, \end{aligned}$$

the last equality being a definition.

Lemma 35-(ii) implies that $\max_i (V_N(i)/\bar{V}_N) \leq \max_i V_N(i)/\min_i V_N(i) \leq 4$ and Lemma 35-(i) implies that $\max_i (\mathcal{L}_N(i)/\bar{\mathcal{L}}_N) \leq \max_i \mathcal{L}_N(i)/\min_i \mathcal{L}_N(i) \leq 3/2$ because $\mathcal{L}_N = A_N^6 \mathbf{1}_N$. Thus $T_N \leq 4 + 3/2 \leq 6$. Next, it holds that $S_N = N \left\| \|A_N^6 \mathbf{1}_N\|_1^{-1} A_N^6 \mathbf{1}_N - \|V_N \mathbf{1}_N\|_1^{-1} V_N \right\|_1$. We thus infer from Lemma 35-(v) that $S_N \leq 3N(2N^{-3/8})^4 = 48N^{-1/2}$. The conclusion follows.

Step 2. We next prove that $\mathbb{E}[\mathbf{1}_{\Omega_N^2} |\mathcal{H}_N - H_N|] \leq CN^{-1/2}$. We first write

$$\begin{aligned} \|\mathcal{L}_N - (\bar{L}_N)^5 L_N\|_2 &= \|A_N^6 \mathbf{1}_N - (\bar{L}_N)^5 A_N \mathbf{1}_N\|_2 \\ &\leq \sum_{k=1}^5 \|(\bar{L}_N)^{5-k} A_N^{k+1} \mathbf{1}_N - (\bar{L}_N)^{6-k} A_N^k \mathbf{1}_N\|_2. \end{aligned}$$

Using that $\bar{L}_N \leq 1$ and $\|A_N\|_2 \leq 1$, we deduce that

$$\|\mathcal{L}_N - (\bar{L}_N)^5 L_N\|_2 \leq 5 \|A_N^2 \mathbf{1}_N - \bar{L}_N A_N \mathbf{1}_N\|_2 = 5 \|A_N L_N - \bar{L}_N L_N\|_2.$$

We thus deduce from point (iv) recalled above that $\mathbb{E}[\|\mathcal{L}_N - (\bar{L}_N)^5 L_N\|_2^2] \leq CN^{-1}$. But it holds that $\|\mathcal{L}_N - (\bar{L}_N)^5 L_N\|_2 = I_N + J_N$, where $I_N = \|(\mathcal{L}_N - \bar{L}_N \mathbf{1}_N) - (\bar{L}_N)^5 (L_N - \bar{L}_N \mathbf{1}_N)\|_2$ and $J_N = \|\bar{L}_N \mathbf{1}_N - (\bar{L}_N)^6 \mathbf{1}_N\|_2 = \sqrt{N} |\bar{L}_N - (\bar{L}_N)^6|$. Consequently, $\mathbb{E}[I_N^2] + \mathbb{E}[J_N^2] \leq CN^{-1}$. Using now that

$$\begin{aligned} H_N &= \frac{\|L_N - \bar{L}_N \mathbf{1}_N\|_2^2}{(\bar{L}_N)^2} = \frac{\|(\bar{L}_N)^5 (L_N - \bar{L}_N \mathbf{1}_N)\|_2^2}{(\bar{L}_N)^{12}} \\ \text{and } \mathcal{H}_N &= \frac{\|\mathcal{L}_N - \bar{L}_N \mathbf{1}_N\|_2^2}{(\bar{L}_N)^2}, \end{aligned}$$

the facts that $\bar{L}_N \geq p^6/192$ and $(\bar{L}_N)^6 \geq p^6/64$ on Ω_N^2 and that the map $x \mapsto x^{-2}$ is globally Lipschitz and bounded on $[p^6/192, \infty)$, we conclude that, still on Ω_N^2 ,

$$\begin{aligned} |H_N - \mathcal{H}_N| &\leq C \left(\|(\bar{L}_N)^5 (L_N - \bar{L}_N \mathbf{1}_N)\|_2^2 |(\bar{L}_N)^6 - \bar{L}_N| \right. \\ &\quad \left. + \left| \|\mathcal{L}_N - \bar{L}_N \mathbf{1}_N\|_2^2 - \|(\bar{L}_N)^5 (L_N - \bar{L}_N \mathbf{1}_N)\|_2^2 \right| \right). \end{aligned}$$

Using now the inequality $|a^2 - b^2| \leq (a - b)^2 + 2a|a - b|$ for $a, b \geq 0$, we deduce that

$$\begin{aligned} |H_N - \mathcal{H}_N| &\leq C \left(\|(\bar{L}_N)^5 (L_N - \bar{L}_N \mathbf{1}_N)\|_2^2 N^{-1/2} J_N + I_N^2 \right. \\ &\quad \left. + \|(\bar{L}_N)^5 (L_N - \bar{L}_N \mathbf{1}_N)\|_2^2 I_N \right) \\ &\leq C \left(\|L_N - \bar{L}_N \mathbf{1}_N\|_2^2 N^{-1/2} J_N + I_N^2 + \|L_N - \bar{L}_N \mathbf{1}_N\|_2^2 I_N \right) \end{aligned}$$

because $\bar{L}_N \leq 1$. Using the Cauchy-Schwarz inequality, that $\mathbb{E}[I_N^2] + \mathbb{E}[J_N^2] \leq CN^{-1}$ and that $\mathbb{E}[\|L_N - \bar{L}_N \mathbf{1}_N\|_2^4] \leq C$ by point (ii) recalled above, we conclude that $\mathbb{E}[\mathbf{1}_{\Omega_N^2} |\mathcal{H}_N - H_N|] \leq CN^{-1/2}$.

Step 3. Here we check that $\mathbb{E}[\mathbf{1}_{\Omega_N^2} |H_N - (1/p - 1)|] \leq CN^{-1/2}$. Since $\bar{L}_N \geq p/2$ on Ω_N^2 and since $x \mapsto x^{-2}$ is bounded and globally Lipschitz continuous on $[p/2, \infty)$, we can write

$$\left| H_N - \left(\frac{1}{p} - 1 \right) \right| = \left| \frac{\|L_N - \bar{L}_N \mathbf{1}_N\|_2^2}{(\bar{L}_N)^2} - \frac{p(1-p)}{p^2} \right|$$

$$\leq C \left(|\bar{L}_N - p|p(1-p) + \left| \|L_N - \bar{L}_N \mathbf{1}_N\|_2^2 - p(1-p) \right| \right).$$

The conclusion follows, since as recalled in points (i) and (iii) above, $\mathbb{E}[|\bar{L}_N - p|] \leq CN^{-1}$ and $\mathbb{E}[\left| \|L_N - \bar{L}_N \mathbf{1}_N\|_2^2 - p(1-p) \right|] \leq CN^{-1/2}$.

Step 4. Gathering Steps 1, 2 and 3, we immediately deduce that $\mathbb{E}[\mathbf{1}_{\Omega_N^2} |\mathcal{U}_\infty^N - (1/p - 1)|] \leq CN^{-1/2}$. Since now $\bar{V}_N \leq 2$ on Ω_N^2 by Lemma 35-(ii), $\|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 = (\bar{V}_N)^2 \mathcal{U}_\infty^N \leq 4\mathcal{U}_\infty^N$, whence of course, $\mathbb{E}[\mathbf{1}_{\Omega_N^2} \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2] \leq C$. \square

5.2. Preliminary analytic estimates

We recall the following lemma, relying on some results of Feller [14] on convolution equations, that can be found in [13, Lemma 26-(b)].

Lemma 41. *Let $\psi : [0, \infty) \mapsto [0, \infty)$ be integrable and such that $\int_0^\infty \psi(t)dt > 1$. Assume also that $t \mapsto \int_0^t |d\psi(s)|$ has at most polynomial growth and set $\Gamma_t = \sum_{n \geq 0} \psi^{*n}(t)$. Consider $\alpha > 0$ such that $\int_0^\infty e^{-\alpha t} \psi(t)dt = 1$. There are $0 < c < C$ such that for all $t \geq 0$, $1 + \Gamma_t \in [ce^{\alpha t}, Ce^{\alpha t}]$.*

Based on this, it is not hard to verify the following result.

Lemma 42. *Assume A. Recall that α_0 was defined in Remark 5 such that $p \int_0^\infty e^{-\alpha_0 t} \varphi(t)dt = 1$ and that ρ_N was defined, for each $N \geq 1$, in Lemma 34. We now set $\Gamma_t^N = \sum_{n \geq 0} \rho_N^n \varphi^{*n}(t)$. For any $\eta > 0$, we can find $N_\eta \geq 1$ and $0 < c_\eta < C_\eta$ (depending only on p, φ and η) such that for all $N \geq N_\eta$, on Ω_N^2 , for all $t \geq 0$, $1 + \Gamma_t^N \in [c_\eta e^{(\alpha_0 - \eta)t}, C_\eta e^{(\alpha_0 + \eta)t}]$.*

Proof. We only prove the result when $\eta \in (0, \alpha_0)$, which of course suffices. We consider $\rho_\eta^+ > p > \rho_\eta^-$ defined respectively by $\int_0^\infty e^{-(\alpha_0 + \eta)t} \varphi(t)dt = 1/\rho_\eta^+$ and $\int_0^\infty e^{-(\alpha_0 - \eta)t} \varphi(t)dt = 1/\rho_\eta^-$. We put $\Gamma_t^{\eta,+} = \sum_{n \geq 0} (\rho_\eta^+)^n \varphi^{*n}(t)$ and $\Gamma_t^{\eta,-} = \sum_{n \geq 0} (\rho_\eta^-)^n \varphi^{*n}(t)$. Applying Lemma 41 with $\psi = \rho_\eta^+ \varphi$ and with $\psi = \rho_\eta^- \varphi$, we deduce that there are some constants $0 < c_\eta < C_\eta$ such that for all $t \geq 0$, $c_\eta e^{(\alpha_0 - \eta)t} \leq 1 + \Gamma_t^{\eta,-} \leq 1 + \Gamma_t^{\eta,+} \leq C_\eta e^{(\alpha_0 + \eta)t}$. But on Ω_N^2 , we know from Lemma 34 that $\rho_N \in [p(1 - N^{-3/8}/2), p(1 + N^{-3/8}/2)]$. Thus for N large enough, we clearly have $\rho_N \in [\rho_\eta^-, \rho_\eta^+]$, so that $\Gamma_t^N \in [\Gamma_t^{\eta,-}, \Gamma_t^{\eta,+}]$. The conclusion follows. \square

We next gather a number of consequences of the above estimate that we will use later.

Lemma 43. *Assume A. Recall that α_0 was defined in Remark 5, that ρ_N was defined in Lemma 34 and that $\Gamma_t^N = \sum_{n \geq 0} \rho_N^n \varphi^{*n}(t)$. We also put $v_t^N = \mu N^{-1/2} \sum_{n \geq 0} \|A_N^n \mathbf{1}_N\|_2 \int_0^t s \varphi^{*n}(t-s)ds$. For any $\eta > 0$, we can find $N_\eta \geq 1$, $t_\eta > 0$ and $0 < c_\eta < C_\eta$ (depending only on p, μ, φ and η) such that for all $N \geq N_\eta$, on Ω_N^2 ,*

- (i) for all $t \geq 0$, $v_t^N \leq C_\eta e^{(\alpha_0 + \eta)t}$,
- (ii) for all $t \geq t_\eta$, $v_t^N \geq c_\eta e^{(\alpha_0 - \eta)t}$,
- (iii) for all $t \geq 0$, $\sum_{n \geq 0} \rho_N^n (2N^{-3/8})^{\lfloor n/2 \rfloor} \int_0^t \varphi^{*n}(t-s)ds \leq C_\eta$,

- (iv) for all $t \geq 0$, $\sum_{n \geq 0} \rho_N^n \int_0^t e^{(\alpha_0 + \eta)s/2} \varphi^{*n}(t-s) ds \leq C_\eta e^{(\alpha_0 + \eta)t}$,
- (v) for all $t \geq 0$, $\sum_{n \geq 0} \rho_N^n \int_0^t s \varphi^{*n}(t-s) ds \leq C_\eta e^{(\alpha_0 + \eta)t}$,
- (vi) for all $t \geq 0$, $\int_0^t \int_0^t \Gamma_{t-r}^N \Gamma_{t-s}^N e^{(\alpha_0 + \eta)(r \wedge s)} dr ds \leq C_\eta e^{2(\alpha_0 + \eta)t}$.

Proof. We fix $\eta > 0$ and work with N large enough and on Ω_N^2 , so that we can use Lemmas 35 and 42.

We start with (i). By Lemma 35-(iii), $\|A_N^n \mathbf{1}_N\|_2 \leq 2\sqrt{N} \rho_N^n$. Hence $v_t^N \leq 2\mu \int_0^t s \Gamma_{t-s}^N ds \leq C_\eta \int_0^t s e^{(\alpha_0 + \eta)(t-s)} ds = C_\eta e^{(\alpha_0 + \eta)t} \int_0^t s e^{-(\alpha_0 + \eta)s} ds \leq C_\eta e^{(\alpha_0 + \eta)t}$.

The LHS of point (iv) is $\int_0^t e^{(\alpha_0 + \eta)s/2} \Gamma_{t-s}^N ds \leq C_\eta \int_0^t e^{(\alpha_0 + \eta)s/2} e^{(\alpha_0 + \eta)(t-s)} ds$, which equals $C_\eta e^{(\alpha_0 + \eta)t} \int_0^t e^{-(\alpha_0 + \eta)s/2} ds \leq C_\eta e^{(\alpha_0 + \eta)t}$.

Point (v) follows from point (iv).

The LHS of (vi) is bounded by

$$C_\eta \int_0^t \int_0^t e^{(\alpha_0 + \eta)(t-r)} e^{(\alpha_0 + \eta)(t-s)} e^{(\alpha_0 + \eta)(r \wedge s)} dr ds,$$

which equals $2C_\eta \int_0^t e^{(\alpha_0 + \eta)(t-s)} \int_0^s e^{(\alpha_0 + \eta)(t-r)} e^{(\alpha_0 + \eta)r} dr ds$, which itself equals $2C_\eta e^{2(\alpha_0 + \eta)t} \int_0^t s e^{-(\alpha_0 + \eta)s} ds \leq C_\eta e^{2(\alpha_0 + \eta)t}$.

Setting $\Lambda = \int_0^\infty \varphi(t) dt$, the LHS of point (iii) is bounded by the quantity $\sum_{n \geq 0} (\Lambda \rho_N)^n (2N^{-3/8})^{\lfloor n/2 \rfloor} \leq \sum_{n \geq 0} (2\Lambda p)^n (2N^{-3/8})^{\lfloor n/2 \rfloor}$ since $\rho_N \leq 2p$ on Ω_N^2 by Lemma 34. This is uniformly bounded, as soon as N is large enough so that $2\Lambda p (2N^{-3/8})^{1/2} \leq 1/2$.

We finally check (ii). We know from Lemma 35-(iii) that, on Ω_N^2 , $\|A_N^n \mathbf{1}_N\|_2 \geq \sqrt{N} \rho_N^n / 2$, whence $v_t^N \geq (\mu/2) \int_0^t s \Gamma_{t-s}^N ds \geq (\mu/2) \int_1^2 s \Gamma_{t-s}^N ds \geq (\mu/2) \int_{t-2}^{t-1} \Gamma_s^N ds$ if $t \geq 2$. By Lemma 42, we thus have $v_t^N \geq (\mu/2) \int_{t-2}^{t-1} (c_\eta e^{(\alpha_0 - \eta)s} - 1) ds \geq (\mu/2) [c_\eta e^{(\alpha_0 - \eta)(t-2)} - 1]$. The conclusion easily follows: we can find $t_\eta \geq 2$ and $c_\eta > 0$ such that for all $t \geq t_\eta$, $v_t^N \geq c_\eta e^{(\alpha_0 - \eta)t}$. \square

5.3. Preliminary stochastic analysis

We now prove a few estimates concerning the processes introduced in Notation 9. We recall that α_0 was defined in Remark 5 and that ρ_N and V_N were defined in Lemma 34. We start from Lemma 11 to write (with as usual $\varphi^{*0}(t-s) ds = \delta_t(ds)$)

$$\mathbb{E}_\theta[\mathbf{Z}_t^N] = \mu \sum_{n \geq 0} \left[\int_0^t s \varphi^{*n}(t-s) ds \right] A_N^n \mathbf{1}_N = v_t^N V_N + \mathbf{I}_t^N, \quad (12)$$

$$\mathbf{U}_t^N = \mathbf{Z}_t^N - \mathbb{E}_\theta[\mathbf{Z}_t^N] = \sum_{n \geq 0} \int_0^t \varphi^{*n}(t-s) A_N^n \mathbf{M}_s^N ds = \mathbf{M}_t^N + \mathbf{J}_t^N, \quad (13)$$

where

$$v_t^N = \mu \sum_{n \geq 0} \frac{\|A_N^n \mathbf{1}_N\|_2}{\sqrt{N}} \int_0^t s \varphi^{*n}(t-s) ds, \quad (14)$$

$$\mathbf{I}_t^N = \mu \sum_{n \geq 0} \left[\int_0^t s \varphi^{*n}(t-s) ds \right] \left[A_N^n \mathbf{1}_N - \frac{\|A_N^n \mathbf{1}_N\|_2}{\sqrt{N}} V_N \right], \quad (15)$$

$$\mathbf{J}_t^N = \sum_{n \geq 1} \int_0^t \varphi^{*n}(t-s) A_N^n \mathbf{M}_s^N ds. \quad (16)$$

As usual, we denote by $I_t^{i,N}$ and $J_t^{i,N}$ the coordinates of \mathbf{I}_t^N and \mathbf{J}_t^N and by \bar{I}_t^N and \bar{J}_t^N their empirical mean. We start with some upperbounds concerning \mathbf{Z}_t^N and \mathbf{U}_t^N .

Lemma 44. *Assume A. For all $\eta > 0$, there are $N_\eta \geq 1$ and $C_\eta > 0$ such that for all $N \geq N_\eta$, all $t \geq 0$, on Ω_N^2 ,*

- (i) $\max_{i=1, \dots, N} \mathbb{E}_\theta[(Z_t^{i,N})^2] \leq C_\eta e^{2(\alpha_0 + \eta)t}$,
- (ii) $\max_{i=1, \dots, N} \mathbb{E}_\theta[(U_t^{i,N})^2] \leq C_\eta (N^{-1} e^{2(\alpha_0 + \eta)t} + e^{(\alpha_0 + \eta)t})$,
- (iii) $\mathbb{E}_\theta[(\bar{U}_t^N)^2] \leq C_\eta N^{-1} e^{2(\alpha_0 + \eta)t}$.

Proof. We fix $\eta > 0$ and work with N large enough and on Ω_N^2 , so that we can use Lemmas 35 and 43.

Step 1. We first verify that $\|\mathbb{E}_\theta[\mathbf{Z}_t^N]\|_\infty \leq C_\eta e^{(\alpha_0 + \eta)t}$. Using (12) and that $\|A_N^n \mathbf{1}_N\|_\infty \leq C \rho_N^n$ for all $n \geq 0$ by Lemma 35-(vii), we see that $\|\mathbb{E}_\theta[\mathbf{Z}_t^N]\|_\infty \leq C \sum_{n \geq 0} \rho_N^n \int_0^t s \varphi^{*n}(t-s) ds$, whence the conclusion by Lemma 43-(v).

Step 2. We next show that for all $i = 1, \dots, N$, $\mathbb{E}_\theta[(J_t^{i,N})^2] \leq C_\eta N^{-1} e^{2(\alpha_0 + \eta)t}$. We start from (16), which gives us

$$\begin{aligned} \mathbb{E}_\theta[(J_t^{i,N})^2] &= \sum_{m, n \geq 1} \int_0^t \int_0^t \varphi^{*m}(t-r) \varphi^{*n}(t-s) \\ &\quad \sum_{j, k=1}^N A_N^m(i, j) A_N^n(i, k) \mathbb{E}_\theta[M_r^{j,N} M_s^{k,N}] dr ds. \end{aligned}$$

But by Remark 10, $\mathbb{E}_\theta[M_r^{j,N} M_s^{k,N}] = \mathbf{1}_{\{j=k\}} \mathbb{E}_\theta[Z_{r \wedge s}^{j,N}] \leq C_\eta \mathbf{1}_{\{j=k\}} e^{(\alpha_0 + \eta)(r \wedge s)}$ thanks to Step 1. Furthermore, $\sum_{j=1}^N A_N^m(i, j) A_N^n(i, j) \leq \|A_N^m e_i\|_2 \|A_N^n e_i\|_2 \leq C N^{-1} \rho_N^{m+n}$ by Lemma 35-(vii) (because $m, n \geq 1$). We thus find, recalling that $\Gamma_t^N = \sum_{n \geq 0} \varphi^{*n}(t)$, that

$$\mathbb{E}_\theta[(J_t^{i,N})^2] = C_\eta N^{-1} \int_0^t \int_0^t \Gamma_{t-r}^N \Gamma_{t-s}^N e^{(\alpha_0 + \eta)(r \wedge s)} dr ds.$$

The conclusion follows from Lemma 43-(vi).

Step 3. Point (ii) follows from the facts that $U_t^{i,N} = M_t^{i,N} + J_t^{i,N}$, that $\mathbb{E}_\theta[(M_t^{i,N})^2] = \mathbb{E}_\theta[Z_t^{i,N}] \leq C_\eta e^{(\alpha_0 + \eta)t}$ by Remark 10 and Step 1 and that $\mathbb{E}_\theta[(J_t^{i,N})^2] \leq C_\eta N^{-1} e^{2(\alpha_0 + \eta)t}$ by Step 2.

Step 4. Since $Z_t^{i,N} = \mathbb{E}_\theta[Z_t^{i,N}] + U_t^{i,N}$, we deduce from Steps 1 and 3 that $\mathbb{E}_\theta[(Z_t^{i,N})^2] \leq C_\eta (e^{2(\alpha_0 + \eta)t} + e^{(\alpha_0 + \eta)t} + N^{-1} e^{2(\alpha_0 + \eta)t}) \leq C_\eta e^{2(\alpha_0 + \eta)t}$, whence point (i).

Step 5. Finally, we write $\bar{U}_t^N = \bar{M}_t^N + \bar{J}_t^N$. It is clear from Step 2 that $\mathbb{E}_\theta[(\bar{J}_t^N)^2] \leq C_\eta N^{-1} e^{2(\alpha_0+\eta)t}$. Moreover, Remark 10 implies that $\mathbb{E}_\theta[(\bar{M}_t^N)^2] = N^{-2} \sum_{i=1}^N \mathbb{E}_\theta[Z_t^{i,N}] \leq C_\eta N^{-1} e^{(\alpha_0+\eta)t}$ by Step 1. Point (iii) is checked. \square

We next show that the term \mathbf{I}_t^N is very small in the present scales.

Lemma 45. *Assume A. For all $\eta > 0$, there are $N_\eta \geq 1$ and $C_\eta > 0$ such that for all $N \geq N_\eta$, all $t \geq 0$, on Ω_N^2 , $\|\mathbf{I}_t^N\|_2 \leq C_\eta N^{1/8} t$.*

Proof. We fix $\eta > 0$ and work with N large enough and on Ω_N^2 , so that we can use Lemmas 35 and 43. Using the Minkowski inequality and then Lemma 35-(iii)-(v), we find

$$\begin{aligned} \|\mathbf{I}_t^N\|_2 &\leq \mu \sum_{n \geq 0} \left[\int_0^t s \varphi^{*n}(t-s) ds \right] \left\| A_N^n \mathbf{1}_N - \frac{\|A_N^n \mathbf{1}_N\|_2}{\sqrt{N}} V_N \right\|_2 \\ &\leq 6\mu t \sum_{n \geq 0} \left[\int_0^t \varphi^{*n}(t-s) ds \right] N^{1/2} \rho_N^n (2N^{-3/8})^{\lfloor n/2 \rfloor + 1} \\ &\leq 12\mu t N^{1/8} \sum_{n \geq 0} \rho_N^n (2N^{-3/8})^{\lfloor n/2 \rfloor} \int_0^t \varphi^{*n}(t-s) ds. \end{aligned}$$

The conclusion follows from Lemma 43-(iii). \square

We now study the empirical variance of \mathbf{J}_t^N .

Lemma 46. *Assume A. For all $\eta > 0$, there are $N_\eta \geq 1$ and $C_\eta > 0$ such that for all $N \geq N_\eta$, all $t \geq 0$, on Ω_N^2 , $\mathbb{E}_\theta[\|\mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N\|_2^2] \leq C_\eta [e^{(\alpha_0+\eta)t} + N^{-1} \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 e^{2(\alpha_0+\eta)t}]$.*

Proof. As usual, we fix $\eta > 0$ and work with N large enough and on Ω_N^2 , so that we can use Lemmas 35 and 43. Starting from (16) and using the Minkowski inequality, we find

$$\mathbb{E}_\theta[\|\mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N\|_2^2]^{1/2} \leq \sum_{n \geq 1} \int_0^t \varphi^{*n}(t-s) \mathbb{E}_\theta[\|A_N^n M_s^N - \overline{A_N^n M_s^N} \mathbf{1}_N\|_2^2]^{1/2} ds.$$

But using Remark 10 and then Lemma 44-(i), we see that

$$\begin{aligned} &\mathbb{E}_\theta[\|A_N^n M_s^N - \overline{A_N^n M_s^N} \mathbf{1}_N\|_2^2] \\ &= \sum_{i=1}^N \mathbb{E}_\theta \left[\left(\sum_{j=1}^N A_N^n(i,j) M_s^{j,N} - \frac{1}{N} \sum_{j,k=1}^N A_N^n(k,j) M_s^{j,N} \right)^2 \right] \\ &= \sum_{i,j=1}^N \left(A_N^n(i,j) - \frac{1}{N} \sum_{k=1}^N A_N^n(k,j) \right)^2 \mathbb{E}_\theta[Z_s^{j,N}] \\ &\leq C_\eta e^{(\alpha_0+\eta)s} \sum_{j=1}^N \|A_N^n e_j - \overline{A_N^n e_j} \mathbf{1}_N\|_2^2. \end{aligned}$$

Using next that, for all $x, y \in \mathbb{R}^N$, $|||x - \bar{x}\mathbf{1}_N|||_2 - |||y - \bar{y}\mathbf{1}_N|||_2 \leq |||x - y|||_2$ (with the notation $\bar{x} = N^{-1} \sum_{i=1}^N x_i$ and $\bar{y} = N^{-1} \sum_{i=1}^N y_i$), we write

$$\begin{aligned} |||A_N^n e_j - \overline{A_N^n e_j} \mathbf{1}_N|||_2 &\leq \left\| A_N^n e_j - \frac{|||A_N^n e_j|||_2}{\sqrt{N}} V_N \right\|_2 + \frac{|||A_N^n e_j|||_2}{\sqrt{N}} |||V_N - \bar{V}_N \mathbf{1}_N|||_2 \\ &\leq |||A_N^n e_j|||_2 \left(12(2N^{-3/8})^{\lfloor n/2 \rfloor} + \frac{|||V_N - \bar{V}_N \mathbf{1}_N|||_2}{\sqrt{N}} \right) \end{aligned}$$

by Lemma 35-(vi). Since $|||A_N^n e_j|||_2 \leq C\rho_N^n/\sqrt{N}$ by Lemma 35-(vii) (because $n \geq 1$), we conclude that

$$\begin{aligned} &\mathbb{E}_\theta[|||A_N^n M_s^N - \overline{A_N^n M_s^N} \mathbf{1}_N|||_2^2]^{1/2} \\ &\leq C_\eta e^{(\alpha_0 + \eta)s/2} \rho_N^n \left((2N^{-3/8})^{\lfloor n/2 \rfloor} + \frac{|||V_N - \bar{V}_N \mathbf{1}_N|||_2}{\sqrt{N}} \right). \end{aligned}$$

Consequently,

$$\begin{aligned} &\mathbb{E}_\theta[|||\mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N|||_2^2]^{1/2} \\ &\leq C_\eta \sum_{n \geq 1} \rho_N^n \left((2N^{-3/8})^{\lfloor n/2 \rfloor} + \frac{|||V_N - \bar{V}_N \mathbf{1}_N|||_2}{\sqrt{N}} \right) \int_0^t \varphi^{*n}(t-s) e^{(\alpha_0 + \eta)s/2} ds \\ &\leq C_\eta e^{(\alpha_0 + \eta)t/2} \sum_{n \geq 1} \rho_N^n (2N^{-3/8})^{\lfloor n/2 \rfloor} \int_0^t \varphi^{*n}(t-s) ds \\ &\quad + C_\eta \frac{|||V_N - \bar{V}_N \mathbf{1}_N|||_2}{\sqrt{N}} \sum_{n \geq 1} \rho_N^n \int_0^t \varphi^{*n}(t-s) e^{(\alpha_0 + \eta)s/2} ds \\ &\leq C_\eta e^{(\alpha_0 + \eta)t/2} + C_\eta \frac{|||V_N - \bar{V}_N \mathbf{1}_N|||_2}{\sqrt{N}} e^{(\alpha_0 + \eta)t} \end{aligned}$$

by Lemma 43-(iii)-(iv). This completes the proof. \square

The last lemma of the subsection concerns the martingale \mathbf{M}_t^N . In point (ii) below, (\cdot, \cdot) stands for the usual scalar product in \mathbb{R}^N .

Lemma 47. *Assume A. For all $\eta > 0$, there are $N_\eta \geq 1$ and $C_\eta > 0$ such that for all $N \geq N_\eta$, all $t \geq 0$, on Ω_N^2 ,*

- (i) $\mathbb{E}_\theta[|||\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N|||_2^2] \leq C_\eta N e^{(\alpha_0 + \eta)t}$,
- (ii) $\mathbb{E}_\theta[(\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N, V_N - \bar{V}_N \mathbf{1}_N)^2] \leq C_\eta |||V_N - \bar{V}_N \mathbf{1}_N|||_2^2 e^{(\alpha_0 + \eta)t}$,
- (iii) $\mathbb{E}_\theta[|X_t^N|] \leq C_\eta \sqrt{N} e^{(\alpha_0 + \eta)t}$, where $X_t^N = |||\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N|||_2^2 - N \bar{Z}_t^N$.

Proof. We fix $\eta > 0$ and work with N large enough and on Ω_N^2 , so that we can use Lemmas 35 and 43.

For point (ii), we write $\mathbb{E}_\theta[(\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N, V_N - \bar{V}_N \mathbf{1}_N)^2] = \mathbb{E}_\theta[(\mathbf{M}_t^N, V_N - \bar{V}_N \mathbf{1}_N)^2] = \mathbb{E}_\theta[(\sum_{i=1}^N (V_N(i) - \bar{V}_N) M_t^{i,N})^2]$. By Remark 10, this is nothing but $\sum_{i=1}^N (V_N(i) - \bar{V}_N)^2 \mathbb{E}_\theta[Z_t^{i,N}]$, which is controlled by $C_\eta |||V_N - \bar{V}_N \mathbf{1}_N|||_2^2 e^{(\alpha_0 + \eta)t}$ by Lemma 44-(i).

For point (iii), we first observe that $X_t^N = Y_t^N - N(\bar{M}_t^N)^2$, where $Y_t^N = \|\mathbf{M}_t^N\|_2^2 - N\bar{Z}_t^N$. Using as usual Remark 10, we deduce that $\mathbb{E}_\theta[N(\bar{M}_t^N)^2] = N^{-1} \sum_{i=1}^N \mathbb{E}_\theta[Z_t^{i,N}] \leq C_\eta e^{(\alpha_0+\eta)t}$ by Lemma 44-(i). Next, we see that $\|\mathbf{M}_t^N\|_2^2 = \sum_{i=1}^N (M_t^{i,N})^2 = 2 \sum_{i=1}^N \int_0^t M_{s-}^{i,N} dM_s^{i,N} + \sum_{i=1}^N Z_t^{i,N}$, since $[M^{i,N}, M^{i,N}]_t = Z_t^{i,N}$, see Remark 10. Thus $Y_t^N = 2 \sum_{i=1}^N \int_0^t M_{s-}^{i,N} dM_s^{i,N}$. But the martingales $\int_0^t M_{s-}^{i,N} dM_s^{i,N}$ are orthogonal and

$$\left[\int_0^\cdot M_{s-}^{i,N} dM_s^{i,N}, \int_0^\cdot M_{s-}^{i,N} dM_s^{i,N} \right]_t = \int_0^t (M_{s-}^{i,N})^2 dZ_s^{i,N} \leq Z_t^{i,N} \sup_{[0,t]} (M_s^{i,N})^2.$$

As a conclusion,

$$\begin{aligned} \mathbb{E}_\theta[(Y_t^N)^2] &= 4 \sum_{i=1}^N \mathbb{E}_\theta \left[\int_0^t (M_{s-}^{i,N})^2 dZ_s^{i,N} \right] \\ &\leq 4 \sum_{i=1}^N \mathbb{E}_\theta \left[(Z_t^{i,N})^2 \right]^{1/2} \mathbb{E}_\theta \left[\sup_{[0,t]} (M_s^{i,N})^4 \right]^{1/2}. \end{aligned}$$

Using again that $[M^{i,N}, M^{i,N}]_t = Z_t^{i,N}$ and the Doob inequality, we see that $\mathbb{E}_\theta[\sup_{[0,t]} (M_s^{i,N})^4] \leq C \mathbb{E}_\theta[(Z_t^{i,N})^2]$. All this shows that we have $\mathbb{E}_\theta[(Y_t^N)^2] \leq C \sum_{i=1}^N \mathbb{E}_\theta[(Z_t^{i,N})^2] \leq C_\eta N e^{2(\alpha_0+\eta)t}$ by Lemma 44-(i). As a consequence, we have $\mathbb{E}_\theta[|Y_t^N|] \leq C_\eta \sqrt{N} e^{(\alpha_0+\eta)t}$ and $\mathbb{E}_\theta[|X_t^N|] \leq \mathbb{E}_\theta[|Y_t^N|] + \mathbb{E}[N(\bar{M}_t^N)^2] \leq C_\eta \sqrt{N} e^{(\alpha_0+\eta)t}$.

Finally, (i) follows from (iii), since $\mathbb{E}_\theta[\|\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N\|_2^2] \leq \mathbb{E}_\theta[|X_t^N|] + N \mathbb{E}_\theta[\bar{Z}_t^N]$ and since $N \mathbb{E}_\theta[\bar{Z}_t^N] \leq C_\eta N e^{(\alpha_0+\eta)t}$ by Lemma 44-(i) again. \square

5.4. Conclusion

We now conclude the proof of Theorem 6. We recall that

$$\begin{aligned} \mathcal{U}_t^N &= \left[\sum_{i=1}^N \left(\frac{Z_t^{i,N} - \bar{Z}_t^N}{\bar{Z}_t^N} \right)^2 - \frac{N}{\bar{Z}_t^N} \right] \mathbf{1}_{\{\bar{Z}_t^N > 0\}} \\ &= \left[\frac{\|\mathbf{Z}_t^N - \bar{Z}_t^N \mathbf{1}_N\|_2^2 - N \bar{Z}_t^N}{(\bar{Z}_t^N)^2} \right] \mathbf{1}_{\{\bar{Z}_t^N > 0\}}, \end{aligned}$$

that V_N was introduced in Lemma 34 and that

$$\mathcal{U}_\infty^N = \sum_{i=1}^N \left(\frac{V_N(i) - \bar{V}_N}{\bar{V}_N} \right)^2 = \frac{\|V_N - \bar{V}_N \mathbf{1}_N\|_2^2}{(\bar{V}_N)^2}.$$

We first proceed to a suitable decomposition of the error.

Remark 48. Assume that $p \in (0, 1]$. We introduce $\mathcal{D}_t^N = |\mathcal{U}_t^N - (1/p - 1)|$ and recall that v_t^N was defined in (14). There is N_0 (depending only on p) such that for all $N \geq N_0$, on the event $\Omega_N^2 \cap \{\bar{Z}_t^N \geq v_t^N/4 > 0\}$,

$$\mathcal{D}_t^N \leq 16 \mathcal{D}_t^{N,1} + 128 \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 \mathcal{D}_t^{N,2} + |\mathcal{U}_\infty^N - (1/p - 1)|,$$

where

$$\mathcal{D}_t^{N,1} = \frac{1}{(v_t^N)^2} \left| \|\mathbf{Z}_t^N - \bar{Z}_t^N \mathbf{1}_N\|_2^2 - N\bar{Z}_t^N - (v_t^N)^2 \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 \right|$$

$$\text{and } \mathcal{D}_t^{N,2} = \left| \frac{\bar{Z}_t^N}{v_t^N} - \bar{V}_N \right|.$$

Proof. We work with N sufficiently large so that we can apply Lemma 35. We obviously have $\mathcal{D}_t^N \leq |\mathcal{U}_t^N - \mathcal{U}_\infty^N| + |\mathcal{U}_\infty^N - (1/p - 1)|$. We next write, on the event $\Omega_N^2 \cap \{\bar{Z}_t^N \geq v_t^N/4 > 0\}$,

$$|\mathcal{U}_t^N - \mathcal{U}_\infty^N| \leq \frac{1}{(\bar{Z}_t^N)^2} \left| \|\mathbf{Z}_t^N - \bar{Z}_t^N \mathbf{1}_N\|_2^2 - N\bar{Z}_t^N - (v_t^N)^2 \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 \right|$$

$$+ \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 \left| \left(\frac{v_t^N}{\bar{Z}_t^N} \right)^2 - \frac{1}{(\bar{V}_N)^2} \right|$$

$$\leq 16\mathcal{D}_t^{N,1} + 128 \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 \mathcal{D}_t^{N,2}.$$

We used that on the present event, $(\bar{Z}_t^N)^{-2} \leq 16(v_t^N)^{-2}$, that $\bar{V}_N \geq 1/2$ (see Lemma 35-(ii)), that $(\bar{Z}_t^N/v_t^N) \geq 1/4$ and that, for all $x, y \geq 1/4$, $|x^{-2} - y^{-2}| \leq 128|x - y|$. \square

We now treat the term $\mathcal{D}_t^{N,2}$.

Lemma 49. *Assume A. For all $\eta > 0$, there are $N_\eta \geq 1$, $t_\eta \geq 0$ and $C_\eta > 0$ such that, for all $N \geq N_\eta$, all $t \geq t_\eta$, on Ω_N^2 ,*

- (i) $\mathbb{E}_\theta[\mathcal{D}_t^{N,2}] \leq C_\eta e^{2\eta t} (N^{-1/2} + e^{-\alpha_0 t})$,
- (ii) $\Pr_\theta(\bar{Z}_t^N \leq v_t^N/4) \leq C_\eta e^{2\eta t} (N^{-1/2} + e^{-\alpha_0 t})$.

Proof. As usual, we fix $\eta > 0$ and consider $N \geq N_\eta$ and $t \geq t_\eta$ and we work on Ω_N^2 so that we can apply Lemmas 35 and 43.

Recalling (12)-(13), we write $\mathbf{Z}_t^N = \mathbb{E}_\theta[\mathbf{Z}_t^N] + \mathbf{U}_t^N = v_t^N V_N + \mathbf{I}_t^N + \mathbf{U}_t^N$, whence $\mathcal{D}_t^{N,2} \leq (v_t^N)^{-1} (|\bar{I}_t^N| + |\bar{U}_t^N|)$. But we infer from Lemma 45 that $|\bar{I}_t^N| \leq N^{-1/2} \|\mathbf{I}_t^N\|_2 \leq C_\eta t N^{1/8-1/2}$, which is obviously bounded by $C_\eta e^{\eta t}$. Next, we know from Lemma 44-(iii) that $\mathbb{E}_\theta[|\bar{U}_t^N|] \leq C_\eta N^{-1/2} e^{(\alpha_0 + \eta)t}$. We deduce that $\mathbb{E}_\theta[\mathcal{D}_t^{N,2}] \leq C_\eta (v_t^N)^{-1} e^{\eta t} [1 + N^{-1/2} e^{\alpha_0 t}]$. But since $t \geq t_\eta$, we know from Lemma 43-(ii) that $v_t^N \geq c_\eta e^{(\alpha_0 - \eta)t}$. This completes the proof of (i).

By Lemma 35-(ii), $\bar{V}_N \geq 1/2$. Thus $\bar{Z}_t^N \leq v_t^N/4$ implies that $\mathcal{D}_t^{N,2} = |\bar{Z}_t^N/v_t^N - \bar{V}_N| \geq 1/4$. Hence $\Pr_\theta(\bar{Z}_t^N \leq v_t^N/4) \leq 4\mathbb{E}_\theta[\mathcal{D}_t^{N,2}]$ and (ii) follows from (i). \square

Lemma 50. *Assume A. For all $\eta > 0$, there are $N_\eta \geq 1$, $t_\eta \geq 0$ and $C_\eta > 0$ such that, for all $N \geq N_\eta$, all $t \geq t_\eta$, on Ω_N^2 ,*

$$\mathbb{E}_\theta[\mathcal{D}_t^{N,1}] \leq C_\eta (1 + \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2) e^{4\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} + \left(\frac{\sqrt{N}}{e^{\alpha_0 t}} \right)^{3/2} \right).$$

Proof. We fix $\eta > 0$ and consider $N \geq N_\eta$ and $t \geq t_\eta$ and we work on Ω_N^2 so that we can apply Lemmas 35 and 43. Recalling (12)-(13), we write $\mathbf{Z}_t^N = v_t^N V_N + \mathbf{I}_t^N + M_t^N + \mathbf{J}_t^N$ and

$$\begin{aligned} \mathcal{D}_t^{N,1} &= \frac{1}{(v_t^N)^2} \left| \|\mathbf{I}_t^N - \bar{I}_t^N \mathbf{1}_N + \mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N\|_2^2 + \|\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N\|_2^2 - N \bar{Z}_t^N \right. \\ &\quad + 2(\mathbf{I}_t^N - \bar{I}_t^N \mathbf{1}_N + \mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N, v_t^N (V_N - \bar{V}_N \mathbf{1}_N) + \mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N) \\ &\quad \left. + 2v_t^N (V_N - \bar{V}_N \mathbf{1}_N, \mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N) \right|. \end{aligned}$$

Recalling that $X_t^N = \|\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N\|_2^2 - N \bar{Z}_t^N$, see Lemma 47, we deduce that

$$\begin{aligned} \mathcal{D}_t^{N,1} &\leq \frac{1}{(v_t^N)^2} \left| 2\|\mathbf{I}_t^N - \bar{I}_t^N \mathbf{1}_N\|_2^2 + 2\|\mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N\|_2^2 + |X_t^N| \right. \\ &\quad + 2(\|\mathbf{I}_t^N - \bar{I}_t^N \mathbf{1}_N\|_2 + \|\mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N\|_2) \\ &\quad \quad \quad \times (v_t^N \|V_N - \bar{V}_N \mathbf{1}_N\|_2 + \|\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N\|_2) \\ &\quad \left. + 2v_t^N |(V_N - \bar{V}_N \mathbf{1}_N, \mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N)| \right|. \end{aligned}$$

We know from Lemma 43-(i)-(ii) that $v_t^N \geq c_\eta e^{(\alpha_0 - \eta)t}$ and $v_t^N \leq C_\eta e^{(\alpha_0 + \eta)t}$ and from Lemma 45 that $\|\mathbf{I}_t^N - \bar{I}_t^N \mathbf{1}_N\|_2 \leq \|\mathbf{I}_t^N\|_2 \leq C_\eta N^{1/8} t \leq C_\eta N^{1/8} e^{\eta t}$. Lemma 46 tells us that that $\mathbb{E}_\theta[\|\mathbf{J}_t^N - \bar{J}_t^N \mathbf{1}_N\|_2^2] \leq C_\eta [e^{(\alpha_0 + \eta)t} + N^{-1} \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2 e^{2(\alpha_0 + \eta)t}]$ and Lemma 47 tells us that $\mathbb{E}_\theta[|X_t^N|] \leq C_\eta \sqrt{N} e^{(\alpha_0 + \eta)t}$, $\mathbb{E}_\theta[\|\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N\|_2^2] \leq C_\eta N e^{(\alpha_0 + \eta)t}$ and also that $\mathbb{E}_\theta[(\mathbf{M}_t^N - \bar{M}_t^N \mathbf{1}_N, V_N - \bar{V}_N \mathbf{1}_N)] \leq C_\eta \|V_N - \bar{V}_N \mathbf{1}_N\|_2 e^{(\alpha_0 + \eta)t/2}$. Using the Cauchy-Schwarz inequality, we find

$$\begin{aligned} \mathbb{E}_\theta[\mathcal{D}_t^{N,1}] &\leq C_\eta (1 + \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2) e^{-2(\alpha_0 - \eta)t} \\ &\quad \times \left(N^{1/4} e^{2\eta t} + [e^{(\alpha_0 + \eta)t} + N^{-1} e^{2(\alpha_0 + \eta)t}] + N^{1/2} e^{(\alpha_0 + \eta)t} \right. \\ &\quad \quad \quad + [N^{1/8} e^{\eta t} + e^{(\alpha_0 + \eta)t/2} + N^{-1/2} e^{(\alpha_0 + \eta)t}] [e^{(\alpha_0 + \eta)t} + N^{1/2} e^{(\alpha_0 + \eta)t/2}] \\ &\quad \left. + e^{3(\alpha_0 + \eta)t/2} \right). \end{aligned}$$

We easily deduce that

$$\begin{aligned} \mathbb{E}_\theta[\mathcal{D}_t^{N,1}] &\leq C_\eta (1 + \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2) e^{4\eta t} \\ &\quad \times \left(N^{-1/2} + e^{-\alpha_0 t/2} + N^{1/2} e^{-\alpha_0 t} + N^{5/8} e^{-3\alpha_0 t/2} \right). \end{aligned}$$

To conclude, it suffices to notice that $e^{-\alpha_0 t/2} \leq N^{-1/2} + N^{1/2} e^{-\alpha_0 t}$ and that $N^{5/8} e^{-3\alpha_0 t/2} \leq N^{3/4} e^{-3\alpha_0 t/2} = (N^{1/2} e^{-\alpha_0 t})^{3/2}$. \square

We now have all the weapons to give the

Proof of Theorem 6. We assume A and fix $\eta > 0$.

Step 1. Starting from Remark 48 and using Lemmas 49 and 50, we deduce that there is $N_\eta \geq 1$, $t_\eta \geq 0$ and $C_\eta > 0$ such that for all $N \geq N_\eta$, all $t \geq t_\eta$,

$$\begin{aligned} \mathbf{1}_{\Omega_N^2} \mathbb{E}_\theta \left[\mathbf{1}_{\{\bar{Z}_t^N \geq v_t^N/4 > 0\}} \left| \mathcal{U}_t^N - \left(\frac{1}{p} - 1\right) \right| \right] &\leq \mathbf{1}_{\Omega_N^2} \left| \mathcal{U}_\infty^N - \left(\frac{1}{p} - 1\right) \right| \\ &+ C_\eta \mathbf{1}_{\Omega_N^2} (1 + \|V_N - \bar{V}_N \mathbf{1}_N\|_2^2) e^{4\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} + \left(\frac{\sqrt{N}}{e^{\alpha_0 t}}\right)^{3/2} \right), \end{aligned}$$

which implies, by Proposition 40, that

$$\mathbb{E} \left[\mathbf{1}_{\Omega_N^2 \cap \{\bar{Z}_t^N \geq v_t^N/4 > 0\}} \left| \mathcal{U}_t^N - \left(\frac{1}{p} - 1\right) \right| \right] \leq C_\eta e^{4\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} + \left(\frac{\sqrt{N}}{e^{\alpha_0 t}}\right)^{3/2} \right)$$

and thus, for all $\varepsilon \in (0, 1)$,

$$\begin{aligned} &\Pr \left(\Omega_N^2, \bar{Z}_t^N \geq v_t^N/4 > 0, \left| \mathcal{U}_t^N - \left(\frac{1}{p} - 1\right) \right| \geq \varepsilon \right) \\ &\leq \frac{C_\eta}{\varepsilon} e^{4\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} + \left(\frac{\sqrt{N}}{e^{\alpha_0 t}}\right)^{3/2} \right) \\ &\leq \frac{C_\eta}{\varepsilon} e^{4\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} \right). \end{aligned}$$

For the last inequality, we used that either $N^{1/2}e^{-\alpha_0 t} \geq 1$ and then the inequality is trivial or $N^{1/2}e^{-\alpha_0 t} \leq 1$ and then $(N^{1/2}e^{-\alpha_0 t})^{3/2} \leq N^{1/2}e^{-\alpha_0 t}$. But we know from Lemma (43)-(ii) that $v_t^N > 0$ on Ω_N^2 (because $t \geq t_\eta$) and from Lemmas 33 and 49-(ii) that

$$\begin{aligned} \Pr((\Omega_N^2)^c \text{ or } \bar{Z}_t^N < v_t^N/4) &\leq C e^{-cN^{1/4}} + C_\eta e^{2\eta t} \left(\frac{1}{\sqrt{N}} + e^{-\alpha_0 t} \right) \\ &\leq C_\eta e^{2\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} \right), \end{aligned}$$

whence finally,

$$\Pr \left(\left| \mathcal{U}_t^N - \left(\frac{1}{p} - 1\right) \right| \geq \varepsilon \right) \leq \frac{C_\eta}{\varepsilon} e^{4\eta t} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} \right).$$

We have proved this inequality only for $N \geq N_\eta$ and $t \geq t_\eta$, but it obviously extends, enlarging C_η is necessary, to any values of $N \geq 1$ and $t \geq 0$.

Step 2. We next recall that $\mathcal{P}_t^N = \Phi(\mathcal{U}_t^N)$, where $\Phi(u) = (1+u)^{-1} \mathbf{1}_{\{u \geq 0\}}$ and observe that $p = \Phi(1/p - 1)$. The function Φ is Lipschitz continuous on $[0, \infty)$ with Lipschitz constant 1. Thus for any $\varepsilon \in (0, 1)$, $|\mathcal{P}_t^N - p| > \varepsilon$ implies that either $|\mathcal{U}_t^N - (1/p - 1)| > \varepsilon$ or $\mathcal{U}_t^N < 0$, so that in any case, $|\mathcal{U}_t^N - (1/p - 1)| > \min\{\varepsilon, (1/p - 1)\}$. We thus conclude from Step 1 that for any $N \geq 1$, any $t \geq 0$, any $\varepsilon \in (0, 1)$,

$$\Pr(|\mathcal{P}_t^N - p| \geq \varepsilon) \leq \frac{C_\eta e^{4\eta t}}{\min\{\varepsilon, (1/p - 1)\}} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} \right) \leq \frac{C_\eta e^{4\eta t}}{\varepsilon} \left(\frac{1}{\sqrt{N}} + \frac{\sqrt{N}}{e^{\alpha_0 t}} \right).$$

The proof is complete. □

Finally, we handle the

Proof of Remark 5. We assume A and fix $\eta > 0$. We know from Lemma 49-(i) that for all $N \geq N_\eta$, $t \geq t_\eta$, $\mathbf{1}_{\Omega_N^2} \mathbb{E}_\theta [|(\bar{Z}_t^N/v_t^N) - \bar{V}_N|] \leq C_\eta e^{2\eta t} (N^{-1/2} +$

$e^{-\alpha_0 t}$), from Lemma 35-(ii) that $\bar{V}_N \in [1/2, 2]$ (on Ω_N^2) and from Lemma 33 that $\Pr(\Omega_N^2) \geq 1 - Ce^{-cN^{1/4}}$. We also know from Lemma 43-(i)-(ii) that on Ω_N^2 , there are $0 < a_\eta < b_\eta$ such that $v_t^N \in [a_\eta e^{(\alpha_0 - \eta)t}, b_\eta e^{(\alpha_0 + \eta)t}]$. We easily deduce that, still for $N \geq N_\eta$ and $t \geq t_\eta$,

$$\Pr(\bar{Z}_t^N \notin [(a_\eta/2)e^{(\alpha_0 - \eta)t}, 2b_\eta e^{(\alpha_0 + \eta)t}]) \leq Ce^{-cN^{1/4}} + C_\eta e^{2\eta t}(N^{-1/2} + e^{-\alpha_0 t}).$$

Hence $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} \Pr(\bar{Z}_t^N \in [(a_\eta/2)e^{(\alpha_0 - \eta)t}, 2b_\eta e^{(\alpha_0 + \eta)t}]) = 1$ for any $\eta \in (0, \alpha_0/2)$. We deduce that $\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} \Pr(\bar{Z}_t^N \in [e^{(\alpha_0 - \eta)t}, e^{(\alpha_0 + \eta)t}]) = 1$ for any $\eta > 0$. \square

6. Detecting subcriticality and supercriticality

Proof of Proposition 7. We first assume $H(1)$. We then know from Lemma 16-(i) with $r = 1$ that, on Ω_N^1 , $\mathbb{E}_\theta[\bar{Z}_t^N] \leq Ct < e^{(\log t)^2}/2$ for all t large enough, say for all $t \geq t_0$. We also know from Lemma 18 that, still on Ω_N^1 , $\mathbb{E}_\theta[|\bar{Z}_t^N - \mathbb{E}_\theta[\bar{Z}_t^N]|] = \mathbb{E}_\theta[|\bar{U}_t^N|] \leq C(t/N)^{1/2}$ and from Lemma 13 that $\Pr[(\Omega_N^1)^c] \leq Ce^{-cN}$. We easily deduce that

$$\begin{aligned} & \Pr(\log(\bar{Z}_t^N) \geq (\log t)^2) \\ & \leq \Pr((\Omega_N^1)^c) + \Pr(\Omega_N^1, \mathbb{E}_\theta[\bar{Z}_t^N] \geq e^{(\log t)^2}/2 \text{ or } |\bar{Z}_t^N - \mathbb{E}_\theta[\bar{Z}_t^N]| \geq e^{(\log t)^2}/2) \\ & \leq Ce^{-cN} + C(t/N)^{1/2} e^{-(\log t)^2} \end{aligned}$$

for all $t \geq t_0$. Enlarging C if necessary, we deduce that $\Pr(\log(\bar{Z}_t^N) \geq (\log t)^2) \leq Ce^{-cN} + Ct^{1/2} e^{-(\log t)^2}$ for all $t \geq 1$,

We next assume A and we fix $\eta \in (0, \alpha_0)$. We know from Lemma 49-(ii) that for all $N \geq N_\eta$ and $t \geq t_\eta$, on Ω_N^2 , $\Pr_\theta(\bar{Z}_t^N \leq v_t^N/4) \leq C_\eta e^{2\eta t}(N^{-1/2} + e^{-\alpha_0 t})$, from Lemma 43-(i)-(ii) that, still on Ω_N^2 , $v_t^N \geq c_\eta e^{(\alpha_0 - \eta)t} \geq 4e^{(\log t)^2}$ (enlarging the value of t_η if necessary). Finally, Lemma 33 tells us that $\Pr((\Omega_N^2)^c) \leq Ce^{-cN^{1/4}}$. We thus see that

$$\begin{aligned} \Pr(\log(\bar{Z}_t^N) \leq (\log t)^2) & \leq \Pr((\Omega_N^2)^c) + \Pr(\Omega_N^2, \bar{Z}_t^N \leq v_t^N/4) \\ & \leq Ce^{-cN^{1/4}} + C_\eta e^{2\eta t}(N^{-1/2} + e^{-\alpha_0 t}) \\ & \leq C_\eta e^{2\eta t}(N^{-1/2} + e^{-\alpha_0 t}). \end{aligned}$$

All this shows that for all $\eta \in (0, \alpha_0)$, we can find C_η and t_η such that for all $t \geq t_\eta$ and all $N \geq N_\eta$, $\Pr(\log(\bar{Z}_t^N) \leq (\log t)^2) \leq C_\eta e^{2\eta t}(N^{-1/2} + e^{-\alpha_0 t})$. We easily conclude that for all $\eta > 0$, there is C_η such that for all $N \geq 1$ and all $t \geq 1$, $\Pr(\log(\bar{Z}_t^N) \leq (\log t)^2) \leq C_\eta e^{2\eta t}(N^{-1/2} + e^{-\alpha_0 t})$ as desired. \square

7. Numerics

We say that we are in the *independent case* when the family $(\theta_{ij})_{1 \leq i, j \leq N}$ is i.i.d. and Bernoulli(p)-distributed, as in the whole paper. We say we are in

the *symmetric case* when the family $(\theta_{ij})_{1 \leq i \leq j \leq N}$ is i.i.d. and Bernoulli(p)-distributed and when $\theta_{ji} = \theta_{ij}$ for all $1 \leq i < j \leq N$. We will see that this does not change much the numerical results (with the very same estimators). Also, we assume that we observe only $(Z_s^{i,N})_{s \in [0,T], i=1, \dots, K}$ for some (large) K smaller than N . The theoretical results of this paper only apply when $K = N$. We adapt the estimators as follows. We introduce $\bar{Z}_t^{N,K} = K^{-1} \sum_{i=1}^K Z_t^{i,N}$ and

$$\mathcal{E}_t^{N,K} = \frac{\bar{Z}_{2t}^{N,K} - \bar{Z}_t^{N,K}}{t}, \quad \mathcal{V}_t^{N,K} = \frac{N}{K} \sum_{i=1}^K \left(\frac{Z_{2t}^{i,N} - Z_t^{i,N}}{t} - \mathcal{E}_t^{N,K} \right)^2 - \frac{N}{t} \mathcal{E}_t^{N,K},$$

$$\mathcal{W}_{\Delta,t}^{N,K} = 2\mathcal{Z}_{2\Delta,t}^{N,K} - \mathcal{Z}_{\Delta,t}^{N,K} \text{ where } \mathcal{Z}_{\Delta,t}^{N,K} = \frac{N}{t} \sum_{k=t/\Delta+1}^{2t/\Delta} \left(\bar{Z}_{k\Delta}^{N,K} - \bar{Z}_{(k-1)\Delta}^{N,K} - \Delta \mathcal{E}_t^{N,K} \right)^2,$$

as well as

$$\mathcal{P}_{\Delta,T}^{sub,N,K} = \Phi_3 \left(\mathcal{E}_{T/2}^{N,K}, \mathcal{V}_{T/2}^{N,K}, \left| \mathcal{W}_{\Delta,T/2}^{N,K} - \frac{N-K}{K} \mathcal{E}_{T/2}^{N,K} \right| \right),$$

with Φ defined in Corollary 4. We added the absolute value around the last argument of Φ_3 for practical reasons: by this way, $\mathcal{P}_{\Delta,T}^{sub,N,K}$ is always well-defined (and seems closer to the reality than Ψ_3 which is 0 when $w \leq 0$). This does not change the theory (since $\mathcal{W}_{\Delta,T/2}^{N,K} - \frac{N-K}{K} \mathcal{E}_{T/2}^{N,K}$ is asymptotically positive, at least when $N = K$). We also put

$$\mathcal{U}_T^{N,K} = \left[\frac{N}{K} \sum_{i=1}^K \left(\frac{Z_T^{i,N} - \bar{Z}_T^{N,K}}{\bar{Z}_T^{N,K}} \right)^2 - \frac{N}{\bar{Z}_T^{N,K}} \right] \mathbf{1}_{\{\bar{Z}_T^{N,K} > 0\}}$$

and $\mathcal{P}_T^{sup,N,K} = \frac{1}{\mathcal{U}_T^{N,K} + 1} \mathbf{1}_{\{\mathcal{U}_T^{N,K} \geq 0\}}.$

We set $\Delta_t = t/(2\lceil t^{9/13} \rceil)$, which corresponds to the (quite arbitrary) choice $q = 12$, and

$$\hat{p}_T^{N,K} = \mathcal{P}_{\Delta_T,T}^{sub,N,K} \mathbf{1}_{\{\log(\bar{Z}_T^{N,K}) < (\log T)^2\}} + \mathcal{P}_T^{sup,N,K} \mathbf{1}_{\{\log(\bar{Z}_T^{N,K}) > (\log T)^2\}}.$$

7.1. Choice of the estimators

Let us explain briefly how we have modified the estimators when observing only $(Z_s^{i,N})_{s \in [0,T], i=1, \dots, K}$. We adopt the notation of Section 2, in particular $A_N(i, j) = N^{-1} \theta_{ij}$, and we follow the considerations therein.

In the subcritical case, we recall that $Q_N = (I - \Lambda A_N)^{-1}$ and that $\ell_N(i) = \sum_{j=1}^N Q_N(i, j)$. Following closely the argumentation of Subsection 2.1, we expect that, for t (and Δ) large and in a suitable regime, we should have $\mathcal{E}_t^{N,K} \simeq \mu \bar{\ell}_N^K$ (where $\bar{\ell}_N^K = K^{-1} \sum_{i=1}^K \ell_N(i)$), $\mathcal{V}_t^{N,K} \simeq \mu^2 (N/K) \sum_{i=1}^K (\ell_N(i) - \bar{\ell}_N^K)^2$, and $\mathcal{W}_{\Delta,t}^{N,K} \simeq \mu (N/K^2) \sum_{j=1}^N (\sum_{i=1}^K Q_N(i, j))^2 \ell_N(j)$. Recalling now that $\ell_N(i) \simeq$

$1 + \Lambda(1 - \Lambda p)^{-1} L_N(i)$ and that NL_N is a vector composed of i.i.d. Binomial(N, p) random variables, we expect that $\mathcal{E}_t^{N,K} \simeq \mu \mathbb{E}[\ell_N(1)] \simeq \mu/(1 - \Lambda p)$ and $\mathcal{V}_t^{N,K} \simeq \mu^2 N \text{Var}(\ell_N(1)) \simeq \mu^2 \Lambda^2 p(1 - p)/(1 - \Lambda p)^2$ for N, K and t large. For the last estimator, one first has to get convinced, following again the arguments of Subsection 2.1, that $\sum_{i=1}^K Q_N(i, j) \simeq 1 + (K/N)\Lambda p/(1 - \Lambda p)$ if $j \in \{1, \dots, K\}$ while $\sum_{i=1}^K Q_N(i, j) \simeq (K/N)(\Lambda p/(1 - \Lambda p))$ if $j \in \{K + 1, \dots, N\}$. Since we still have $\ell_N(j) \simeq 1 + \Lambda p/(1 - \Lambda p) = 1/(1 - \Lambda p)$, we find that

$$\begin{aligned} \mathcal{W}_{\Delta,t}^{N,K} &\simeq \frac{\mu N}{K^2(1 - \Lambda p)} \left(K \left[1 + \frac{K\Lambda p}{N(1 - \Lambda p)} \right]^2 + (N - K) \left[\frac{K\Lambda p}{N(1 - \Lambda p)} \right]^2 \right) \\ &= \frac{\mu}{(1 - \Lambda p)^3} + \frac{(N - K)\mu}{K(1 - \Lambda p)}. \end{aligned}$$

Recalling that $\mathcal{E}_t^{N,K} \simeq \mu/(1 - \Lambda p)$, we conclude that $\mathcal{W}_{\Delta,t}^{N,K} - (N - K)\mathcal{E}_t^{N,K}/K \simeq \mu/(1 - \Lambda p)^3$. For N, K, t and Δ large, we thus should have

$$\begin{aligned} &\Phi_3 \left(\mathcal{E}_t^{N,K}, \mathcal{V}_t^{N,K}, \left| \mathcal{W}_{\Delta,t}^{N,K} - \frac{N - K}{K} \mathcal{E}_t^{N,K} \right| \right) \\ &\simeq \Phi_3 \left(\frac{\mu}{1 - \Lambda p}, \frac{\mu^2 \Lambda^2 p(1 - p)}{(1 - \Lambda p)^2}, \frac{\mu}{(1 - \Lambda p)^3} \right) = p. \end{aligned}$$

We introduce the conjectured limit of $\mathcal{P}_{\Delta,t}^{sub,N,K}$ as $t \rightarrow \infty$:

$$\begin{aligned} \mathcal{P}_{\infty,\infty}^{sub,N,K} &= \Phi_3 \left(\mu \bar{\ell}_N^K, \frac{\mu^2 N}{K} \sum_{i=1}^K (\ell_N(i) - \bar{\ell}_N^K)^2, \right. \\ &\quad \left. \left| \frac{\mu N}{K^2} \sum_{j=1}^N \left(\sum_{i=1}^K Q_N(i, j) \right)^2 \ell_N(j) - \frac{N - K}{K} \mu \bar{\ell}_N^K \right| \right). \end{aligned}$$

In the supercritical case, we follow Subsection 2.2 and deduce that for t large, we should have $\mathcal{U}_t^{N,K} \simeq (N/K)(\bar{V}_N^K)^{-2} \sum_{i=1}^K (V_N(i) - \bar{V}_N^K)^2$, where V_N is the Perron-Frobenius eigenvector of A_N and where $\bar{V}_N^K = K^{-1} \sum_{i=1}^K V_N(i)$. Recalling now that V_N is almost colinear to L_N and that NL_N is a vector composed of i.i.d. Binomial(N, p) random variables, we conclude that indeed, it should hold that $\mathcal{U}_t^{N,K} \simeq N(\mathbb{E}[V_N(1)])^{-2} \text{Var}(V_N(1)) \simeq 1/p - 1$ for N, K and t large, whence $\mathcal{P}_t^{sup,N,K} \simeq p$. Here we introduce the conjectured limit of $\mathcal{P}_t^{sup,N,K}$ as $t \rightarrow \infty$:

$$\mathcal{P}_{\infty}^{sup,N,K} = \left(1 + \frac{N}{K(\bar{V}_N^K)^2} \sum_{i=1}^K (V_N(i) - \bar{V}_N^K)^2 \right)^{-1}.$$

7.2. Numerical results

From now on, we assume that $\varphi(t) = a \exp(-bt)$ for some $a, b > 0$, which satisfies all our assumptions and is easy to simulate. We also always assume that $a = 2$ and $b = 1$, whence $\Lambda = 2$. We did not find interesting different behaviors when using other values.

On all the pictures below but the two last ones, we plot the time evolution of the three quartiles, using 1000 simulations, of $\hat{p}_t^{N,K} - p$, as a function of time $t \in [0, T]$. We always choose T in such a way that $\bar{Z}_T^N \simeq 3000$, so that on the right of all the pictures below, we always have more or less the same quantity of data (for a given value of K). The median is plotted in black and the two other quartiles (25% and 75%) are plotted in red.

Except in the case where $p = 0$ and $\mu = 1$, we always use the same range for the y coordinate, from -0.1 to 0.1 for the This explains why some curves are off the chart. An exception is the case where $p = 0$.


For a given simulation, we say that the choice is *good* when $\hat{p}_t^{N,K} = \mathcal{P}_{\Delta_t,t}^{sub,N,K}$ and $\Lambda p < 1$ or $\hat{p}_t^{N,K} = \mathcal{P}_t^{sup,N,K}$ and $\Lambda p > 1$. When the choice is almost always good (that is, for a large proportion of the simulations), we also indicate below the picture the three quartiles of $\hat{p}_\infty^{N,K} - p$, where $\hat{p}_\infty^{N,K}$ is the conjectured limit as $t \rightarrow \infty$ of $\hat{p}_t^{N,K}$, given by $\hat{p}_\infty^{N,K} = \mathcal{P}_{\infty,\infty}^{sub,N,K}$ when $\Lambda p < 1$ and $\hat{p}_\infty^{N,K} = \mathcal{P}_\infty^{sup,N,K}$ when $\Lambda p > 1$.

7.2.1. Estimation of p in the independent case


We start with the independent case. As the pictures below show, the estimation of p is more precise when p is large. Also, around the critical case, $\hat{p}_t^{N,K}$ is far from always making the good choice, but this does not, however, produce too bad results. On the contrary, the estimation of p when p is very small does not work very well. We did not investigate the case where p is very small and $\Lambda p > 1$ and we do not know what would happen. This would presumably require a lot of work.

Observe that the results with $N = K = 1000$ are most often not better than those with $N = K = 250$. This is not so surprising for a given value of T , since our rate of convergence resembles $T^{-1}N^{1/2} + N^{-1/2}$ in the subcritical case and something similar in the supercritical case.

Finally, in all the trials below, it seems that $|\hat{p}_\infty^{N,K} - p|$ is much smaller than $|\hat{p}_t^{N,K} - p|$.




$p = 0.65, \mu = 1$ (supercritical). The choice between $\mathcal{P}_{\Delta_t,t}^{sub,N,K}$ and $\mathcal{P}_t^{sup,N,K}$ is always bad for $t \in [14, 19]$.




However, the error (using the wrong estimator) does not seem so large. The jumps correspond to jumps of $t \mapsto \Delta_t$. This is particularly visible on these pictures because the time interval is short. Let us mention that the choice becomes good around $t = 22$.

$p = 0.51, \mu = 1$ (slightly supercritical). The choice between $\mathcal{P}_{\Delta_t,t}^{sub,N,K}$ and $\mathcal{P}_t^{sup,N,K}$ is always bad for $t \in [9, 62]$.




However, the error (using the wrong estimator) does not seem so large.

$p = 0.48, \mu = 20$ (slightly subcritical). The choice between $\mathcal{P}_{\Delta_t,t}^{sub,N,K}$ and $\mathcal{P}_t^{sup,N,K}$ is always bad for $t \in [1, 15]$ and always good for $t \in [17, 20]$.




We clearly see the change of choice around $t = 16$.

$p = 0.35, \mu = 1$ (fairly subcritical). The choice between $\mathcal{P}_{\Delta_t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always good for $t \in (0, 900]$.




These results show that the bias is rather large, of the same order as the standard deviation.

$p = 0.1, \mu = 1$ (fairly subcritical). The choice between $\mathcal{P}_{\Delta_t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always good.



These pictures illustrate that this situation (p small) is not quite favorable.

$p = 0, \mu = 1$ (subcritical). The choice between $\mathcal{P}_{\Delta_t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always good.




This is catastrophic. The 25% quartile is below 0. One can use $\hat{p}_t^{N, K} \vee 0$ instead of $\hat{p}_t^{N, K}$.


7.2.2. Estimation of p in the symmetric case

We obtain very similar numerical results. When $p = 0$, the symmetric and independent cases are precisely the same so that we do not plot it again.


$p = 0.85, \mu = 1$ (fairly supercritical). The choice between $\mathcal{P}_{\Delta_t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always good for $t \in [1, 9.7]$.




$p = 0.65, \mu = 1$ (supercritical). The choice between $\mathcal{P}_{\Delta_t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always bad for $t \in [14, 19]$.




$p = 0.51, \mu = 1$ (slightly supercritical). The choice between $\mathcal{P}_{\Delta_t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always bad for $t \in [9, 62]$.




$p = 0.48, \mu = 20$ (slightly subcritical but large μ). The choice between $\mathcal{P}_{\Delta t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always bad for $t \in [1, 15]$ and always good for $t \in [17, 20]$.



$p = 0.35, \mu = 1$ (fairly subcritical). The choice between $\mathcal{P}_{\Delta t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always good for $t \in (0, 900]$.




$p = 0.1, \mu = 1$ (fairly subcritical). The choice between $\mathcal{P}_{\Delta t, t}^{sub, N, K}$ and $\mathcal{P}_t^{sup, N, K}$ is always good.



7.2.3. Practical choice of Δ

Independent case, $\mu = 1$, $p = 0.35$ (fairly subcritical), with $T = 900$ and $N = K = 1000$. On the left, we have plotted $\mathcal{P}_{\Delta,T}^{sub,N,K} - p$ as a function of $\Delta \in [1, 15]$ obtained with one simulation. On the right, we have plotted the quartiles of the same quantity using 1000 simulations.



Our (arbitrary) choice $\Delta_T = T/(2\lfloor T^{9/13} \rfloor) \simeq 4.1$ seems rather suitable: we see on the right picture that the “optimal” Δ lies between 4 and 6. This is mainly due to chance and probably depends strongly on the parameters of the model. We see on the left picture that given one set of data, $\mathcal{P}_{\Delta,T}^{sub,N,K}$ varies a lot.

References

- [1] Y. Aït-Sahalia, J. Cacho-Diaz, R. J. A. Laeven, *Modeling financial contagion using mutually exciting jump processes*, to appear in the Journal of Financial Economics.
- [2] E. Bacry, S. Delattre, M. Hoffmann, J. F. Muzy, *Modeling microstructure noise with mutually exciting point processes*, Quantitative Finance 13 (2013), 65–77. [MR3005350](#)
- [3] E. Bacry, S. Delattre, M. Hoffmann, J. F. Muzy, *Some limit theorems for Hawkes processes and applications to financial statistics*, Stoch. Processes Appl. 123 (2013), 2475–2499. [MR3054533](#)
- [4] E. Bacry and J. F. Muzy, *Hawkes model for price and trades high-frequency dynamics*, Quantitative Finance 14 (2014), 1147–1166. [MR3219705](#)
- [5] E. Bacry and J. F. Muzy, *Second order statistics characterization of Hawkes processes and non-parametric estimation*, arXiv:1401.0903.
- [6] L. Bauwens, N. Hautsch, *Modeling financial high frequency data using point processes*, ser. In T. Mikosch, J.-P. Kreiss, R. A. Davis, and T. G. Andersen, editors, Handbook of Financial Time Series. Springer, 2009.
- [7] G. Birkhoff, *Extensions of Jentzsch’s theorem*, Trans. Am. Math. Soc. 85 (1957), 219–227. [MR0087058](#)
- [8] C. Blundell, K. A. Heller, J. F. Beck, *Modeling reciprocating relationships with Hawkes processes*, Neural Information Processing Systems 2012.
- [9] P. Brémaud, L. Massoulié, *Stability of nonlinear Hawkes processes*, Ann. Probab. 24 (1996), 1563–1588. [MR1411506](#)

- [10] P. Brémaud, G. Nappo, G. L. Torrisi, *Rate of convergence to equilibrium of marked Hawkes processes*, J. Appl. Probab. 39 (2002), 123–136. [MR1895148](#)
- [11] R. Cavazos-Cadena, *An alternative derivation of Birkhoff's formula for the contraction coefficient of a positive matrix*, Linear Algebra Appl. 375 (2003), 291–297. [MR2013472](#)
- [12] D. J. Daley, D. Vere-Jones, *An introduction to the theory of point processes*, Vol. I. Probability and its Applications. Springer-Verlag, second edition, 2003. [MR1950431](#)
- [13] S. Delattre, N. Fournier, M. Hoffmann, *Hawkes processes on large networks*, to appear in Ann. Appl. Probab. [MR3449317](#)
- [14] W. Feller, *On the integral equation of renewal theory*, Ann. Math. Statistics 12 (1941), 243–267. [MR0005419](#)
- [15] S. Grün, M. Diedsmann, A. M. Aertsen, *Unitary events analysis*, in *Analysis of parallel spike trains*, S. Grün and S. Rotter, Springer series in computational neurosciences, 2010.
- [16] N. R. Hansen, P. Reynaud-Bouret, V. Rivoirard, *Lasso and probabilistic inequalities for multivariate point processes*, to appear in Bernoulli. [MR3322314](#)
- [17] A. Hawkes, *Spectra of some self-exciting and mutually exciting point processes*, Biometrika 58 (1971), 83–90. [MR0278410](#)
- [18] A. Hawkes, D. Oakes, *A cluster process representation of a self-exciting process*, J. Appl. Probability 11 (1974), 493–503. [MR0378093](#)
- [19] A. Helmstetter and D. Sornette, *Subcritical and supercritical regimes in epidemic models of earthquake aftershocks*, Journal of geophysical research, 107 (2002), 2237.
- [20] P. Hewlett *Clustering of order arrivals, price impact and trade path optimisation*, In Workshop on Financial Modeling with Jump processes. Ecole Polytechnique, 2006.
- [21] W. Hoeffding, *Probability inequalities for sums of bounded random variables*, J. Amer. Statist. Assoc. 58 (1963), 13–30. [MR0144363](#)
- [22] J. Jacod, A. N. Shiryaev, *Limit theorems for stochastic processes*, Second edition. Springer-Verlag, 2003.
- [23] Y. Y. Kagan, *Statistical distributions of earthquake numbers: consequence of branching process*, Geophysical Journal International 180 (2010), 1313–1328.
- [24] L. Massoulié, *Stability results for a general class of interacting point processes dynamics, and applications*, Stochastic Process. Appl. 75 (1998), 1–30. [MR1629010](#)
- [25] Y. Ogata, *The asymptotic behaviour of maximum likelihood estimators for stationary point processes*, Ann. Instit. Math. Statist. 30 (1978), 243–261. [MR0514494](#)
- [26] Y. Ogata, *Seismicity analysis through point-process modeling: a review*, Pure and Applied Geophysics 155 (1999), 471–507.
- [27] M. Okatan, M. A. Wilson, E. N. Brown, *Analyzing functional connectivity using a network likelihood model of ensemble neural spiking activity*, Neural Computation 17 (2005), 1927–1961.

- [28] J. W. Pillow, J. Shlens, L. Paninski, A. Scher, A. M. Litke, E. J. Chichilnisky, E. P. Simoncelli, *Spatio-temporal correlations and visual signalling in a complete neuronal population*, Nature 454 (2008), 995–999.
- [29] J. G. Rasmussen, *Bayesian inference for Hawkes processes*, Methodol. Comput. Appl. Probab. 15 (2013), 623–642. [MR3085883](#)
- [30] P. Reynaud-Bouret and S. Schbath, *Adaptive estimation for Hawkes processes: application to genome analysis*, Ann. Statist. 38 (2010), 2781–2822. [MR2722456](#)
- [31] P. Reynaud-Bouret, V. Rivoirard, F. Grammont, C. Tuleau-Malot, *Goodness-of-fit tests and nonparametric adaptive estimation for spike train analysis*, Journal of Math. Neuroscience 4:3 (2014). [MR3197017](#)
- [32] P. Reynaud-Bouret, V. Rivoirard, C. Tuleau-Malot, *Inference of functional connectivity in Neurosciences via Hawkes processes*, 1st IEEE Global Conference on Signal and Information Processing, 2013.
- [33] A. S. Sznitman, *Topics in propagation of chaos*, Ecole d’Été de Probabilités de Saint-Flour XIX-1989, Vol. 1464 of Lecture Notes in Math. Springer, 1991, 165–251. [MR1108185](#)
- [34] K. Zhou, H. Zha, L. Song, *Learning triggering kernels for multi-dimensional Hawkes processes*, Proceedings of the 30th International Conference on Machine Learning, 2013.
- [35] L. Zhu, *Central limit theorem for nonlinear Hawkes processes*, J. App. Probab. 50 (2013), 760–771. [MR3102513](#)
- [36] L. Zhu, *Large deviations for Markovian nonlinear Hawkes processes*, Ann. App. Probab. 25 (2015), 548–581. [MR3313748](#)