

The expected Euler characteristic approximation to excursion probabilities of smooth Gaussian random fields with general variance functions*

Dan Cheng[†]

Abstract

Consider a centered smooth Gaussian random field $\{X(t), t \in T\}$ with a general (nonconstant) variance function. In this work, we demonstrate that as $u \rightarrow \infty$, the excursion probability $\mathbb{P}\{\sup_{t \in T} X(t) \geq u\}$ can be accurately approximated by $\mathbb{E}\{\chi(A_u)\}$ such that the error decays at a super-exponential rate. Here, $A_u = \{t \in T : X(t) \geq u\}$ represents the excursion set above u , and $\mathbb{E}\{\chi(A_u)\}$ is the expectation of its Euler characteristic $\chi(A_u)$. This result substantiates the expected Euler characteristic heuristic for a broad class of smooth Gaussian random fields with diverse covariance structures. In addition, we employ the Laplace method to derive explicit approximations to the excursion probabilities.

Keywords: Gaussian random fields; excursion probability; excursion set; Euler characteristic; nonconstant variance; asymptotics; Laplace method; super-exponentially small.

MSC2020 subject classifications: 60G15; 60G60; 60G70.

Submitted to EJP on September 12, 2023, final version accepted on April 24, 2024.

1 Introduction

Let $X = \{X(t), t \in T\}$ represent a real-valued Gaussian random field defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where T denotes the parameter space. The study of excursion probabilities, denoted as $\mathbb{P}\{\sup_{t \in T} X(t) \geq u\}$, is a classical and fundamental problem in both probability and statistics. It finds extensive applications across numerous domains, including p -value computations, risk control and extreme event analysis, etc.

In the field of statistics, excursion probabilities play a critical role in tasks such as controlling family-wise error rates [14, 15], constructing confidence bands [11], and detecting signals in noisy data [9, 14]. However, except for only a few examples, computing the exact values of these probabilities is almost impossible. To address this challenge, many researchers have developed various methods for precise approximations

*The author acknowledges support from NSF Grants DMS-1902432 and DMS-2220523, as well as the Simons Foundation Collaboration Grant 854127.

[†]Arizona State University, U.S.A. E-mail: cheng.stats@gmail.com

of $\mathbb{P}\{\sup_{t \in T} X(t) \geq u\}$. These methods encompass techniques like the double sum method [7, 6], the tube method [10] and the Rice method [3, 4]. For comprehensive theoretical insights and related applications, we refer readers to the survey by Adler [1] and the monographs by Piterbarg [7], Adler and Taylor [2], and Azaïs and Wschebor [4], as well as the references therein.

In recent years, the expected Euler characteristic (EEC) method has emerged as a powerful tool for approximating excursion probabilities. This method, originating from the works of Taylor et al. [13] and Adler and Taylor [2], provides the following approximation:

$$\mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} = \mathbb{E}\{\chi(A_u)\} + \text{error}, \quad \text{as } u \rightarrow \infty, \quad (1.1)$$

where $\chi(A_u)$ represents the Euler characteristic of the excursion set $A_u = \{t \in T : X(t) \geq u\}$. This approximation (1.1) is highly elegant and accurate, primarily due to the fact that the principle term $\mathbb{E}\{\chi(A_u)\}$ is computable and the error term decays exponentially faster than the major component. However, it is essential to note that this method assumes a Gaussian field with constant variance, limiting its applicability in various scenarios.

Inspired by [5] where Gaussian fields with stationary increments are considered, we improve and generalize the techniques therein to establish the EEC approximation to accommodate smooth Gaussian random fields with general (nonconstant) variance functions in this paper. Our main objective is to demonstrate that the EEC approximation (1.1) remains valid under these conditions, with the error term exhibiting super-exponential decay. For a precise description of our findings, please refer to Theorem 3.1 below. Our derived approximation result shows that the maximum variance of $X(t)$, denoted by σ_T^2 (see (2.1) below), plays a pivotal role in both $\mathbb{E}\{\chi(A_u)\}$ and the super-exponentially small error. In our analysis, we observe that the points where σ_T^2 is attained make the most substantial contributions to $\mathbb{E}\{\chi(A_u)\}$. Building on this observation, we establish two simpler approximations: one in Theorem 3.2, which incorporates boundary conditions on nonzero derivatives of the variance function over points where σ_T^2 is attained, and another in Theorem 3.3, assuming only a single point attains σ_T^2 .

In general, the EEC approximation can be expressed as an integral using the Kac-Rice formula, as outlined in (3.2) in Theorem 3.1. While [13, 2] provided an elegant expression for $\mathbb{E}\{\chi(A_u)\}$ termed the Gaussian kinematic formula, this expression heavily relies on the assumption of unit variance, which simplifies the calculation. In our case, where the variance function of $X(t)$ varies across T , deriving an explicit expression for $\mathbb{E}\{\chi(A_u)\}$ becomes challenging. Instead, we apply the Laplace method to extract the term with the leading order of u from the integral, leaving a remaining error that is $\mathbb{E}\{\chi(A_u)\}o(1/u)$. For a more detailed explanation, we offer specific calculations in Sections 4 and 5. To intuitively grasp the EEC approximation, one can roughly consider the major term as $g(u)e^{-u^2/(2\sigma_T^2)}$, while the error term diminishes as $o(e^{-u^2/(2\sigma_T^2) - \alpha u^2})$, where $g(u)$ is a polynomial in u , and $\alpha > 0$ is a constant.

In terms of statistical applications, the EEC approximation is especially useful in estimating the EEC curve and hence the excursion probabilities; see a recent reference [16]. The derived EEC approximation in this paper verifies the validity of the method on estimating the EEC curve (such as taking the nonparametric average of Euler characteristic curves) to approximate the excursion probabilities for smooth Gaussian fields with general nonconstant variance functions. However, the method of Hermite projection for estimating Lipschitz-Killing curvatures introduced in [16] seems challenging for nonconstant variances due to the complexity of the EEC expression mentioned above.

The structure of this paper is as follows: We begin by introducing the notations

and assumptions in Section 2. In Section 3, we present our main results, including Theorems 3.1, 3.2, and 3.3. In Section 4, we apply the Laplace method to derive explicit approximations (Theorems 4.1 and 4.2) for cases where a unique maximum point of the variance is present. We then demonstrate several examples in Section 5 and illustrate the evaluation of EEC and the subsequent approximation of excursion probabilities, including the case when the maximum of the variance is achieved on a line. To understand our approach, we outline the main ideas in Section 6 and delve into the analysis of super-exponentially small errors in Sections 7 and 8. Finally, we provide the proofs of our main results and of the results for the unique maximum point in Sections 9 and 10, respectively.

2 Notations and assumptions

Let $\{X(t), t \in T\}$ be a real-valued and centered Gaussian random field, where T is a compact rectangle in \mathbb{R}^N . We define

$$\nu(t) = \sigma_t^2 = \text{Var}(X(t)) \quad \text{and} \quad \sup_{t \in T} \nu(t) = \sigma_T^2. \tag{2.1}$$

Here, $\nu(\cdot)$ represents the variance function of the field and σ_T^2 is the maximum variance over T . For a function $f(\cdot) \in C^2(\mathbb{R}^N)$ and $t \in \mathbb{R}^N$, we introduce the following notations on derivatives:

$$\begin{aligned} f_i(t) &= \frac{\partial f(t)}{\partial t_i}, \quad f_{ij}(t) = \frac{\partial^2 f(t)}{\partial t_i \partial t_j}, \quad \forall i, j = 1, \dots, N; \\ \nabla f(t) &= (f_1(t), \dots, f_N(t))^T, \quad \nabla^2 f(t) = (f_{ij}(t))_{i,j=1, \dots, N}. \end{aligned} \tag{2.2}$$

Let $B \prec 0$ (negative definite) and $B \preceq 0$ (negative semi-definite) denote that a symmetric matrix B has all negative or nonpositive eigenvalues, respectively. Additionally, we use $\text{Cov}(\xi_1, \xi_2)$ and $\text{Corr}(\xi_1, \xi_2)$ to represent the covariance and correlation between two random variables ξ_1 and ξ_2 . The density of the standard Normal distribution is denoted as $\phi(x)$, and its tail probability is $\Psi(x) = \int_x^\infty \phi(y) dy$. Let \mathbb{S}^j be the j -dimensional unit sphere.

Consider the domain $T = \prod_{i=1}^N [a_i, b_i]$, where $-\infty < a_i < b_i < \infty$. We draw from the notation established by Adler and Taylor in [2] to demonstrate that T can be decomposed into the union of its interior and lower-dimensional faces. This decomposition forms the basis for calculating the Euler characteristic of the excursion set A_u , as elaborated in Section 3.

Let $k \in \{0, 1, \dots, N\}$. Each face K of dimension k is defined by fixing a subset $\tau(K) \subset \{1, \dots, N\}$ of size k and a subset $\varepsilon(K) = \{\varepsilon_j, j \notin \tau(K)\} \subset \{0, 1\}^{N-k}$ of size $N - k$ so that

$$\begin{aligned} K &= \{t = (t_1, \dots, t_N) \in T : a_j < t_j < b_j \text{ if } j \in \tau(K), \\ &\quad t_j = (1 - \varepsilon_j)a_j + \varepsilon_j b_j \text{ if } j \notin \tau(K)\}. \end{aligned}$$

Denote by $\partial_k T$ the collection of all k -dimensional faces in T . The interior of T is designated as $\overset{\circ}{T} = \partial_N T$, while the boundary of T is formulated as $\partial T = \cup_{k=0}^{N-1} \cup_{K \in \partial_k T} K$. This allows us to partition T in the following manner:

$$T = \bigcup_{k=0}^N \bigcup_{K \in \partial_k T} K.$$

For each $t \in T$, let

$$\begin{aligned} \nabla X|_K(t) &= (X_{i_1}(t), \dots, X_{i_k}(t))_{i_1, \dots, i_k \in \tau(K)}^T, \quad \nabla^2 X|_K(t) = (X_{mn}(t))_{m, n \in \tau(K)}, \\ \Sigma(t) &= \mathbb{E}\{X(t)\nabla^2 X(t)\} = (\mathbb{E}\{X(t)X_{ij}(t)\})_{1 \leq i, j \leq N}, \\ \Sigma_K(t) &= \mathbb{E}\{X(t)\nabla^2 X|_K(t)\} = (\mathbb{E}\{X(t)X_{ij}(t)\})_{i, j \in \tau(K)}, \\ \Lambda(t) &= \text{Cov}(\nabla X(t)) = (\mathbb{E}\{X_i(t)X_j(t)\})_{1 \leq i, j \leq N}, \\ \Lambda_K(t) &= \text{Cov}(\nabla X|_K(t)) = (\mathbb{E}\{X_i(t)X_j(t)\})_{i, j \in \tau(K)}. \end{aligned} \tag{2.3}$$

For each $K \in \partial_k T$, we define the *number of extended outward maxima above u on face K* as

$$M_u^E(K) := \#\{t \in K : X(t) \geq u, \nabla X|_K(t) = 0, \nabla^2 X|_K(t) \prec 0, \varepsilon_j^* X_j(t) \geq 0, \forall j \notin \tau(K)\},$$

where $\varepsilon_j^* = 2\varepsilon_j - 1$, and define the *number of local maxima above u on face K* as

$$M_u(K) := \#\{t \in K : X(t) \geq u, \nabla X|_K(t) = 0, \nabla^2 X|_K(t) \prec 0\}.$$

Clearly, $M_u^E(K) \leq M_u(K)$.

For each $t \in T$ with $\nu(t) = \sigma_T^2$, we define the index set $\mathcal{I}(t) = \{\ell : \nu_\ell(t) = 0\}$ representing the directions along which the partial derivatives of $\nu(t)$ vanish. If $t \in K \in \partial_k T$ with $\nu(t) = \sigma_T^2$, then we have $\tau(K) \subset \mathcal{I}(t)$ since $\nu_\ell(t) = 0$ for all $\ell \in \tau(K)$. It is worth noting that since $\nu_i(t) = 2\mathbb{E}\{X_i(t)X(t)\}$, we can also express this index set as $\mathcal{I}(t) = \{\ell : \mathbb{E}\{X(t)X_\ell(t)\} = 0\}$.

Our analytical framework relies on the following conditions for smoothness **(H1)** and regularity **(H2)**, in addition to curvature conditions **(H3)** or **(H3')**.

(H1) $X \in C^2(\mathbb{R}^N)$ almost surely and the second derivatives satisfy the *uniform mean-square Hölder condition*: there exist constants $C, \delta > 0$ such that

$$\mathbb{E}(X_{ij}(t) - X_{ij}(t'))^2 \leq C\|t - t'\|^{2\delta}, \quad \forall t, t' \in T, \quad i, j = 1, \dots, N.$$

(H2) For every pair $(t, t') \in T^2$ with $t \neq t'$, the Gaussian vector

$$(X(t), \nabla X(t), X_{ij}(t), X(t'), \nabla X(t'), X_{ij}(t'), 1 \leq i \leq j \leq N)$$

is non-degenerate.

(H3) For every $t \in K \in \partial_k T$, $0 \leq k \leq N - 2$, such that $\nu(t) = \sigma_T^2$ and $\mathcal{I}(t)$ contains at least two indices, we have

$$(\mathbb{E}\{X(t)X_{ij}(t)\})_{i, j \in \mathcal{I}(t)} \prec 0. \tag{2.4}$$

(H3') For every $t \in K \in \partial_k T$, $0 \leq k \leq N - 2$, such that $\nu(t) = \sigma_T^2$ and $\mathcal{I}(t)$ contains at least two indices, we have

$$(\nu_{ij}(t))_{i, j \in \mathcal{I}(t)} \preceq 0. \tag{2.5}$$

The smoothness condition **(H1)** and regularity condition **(H2)** imply the validity of Corollary 11.3.2 in [2], showing that X is almost surely a Morse function on T . Additionally, the conditions required for Kac-Rice formulas in Theorems 11.2.1 and 11.5.1 in [2] are satisfied, so that we can apply them to compute moments of the number of critical points such as $\mathbb{E}\{M_u^E(K)\}$, $\mathbb{E}\{M_u^E(K)M_u^E(K')\}$ and $\mathbb{E}\{M_u^E(K)(M_u^E(K) - 1)\}$, where K and K' are different faces of T ; see also [8].

Conditions **(H3)** and **(H3')** involve the behavior of the variance function $\nu(t)$ at critical points, and they are closely related, as shown in Proposition 2.1 below. Here we provide

some additional insights into $(\mathbf{H3}')$. Despite its initially technical appearance, $(\mathbf{H3}')$ is in fact a mild condition that specifically applies to lower-dimensional boundary points t where $\nu(t) = \sigma_T^2$. In essence, it indicates that the variance function should possess a negative semi-definite Hessian matrix at these boundary critical points where $\nu(t) = \sigma_T^2$ while concurrently exhibiting at least two zero partial derivatives. Conditions $(\mathbf{H3})$ and $(\mathbf{H3}')$ generalize the nondegeneracy condition of $\mathbb{E}\{(X(t) - X(0))\nabla^2 X(t)\}$ which is condition $(\mathbf{H2})$ in [5] from stationary increments to general nonconstant variances.

For example, in the 1D case, since $\mathcal{I}(t)$ contains at most one index, there is no need to check $(\mathbf{H3}')$. Similarly, in the 2D case, we only need to check $(\mathbf{H3}')$ or (2.5) when σ_T^2 is achieved at corner points $t \in \partial_0 T$ with $\mathcal{I}(t) = \{1, 2\}$. Moreover, if the variance function $\nu(t)$ demonstrates strict monotonicity in all directions across \mathbb{R}^N , then $\mathcal{I}(t) = \emptyset$ and there is no need to verify $(\mathbf{H3}')$.

Proposition 2.1. *The condition $(\mathbf{H3}')$ implies $(\mathbf{H3})$. In addition, $(\mathbf{H3})$ implies that*

$$(\mathbb{E}\{X(t)X_{ij}(t)\})_{i,j \in \mathcal{I}(t)} < 0, \quad \forall t \in T \text{ with } \nu(t) = \sigma_T^2. \quad (2.6)$$

Proof. Taking the second derivative on both sides of $\nu(t) = \mathbb{E}\{X(t)^2\}$, we obtain $\nu_{ij}(t)/2 = \mathbb{E}\{X(t)X_{ij}(t)\} + \mathbb{E}\{X_i(t)X_j(t)\}$, implying

$$(\mathbb{E}\{X(t)X_{ij}(t)\})_{i,j \in \mathcal{I}(t)} = \frac{1}{2}(\nu_{ij}(t))_{i,j \in \mathcal{I}(t)} - (\mathbb{E}\{X_i(t)X_j(t)\})_{i,j \in \mathcal{I}(t)}. \quad (2.7)$$

Note that, as a covariance matrix, $(\mathbb{E}\{X_i(t)X_j(t)\})_{i,j \in \mathcal{I}(t)}$ is positive definite by $(\mathbf{H2})$. Therefore, (2.5) implies (2.4), or equivalently $(\mathbf{H3}')$ implies $(\mathbf{H3})$.

Next we demonstrate that $(\mathbf{H3})$ implies (2.6). It suffices to show (2.4) for $k = N - 1$ and $k = N$, and for the case that $\mathcal{I}(t)$ contains at most one index, which complement those cases in $(\mathbf{H3})$.

(i) If $k = N$, then t becomes a maximum point of ν within the interior of T and $\mathcal{I}(t) = \tau(K) = \{1, \dots, N\}$, implying (2.5), and hence (2.4) holds by (2.7).

(ii) For $k = N - 1$, we consider two scenarios. If $\mathcal{I}(t) = \tau(K)$, then t becomes a maximum point of ν restricted on K , hence (2.4) is satisfied as discussed above. If $\mathcal{I}(t) = \{1, \dots, N\}$, then it follows from Taylor's formula that

$$\nu(t') = \nu(t) + (t' - t)^T \nabla^2 \nu(t)(t' - t) + o(\|t' - t\|^2), \quad t' \in T.$$

Notice that $\{(t' - t)/\|t' - t\| : t' \in T\}$ contains all directions in \mathbb{R}^N since $t \in K \in \partial_{N-1} T$, together with the fact $\nu(t) = \sigma_T^2$, we see that $\nabla^2 \nu(t)$ cannot have any positive eigenvalue, thus (2.5) and hence (2.4) hold.

(iii) Finally, it's evident from the 1D Taylor's formula that (2.5) is valid when $\mathcal{I}(t)$ contains only one index. \square

The condition (2.6) established in Proposition 2.1 serves as the fundamental requirement for our main results, as demonstrated in Theorems 3.1, 3.2 and 3.3 below. As seen from Proposition 2.1, we can simplify (2.6) to condition $(\mathbf{H3})$. Thus our main results will be presented under the assumption of condition $(\mathbf{H3})$.

Furthermore, it is worth highlighting that, in practical applications, verifying $(\mathbf{H3}')$ can often be a more straightforward process. This condition directly pertains to the variance function $\nu(t)$, making it easier to assess. Thus, Proposition 2.1 provides the flexibility to check $(\mathbf{H3}')$ instead of $(\mathbf{H3})$. This insight simplifies the verification procedure, enhancing the practical applicability of our results.

3 Main results

Here, we will present our main results Theorems 3.1, 3.2 and 3.3, whose proofs are given in Section 9. Define the *number of extended outward critical points of index i above level u on the face K* be

$$\begin{aligned} \mu_i(K) &:= \#\{t \in K : X(t) \geq u, \nabla X|_K(t) = 0, \text{index}(\nabla^2 X|_K(t)) = i, \\ &\quad \varepsilon_j^* X_j(t) \geq 0 \text{ for all } j \notin \tau(K)\}. \end{aligned}$$

Recall that $\varepsilon_j^* = 2\varepsilon_j - 1$ and the index of a matrix is defined as the number of its negative eigenvalues. It is evident to observe that $\mu_N(K) = M_u^E(K)$. Here, by convention, if $K \in \partial_0 T$, then the terms on $\nabla X|_K(t)$ and $\nabla^2 X|_K(t)$ in the definition above vanish. It follows from **(H1)**, **(H2)** and the Morse theorem (see Corollary 9.3.5 or pages 211–212 in [2]) that the Euler characteristic of the excursion set A_u can be represented as

$$\chi(A_u) = \sum_{k=0}^N \sum_{K \in \partial_k T} (-1)^k \sum_{i=0}^k (-1)^i \mu_i(K). \tag{3.1}$$

Now we state the following general result on the EEC approximation for the excursion probability.

Theorem 3.1. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying **(H1)**, **(H2)** and **(H3)**. Then there exists a constant $\alpha > 0$ such that as $u \rightarrow \infty$,*

$$\mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} = \mathbb{E}\{\chi(A_u)\} + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right),$$

where the EEC is expressed as

$$\begin{aligned} \mathbb{E}\{\chi(A_u)\} &= \sum_{k=0}^N \sum_{K \in \partial_k T} (-1)^k \int_K \mathbb{E}\{\det \nabla^2 X|_K(t) \mathbb{1}_{\{X(t) \geq u, \varepsilon_\ell^* X_\ell(t) \geq 0 \text{ for all } \ell \notin \tau(K)\}} \mid \\ &\quad \nabla X|_K(t) = 0\} p_{\nabla X|_K(t)}(0) dt. \end{aligned} \tag{3.2}$$

It is worth noting that, the principle term $\mathbb{E}\{\chi(A_u)\}$ can be roughly treated as $g(u)e^{-u^2/(2\sigma_T^2)}$, where $g(u)$ is a polynomial in u . So the error term decays exponentially faster than $\mathbb{E}\{\chi(A_u)\}$. In general, computing the EEC approximation $\mathbb{E}\{\chi(A_u)\}$ is a challenging task because it involves conditional expectations over the joint covariance of the Gaussian field and its Hessian, given zero gradient, which vary across T . However, one can apply the Laplace method to extract the term with the largest order of u from $\mathbb{E}\{\chi(A_u)\}$ such that the remaining error is $o(1/u)\mathbb{E}\{\chi(A_u)\}$. Examples demonstrating the Laplace method are presented in Section 5.

It is important to note that in the expression (3.2), when $k = 0$, all terms involving $\nabla X|_K(t)$ and $\nabla^2 X|_K(t)$ vanish. Consequently, if $k = 0$, we treat the integral in (3.2) as the usual Gaussian tail probabilities. This notation is also adopted in the results presented in Theorems 3.2 and 3.3 below.

The proof of Theorem 3.1 reveals that the points where the maximum variance σ_T^2 is attained make the most significant contribution to $\mathbb{E}\{\chi(A_u)\}$. Therefore, in many cases, the general EEC approximation $\mathbb{E}\{\chi(A_u)\}$ can be simplified. The following result is based on the boundary condition (3.3) and is applicable at boundary points where nonzero partial derivatives of the variance function occur when σ_T^2 is reached.

Theorem 3.2. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying **(H1)**, **(H2)** and the following boundary condition*

$$\left\{t \in J : \nu(t) = \sigma_T^2, \prod_{i \notin \tau(J)} \nu_i(t) = 0\right\} = \emptyset, \quad \forall \text{ face } J \subset T. \tag{3.3}$$

Then there exists a constant $\alpha > 0$ such that as $u \rightarrow \infty$,

$$\begin{aligned} \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} &= \sum_{k=0}^N \sum_{K \in \partial_k T} (-1)^k \int_K \mathbb{E} \left\{ \det \nabla^2 X|_K(t) \mathbb{1}_{\{X(t) \geq u\}} \mid \nabla X|_K(t) = 0 \right\} \\ &\quad \times p_{\nabla X|_K(t)}(0) dt + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right). \end{aligned}$$

In other words, the boundary condition (3.3) indicates that, for any point $t \in J$ attaining the maximum variance σ_T^2 , there must be $\nu_i(t) \neq 0$ for all $i \notin \tau(J)$. In particular, as an important property, we observe that (3.3) implies the condition **(H3')** and hence **(H3)**. The following result provides an asymptotic approximation for the special case where the variance function attains its maximum σ_T^2 only at a unique point.

Theorem 3.3. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying **(H1)**, **(H2)** and **(H3)**. Suppose $\nu(t)$ attains its maximum σ_T^2 only at a single point $t^* \in K$, where $K \in \partial_k T$ with $k \geq 0$. Then there exists a constant $\alpha > 0$ such that as $u \rightarrow \infty$,*

$$\begin{aligned} &\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &= \sum_J (-1)^{\dim(J)} \int_J \mathbb{E} \left\{ \det \nabla^2 X|_J(t) \mathbb{1}_{\{X(t) \geq u, \varepsilon_\ell^* X_\ell(t) \geq 0 \text{ for all } \ell \in \mathcal{I}(t^*) \setminus \tau(J)\}} \mid \nabla X|_J(t) = 0 \right\} \\ &\quad \times p_{\nabla X|_J(t)}(0) dt + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right), \end{aligned}$$

where the sum is taken over all faces J of T such that $t^* \in \bar{J}$ and $\tau(J) \subset \mathcal{I}(t^*)$.

Employing the Laplace method, we will provide refined explicit approximation results in Section 4 under the assumptions in Theorem 3.3. Furthermore, we demonstrate several examples that illustrate the evaluation of approximating excursion probabilities in Section 5, including the case that the maximum of the variance function is achieved on a line.

4 Gaussian fields with a unique maximum point of the variance

In this section, we delve deeper into EEC approximations when the variance function $\nu(t)$ reaches its maximum value σ_T^2 at a solitary point t^* . While Theorem 3.3 provides an implicit formula for such scenarios, our objective here is to obtain explicit formulae by employing integral approximation techniques based on the Kac-Rice formula. The proofs are given in Section 10.

There are some existing references on approximating the excursion probabilities for Gaussian fields with a unique maximum point of the variance function, such as [7, 6], where the double sum method was employed but the error rate was hard to obtain (only stated as $o(1)$ multiplying the major term). Our derived EEC approximations show a super-exponentially small error for the smooth case, and one can apply the Laplace method to derive a specific approximation. Meanwhile, we provide the approximation for the case when the maximum point t^* is on the boundary. This boundary case turns to be difficult for the double sum method and is usually ignored.

4.1 Gaussian fields satisfying the boundary condition (3.3)

The following result provides explicit approximations to the excursion probabilities when the maximum of the variance is reached only at a single point and the boundary condition (3.3) is satisfied.

Theorem 4.1. Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1) and (H2). Suppose ν attains its maximum σ_T^2 only at $t^* \in K \in \partial_k T$, $\nu_i(t^*) \neq 0$ for all $i \notin \tau(K)$, and $\nabla^2 \nu|_K(t^*) \prec 0$. Then, as $u \rightarrow \infty$,

$$\begin{aligned} \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} &= \Psi \left(\frac{u}{\sigma_T} \right) + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right) \text{ for some } \alpha > 0, \quad \text{if } k = 0, \\ \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} &= \sqrt{\frac{\det(\Sigma_K(t^*))}{\det(\Lambda_K(t^*) + \Sigma_K(t^*))}} \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)), \quad \text{if } k \geq 1, \end{aligned} \tag{4.1}$$

where $\Lambda_K(t^*)$ and $\Sigma_K(t^*)$ are defined in (2.3).

Now we apply Theorem 4.1 to the 1D case when $T = [a, b]$. If $t^* = a$ or $t^* = b$, then it is a direct application of the first line in (4.1). If $t^* \in (a, b)$, then it follows from (4.1) that

$$\mathbb{P} \left\{ \sup_{t \in [a, b]} X(t) \geq u \right\} = \sqrt{\frac{\mathbb{E}\{X(t^*)X''(t^*)\}}{\text{Var}(X'(t^*)) + \mathbb{E}\{X(t^*)X''(t^*)\}}} \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)).$$

4.2 Gaussian fields not satisfying the boundary condition (3.3)

We consider here the other case when $\nu_i(t^*) \neq 0$ for some $i \notin \tau(K)$. For a symmetric matrix $B = (B_{ij})_{1 \leq i, j \leq N}$, we call $(B_{ij})_{i, j \in \mathcal{I}}$ the matrix B with indices restricted on \mathcal{I} .

Theorem 4.2. Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1) and (H2). Suppose ν attains its maximum σ_T^2 only at $t^* \in K \in \partial_k T$ such that $\mathcal{I}(t^*) \setminus \tau(K)$ contains $m \geq 1$ indices and $(\nu_{i i'}(t^*))_{i, i' \in \mathcal{I}(t^*)} \prec 0$. Then, as $u \rightarrow \infty$,

$$\begin{aligned} &\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &= \sum_J \sqrt{\frac{\det(\Sigma_J(t^*))}{\det(\Lambda_J(t^*) + \Sigma_J(t^*))}} \mathbb{P}\{(Z_{J'_1}, \dots, Z_{J'_{j-k}}) \in E'(J)\} \\ &\quad \times \mathbb{P}\{(X_{J_1}(t^*), \dots, X_{J_{k+m-j}}(t^*)) \in E(J) \mid \nabla X|_J(t^*) = 0\} \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)), \end{aligned} \tag{4.2}$$

where the sum is taken over all faces J such that $t^* \in \bar{J}$ and $\tau(J) \subset \mathcal{I}(t^*)$, $j = \dim(J)$,

$$\begin{aligned} (J_1, \dots, J_{k+m-j}) &= \mathcal{I}(t^*) \setminus \tau(J), \quad (J'_1, \dots, J'_{j-k}) = \tau(J) \setminus \tau(K), \\ E(J) &= \{(y_{J_1}, \dots, y_{J_{k+m-j}}) \in \mathbb{R}^{k+m-j} : \varepsilon_{J_\ell}^*(J) y_{J_\ell} \geq 0, \forall \ell = 1, \dots, k+m-j\}, \\ E'(J) &= \{(y_{J'_1}, \dots, y_{J'_{j-k}}) \in \mathbb{R}^{j-k} : \varepsilon_{J'_\ell}^*(K) y_{J'_\ell} \geq 0, \forall \ell = 1, \dots, j-k\}, \end{aligned}$$

$\varepsilon_{J_\ell}^*(J)$ and $\varepsilon_{J'_\ell}^*(K)$ are the ε^* numbers for faces J and K respectively, $(Z_{J'_1}, \dots, Z_{J'_{j-k}})$ is a centered Gaussian vector having covariance matrix $\Sigma(t^*) + \Sigma(t^*)\Lambda^{-1}(t^*)\Sigma(t^*)$ with indices restricted on $\tau(J) \setminus \tau(K)$, and $\Lambda_J(t^*)$ and $\Sigma_J(t^*)$ are defined in (2.3). In particular, for $k = 0$, the term inside the sum in (4.2) with $J = K = \{t^*\}$ is given by

$$\mathbb{P}\{(X_{J_1}(t^*), \dots, X_{J_m}(t^*)) \in E(J)\} \Psi \left(\frac{u}{\sigma_T} \right).$$

Now we apply Theorem 4.2 to the 1D case when $T = [a, b]$. Without loss of generality,

assume $t^* = b$ and $\nu'(t^*) = 0$. Then it follows from Theorem 4.2 that

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{t \in [a, b]} X(t) \geq u \right\} \\ &= \left(\mathbb{P}\{X'(t^*) > 0\} + \sqrt{\frac{\mathbb{E}\{X(t^*)X''(t^*)\}}{\text{Var}(X'(t^*)) + \mathbb{E}\{X(t^*)X''(t^*)\}}} \mathbb{P}\{Z > 0\} \right) \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)) \\ &= \frac{1}{2} \left(1 + \sqrt{\frac{\mathbb{E}\{X(t^*)X''(t^*)\}}{\text{Var}(X'(t^*)) + \mathbb{E}\{X(t^*)X''(t^*)\}}} \right) \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)), \end{aligned}$$

where Z is a centered Gaussian variable.

Denote by $\mathbb{R}_+^n = (0, \infty)^n$. To simplify the statement in Theorem 4.2, we present below another version with less notations on faces.

Corollary 4.3. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1), (H2) and (H3). Suppose ν attains its maximum σ_T^2 only at $t^* \in K \in \partial_k T$ with $\tau(K) = \{1, \dots, k\}$ such that $\mathcal{I}(t^*) = \{1, \dots, k, k+1, \dots, k+m\}$ with $m \geq 1$ and $(\nu_{i,i'}(t^*))_{1 \leq i, i' \leq k+m} \prec 0$. Then, as $u \rightarrow \infty$,*

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &= \sum_{j=k}^{k+m} \sum_{J \in \partial_j T: t^* \in \bar{J}} \sqrt{\frac{\det(\Sigma_J(t^*))}{\det(\Lambda_J(t^*) + \Sigma_J(t^*))}} \mathbb{P}\{(Z_1, \dots, Z_{j-k}) \in \mathbb{R}_+^{j-k}\} \\ & \quad \times \mathbb{P}\{(X_{j+1}(t^*), \dots, X_{k+m}(t^*)) \in \mathbb{R}_+^{k+m-j} | \nabla X|_J(t^*) = 0\} \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)), \end{aligned} \tag{4.3}$$

where (Z_1, \dots, Z_{j-k}) is a centered Gaussian random vector having covariance $\Sigma(t^*) + \Sigma(t^*)\Lambda^{-1}(t^*)\Sigma(t^*)$ with indices restricted on $\{k+1, \dots, j\}$, and $\Lambda_J(t^*)$ and $\Sigma_J(t^*)$ are defined in (2.3). In particular, for $k = 0$, the term inside the sum in (4.3) with $J = K = \{t^*\}$ is

$$\mathbb{P}\{(X_1(t^*), \dots, X_m(t^*)) \in \mathbb{R}_+^m\} \Psi \left(\frac{u}{\sigma_T} \right).$$

5 Examples

Throughout this section, we consider a centered Gaussian random field $\{X(t), t \in T\}$ satisfying (H1), (H2) and (H3), where $T = [a_1, b_1] \times [a_2, b_2] \subset \mathbb{R}^2$.

5.1 Examples with a unique maximum point of the variance

Suppose $\nu(t_1, t_2)$ attains the maximum σ_T^2 only at a single point $t^* = (t_1^*, t_2^*)$; and the assumptions in Theorems 4.1 or 4.2 are satisfied.

Case 1: $t^* = (b_1, b_2)$ and $\nu_1(t^*)\nu_2(t^*) \neq 0$. It follows directly from Theorem 4.1 that

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} = \Psi \left(\frac{u}{\sigma_T} \right) + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right).$$

Case 2: $t^* = (b_1, b_2)$, $\nu_1(t^*) = 0$ and $\nu_2(t^*) \neq 0$. It follows from Corollary 4.3 that

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &= \left(\mathbb{P}\{X_1(t^*) > 0\} + \sqrt{\frac{\mathbb{E}\{X(t^*)X_{11}(t^*)\}}{\text{Var}(X_1(t^*)) + \mathbb{E}\{X(t^*)X_{11}(t^*)\}}} \mathbb{P}\{Z > 0\} \right) \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)) \\ &= \frac{1}{2} \left(1 + \sqrt{\frac{\mathbb{E}\{X(t^*)X_{11}(t^*)\}}{\text{Var}(X_1(t^*)) + \mathbb{E}\{X(t^*)X_{11}(t^*)\}}} \right) \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)), \end{aligned}$$

where Z is a centered Gaussian variable.

Case 3: $t^* = (b_1, b_2)$ and $\nu_1(t^*) = \nu_2(t^*) = 0$. Applying Corollary 4.3 and noting the calculations in Case 2 above, we obtain

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &= \left(\mathbb{P}\{X_1(t^*) > 0, X_2(t^*) > 0\} + \frac{1}{2} \sqrt{\frac{\mathbb{E}\{X(t^*)X_{11}(t^*)\}}{\text{Var}(X_1(t^*)) + \mathbb{E}\{X(t^*)X_{11}(t^*)\}}} \right. \\ & \quad \left. + \frac{1}{2} \sqrt{\frac{\mathbb{E}\{X(t^*)X_{22}(t^*)\}}{\text{Var}(X_2(t^*)) + \mathbb{E}\{X(t^*)X_{22}(t^*)\}}} \right. \\ & \quad \left. + \mathbb{P}\{Z_1 > 0, Z_2 > 0\} \sqrt{\frac{\det(\Sigma(t^*))}{\det(\Lambda(t^*) + \Sigma(t^*))}} \right) \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)), \end{aligned}$$

where (Z_1, Z_2) is a centered Gaussian vector with covariance $\Sigma(t^*) + \Sigma(t^*)\Lambda^{-1}(t^*)\Sigma(t^*)$.

Case 4: $t^* = (t_1^*, b_2)$, where $t_1^* \in (a_1, b_1)$ and $\nu_2(t^*) \neq 0$. It follows directly from Theorem 4.1 that

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} = \sqrt{\frac{\mathbb{E}\{X(t^*)X_{11}(t^*)\}}{\text{Var}(X_1(t^*)) + \mathbb{E}\{X(t^*)X_{11}(t^*)\}}} \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)).$$

Case 5: $t^* = (t_1^*, b_2)$, where $t_1^* \in (a_1, b_1)$ and $\nu_2(t^*) = 0$. Applying Corollary 4.3 and noting the calculations in Case 2 above, we obtain

$$\begin{aligned} & \mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &= \frac{1}{2} \left(\sqrt{\frac{\mathbb{E}\{X(t^*)X_{11}(t^*)\}}{\text{Var}(X_1(t^*)) + \mathbb{E}\{X(t^*)X_{11}(t^*)\}}} + \sqrt{\frac{\det(\Sigma(t^*))}{\det(\Lambda(t^*) + \Sigma(t^*))}} \right) \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)). \end{aligned}$$

Case 6: $a_1 < t_1^* < b_1$ and $a_2 < t_2^* < b_2$. It follows directly from Theorem 4.1 that

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} = \sqrt{\frac{\det(\Sigma(t^*))}{\det(\Lambda(t^*) + \Sigma(t^*))}} \Psi \left(\frac{u}{\sigma_T} \right) (1 + o(1)).$$

5.2 Examples with the maximum of the variance achieved on a line

Consider the Gaussian random field $X(t)$ defined as:

$$X(t) = \xi_1 \cos t_1 + \xi_1' \sin t_1 + t_2(\xi_2 \cos t_2 + \xi_2' \sin t_2),$$

where $t = (t_1, t_2) \in T = [a_1, b_1] \times [a_2, b_2] \subset (0, 2\pi)^2$, and $\xi_1, \xi_1', \xi_2, \xi_2'$ are independent standard Gaussian random variables. This is a Gaussian random field on \mathbb{R}^2 generated

from the cosine field, with an additional product of t_2 along the vertical direction. The constraint on the parameter space within $(0, 2\pi)^2$ is imposed to prevent degeneracy in derivatives. For this field, we have $\nu(t) = 1 + t_2^2$, which reaches the maximum $\sigma_T^2 = 1 + b_2^2$ on the entire real line $L := \{(t_1, b_2) : a_1 \leq t_1 \leq b_1\}$. Furthermore,

$$\nu_1(t)|_{t \in L} = 0, \quad \nu_2(t)|_{t \in L} = 2b_2 > 0, \quad \forall t \in L.$$

By employing similar reasoning in the proofs of Theorems 3.1 and 3.2, we see that, in the EEC approximation $\mathbb{E}\{\chi(A_u)\}$, all integrals (derived from the Kac-Rice formula) over faces not contained within \bar{L} are super-exponentially small. Thus, there exists $\alpha > 0$ such that as $u \rightarrow \infty$,

$$\begin{aligned} \mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} &= \mathbb{P}\{X(a_1, b_2) \geq u, X_1(a_1, b_2) < 0\} + \mathbb{P}\{X(b_1, b_2) \geq u, X_1(b_1, b_2) > 0\} \\ &\quad + I(u) + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right) \\ &= \Psi\left(\frac{u}{\sqrt{1 + b_2^2}}\right) + I(u) + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right), \end{aligned} \tag{5.1}$$

where

$$I(u) = - \int_{a_1}^{b_1} \mathbb{E}\{X_{11}(t_1, b_2) \mathbb{1}_{\{X(t_1, b_2) \geq u\}} | X_1(t_1, b_2) = 0\} p_{X_1(t_1, b_2)}(0) dt_1.$$

Since $X_1(t_1, b_2) = -\xi_1 \sin t_1 + \xi'_1 \cos t_1$ and $X_{11}(t_1, b_2) = -\xi_1 \cos t_1 - \xi'_1 \sin t_1$, one has

$$\text{Cov}(X(t_1, b_2), X_1(t_1, b_2), X_{11}(t_1, b_2)) = \begin{pmatrix} 1 + b_2^2 & 0 & -1 \\ 0 & 1 & 0 \\ -1 & 0 & 1 \end{pmatrix},$$

which does not depend on t_1 . Particularly, $X_1(t_1, b_2)$ is independent of both $X(t_1, b_2)$ and $X_{11}(t_1, b_2)$. Thus

$$\begin{aligned} I(u) &= -\frac{b_1 - a_1}{\sqrt{2\pi}} \mathbb{E}\{X_{11}(t_1, b_2) \mathbb{1}_{\{X(t_1, b_2) \geq u\}}\} \\ &= -\frac{b_1 - a_1}{\sqrt{2\pi}} \int_u^\infty \mathbb{E}\{X_{11}(t_1, b_2) | X(t_1, b_2) = x\} \phi\left(\frac{x}{\sqrt{1 + b_2^2}}\right) dx \\ &= \frac{b_1 - a_1}{\sqrt{2\pi}} \int_u^\infty \frac{x}{1 + b_2^2} \phi\left(\frac{x}{\sqrt{1 + b_2^2}}\right) dx \\ &= \frac{b_1 - a_1}{\sqrt{2\pi}} \phi\left(\frac{u}{\sqrt{1 + b_2^2}}\right). \end{aligned}$$

Substituting this expression into (5.1), we arrive at the following refined approximation:

$$\mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} = \Psi\left(\frac{u}{\sqrt{1 + b_2^2}}\right) + \frac{b_1 - a_1}{\sqrt{2\pi}} \phi\left(\frac{u}{\sqrt{1 + b_2^2}}\right) + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right),$$

which has a super-exponentially small error.

6 Outline of the proofs of main results

Here we show the main idea for proving the main results above. The main idea below is similar to that introduced in [5] by estimating the moments of the number of critical points. Let f be a smooth real-valued function, then $\sup_{t \in T} f(t) \geq u$ if and only if there exists at least one extended outward local maximum above u on some face of T . Thus, under conditions **(H1)** and **(H2)**, the following relation holds for each $u \in \mathbb{R}$:

$$\left\{ \sup_{t \in T} X(t) \geq u \right\} = \bigcup_{k=0}^N \bigcup_{K \in \partial_k T} \{M_u^E(K) \geq 1\} \quad \text{a.s.} \quad (6.1)$$

This implies that the probability of the supremum of the Gaussian random field exceeding u is equal to the probability that there exists at least one extended outward local maximum above u on some face K of T . Therefore, we obtain the following upper bound for the excursion probability:

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \leq \sum_{k=0}^N \sum_{K \in \partial_k T} \mathbb{P}\{M_u^E(K) \geq 1\} \leq \sum_{k=0}^N \sum_{K \in \partial_k T} \mathbb{E}\{M_u^E(K)\}. \quad (6.2)$$

On the other hand, notice that

$$\begin{aligned} \mathbb{E}\{M_u^E(K)\} - \mathbb{P}\{M_u^E(K) \geq 1\} &= \sum_{i=1}^{\infty} (i-1) \mathbb{P}\{M_u^E(K) = i\} \\ &\leq \sum_{i=1}^{\infty} i(i-1) \mathbb{P}\{M_u^E(K) = i\} = \mathbb{E}\{M_u^E(K)[M_u^E(K) - 1]\} \end{aligned}$$

and

$$\mathbb{P}\{M_u^E(K) \geq 1, M_u^E(K') \geq 1\} \leq \mathbb{E}\{M_u^E(K)M_u^E(K')\}.$$

Applying the Bonferroni inequality to (6.1) and combining these two inequalities, we obtain the following lower bound for the excursion probability:

$$\begin{aligned} &\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} \\ &\geq \sum_{k=0}^N \sum_{K \in \partial_k T} \mathbb{P}\{M_u^E(K) \geq 1\} - \sum_{K \neq K'} \mathbb{P}\{M_u^E(K) \geq 1, M_u^E(K') \geq 1\} \\ &\geq \sum_{k=0}^N \sum_{K \in \partial_k T} (\mathbb{E}\{M_u^E(K)\} - \mathbb{E}\{M_u^E(K)[M_u^E(K) - 1]\}) - \sum_{K \neq K'} \mathbb{E}\{M_u^E(K)M_u^E(K')\}, \end{aligned} \quad (6.3)$$

where the last sum is taken over all possible pairs of different faces (K, K') .

Remark 6.1. Note that, following the same arguments above, we have that the expectations on the number of extended outward maxima $M_u^E(\cdot)$ in both (6.2) and (6.3) can be replaced by the expectations on the number of local maxima $M_u(\cdot)$.

We call a function $h(u)$ *super-exponentially small* [when compared with the excursion probability $\mathbb{P}\{\sup_{t \in T} X(t) \geq u\}$ or $\mathbb{E}\{\chi(A_u)\}$], if there exists a constant $\alpha > 0$ such that $h(u) = o(e^{-u^2/(2\sigma_T^2)} - \alpha u^2)$ as $u \rightarrow \infty$. The main idea for proving the EEC approximation Theorem 3.1 consists of the following two steps: (i) show that, except for the upper bound in (6.2), all terms in the lower bound in (6.3) are super-exponentially small; and

(ii) demonstrate that the difference between the upper bound in (6.2) and $\mathbb{E}\{\chi(A_u)\}$ is also super-exponentially small. The proofs for Theorems 3.2 and 3.3 follow the same ideas, aiming to establish super-exponential smallness for the terms involved in the lower bounds, as well as for the difference between the upper bound and EEC.

Remark 6.2. In terms of the detailed proofs in Sections 7 and 8 below, the methodologies are similar to those in [5]. The main difference is due to the condition **(H3)** or **(H3')**, which generalizes the condition **(H2)** in [5] where Gaussian fields with stationary increments are considered. It is important to recognize that in our current framework, there is no assumption of stationary increments; and the condition **(H3)** or **(H3')** is imposed only on the points attaining the maximum of the variance. Such general assumptions make the proofs more challenging, requiring improved and refined techniques on estimating the super-exponentially small errors, especially on estimating the adjacent faces in Proposition 7.4 and approximating the major term in Proposition 8.1. Meanwhile, due to the absence of stationary increments, we employ the Laplace method to derive general explicit approximations to excursion probabilities for smooth Gaussian fields with the maximum of variance achieved at a single point in Section 4, and to derive the approximation for an example with the maximum of variance achieved on a line in Section 5.

7 Estimation of super-exponential smallness for terms in the lower bound

7.1 Factorial moments

We first state the following result, which is a modified version (restricted on a face K) of Lemma 4 in Piterbarg [8], characterizing the decaying rate for factorial moments of the number of critical points exceeding a high level for Gaussian fields.

Lemma 7.1. *Assume **(H1)** and **(H2)**. Then there exists a positive constant C such that for any $\varepsilon > 0$ one can find a number $\varepsilon_1 > 0$ such that for any $K \in \partial_k T$,*

$$\mathbb{E}\{M_u(K)(M_u(K) - 1)\} \leq C u^{2k+1} \exp\left\{-\frac{u^2}{2\beta_K^2 + \varepsilon}\right\} + C u^{4k+2} \exp\left\{-\frac{u^2}{2\sigma_K^2 - \varepsilon_1}\right\}, \tag{7.1}$$

where

$$\beta_K^2 = \sup_{t \in K} \sup_{e \in \mathbb{S}^{k-1}} \text{Var}(X(t)|\nabla X|_K(t), \nabla^2 X|_K(t)e), \quad \sigma_K^2 = \sup_{t \in K} \text{Var}(X(t)).$$

The following result shows that the factorial moments in (6.3) are super-exponentially small under our assumptions.

Proposition 7.2. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying **(H1)**, **(H2)** and **(H3)**. Then there exists $\alpha > 0$ such that as $u \rightarrow \infty$,*

$$\sum_{k=0}^N \sum_{K \in \partial_k T} \mathbb{E}\{M_u(K)(M_u(K) - 1)\} = o\left(e^{-u^2/(2\sigma_T^2) - \alpha u^2}\right). \tag{7.2}$$

Proof. Due to Lemma 7.1, it suffices to show that for each $K \in \partial_k T$, $\beta_K^2 < \sigma_T^2$, which is equivalent to $\text{Var}(X(t)|\nabla X|_K(t), \nabla^2 X|_K(t)e) < \sigma_T^2$ for all $t \in \bar{K} = K \cup \partial K$ and $e \in \mathbb{S}^{k-1}$. Suppose $\text{Var}(X(t)|\nabla X|_K(t), \nabla^2 X|_K(t)e) = \sigma_T^2$ for some $t \in K$, then

$$\sigma_T^2 = \text{Var}(X(t)|\nabla X|_K(t), \nabla^2 X|_K(t)e) \leq \text{Var}(X(t)|\nabla^2 X|_K(t)e) \leq \text{Var}(X(t)) \leq \sigma_T^2.$$

Note that

$$\text{Var}(X(t)|\nabla^2 X|_K(t)e) = \text{Var}(X(t)) \Leftrightarrow \mathbb{E}\{X(t)(\nabla^2 X|_K(t)e)\} = 0 \Leftrightarrow \Sigma_K(t)e = 0.$$

But t is a point with $\nu(t) = \sigma_T^2$, thus $\Sigma_K(t) \prec 0$ by Proposition 2.1, implying $\Sigma_K(t)e \neq 0$ for all $e \in \mathbb{S}^{k-1}$ and causing a contradiction.

On the other hand, suppose $\text{Var}(X(t)|\nabla X|_K(t), \nabla^2 X|_K(t)e) = \sigma_T^2$ for some $t \in \partial K$, then $\text{Var}(X(t)|\nabla X|_K(t)) = \sigma_T^2$ and hence $\nu_i(t) = 0$ for all $i \in \tau(K)$, implying $\Sigma_K(t) \prec 0$ by Proposition 2.1. Similarly to the previous arguments, this will lead to a contradiction. The proof is completed. \square

7.2 Non-adjacent faces

For two sets $D, D' \subset \mathbb{R}^N$, let $d(D, D') = \inf\{\|t - t'\| : t \in D, t' \in D'\}$ denote their distance. The following result demonstrates that the last two sums involving the joint moment of two non-adjacent faces in (6.3) are super-exponentially small.

Proposition 7.3. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1) and (H2). Then there exists $\alpha > 0$ such that as $u \rightarrow \infty$,*

$$\mathbb{E}\{M_u(K)M_u(K')\} = o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right), \tag{7.3}$$

where K and K' are different faces of T with $d(K, K') > 0$.

Proof. Consider first the case where $\dim(K) = k \geq 1$ and $\dim(K') = k' \geq 1$. Applying the Kac-Rice formula in [2, Theorem 11.2.1] with $f = (\nabla X|_K(t), \nabla X|_{K'}(t'))$, we obtain

$$\begin{aligned} & \mathbb{E}\{M_u(K)M_u(K')\} \\ &= \int_K dt \int_{K'} dt' \mathbb{E}\{|\det \nabla^2 X|_K(t)| |\det \nabla^2 X|_{K'}(t')| \mathbb{1}_{\{X(t) \geq u, X(t') \geq u\}} \\ & \quad \times \mathbb{1}_{\{\nabla^2 X|_K(t) \prec 0, \nabla^2 X|_{K'}(t') \prec 0\}} | \nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0\} p_{\nabla X|_K(t), \nabla X|_{K'}(t')}(0, 0) \\ &\leq \int_K dt \int_{K'} dt' \int_u^\infty dx \int_u^\infty dx' p_{X(t), X(t')}(x, x') p_{\nabla X|_K(t), \nabla X|_{K'}(t')}(0, 0 | X(t) = x, X(t') = x') \\ & \quad \times \mathbb{E}\{|\det \nabla^2 X|_K(t)| |\det \nabla^2 X|_{K'}(t')| | X(t) = x, X(t') = x', \nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0\}. \end{aligned} \tag{7.4}$$

Notice that the following two inequalities hold: for constants a_{i_1} and b_{i_2} ,

$$\prod_{i_1=1}^k |a_{i_1}| \prod_{i_2=1}^{k'} |b_{i_2}| \leq \frac{\sum_{i_1=1}^k |a_{i_1}|^{k+k'} + \sum_{i_2=1}^{k'} |b_{i_2}|^{k+k'}}{k+k'};$$

and for any Gaussian variable ξ and positive integer m , by Jensen's inequality,

$$\begin{aligned} \mathbb{E}|\xi|^m &\leq \mathbb{E}(|\mathbb{E}\xi| + |\xi - \mathbb{E}\xi|)^m \leq 2^{m-1}(\mathbb{E}|\xi|^m + \mathbb{E}|\xi - \mathbb{E}\xi|^m) \\ &= 2^{m-1}(\mathbb{E}|\xi|^m + B_m(\text{Var}(\xi))^{m/2}), \end{aligned}$$

where B_m is some constant depending only on m . Combining these two inequalities with the well-known conditional formula for Gaussian variables, we obtain that there exist

We first consider the case when $k \geq 1$ and $l \geq 1$. By the Kac-Rice formula,

$$\begin{aligned} & \mathbb{E}\{M_u^E(K)M_u^E(K')\} \\ & \leq \int_{K \times K'} dt dt' \int_u^\infty dx \int_u^\infty dx' \int_0^\infty dz_{k+1} \cdots \int_0^\infty dz_{k+k'-m} \int_0^\infty dw_{m+1} \cdots \int_0^\infty dw_k \\ & \quad \mathbb{E}\{|\det \nabla^2 X|_K(t)|\det \nabla^2 X|_{K'}(t')|X(t) = x, X(t') = x', \nabla X|_K(t) = 0, X_{k+1}(t) = z_{k+1}, \\ & \quad \dots, X_{k+k'-m}(t) = z_{k+k'-m}, \nabla X|_{K'}(t') = 0, X_{m+1}(t') = w_{m+1}, \dots, X_k(t') = w_k\} \\ & \quad \times p_{t,t'}(x, x', 0, z_{k+1}, \dots, z_{k+k'-m}, 0, w_{m+1}, \dots, w_k) \\ & := \int_{K \times K'} A(t, t', u) dt dt', \end{aligned} \tag{7.9}$$

where $p_{t,t'}(x, x', 0, z_{k+1}, \dots, z_{k+k'-m}, 0, w_{m+1}, \dots, w_k)$ is the density of the joint distribution of the variables involved in the given condition. We define

$$\mathcal{M}_0 := \{t \in I : \nu(t) = \sigma_T^2, \nu_i(t) = 0, \forall i = 1, \dots, k + k' - m\}, \tag{7.10}$$

and consider two cases for \mathcal{M}_0 .

Case (i): $\mathcal{M}_0 = \emptyset$. Under this case, since I is a compact set, by the uniform continuity of conditional variance, there exist constants $\varepsilon_1, \delta_1 > 0$ such that

$$\sup_{t \in B(I, \delta_1), t' \in B'(I, \delta_1)} \text{Var}(X(t)|\nabla X|_K(t), \nabla X|_{K'}(t')) \leq \sigma_T^2 - \varepsilon_1, \tag{7.11}$$

where $B(I, \delta_1) = \{t \in K : d(t, I) \leq \delta_1\}$ and $B'(I, \delta_1) = \{t' \in K' : d(t', I) \leq \delta_1\}$. By partitioning $K \times K'$ into $B(I, \delta_1) \times B'(I, \delta_1)$ and $(K \times K') \setminus (B(I, \delta_1) \times B'(I, \delta_1))$ and applying the Kac-Rice formula, we obtain

$$\begin{aligned} & \mathbb{E}\{M_u(K)M_u(K')\} \\ & \leq \int_{(K \times K') \setminus (B(I, \delta_1) \times B'(I, \delta_1))} dt dt' p_{\nabla X|_K(t), \nabla X|_{K'}(t')}(0, 0) \\ & \quad \times \mathbb{E}\{|\det \nabla^2 X|_K(t)|\det \nabla^2 X|_{K'}(t')\mathbb{1}_{\{X(t) \geq u, X(t') \geq u\}}|\nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0\} \\ & + \int_{B(I, \delta_1) \times B'(I, \delta_1)} dt dt' p_{\nabla X|_K(t), \nabla X|_{K'}(t')}(0, 0) \\ & \quad \times \mathbb{E}\{|\det \nabla^2 X|_K(t)|\det \nabla^2 X|_{K'}(t')\mathbb{1}_{\{X(t) \geq u, X(t') \geq u\}}|\nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0\} \\ & := I_1(u) + I_2(u). \end{aligned} \tag{7.12}$$

Note that

$$\begin{aligned} (K \times K') \setminus (B(I, \delta_1) \times B'(I, \delta_1)) & = \left((K \setminus B(I, \delta_1)) \times B'(I, \delta_1)\right) \cup \left(B(I, \delta_1) \times (K \setminus B(I, \delta_1))\right) \\ & \quad \cup \left((K \setminus B(I, \delta_1)) \times (K \setminus B(I, \delta_1))\right), \end{aligned}$$

where each product on the right hand side consists of two sets with a positive distance. It then follows from Proposition 7.3 that $I_1(u)$ is super-exponentially small. On the other hand, since $\mathbb{1}_{\{X(t) \geq u, X(t') \geq u\}} \leq \mathbb{1}_{\{[X(t)+X(t')]/2 \geq u\}}$, one has

$$\begin{aligned} I_2(u) & \leq \int_{B(I, \delta_1) \times B'(I, \delta_1)} dt dt' \int_u^\infty dx p_{\frac{X(t)+X(t')}{2}}(x|\nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0) \\ & \quad \times \mathbb{E}\{|\det \nabla^2 X|_K(t)|\det \nabla^2 X|_{K'}(t')\mathbb{1}_{\{[X(t)+X(t')]/2 = x, \\ & \quad \nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0\}}p_{\nabla X|_K(t), \nabla X|_{K'}(t')}(0, 0)\}. \end{aligned} \tag{7.13}$$

Combining this with (7.11), we conclude that $I_2(u)$ and hence $\mathbb{E}\{M_u^E(X, K)M_u^E(X, K')\}$ are super-exponentially small.

Case (ii): $\mathcal{M}_0 \neq \emptyset$. Let

$$B(\mathcal{M}_0, \delta_2) := \{(t, t') \in K \times K' : d(t, \mathcal{M}_0) \vee d(t', \mathcal{M}_0) \leq \delta_2\},$$

where δ_2 is a small positive number to be specified. Note that, by the definitions of \mathcal{M}_0 and $B(\mathcal{M}_0, \delta_2)$, there exists $\varepsilon_2 > 0$ such that

$$\sup_{(t, t') \in (K \times K') \setminus B(\mathcal{M}_0, \delta_2)} \text{Var}([X(t) + X(t')]/2 | \nabla X|_K(t), \nabla X|_{K'}(t')) \leq \sigma_T^2 - \varepsilon_2. \tag{7.14}$$

Similarly to (7.13), we obtain that $\int_{(K \times K') \setminus B(\mathcal{M}_0, \delta_2)} A(t, t', u) dt dt'$ is super-exponentially small. It suffices to show below that $\int_{B(\mathcal{M}_0, \delta_2)} A(t, t', u) dt dt'$ is super-exponentially small.

Due to (H3) and Proposition 2.1, we can choose δ_2 small enough such that for all $(t, t') \in B(\mathcal{M}_0, \delta_2)$,

$$\Lambda_{K \cup K'}(t) := -\mathbb{E}\{X(t) \nabla^2 X|_{K \cup K'}(t)\} = -(\mathbb{E}\{X(t) X_{ij}(t)\})_{i, j=1, \dots, k+k'-m}$$

are positive definite. Let $\{e_1, e_2, \dots, e_N\}$ be the standard orthonormal basis of \mathbb{R}^N . For $t \in K$ and $t' \in K'$, let $e_{t, t'} = (t' - t) / \|t' - t\|$ and $\alpha_i(t, t') = \langle e_i, \Lambda_{K \cup K'}(t) e_{t, t'} \rangle$. Then

$$\Lambda_{K \cup K'}(t) e_{t, t'} = \sum_{i=1}^N \langle e_i, \Lambda_{K \cup K'}(t) e_{t, t'} \rangle e_i = \sum_{i=1}^N \alpha_i(t, t') e_i \tag{7.15}$$

and there exists $\alpha_0 > 0$ such that for all $(t, t') \in B(\mathcal{M}_0, \delta_2)$,

$$\langle e_{t, t'}, \Lambda_{K \cup K'}(t) e_{t, t'} \rangle \geq \alpha_0. \tag{7.16}$$

Since all elements in $\varepsilon(K)$ and $\varepsilon(K')$ are 1, we may write

$$\begin{aligned} t &= (t_1, \dots, t_m, t_{m+1}, \dots, t_k, b_{k+1}, \dots, b_{k+k'-m}, 0, \dots, 0), \\ t' &= (t'_1, \dots, t'_m, b_{m+1}, \dots, b_k, t'_{k+1}, \dots, t'_{k+k'-m}, 0, \dots, 0), \end{aligned}$$

where $t_i \in (a_i, b_i)$ for $i \in \tau(K)$ and $t'_j \in (a_j, b_j)$ for $j \in \tau(K')$. Therefore,

$$\begin{aligned} \langle e_i, e_{t, t'} \rangle &\geq 0, \quad \forall m+1 \leq i \leq k, \\ \langle e_i, e_{t, t'} \rangle &\leq 0, \quad \forall k+1 \leq i \leq k+k'-m, \\ \langle e_i, e_{t, t'} \rangle &= 0, \quad \forall k+k'-m < i \leq N. \end{aligned} \tag{7.17}$$

Let

$$\begin{aligned} D_i &= \{(t, t') \in B(\mathcal{M}_0, \delta_2) : \alpha_i(t, t') \geq \beta_i\}, \quad \text{if } m+1 \leq i \leq k, \\ D_i &= \{(t, t') \in B(\mathcal{M}_0, \delta_2) : \alpha_i(t, t') \leq -\beta_i\}, \quad \text{if } k+1 \leq i \leq k+k'-m, \\ D_0 &= \left\{ (t, t') \in B(\mathcal{M}_0, \delta_2) : \sum_{i=1}^m \alpha_i(t, t') \langle e_i, e_{t, t'} \rangle \geq \beta_0 \right\}, \end{aligned} \tag{7.18}$$

where $\beta_0, \beta_1, \dots, \beta_{k+k'-m}$ are positive constants such that $\beta_0 + \sum_{i=m+1}^{k+k'-m} \beta_i < \alpha_0$. It follows from (7.17) and (7.18) that, if (t, s) does not belong to any of $D_0, D_{m+1}, \dots, D_{k+k'-m}$, then by (7.15),

$$\langle \Lambda_{K \cup K'}(t) e_{t, t'}, e_{t, t'} \rangle = \sum_{i=1}^N \alpha_i(t, t') \langle e_i, e_{t, t'} \rangle \leq \beta_0 + \sum_{i=m+1}^{k+k'-m} \beta_i < \alpha_0,$$

which contradicts (7.16). Thus $D_0 \cup (\cup_{i=m+1}^{k+k'-m} D_i)$ is a covering of $B(\mathcal{M}_0, \delta_2)$. By (7.9),

$$\mathbb{E}\{M_u^E(K)M_u^E(K')\} \leq \int_{D_0} A(t, t', u) dt dt' + \sum_{i=m+1}^{k+k'-m} \int_{D_i} A(t, t', u) dt dt'.$$

By the Kac-Rice metatheorem and the fact $\mathbb{1}_{\{X(t) \geq u, Y(s) \geq u\}} \leq \mathbb{1}_{\{X(t) \geq u\}}$, we obtain

$$\begin{aligned} & \int_{D_0} A(t, t', u) dt dt' \\ & \leq \int_{D_0} dt dt' \int_u^\infty dx p_{\nabla X|_K(t), \nabla X|_{K'}(t')}(0, 0) p_{X(t)}(x | \nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0) \\ & \quad \times \mathbb{E}\{|\det \nabla^2 X|_K(t)| |\det \nabla^2 X|_{K'}(t') | X(t) = x, \nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0\}, \end{aligned} \tag{7.19}$$

and that for $i = m + 1, \dots, k$,

$$\begin{aligned} & \int_{D_i} A(t, t', u) dt dt' \\ & \leq \int_{D_i} dt dt' \int_u^\infty dx \int_0^\infty dw_i p_{X(t), \nabla X|_K(t), X_i(t'), \nabla X|_{K'}(t')}(x, 0, w_i, 0) \\ & \quad \times \mathbb{E}\{|\det \nabla^2 X|_K(t)| |\det \nabla^2 X|_{K'}(t') | X(t) = x, \nabla X|_K(t) = 0, X_i(t') = w_i, \nabla X|_{K'}(t') = 0\}. \end{aligned} \tag{7.20}$$

Comparing (7.19) and (7.20) with Eqs. (4.33) and (4.36) respectively in the proof of Theorem 4.8 in Cheng and Xiao [5], one can employ the same reasoning therein to show that $\text{Var}(X(t) | \nabla X|_K(t), \nabla X|_{K'}(t')) < \sigma_T^2$ uniformly on D_0 and $\mathbb{P}(X(t) > u, X_i(t') > 0 | \nabla X|_K(t) = 0, \nabla X|_{K'}(t') = 0) = o(e^{-u^2/(2\sigma_T^2)} - \alpha u^2)$ uniformly on D_i , and deduce that $\int_{D_0} A(t, t', u) dt dt'$ and $\int_{D_i} A(t, t', u) dt dt'$ ($i = m + 1, \dots, k$) are super-exponentially small.

It is similar to show that $\int_{D_i} A(t, t', u) dt dt'$ are super-exponentially small for $i = k + 1, \dots, k + k' - m$. For the case $k = 0$ or $l = 0$, the argument is even simpler when applying the Kac-Rice formula; the details are omitted here. The proof is finished. \square

In the proof of Proposition 7.4, we have shown in (7.12) that, if $\mathcal{M}_0 = \emptyset$, then the moment $\mathbb{E}\{M_u(X, K)M_u(X, K')\}$ is super-exponentially small. It is important to note that, the boundary condition (3.3) implies (and generalizes) the condition $\mathcal{M}_0 = \emptyset$, yielding the following result.

Proposition 7.5. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1), (H2) and the boundary condition (3.3). Then there exists $\alpha > 0$ such that as $u \rightarrow \infty$,*

$$\mathbb{E}\{M_u(K)M_u(K')\} = o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right),$$

where K and K' are different faces of T with $d(K, K') = 0$.

8 Estimation of the difference between EEC and the upper bound

In this section, we demonstrate that the difference between $\mathbb{E}\{\chi(A_u)\}$ and the expected number of extended outward local maxima, i.e. the upper bound in (6.2), is super-exponentially small.

Proposition 8.1. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1),*

(H2) and (H3). Then there exists $\alpha > 0$ such that for any $K \in \partial_k T$ with $k \geq 0$, as $u \rightarrow \infty$,

$$\begin{aligned} \mathbb{E}\{M_u^E(K)\} &= (-1)^k \int_K \mathbb{E}\{\det \nabla^2 X|_K(t) \mathbb{1}_{\{X(t) \geq u, \varepsilon_\ell^* X_\ell(t) \geq 0 \text{ for all } \ell \notin \tau(K)\}} | \nabla X|_K(t) = 0\} \\ &\quad \times p_{\nabla X|_K(t)}(0) dt + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right) \\ &= (-1)^k \mathbb{E}\left\{\left(\sum_{i=0}^k (-1)^i \mu_i(K)\right)\right\} + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right). \end{aligned} \tag{8.1}$$

Proof. The second equality in (8.1) arises from the application of the Kac-Rice formula:

$$\begin{aligned} &\mathbb{E}\left\{\left(\sum_{i=0}^k (-1)^i \mu_i(K)\right)\right\} \\ &= \sum_{i=0}^k (-1)^i \int_K \mathbb{E}\{|\det \nabla^2 X|_K(t)| \mathbb{1}_{\{\text{index}(\nabla^2 X|_K(t))=i\}} \\ &\quad \times \mathbb{1}_{\{X(t) \geq u, \varepsilon_\ell^* X_\ell(t) \geq 0 \text{ for all } \ell \notin \tau(K)\}} | \nabla X|_K(t) = 0\} p_{\nabla X|_K(t)}(0) dt \\ &= \int_K \mathbb{E}\{\det \nabla^2 X|_K(t) \mathbb{1}_{\{X(t) \geq u, \varepsilon_\ell^* X_\ell(t) \geq 0 \text{ for all } \ell \notin \tau(K)\}} | \nabla X|_K(t) = 0\} p_{\nabla X|_K(t)}(0) dt. \end{aligned}$$

To prove the first approximation in (8.1) and convey the main idea, we start with the case when the face K represents the interior of T .

Case (i): $k = N$. By the Kac-Rice formula, we have

$$\begin{aligned} \mathbb{E}\{M_u^E(K)\} &= \int_K p_{\nabla X(t)}(0) dt \int_u^\infty p_{X(t)}(x | \nabla X(t) = 0) \\ &\quad \times \mathbb{E}\{\det \nabla^2 X(t) \mathbb{1}_{\{\nabla^2 X(t) < 0\}} | X(t) = x, \nabla X(t) = 0\} dx \\ &:= \int_K p_{\nabla X(t)}(0) dt \int_u^\infty A(t, x) dx. \end{aligned}$$

Let

$$\begin{aligned} \mathcal{M}_1 &= \{t \in \bar{K} = T : \nu(t) = \sigma_T^2, \nabla \nu(t) = 2\mathbb{E}\{X(t)\nabla X(t)\} = 0\}, \\ B(\mathcal{M}_1, \delta_1) &= \{t \in K : d(t, \mathcal{M}_1) \leq \delta_1\}, \end{aligned} \tag{8.2}$$

where δ_1 is a small positive number to be specified. Then, we only need to estimate

$$\int_{B(\mathcal{M}_1, \delta_1)} p_{\nabla X(t)}(0) dt \int_u^\infty A(t, x) dx, \tag{8.3}$$

since the integral above with $B(\mathcal{M}_1, \delta_1)$ replaced by $K \setminus B(\mathcal{M}_1, \delta_1)$ becomes super-exponentially small due to the fact

$$\sup_{t \in K \setminus B(\mathcal{M}_1, \delta_1)} \text{Var}(X(t) | \nabla X(t) = 0) < \sigma_T^2.$$

Notice that, by Proposition 2.1, $\mathbb{E}\{X(t)\nabla^2 X(t)\} \prec 0$ for all $t \in \mathcal{M}_1$. Thus there exists δ_1 small enough such that $\mathbb{E}\{X(t)\nabla^2 X(t)\} \prec 0$ for all $t \in B(\mathcal{M}_1, \delta_1)$. In particular, let λ_0 be the largest eigenvalue of $\mathbb{E}\{X(t)\nabla^2 X(t)\}$ over $B(\mathcal{M}_1, \delta_1)$, then $\lambda_0 < 0$ by the uniform continuity. Also note that $\mathbb{E}\{X(t)\nabla X(t)\}$ tends to 0 as $\delta_1 \rightarrow 0$. Therefore, as $\delta_1 \rightarrow 0$,

$$\begin{aligned} &\mathbb{E}\{X_{ij}(t) | X(t) = x, \nabla X(t) = 0\} \\ &= (\mathbb{E}\{X_{ij}(t)X(t)\}, \mathbb{E}\{X_{ij}(t)X_1(t)\}, \dots, \mathbb{E}\{X_{ij}(t)X_N(t)\}) \cdot [\text{Cov}(X(t), \nabla X(t))]^{-1} \\ &\quad \cdot (x, 0, \dots, 0)^T \\ &= \frac{\mathbb{E}\{X_{ij}(t)X(t)\}x}{\sigma_T^2} (1 + o(1)). \end{aligned}$$

Thus, for all $x \geq u$ and $t \in B(\mathcal{M}_1, \delta_1)$ with δ_1 small enough,

$$\Sigma_1(t, x) := \mathbb{E}\{\nabla^2 X(t) | X(t) = x, \nabla X(t) = 0\} \prec 0.$$

Let $\Delta_1(t, x) = \nabla^2 X(t) - \Sigma_1(t, x)$. We have

$$\begin{aligned} \int_u^\infty A(t, x) dx &= \int_u^\infty p_{X(t)}(x | \nabla X(t) = 0) \mathbb{E}\{\det(\Delta_1(t, x) + \Sigma_1(t, x)) \\ &\quad \times \mathbb{1}_{\{\Delta_1(t, x) + \Sigma_1(t, x) \prec 0\}} | X(t) = x, \nabla X(t) = 0\} dx \\ &:= \int_u^\infty p_{X(t)}(x | \nabla X(t) = 0) E(t, x) dx. \end{aligned} \tag{8.4}$$

Note that the following is a centered Gaussian random matrix not depending on x :

$$\Omega(t) = (\Omega_{ij}(t))_{1 \leq i, j \leq N} = (\Delta_1(t, x) | X(t) = x, \nabla X(t) = 0).$$

Let $h_t(v)$ denote the density of the Gaussian random vector $((\Omega_{ij}(t))_{1 \leq i \leq j \leq N})$ with $v = (v_{ij})_{1 \leq i \leq j \leq N} \in \mathbb{R}^{N(N+1)/2}$. Then

$$\begin{aligned} E(t, x) &= \mathbb{E}\{\det(\Omega(t) + \Sigma_1(t, x)) \mathbb{1}_{\{\Omega(t) + \Sigma_1(t, x) \prec 0\}}\} \\ &= \int_{v: (v_{ij}) + \Sigma_1(t, x) \prec 0} \det((v_{ij}) + \Sigma_1(t, x)) h_t(v) dv, \end{aligned} \tag{8.5}$$

where (v_{ij}) is the abbreviation of the matrix $v = (v_{ij})_{1 \leq i, j \leq N}$. There exists a constant $c > 0$ such that for δ_1 small enough and all $t \in B(\mathcal{M}_1, \delta_1)$, and $x \geq u$, we have

$$(v_{ij}) + \Sigma_1(t, x) \prec 0, \quad \forall \|(v_{ij})\| := \left(\sum_{i,j=1}^N v_{ij}^2\right)^{1/2} < cu.$$

This implies $\{v : (v_{ij}) + \Sigma_1(t, x) \not\prec 0\} \subset \{v : \|(v_{ij})\| \geq cu\}$. Consequently, the integral in (8.5) with the domain of integration replaced by $\{v : (v_{ij}) + \Sigma_1(t, x) \not\prec 0\}$ is $o(e^{-\alpha' u^2})$ uniformly for all $t \in B(\mathcal{M}_1, \delta_1)$, where α' is a positive constant. As a result, we conclude that, uniformly for all $t \in B(\mathcal{M}_1, \delta_1)$ and $x \geq u$,

$$E(t, x) = \int_{\mathbb{R}^{N(N+1)/2}} \det((v_{ij}) + \Sigma_1(t, x)) h_t(v) dv + o(e^{-\alpha' u^2}).$$

By substituting this result into (8.4), we observe that the indicator function $\mathbb{1}_{\{\nabla^2 X(t) \prec 0\}}$ in (8.3) can be eliminated, causing only a super-exponentially small error. Thus, for sufficiently large u , there exists $\alpha > 0$ such that

$$\begin{aligned} \mathbb{E}\{M_u^E(K)\} &= \int_K p_{\nabla X(t)}(0) dt \int_u^\infty \mathbb{E}\{\det \nabla^2 X(t) | X(t) = x, \nabla X(t) = 0\} \\ &\quad \times p_{X(t)}(x | \nabla X(t) = 0) dx + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right). \end{aligned}$$

Case (ii): $k \geq 0$. It is worth noting that when $k = 0$, the terms in (8.1) related to the Hessian will vanish, simplifying the proof. Therefore, without loss of generality, let $k \geq 1$, $\tau(K) = \{1, \dots, k\}$ and assume all the elements in $\varepsilon(K)$ are 1. By the Kac-Rice formula,

$$\begin{aligned} \mathbb{E}\{M_u^E(K)\} &= (-1)^k \int_K p_{\nabla X_{|K}(t)}(0) dt \int_u^\infty p_{X(t)}(x | \nabla X_{|K}(t) = 0) \mathbb{E}\{\det \nabla^2 X_{|K}(t) \\ &\quad \times \mathbb{1}_{\{\nabla^2 X_{|K}(t) \prec 0\}} \mathbb{1}_{\{X_{k+1}(t) > 0, \dots, X_N(t) > 0\}} | X(t) = x, \nabla X_{|K}(t) = 0\} dx \\ &:= (-1)^k \int_K p_{\nabla X_{|K}(t)}(0) dt \int_u^\infty A'(t, x) dx. \end{aligned}$$

Let

$$\begin{aligned} \mathcal{M}_2 &= \{t \in \bar{K} : \nu(t) = \sigma_T^2, \nabla \nu|_K(t) = 2\mathbb{E}\{X(t)\nabla X|_K(t)\} = 0\}, \\ B(\mathcal{M}_2, \delta_2) &= \{t \in K : d(t, \mathcal{M}_2) \leq \delta_2\}, \end{aligned} \tag{8.6}$$

where δ_2 is another small positive number to be specified. Here, we only need to estimate

$$\int_{B(\mathcal{M}_2, \delta_2)} p_{\nabla X|_K(t)}(0) dt \int_u^\infty A'(t, x) dx, \tag{8.7}$$

since the integral above with $B(\mathcal{M}_2, \delta_2)$ replaced by $K \setminus B(\mathcal{M}_2, \delta_2)$ is super-exponentially small due to the fact

$$\sup_{t \in K \setminus B(\mathcal{M}_2, \delta_2)} \text{Var}(X(t) | \nabla X(t) = 0) < \sigma_T^2.$$

On the other hand, following similar arguments in the proof for Case (i), we have that removing the indicator functions $\mathbb{1}_{\{\nabla^2 X|_K(t) < 0\}}$ in (8.7) will only cause a super-exponentially small error. Combining these results, we conclude that the first approximation in (8.1) holds, thus completing the proof. \square

From the proof of Proposition 8.1, it is evident that the same line of reasoning and arguments can be readily extended to $\mathbb{E}\{M_u(X, K)\}$, leading to the following result.

Proposition 8.2. *Let $\{X(t), t \in T\}$ be a centered Gaussian random field satisfying (H1), (H2) and (H3). Then there exists a constant $\alpha > 0$ such that for any $K \in \partial_k T$, as $u \rightarrow \infty$,*

$$\begin{aligned} \mathbb{E}\{M_u(K)\} &= (-1)^k \int_K \mathbb{E}\{\det \nabla^2 X|_K(t) \mathbb{1}_{\{X(t) \geq u\}} | \nabla X|_K(t) = 0\} p_{\nabla X|_K(t)}(0) dt \\ &\quad + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right). \end{aligned}$$

9 Proofs of the main results

Proof of Theorem 3.1. By Propositions 7.2, 7.3 and 7.4, together with the fact $M_u^E(K) \leq M_u(K)$, we obtain that the factorial moments and the last two sums in (6.3) are super-exponentially small. Therefore, from (6.2) and (6.3), it follows that there exists a constant $\alpha > 0$ such that as $u \rightarrow \infty$,

$$\mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} = \sum_{k=0}^N \sum_{K \in \partial_k T} \mathbb{E}\{M_u^E(K)\} + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right).$$

This desired result follows as an immediate consequence of Proposition 8.1. \square

Proof of Theorem 3.2. Remark 6.1 indicates that both inequalities (6.2) and (6.3) hold with $M_u^E(\cdot)$ replaced by $M_u(\cdot)$. Therefore, the corresponding factorial moments and the last two sums in (6.3) with $M_u^E(\cdot)$ replaced by $M_u(\cdot)$ are super-exponentially small by Propositions 7.2, 7.3 and 7.5. Consequently, there exists a constant $\alpha > 0$ such that as $u \rightarrow \infty$,

$$\mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} = \sum_{k=0}^N \sum_{K \in \partial_k T} \mathbb{E}\{M_u(K)\} + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right).$$

The desired result follows directly from Proposition 8.2. \square

Proof of Theorem 3.3. Note that, in the proof of Theorem 3.1, we have seen that the points in \mathcal{M}_2 defined in (8.6) make major contribution to the excursion probability. That is, with up to a super-exponentially small error, we can focus only on those faces, say J , whose closure \bar{J} contains the unique point t^* with $\nu(t^*) = \sigma_T^2$ and satisfying $\tau(J) \subset \mathcal{I}(t^*)$ (i.e., the partial derivatives of ν are 0 at t^* restricted on J). To formalize this concept, we define a set of faces T^* as follows:

$$T^* = \{J \in \partial_k T : t^* \in \bar{J}, \tau(J) \subset \mathcal{I}(t^*), k = 0, \dots, N\}.$$

For each $J \in T^*$, let

$$M_u^{E^*}(J) := \#\{t \in J : X(t) \geq u, \nabla X|_J(t) = 0, \nabla^2 X|_J(t) \prec 0, \varepsilon_j^* X_j(t) \geq 0 \text{ for all } j \in \mathcal{I}(t^*) \setminus \tau(J)\}.$$

Note that, both inequalities (6.2) and (6.3) remain valid when we replace $M_u^E(J)$ with $M_u^{E^*}(J)$ for faces J belonging to T^* , and replace $M_u^E(J)$ with $M_u(J)$ otherwise. Employing analogous reasoning as used in the derivation of Theorems 3.1 and 3.2, we obtain that, there exists $\alpha > 0$ such that as $u \rightarrow \infty$,

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} = \sum_{J \in T^*} \mathbb{E}\{M_u^{E^*}(J)\} + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right).$$

This desired result is then deduced from Proposition 8.1. □

10 Proofs on results with a unique maximum point of the variance

We begin by presenting the following two auxiliary results on the Laplace method for integral approximations. Lemma 10.1 can be found in many books on the approximations of integrals; here we refer to Wong [17]. Lemma 10.2 can be derived by following similar arguments in the proof of the Laplace method for the case of boundary points in [17].

Lemma 10.1. [Laplace method for interior points] *Let t_0 be an interior point of T . Suppose the following conditions hold: (i) $g(t) \in C(T)$ and $g(t_0) \neq 0$; (ii) $h(t) \in C^2(T)$ and attains its minimum only at t_0 ; and (iii) $\nabla^2 h(t_0)$ is positive definite. Then as $u \rightarrow \infty$,*

$$\int_T g(t) e^{-uh(t)} dt = \frac{(2\pi)^{N/2}}{u^{N/2} (\det \nabla^2 h(t_0))^{1/2}} g(t_0) e^{-uh(t_0)} (1 + o(1)).$$

Lemma 10.2. [Laplace method for boundary points] *Let $t_0 \in K \in \partial_k T$ with $0 \leq k \leq N-1$. Suppose that conditions (i), (ii) and (iii) in Lemma 10.1 hold, and additionally $\nabla h(t_0) = 0$. Then as $u \rightarrow \infty$,*

$$\int_T g(t) e^{-uh(t)} dt = \frac{(2\pi)^{N/2} \mathbb{P}\{Z_{i_\ell} \varepsilon_{i_\ell}^* > 0, \forall i_\ell \notin \tau(K)\}}{u^{N/2} (\det \nabla^2 h(t_0))^{1/2}} g(t_0) e^{-uh(t_0)} (1 + o(1)),$$

where $(Z_{i_1}, \dots, Z_{i_{N-k}})$ is a centered $(N-k)$ -dimensional Gaussian vector with covariance matrix $(h_{i_\ell i_{\ell'}}(t_0))_{i_\ell, i_{\ell'} \notin \tau(K)}$ and $\tau(K)$ and $\varepsilon_{i_\ell}^*$ are defined in Section 2.

We now provide below the proofs of Theorems 4.1 and 4.2.

Proof of Theorem 4.1. For $t \in T$, we define the following notation for conditional variance $\tilde{\nu}|_K(t) = \text{Var}(X(t) | \nabla X|_K(t) = 0)$. If $k = 0$, then $\nu_i(t^*) \neq 0$ for all $i \geq 1$, and hence $\mathcal{I}(t^*) = \emptyset$. The first line of (4.1) follows from Theorem 3.3 that

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} = \mathbb{P}\{X(t^*) \geq u\} + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right).$$

Now, let us consider the case when $k \geq 1$. Note that the assumption on partial derivatives of $\nu(t)$ implies $\mathcal{I}(t^*) = \tau(K)$. By Theorem 3.3, we have

$$\mathbb{P} \left\{ \sup_{t \in T} X(t) \geq u \right\} = (-1)^k I(u, K) + o \left(\exp \left\{ -\frac{u^2}{2\sigma_T^2} - \alpha u^2 \right\} \right), \quad (10.1)$$

where

$$\begin{aligned} I(u, K) &= \int_K \mathbb{E} \{ \det \nabla^2 X_{|K}(t) \mathbb{1}_{\{X(t) \geq u\}} | \nabla X_{|K}(t) = 0 \} p_{\nabla X_{|K}(t)}(0) dt \\ &= \int_u^\infty \int_K \frac{(2\pi)^{-(k+1)/2}}{\sqrt{\tilde{\nu}_{|K}(t)} \det(\Lambda_K(t))} \mathbb{E} \{ \det \nabla^2 X_{|K}(t) | X(t) = x, \nabla X_{|K}(t) = 0 \} \\ &\quad \times e^{-\frac{x^2}{2\nu(t)}} dt dx. \end{aligned}$$

Applying the Laplace method in Lemma 10.1 with

$$\begin{aligned} g(t) &= \frac{1}{\sqrt{\tilde{\nu}_{|K}(t)} \det(\Lambda_K(t))} \mathbb{E} \{ \det \nabla^2 X_{|K}(t) | X(t) = x, \nabla X_{|K}(t) = 0 \}, \\ h(t) &= \frac{1}{2\tilde{\nu}_{|K}(t)}, \quad u = x^2, \end{aligned}$$

and noting that the Hessian matrix of $1/(2\tilde{\nu}_{|K}(t))$ evaluated at t^* is

$$-\frac{1}{2\tilde{\nu}_{|K}^2(t^*)} (\tilde{\nu}_{ij}(t^*))_{i,j \in \tau(K)} = -\frac{1}{2\sigma_T^4} \nabla^2 \tilde{\nu}_{|K}(t^*) \succ 0, \quad (10.2)$$

we obtain

$$I(u, K) = \frac{(2\sigma_T^4)^{k/2}}{\sqrt{2\pi\sigma_T^2 \det(\Lambda_K(t^*))} \sqrt{|\det \nabla^2 \tilde{\nu}_{|K}(t^*)|}} I(u)(1 + o(1)), \quad (10.3)$$

where

$$\begin{aligned} I(u) &= \int_u^\infty \mathbb{E} \{ \det \nabla^2 X_{|K}(t^*) | X(t^*) = x, \nabla X_{|K}(t^*) = 0 \} x^{-k} e^{-\frac{x^2}{2\sigma_T^2}} dx \\ &= \det(\Sigma_K(t^*)) \int_u^\infty \mathbb{E} \{ \det(Q \nabla^2 X_{|K}(t^*) Q) | X(t^*) = x, \nabla X_{|K}(t^*) = 0 \} x^{-k} e^{-\frac{x^2}{2\sigma_T^2}} dx. \end{aligned} \quad (10.4)$$

Here, noting that $\Sigma_K(t^*) = \mathbb{E} \{ X(t) \nabla^2 X_{|K}(t^*) \} \prec 0$ by Proposition 2.1, we let Q in (10.4) be a $k \times k$ positive definite matrix such that $Q(-\Sigma_K(t^*))Q = I_k$, where I_k is the size- k identity matrix. Then

$$\mathbb{E} \{ X(t) (Q \nabla^2 X_{|K}(t^*) Q) \} = Q \Sigma_K(t^*) Q = -I_k.$$

Notice that $\mathbb{E} \{ X(t^*) \nabla X_{|K}(t^*) \} = 0$ due to $\nu_{|K}(t^*) = 0$, we have

$$\mathbb{E} \{ Q \nabla^2 X_{|K}(t^*) Q | X(t^*) = x, \nabla X_{|K}(t^*) = 0 \} = -\frac{x}{\sigma_T^2} I_k.$$

One can write

$$\mathbb{E} \{ \det(Q \nabla^2 X_{|K}(t^*) Q) | X(t^*) = x, \nabla X_{|K}(t^*) = 0 \} = \mathbb{E} \{ \det(\Delta(t^*) - (x/\sigma_T^2) I_k) \},$$

where $\Delta(t^*)$ is a centered Gaussian random matrix with covariance independent of x . According to the Laplace expansion of determinant, $\mathbb{E} \{ \det(\Delta(t^*) - (x/\sigma_T^2) I_k) \}$ is a

polynomial in x with the highest-order term being $(-1)^k \sigma_T^{-2k} x^k$. Plugging this into (10.4) and (10.3), we obtain

$$I(u, K) = \frac{(-1)^k 2^{k/2} |\det(\Sigma_K(t^*))|}{\sqrt{\det(\Lambda_K(t^*))} \sqrt{|\det(\nabla^2 \tilde{\nu}_{|K}(t^*))|}} \Psi\left(\frac{u}{\sigma_T}\right) (1 + o(1)).$$

Finally, note that

$$\tilde{\nu}_{|K}(t) = \mathbb{E}\{X(t)^2\} - \mathbb{E}\{X(t)\nabla X_{|K}(t)\}^T \Lambda_K^{-1}(t) \mathbb{E}\{X(t)\nabla X_{|K}(t)\},$$

we have

$$\begin{aligned} \nabla^2 \tilde{\nu}_{|K}(t^*) &= 2[\Lambda_K(t^*) + \Sigma_K(t^*)] - 2[\Lambda_K(t^*) + \Sigma_K(t^*)] \Lambda_K^{-1}(t^*) [\Lambda_K(t^*) + \Sigma_K(t^*)] \\ &= -2\Sigma_K(t^*) [I_k + \Lambda_K^{-1}(t^*) \Sigma_K(t^*)]. \end{aligned} \tag{10.5}$$

Therefore,

$$I(u, K) = (-1)^k \sqrt{\frac{\det(\Sigma_K(t^*))}{\det(\Lambda_K(t^*) + \Sigma_K(t^*))}} \Psi\left(\frac{u}{\sigma_T}\right) (1 + o(1)),$$

where $\Sigma_K(t^*) \prec 0$ by Proposition 2.1 and $\Lambda_K(t^*) + \Sigma_K(t^*) = \nabla^2 \nu_{|K}(t^*)/2 \prec 0$ by assumption. Plugging this into (10.1) yields the desired result. \square

Proof of Theorem 4.2. We first prove the case when $k \geq 1$. By Theorem 3.3, we have

$$\mathbb{P}\left\{\sup_{t \in T} X(t) \geq u\right\} = \sum_J (-1)^j I(u, J) + o\left(\exp\left\{-\frac{u^2}{2\sigma_T^2} - \alpha u^2\right\}\right), \tag{10.6}$$

where $j = \dim(J)$, the sum is taken over all faces J such that $t^* \in \bar{J}$ and $\tau(J) \subset \mathcal{I}(t^*)$, and

$$\begin{aligned} I(u, J) &= \int_J \mathbb{E}\{\det \nabla^2 X_{|J}(t) \mathbb{1}_{\{X(t) \geq u\}} \mathbb{1}_{\{\varepsilon_\ell^* X_\ell(t) \geq 0, \forall \ell \in \mathcal{I}(t^*) \setminus \tau(J)\}} | \nabla X_{|J}(t) = 0\}} \\ &\quad \times p_{\nabla X_{|J}(t)}(0) dt \\ &= \int_u^\infty \int_J \frac{(2\pi)^{-(j+1)/2}}{\sqrt{\tilde{\nu}_{|J}(t) \det(\Lambda_J(t))}} \mathbb{E}\{\det \nabla^2 X_{|K}(t) \mathbb{1}_{\{\varepsilon_\ell^* X_\ell(t) \geq 0, \forall \ell \in \mathcal{I}(t^*) \setminus \tau(J)\}} | X(t) = x, \\ &\quad \nabla X_{|J}(t) = 0\} e^{-\frac{x^2}{2\tilde{\nu}_{|J}(t)}} dt dx. \end{aligned}$$

Applying the Laplace method in Lemma 10.2 with

$$\begin{aligned} g(t) &= \frac{\mathbb{E}\{\det \nabla^2 X_{|J}(t) \mathbb{1}_{\{\varepsilon_\ell^* X_\ell(t) \geq 0, \forall \ell \in \mathcal{I}(t^*) \setminus \tau(J)\}} | X(t) = x, \nabla X_{|J}(t) = 0\}}{\sqrt{\tilde{\nu}_{|J}(t) \det(\Lambda_J(t))}}, \\ h(t) &= \frac{1}{2\tilde{\nu}_{|J}(t)}, \quad u = x^2, \end{aligned}$$

we obtain

$$I(u, J) = \frac{(2\sigma_T^4)^{j/2} \mathbb{P}\{(Z_{J'_1}, \dots, Z_{J'_{j-k}}) \in E'(J)\}}{\sqrt{2\pi\sigma_T^2 \det(\Lambda_J(t^*))} \sqrt{|\det \nabla^2 \tilde{\nu}_{|J}(t^*)|}} I(u) (1 + o(1)),$$

where $(Z_{J_1}, \dots, Z_{J_{j-k}})$ is a centered $(j - k)$ -dimensional Gaussian vector having covariance matrix $\nabla^2 h(t^*)$ with indices restricted on $\tau(J) \setminus \tau(K)$, and

$$\begin{aligned} I(u) &= \int_u^\infty \mathbb{E}\{\det \nabla^2 X_{|J}(t^*) \mathbb{1}_{\{\varepsilon_\ell^* X_\ell(t^*) \geq 0, \forall \ell \in \mathcal{I}(t^*) \setminus \tau(J)\}} | X(t^*) = x, \nabla X_{|J}(t^*) = 0\} \\ &\quad \times x^{-j} e^{-\frac{x^2}{2\sigma_T^2}} dx \\ &= \det(\Sigma_J(t^*)) \int_u^\infty \int_{E(J)} \mathbb{E}\{\det(Q \nabla^2 X_{|J}(t^*) Q) | X(t^*) = x, \nabla X_{|J}(t^*) = 0, \\ &\quad X_{J_1}(t^*) = y_{J_1}, \dots, X_{J_{k+m-j}}(t^*) = y_{J_{k+m-j}}\} p(y_{J_1}, \dots, y_{J_{k+m-j}} | x, 0) \\ &\quad \times x^{-j} e^{-\frac{x^2}{2\sigma_T^2}} dy_{J_1} \cdots dy_{J_{k+m-j}} dx. \end{aligned} \tag{10.7}$$

Here $p(y_{J_1}, \dots, y_{J_{k+m-j}} | x, 0)$ is the pdf of $(X_{J_1}(t^*), \dots, X_{J_{k+m-j}}(t^*) | X(t^*) = x, \nabla X_{|J}(t^*) = 0)$, and Q is a $j \times j$ positive definite matrix such that $Q(-\Sigma_J(t^*))Q = I_j$. Then, similarly to the arguments in the proof of Theorem 4.1, one can write the last expectation in (10.7) as

$$\mathbb{E}\{\det(\Delta(t^*, y_{J_1}, \dots, y_{J_{k+m-j}}) - (x/\sigma_T^2)I_k)\},$$

where $\Delta(t^*, y_{J_1}, \dots, y_{J_{k+m-j}})$ is a centered Gaussian random matrix with covariance independent of x , and hence the highest-order term in x is $(-1)^j x^j / \sigma_T^{2j}$. Noting that $\mathbb{E}\{X(t^*)X_i(t^*)\} = 0$ for all $i \in \mathcal{I}(t^*)$ and following similar arguments in the proof of Theorem 4.1, we obtain

$$\begin{aligned} I(u, J) &= (-1)^j \sqrt{\frac{\det(\Sigma_J(t^*))}{\det(\Lambda_J(t^*) + \Sigma_J(t^*))}} \mathbb{P}\{(Z_{J_1}, \dots, Z_{J_{j-k}}) \in E'(J)\} \\ &\quad \times \mathbb{P}\{(X_{J_1}(t^*), \dots, X_{J_{k+m-j}}(t^*)) \in E(J) | \nabla X_{|J}(t^*) = 0\} \Psi\left(\frac{u}{\sigma_T}\right) (1 + o(1)), \end{aligned}$$

which yields the desired result together with (10.6). In particular, by (10.5), one can treat $(Z_{J_1}, \dots, Z_{J_{j-k}})$ having covariance $\Sigma(t^*) + \Sigma(t^*)\Lambda^{-1}(t^*)\Sigma(t^*)$ with indices restricted on $\tau(J) \setminus \tau(K)$ while not changing the probability that it falls in $E(J)$. Lastly, the case when $k = 0$ can be shown similarly. \square

References

- [1] Adler, R. J. (2000). On excursion sets, tube formulas and maxima of random fields. *Ann. Appl. Probab.* **10**, 1–74. MR1765203
- [2] Adler, R. J. and Taylor, J. E. (2007). *Random fields and geometry*. Springer, New York. MR2319516
- [3] Azaïs, J. M. and Delmas, C. (2002). Asymptotic expansions for the distribution of the maximum of Gaussian random fields. *Extremes.* **5**, 181–212. MR1965978
- [4] Azaïs, J. M. and Wschebor, M. (2009). *Level Sets and Extrema of Random Processes and Fields*. John Wiley & Sons, Hoboken, NJ. MR2478201
- [5] Cheng, D. and Xiao, Y. (2016). The mean Euler characteristic and excursion probability of Gaussian random fields with stationary increments. *Ann. Appl. Probab.* **26**, 722–759. MR3476623
- [6] Kobelkov, S. G. and Piterbarg, V. I., (2019). On maximum of Gaussian random field having unique maximum point of its variance. *Extremes.* **22**, 413–432. MR3984356
- [7] Piterbarg, V. I. (1996a). *Asymptotic Methods in the Theory of Gaussian Processes and Fields. Translations of Mathematical Monographs 148*. Amer. Math. Soc., Providence, RI. MR1361884

- [8] Piterbarg, V. I. (1996b). Rice's method for large excursions of Gaussian random fields. Technical Report NO. 478, Center for Stochastic Processes, Univ. North Carolina.
- [9] Siegmund, D. and Worsley, K. J. (1995). Testing for a signal with unknown location and scale in a stationary Gaussian random field. *Ann. Statist.* **23**, 608–639. MR1332585
- [10] Sun, J. (1993). Tail probabilities of the maxima of Gaussian random fields. *Ann. Probab.* **21**, 34–71. MR1207215
- [11] Sun, J. (2001). Multiple comparisons for a large number of parameters. *Biometrical Journal* **43**, 627–643. MR1859377
- [12] Taylor, J. E. and Adler, R. J. (2003). Euler characteristics for Gaussian fields on manifolds. *Ann. Probab.* **31**, 533–563. MR1964940
- [13] Taylor, J. E., Takemura, A. and Adler, R. J. (2005). Validity of the expected Euler characteristic heuristic. *Ann. Probab.* **33**, 1362–1396. MR2150192
- [14] Taylor, J. E. and Worsley, K. J. (2007). Detecting sparse signals in random fields, with an application to brain mapping. *J. Amer. Statist. Assoc.* **102**, 913–928. MR2354405
- [15] Taylor, J. E. and Worsley, K. J. (2008). Random fields of multivariate test statistics, with applications to shape analysis. *Ann. Statist.* **36**, 1–27. MR2387962
- [16] Telschow, F., Cheng, D., Pranav, P. and Schwartzman, A. (2023). Estimation of expected Euler characteristic curves of nonstationary smooth random fields. *Ann. Statist.* **51**, 2272–2297. MR4678804
- [17] Wong, R. (2001). *Asymptotic Approximations of Integrals*. SIAM, Philadelphia, PA. MR1851050

Acknowledgments. The author is grateful to two referees for their efforts and valuable comments, which have led to great improvement for this paper.