

ON THE OCCUPATION TIME OF BROWNIAN EXCURSION

GERARD HOOGHIEMSTRA

Faculty ITS, Technical University Delft, Holland.

email: G.Hooghiemstra@its.tudelft.nl

submitted January 25, 1999; *revised* August 4, 1999AMS subject classification: 60J65
Brownian excursion, occupation time*Abstract*

Recently, Kalvin M. Jansons derived in an elegant way the Laplace transform of the time spent by an excursion above a given level $a > 0$. This result can also be derived from previous work of the author on the occupation time of the excursion in the interval $(a, a + b]$, by sending $b \rightarrow \infty$. Several alternative derivations are included.

1 Introduction

In [5], the author derives in an elegant way the Laplace transform of the time spent by an excursion above a given level $a > 0$. This result can also be derived from the occupation time of the excursion in the interval $(a, a + b]$, by sending $b \rightarrow \infty$ (cf. [2] or [4]).

2 Occupation times

Introduce for α, β complex and $a \geq 0$,

$$\psi(\alpha, \beta, a) = \left[\frac{\alpha \cosh(a\beta) + \beta \sinh(a\beta)}{\alpha \sinh(a\beta) + \beta \cosh(a\beta)} \right].$$

Denote by W_0^+ , Brownian excursion with time parameter $t \in [0, 1]$, see [4], I.2 for a precise definition. According to p. 117 and p. 120 of [4], or Theorem 5.1 of [2], the Laplace transform of the occupation time $T(a, a + b) = \int_0^1 1_{(a, a+b]}(W_0^+(t)) dt$, is given by:

$$E e^{-\beta T(a, a+b)} = \frac{\sqrt{2\pi}}{a^3} \sum_{k=1}^{\infty} k^2 \pi^2 e^{-k^2 \pi^2 / 2a^2} + \frac{1}{i\sqrt{\pi}} \int_S \frac{\alpha e^\alpha}{\sinh\{a\sqrt{2\alpha}\}} \times [\sqrt{\alpha} \cosh\{a\sqrt{2\alpha}\} + (\alpha + \beta)^{1/2} \psi(\sqrt{\alpha}, \sqrt{\alpha + \beta}, b\sqrt{2}) \sinh\{a\sqrt{2\alpha}\}]^{-1} d\alpha, \quad (1)$$

where the path S is defined by

$$S = \{\alpha : \alpha = iy, |y| \geq \xi\} \cup \{\alpha : \alpha = \xi e^{i\eta}, -\pi/2 \leq \eta \leq \pi/2\},$$

for some $\xi > 0$.

In order to write the first term on the right side of (1), which term is equal to the distribution function of the supremum of Brownian excursion,¹ as a complex integral we introduce the path:

$$\Gamma = \{\alpha : \alpha = ye^{\pm i\phi}, y \geq \xi\} \cup \{\alpha : \alpha = \xi e^{i\eta}, -\phi \leq \eta \leq \phi\},$$

with $\pi/2 < \phi < \pi$, $\xi > 0$ and the orientation counterclockwise. We choose the angle ϕ in such a way that all singularities of the integrand in (1) remain on the left of the path Γ . Then

$$\frac{\sqrt{2\pi}}{a^3} \sum_{k=1}^{\infty} k^2 \pi^2 e^{-k^2 \pi^2 / 2a^2} = -\frac{1}{i\sqrt{\pi}} \int_{\Gamma} \sqrt{\alpha} e^{\alpha} \frac{\cosh\{a\sqrt{2\alpha}\}}{\sinh\{a\sqrt{2\alpha}\}} d\alpha, \quad (2)$$

since the integrand has only simple poles at $\alpha_k = -k^2 \pi^2 / 2a^2$, $k \geq 1$. Combining (1) and (2) and deforming the path S into the path Γ (again using Cauchy's theorem), yields

$$Ee^{-\beta T(a,a+b)} = -\frac{1}{i\sqrt{\pi}} \int_{\Gamma} \sqrt{\alpha} e^{\alpha} d\alpha \quad (3)$$

$$\times \left[\frac{\sqrt{\alpha} \sinh\{a\sqrt{2\alpha}\} + (\alpha + \beta)^{1/2} \psi(\sqrt{\alpha}, \sqrt{\alpha + \beta}, b\sqrt{2}) \cosh\{a\sqrt{2\alpha}\}}{\sqrt{\alpha} \cosh\{a\sqrt{2\alpha}\} + (\alpha + \beta)^{1/2} \psi(\sqrt{\alpha}, \sqrt{\alpha + \beta}, b\sqrt{2}) \sinh\{a\sqrt{2\alpha}\}} \right].$$

By taking the limit for $b \rightarrow \infty$, $(\psi(\cdot, \cdot, b\sqrt{2}) \rightarrow 1$, uniformly on compacta of Γ) we obtain for the Laplace transform of the occupation time $T(a) = T(a, \infty)$,

$$Ee^{-\beta T(a)} = -\frac{1}{i\sqrt{\pi}} \int_{\Gamma} \sqrt{\alpha} e^{\alpha} \psi(\sqrt{\alpha + \beta}, \sqrt{\alpha}, a\sqrt{2}) d\alpha. \quad (4)$$

Alternatively, one could take the limit for $a \downarrow 0$ in (3), resulting in the transform: $Ee^{-\beta(1-T(b))}$. For the occupation time $T_t(a)$ of the excursion straddling t , we have

$$T_t(a) \stackrel{d}{=} (L_t)^{1/2} T(a(L_t)^{-1/2}), \quad (5)$$

with $T(a)$ and L_t independent, and where L_t denotes the length of the excursion. It is readily verified from the density of L_t , see [1], (4.4), that for integrable φ ,

$$\int_0^{\infty} e^{-\alpha t} E\varphi(L_t) dt = \frac{1}{2\sqrt{\pi\alpha^3}} \int_0^{\infty} \varphi(y)(1 - e^{-\alpha y}) dy. \quad (6)$$

¹According to the Poisson-summation formula

$$\frac{\sqrt{2\pi}}{a^3} \sum_{k=1}^{\infty} k^2 \pi^2 e^{-k^2 \pi^2 / 2a^2} = 1 + 2 \sum_{k=1}^{\infty} (1 - 4k^2 a^2) e^{-2k^2 a^2},$$

which is the more familiar form of this distribution function.

Hence, using (5) and (6), the Laplace transform (4) yields the double Laplace transform:

$$\begin{aligned} & \int_0^\infty e^{-\alpha t} E e^{-\beta T_t(a)} dt \tag{7} \\ &= \frac{1}{\alpha} \psi(\sqrt{\alpha + \beta}, \sqrt{\alpha}, a\sqrt{2}) - \frac{1}{\alpha^{3/2}} \lim_{\alpha \downarrow 0} \sqrt{\alpha} \psi(\sqrt{\alpha + \beta}, \sqrt{\alpha}, a\sqrt{2}) \\ &= \frac{1}{\alpha} \left[\frac{\sqrt{\alpha} \sinh\{a\sqrt{2\alpha}\} + \sqrt{\alpha + \beta} \cosh\{a\sqrt{2\alpha}\}}{\sqrt{\alpha} \cosh\{a\sqrt{2\alpha}\} + \sqrt{\alpha + \beta} \sinh\{a\sqrt{2\alpha}\}} \right] - \frac{1}{\alpha^{3/2}} \frac{\sqrt{\beta}}{1 + a\sqrt{2\beta}}. \end{aligned}$$

This result can also be derived starting from reflected Brownian motion $|W|$ (cf. [3], p. 92, Remark (3.20)).

Perhaps the most elegant formulation of the Laplace transform of the occupation time is that for β strictly positive

$$\begin{aligned} & \frac{1}{\sqrt{2\pi}} \int_0^\infty \frac{e^{-\alpha x}}{x^{3/2}} \left[1 - E e^{-\beta x T(x^{-1/2})} \right] dx \tag{8} \\ &= \frac{2\sqrt{2\alpha}(\sqrt{\alpha + \beta} - \sqrt{\alpha})}{(\sqrt{\alpha} + \sqrt{\alpha + \beta})e^{2\sqrt{2\alpha}} + (\sqrt{\alpha} - \sqrt{\alpha + \beta})}. \end{aligned}$$

Equation (8) can be derived as follows. On the path Γ we have:

$$\begin{aligned} 1 - E e^{-\beta T(a)} &= -\frac{1}{i\sqrt{\pi}} \int_\Gamma \sqrt{\alpha} e^\alpha d\alpha + \frac{1}{i\sqrt{\pi}} \int_\Gamma \sqrt{\alpha} e^\alpha \psi(\sqrt{\alpha + \beta}, \sqrt{\alpha}, a\sqrt{2}) d\alpha \\ &= \frac{1}{i\sqrt{\pi}} \int_\Gamma \sqrt{\alpha} e^\alpha \left[\frac{2(\sqrt{\alpha + \beta} - \sqrt{\alpha})e^{-a\sqrt{2\alpha}}}{(\sqrt{\alpha} + \sqrt{\alpha + \beta})e^{a\sqrt{2\alpha}} + (\sqrt{\alpha} - \sqrt{\alpha + \beta})e^{-a\sqrt{2\alpha}}} \right] d\alpha. \end{aligned}$$

Now for $a > 0$ the integral over the path Γ may be replaced by integration over the line $(c - i\infty, c + i\infty)$, where $c > 0$ is arbitrary. Hence after the substitution $\alpha = xz$, with x positive and replacement of the path $(c/x - i\infty, c/x + i\infty)$ by the path $(c - i\infty, c + i\infty)$, we obtain

$$\begin{aligned} & x^{-3/2} \left(1 - E e^{-\beta x T(x^{-1/2})} \right) \\ &= \frac{1}{i\sqrt{\pi}} \int_{c-i\infty}^{c+i\infty} \sqrt{z} e^{xz} \left[\frac{2(\sqrt{z + \beta} - \sqrt{z})e^{-\sqrt{2z}}}{(\sqrt{z} + \sqrt{z + \beta})e^{\sqrt{2z}} + (\sqrt{z} - \sqrt{z + \beta})e^{-\sqrt{2z}}} \right] dz. \end{aligned}$$

Taking Laplace transforms on both sides gives (8).

Each of the representations (4), (7) or (8) is equivalent to Theorem 1 of [5], where the duration of the excursion was scaled with a gamma($\frac{1}{2}, \frac{1}{2}\nu^2$) density. In particular, Theorem 1 of [5] can be obtained from (8) by differentiating both sides with respect to α and using that

$$\int_0^\infty x^{-1/2} e^{-\alpha x} dx = \sqrt{\pi/\alpha}.$$

References

- [1] K.L. CHUNG *Excursions in Brownian motion*. Ark. Math. **14**, 155-177, 1976.

- [2] J.W. COHEN AND HOOGHIEMSTRA, G. *Brownian excursion, the M/M/1 queue and their occupation times*. Math. Oper. Res. **6**, 608-629, 1981.
- [3] R.K. GETTOOR AND SHARPE, M.J. *Excursions of Brownian motion and Bessel processes*. Z. Wahrscheinlichkeitstheorie un Verw. Gebiete **47**, 83-106, 1979.
- [4] G. HOOGHIEMSTRA *Brownian Excursion and Limit Theorems for the M/G/1 queue*. Ph.D. thesis University Utrecht, 1979.
- [5] K.M. JANSON *The distribution of time spent by a standard excursion above a given level, with applications to ring polymers near a discontinuity in potential*. Elect. Comm. in Probab. **2**, 53-58, 1997.