



Vol. 10 (2005), Paper no. 29, pages 962-987.

Journal URL

<http://www.math.washington.edu/~ejpecp/>

EQUILIBRIUM FLUCTUATIONS FOR A ONE-DIMENSIONAL INTERFACE
IN THE SOLID ON SOLID APPROXIMATION

GUSTAVO POSTA*

Abstract. An unbounded one-dimensional solid-on-solid model with integer heights is studied. Unbounded here means that there is no *a priori* restrictions on the discrete gradient of the interface. The interaction Hamiltonian of the interface is given by a finite range part, proportional to the sum of height differences, plus a part of exponentially decaying long range potentials. The evolution of the interface is a reversible Markov process. We prove that if this system is started in the center of a box of size L after a time of order L^3 it reaches, with a very large probability, the top or the bottom of the box.

Keywords: Solid on solid, SOS, interface dynamics, spectral gap.

Mathematics subject classification: 60K35, 82C22.

Submitted to EJP on January 19, 2004. Final version accepted on May 24, 2005.

* Dipartimento di Matematica F. Brioschi, Politecnico di Milano, Piazza Leonardo da Vinci 32, I-20133 Milano, Italy. e-mail: gustavo.posta@polimi.it

1. Introduction

The rigorous analysis of Glauber dynamics for classical spin systems when the inverse temperature β is such that the static system does not undergo phase transition in the thermodynamic limit has been in the last years the argument of many important works. In particular we refer to [12], [8], [9] and references in these papers.

A natural question is what happens when the thermodynamic parameters are such that there is a phase transition. To be concrete consider the stochastic ferromagnetic Ising model in $\Lambda_L \equiv [1, L] \times [1, L] \cap \mathbb{Z}^2$ in absence of external field with free boundary conditions and let us suppose that the inverse temperature β is much larger than the critical one. Then, as well known (see [4]), any associated infinite volume (*i.e.* $L = +\infty$) Glauber dynamics is not ergodic. However, for $L < +\infty$, every associated Glauber dynamics is ergodic, because it is an irreducible finite-state Markov chain. The problem of how this system behaves at the equilibrium is discussed in [6]. In that paper is proved that the system fluctuates between the “+” phase and the “−” on two distinct time scales. In a first time interval the system creates a layer of the opposite phase, separated from the initial one by a one-dimensional interface. In a second time interval the interface moves until the new phase invades the system. The time the system spends for this transition is exponential in L . In this paper we will study the motion of the interface in the solid-on-solid (SOS) approximation. In particular we will show that after the formation of the interface, a time of order L^3 is sufficient to reach the opposite phase.

The SOS model studied here is a one-dimensional random interface (or surface) with integer heights. There is no restriction on the discrete gradient of the surface (unbounded SOS), thus the configuration space is \mathbb{Z}^L . The interaction Hamiltonian of the interface is given by the usual energy proportional to the sum of height differences plus a part of exponentially decaying potential which mimics the long-range dependence that the interface feels due to the surrounding bulk phases. This long-range interaction is small in the regime studied, but the potentials involved are of unbounded range, so it is not completely trivial to handle. We restrict the interface to stay in a finite box $[1, L] \times [-M, M]$. This gives (see Section 2 for more details) the Gibbs measure on \mathbb{Z}^L

$$\mu_L^M(\phi) \equiv \frac{\mathbf{1}(\|\phi\|_\infty \leq M) e^{-\beta H_L(\phi) - W_L^M(\beta, \phi)}}{Z_L^M(\beta)}.$$

The evolution of the interface is described by a reversible Markov process with generator

$$(G_L^M f)(\phi) \equiv \sum_{\psi} c_L^M(\phi, \psi) [f(\psi) - f(\phi)],$$

where the jump rates are bounded and such that only transitions of the form $(\phi_1, \dots, \phi_L) \mapsto (\phi_1, \dots, \phi_k \pm 1, \dots, \phi_L)$, for some $k = 1, \dots, L$, are allowed. We study this process for $M = L/2$, *i.e.* in a “box” of size L and we prove, in a sense given precisely by Theorem 2.1, that if the interface

is started in the center of the box then in a time of order L^3 it reaches the bottom or the top of the box.

The technique we use to get this result is a mix of analytical and probabilistic tools. In fact it is standard to obtain estimates on the exit time distribution of a reversible Markov process from a region if one has an estimate from below of the spectral gap of its generator with Dirichlet boundary conditions. We will not give bounds on the spectral gap of $G_L^{L/2}$ but on the generator of an auxiliary simpler process and we conclude with a coupling argument.

This paper completes the study of the asymptotic properties of the solid-on-solid model started in [10] where a similar model, in the presence of boundary conditions, is studied.

Acknowledgments: I would like to thank Fabio Martinelli who posed this problem to me and helped me with many constructive discussions.

2. Notation and Results

Our sample space is $\Omega_L \equiv \mathbb{Z}^L$ for fixed $L \in \mathbb{N}$. Configurations, *i.e.* elements of the sample space will be denoted by Greek letters, *e.g.* $\phi = (\phi_1, \dots, \phi_L) \in \Omega_L$. Given $M \in \mathbb{N} \cup \{+\infty\}$, and $\beta > 0$ one defines the *energy* associated with the configuration $\phi \in \Omega_L$ as:

$$\beta H_L(\phi) + W_L^M(\beta, \phi).$$

Here H_L is a local interaction:

$$H_L(\phi) \equiv \sum_{k=1}^{L-1} |\phi_{k+1} - \phi_k|, \tag{2.1}$$

while $W_L^M(\beta, \phi)$ is a long range interaction that will be defined below.

Consider the lattices \mathbb{Z}^2 and $(\mathbb{Z}^2)^* \equiv (1/2, 1/2) + \mathbb{Z}^2$ as graphs embedded in \mathbb{R}^2 equipped with the usual Euclidean metric denoted with $\text{dist}(\cdot, \cdot)$:

$$\text{dist}((x_1, y_1), (x_2, y_2)) \equiv \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}.$$

For every $\phi \in \Omega_L$ the *contour associated with* ϕ is the subset of \mathbb{R}^2 defined by:

$$\Gamma(\phi) \equiv \left[\bigcup_{i=1}^L \{(x, y) : x \in (i-1, i), y = \phi_i\} \right] \cup \left[\bigcup_{i=1}^{L-1} \{(x, y) : x = i, y \in [\phi_i \wedge \phi_{i+1}, \phi_i \vee \phi_{i+1}]\} \right].$$

For $A \subset \mathbb{R}^2$ and $p \in \mathbb{R}^2$, the distance of x from A is defined as $\text{dist}(p, A) \equiv \inf\{\text{dist}(x, y) : y \in A\}$. If we denote with

$$\Delta(\phi) \equiv \left\{ p^* \in (\mathbb{Z}^2)^* : \text{dist}(p^*, \Gamma(\phi)) = \frac{1}{2}, \text{ or } \text{dist}(p^*, \Gamma(\phi)) = \frac{1}{\sqrt{2}} \right\}$$

the set of sites *attached* to the contour $\Gamma(\phi)$ and define

$$V_L^M \equiv ([-1/2, L + 1/2] \times [-(M + 1)/2, (M + 1)/2]) \cap (\mathbb{Z}^2)^*, \quad (2.2)$$

the long range interaction may be written as:

$$W_L^M(\beta, \phi) \equiv \sum_{\substack{\Lambda \cap \Delta(\phi) \neq \emptyset \\ \Lambda \subset V_L^M}} \Phi(\beta, \Lambda).$$

The sum is over all $\Lambda \subset V_L^M$ connected in the sense of the dual graph $(\mathbb{Z}^2)^*$. The *potential* $\Phi(\cdot, \cdot)$ is a function satisfying (see. [2]):

i) there exists $\bar{\beta} > 0$ such that for every $\beta > \bar{\beta}$ we have:

$$\sum_{\substack{\Lambda \ni p^* \\ \text{diam}(\Lambda) \geq k}} |\Phi(\beta, \Lambda)| \leq e^{-m(\beta)k} \quad (2.3)$$

for any $k > 0$ and $p^* \in (\mathbb{Z}^2)^*$. Here the sum is over all $\Lambda \subset (\mathbb{Z}^2)^*$ connected and such that $\Lambda \ni p^*$, $m(\beta)$ is a positive function such that $m(\beta) \rightarrow +\infty$ for $\beta \rightarrow +\infty$ and $\text{diam}(\Lambda)$ is the Euclidean diameter of the set Λ .

ii) For every $p^* \in (\mathbb{Z}^2)^*$

$$\Phi(\beta, \Lambda + p^*) = \Phi(\beta, \Lambda). \quad (2.4)$$

It is easy to check that for every $\beta > 0$ and $M \in (0, +\infty]$ there exists finite the *partition function*:

$$Z_L^M(\beta) \equiv \sum_{\phi \in \Omega_L} \mathbf{1}(\|\phi\|_\infty \leq M) e^{-\beta H_L(\phi) - W_L^M(\beta, \phi)}, \quad (2.5)$$

where $\|\phi\|_\infty \equiv \max_{1 \leq i \leq L} |\phi_i|$. Thus it is possible to define on Ω_L the *Gibbs measure*:

$$\mu_L^M(\phi) \equiv \frac{\mathbf{1}(\|\phi\|_\infty \leq M) e^{-\beta H_L(\phi) - W_L^M(\beta, \phi)}}{Z_L^M(\beta)}. \quad (2.6)$$

This measure represents the equilibrium of the system. The dynamics of the system is a continuous time Markov chain with values in Ω_L and stationary measure μ_L^M . This process will be defined by means of its generator.

For every $\phi \in \Omega_L$, $k = 1, \dots, L$, define $\phi \pm \delta_k = (\phi_1, \dots, \phi_k \pm 1, \dots, \phi_L)$, and the *jump rates*:

$$c_L^M(\phi, \psi) = \begin{cases} \left(\frac{\mu_L^M(\psi)}{\mu_L^M(\phi)} \right)^{\frac{1}{2}} & \text{if } \mu_L^M(\phi) > 0 \text{ and } \psi = \phi \pm \delta_k \text{ for some } k = 1, \dots, L; \\ 0 & \text{otherwise.} \end{cases} \quad (2.7)$$

It is simple to prove that there exists a unique Markov process $\Phi \equiv \{\Phi(t) : t \geq 0\}$ with generator

$$(G_L^M f)(\phi) \equiv \sum_{\psi} c_L^M(\phi, \psi) [f(\psi) - f(\phi)]. \quad (2.8)$$

Moreover G_L^M is self-adjoint in $L^2(\mu_L^M)$, *i.e.* Φ is *reversible* and G_L^M is negative semidefinite. The absolute value of the largest negative eigenvalue of G_L^M is denoted by $\lambda_1(G_L^M)$ and it is called *spectral gap* of G_L^M .

We will give a direct construction of Φ in Section 4. More precisely (see Proposition 3.1) we will define a measurable space (Ω_L, \mathbf{F}_L) and a family of probability measures on it $\{\mathbb{P}_\phi : \phi \in \Omega_L\}$ such that:

- i)* for every $\phi \in \Omega_L$ the process Φ is a Markov process on $(\Omega_L, \mathbf{F}_L, \mathbb{P}_\phi)$ with generator $G_L^{\emptyset, M, W}$;
- ii)* $\mathbb{P}_\phi(\Phi(0) = \phi) = 1$.

For every measurable set $A \subset \Omega_L$ define the *first exit time* from A as:

$$\tau(A^c) \equiv \inf \{t \geq 0 : \Phi(t) \in A^c\}.$$

The main result of this paper is:

Theorem 2.1. *Let Φ be the process associated with the generator $G_L^{L/2}$. Fix $\alpha \in (0, 1/4)$, $\varepsilon \in (0, 1/100)$ and define $A \equiv \{\phi \in \Omega_L : \|\phi\|_\infty \leq (1 - \varepsilon)L/2\}$. Then there exists $\bar{\beta} > 0$ and for every $\beta > \bar{\beta}$ constants $K_1(\beta)$, $K_2(\beta)$, $K_3(\beta)$ and $K_4(\beta) > 0$ such that:*

$$\int_{\Omega_L} d\mu_L^{\emptyset, L/2, W}(\phi | \|\phi\|_\infty \leq \alpha L) \mathbb{P}_\phi(\tau(A^c) > t) \leq \alpha^{-1} K_1 e^{-K_2(\frac{t}{L^3} \wedge L)} + \alpha^{-1} K_3 e^{-K_4 \varepsilon L} \quad (2.9)$$

for any $L > 0$.

This result can be read in the following way: starting the interface in a square box of size L , from an initial condition randomly chosen under $\mu_L^{\emptyset, L/2, W}(\cdot | \|\cdot\|_\infty \leq \alpha L)$ (*i.e.* the interface is forced to stay at least αL far away from the top or the bottom of the box), the probability of reaching within εL of the top or the bottom of the box in time bigger than t is exponentially small in t/L^3 .

Remark 2.2. In what follows we use constants K_1, K_2, \dots in the statement of theorems, propositions and so on, while we use constants C_1, C_2, \dots inside proofs. The reader should be warned that the use of constants is coherent only inside the same structure. This means, *e.g.*, that constants which appears in the proof or in the statement of a proposition may differ from constants, with the same name, which appears in the proof or in the statement of a different proposition.

3. Proof of Main Result

In this section we will prove our main result Theorem 2.1. The technique we will use is the following. We can estimate the first exit time of a reversible Markov process from a region A by bounding from below the spectral gap of the generator of the process. Actually we will not bound the spectral gap of the process Φ . Instead we will estimate the spectral gap of a simpler auxiliary process $\bar{\Phi}$ that in the region A is similar to Φ . Then a *coupling* argument (Proposition 3.1) concludes the proof.

We begin this section introducing the auxiliary above mentioned process. Fix $L \in \mathbb{N}$, β and $M > 0$ and define on Ω_L the probability measure

$$\bar{\mu}_L^M(\phi) \equiv \mathbf{1}(|\phi_1| \leq M) \frac{e^{-\beta H_L(\phi) - W_L^\infty(\beta, \phi)}}{\bar{Z}_L^M(\beta)}, \quad (3.1)$$

where

$$\bar{Z}_L^M(\beta) \equiv \sum_{\phi} \mathbf{1}(|\phi_1| \leq M) e^{-\beta H_L(\phi) - W_L^\infty(\beta, \phi)}$$

and (see Section 2)

$$W_L^\infty(\beta, \phi) \equiv \sum_{\substack{\Lambda \cap \Delta(\phi) \neq \emptyset \\ \Lambda \subset V_L^\infty}} \Phi(\beta, \Lambda).$$

The process $\bar{\Phi}$ is defined by means of its generator on $L^2(\Omega_L, \bar{\mu}_L^M)$

$$(\bar{G}_L^M f)(\phi) \equiv \sum_{\psi} \bar{c}_L^M(\phi, \psi) [f(\psi) - f(\phi)], \quad (3.2)$$

where

$$\bar{c}_L^M(\phi, \psi) \equiv \begin{cases} \left(\frac{\bar{\mu}_L^M(\psi)}{\bar{\mu}_L^M(\phi)} \right)^{\frac{1}{2}} & \text{if } \bar{\mu}_L^M(\phi) > 0 \text{ and } \psi = \phi \pm \delta_k \text{ for some } k = 1, \dots, L; \\ 0 & \text{otherwise.} \end{cases} \quad (3.3)$$

It simple to check that \bar{G}_L^M is a self-adjoint Markov generator which defines a unique Markov process. Because for $\phi \in A$ (A is defined in Theorem 2.1) the jump rates of \bar{G}_L^M are very close to the jump rates of $G_L^{L/2}$ (see (2.7)), the processes $\bar{\Phi}$ and Φ evolve in a similar way as long as they remains within A . This fact is formally proved in the following result which gives also a direct construction of the processes.

Proposition 3.1. *It is possible to construct a family of probability spaces $(\Omega_L, \mathbf{F}_L, \mathbb{P}_{\phi, \bar{\phi}})$ and a process $\{(\Phi(t), \bar{\Phi}(t)) : t \geq 0\}$ taking values in $\Omega_L \times \Omega_L$, with $\mathbb{P}_{\phi, \bar{\phi}}(\Phi(0) = \phi, \bar{\Phi}(0) = \bar{\phi}) = 1$ for any $(\phi, \bar{\phi}) \in \Omega_L \times \Omega_L$, and such that:*

- i) Φ and $\bar{\Phi}$ are Markov processes with generators $G_L^{L/2}$ and $\bar{G}_L^{L/2}$ respectively;
ii) if we define $\sigma \equiv \inf \{t \geq 0 : \Phi(t) \neq \bar{\Phi}(t)\}$ and $\bar{\tau}(A^c) \equiv \inf \{t \geq 0 : \bar{\Phi}(t) \in A^c\}$, where A is defined in Theorem 2.1, then there exist $K_1(\beta)$ and $K_2(\beta) > 0$, with $K_2(\beta) \rightarrow +\infty$ for $\beta \rightarrow +\infty$, such that for every $\phi \in A$:

$$\mathbb{P}_{\phi,\phi}(\sigma \leq t, \sigma \leq \bar{\tau}) \leq K_1 L t e^{-K_2 \varepsilon L}. \quad (3.4)$$

This proposition will be proved in Section 4.

The advantage in considering the process $\bar{\Phi}$ instead of Φ is that the first one is simpler to study because it has no interaction with the top and the bottom of the box V_L^M . In particular in Section 5 we will prove the following result on the spectral gap $\lambda_1(\bar{G}_L^M)$:

Proposition 3.2. *There exists $\bar{\beta} > 0$ such that for every $\beta \geq \bar{\beta}$ there exists $K_1(\beta) > 0$, so that:*

$$\lambda_1(\bar{G}_L^M) \geq K_1 L^{-1} (L^{-1} \wedge M^{-2}) \quad (3.5)$$

for every L and $M > 0$.

We are now in a position to prove Theorem 2.1.

Proof of Theorem 2.1.

In this proof we simplify notation writing $G \equiv G_L^{L/2}$, $\bar{G} \equiv \bar{G}_L^{L/2}$, $\mu \equiv \mu_L^{L/2}$, $\bar{\mu} \equiv \bar{\mu}_L^{L/2}$ and

$$\begin{aligned} \tau &\equiv \tau(A^c) = \inf \{t \geq 0 : \|\Phi(t)\|_\infty > (1 - \varepsilon)L/2\}, \\ \bar{\tau} &\equiv \bar{\tau}(A^c) = \inf \{t \geq 0 : \|\bar{\Phi}(t)\|_\infty > (1 - \varepsilon)L/2\}. \end{aligned}$$

Recall that σ was defined in Proposition 3.1 as the first time such that $\bar{\Phi}(\sigma) \neq \Phi(\sigma)$ and suppose that $\phi \in A$. Then for any $t > 0$

$$\mathbb{P}_{\phi,\phi}(\tau > t) = \mathbb{P}_{\phi,\phi}(\tau > t, \tau = \bar{\tau}) + \mathbb{P}_{\phi,\phi}(\tau > t, \tau \neq \bar{\tau}) \leq \mathbb{P}_{\phi,\phi}(\bar{\tau} > t) + \mathbb{P}_{\phi,\phi}(\sigma \leq \bar{\tau}),$$

while for any $s > 0$

$$\begin{aligned} \mathbb{P}_{\phi,\phi}(\sigma \leq \bar{\tau}) &= \mathbb{P}_{\phi,\phi}(\sigma \leq \bar{\tau}, \sigma \leq s) + \mathbb{P}_{\phi,\phi}(\sigma \leq \bar{\tau}, \sigma > s) \leq \\ &\leq \mathbb{P}_{\phi,\phi}(\sigma \leq \bar{\tau}, \sigma \leq s) + \mathbb{P}_{\phi,\phi}(\bar{\tau} > s). \end{aligned}$$

In conclusion for any $t, s > 0$

$$\mathbb{P}_{\phi,\phi}(\tau > t) \leq 2\mathbb{P}_{\phi,\phi}(\bar{\tau} > t \wedge s) + \mathbb{P}_{\phi,\phi}(\sigma \leq \bar{\tau}, \sigma \leq s). \quad (3.6)$$

Define $B \equiv \{\phi \in \Omega_L : \|\phi\|_\infty \leq \alpha L\}$, where $\alpha \in (0, 1/4)$. By (3.6) we obtain:

$$\int_{\Omega_L} d\mu(\phi|B) \mathbb{P}_{\phi,\phi}(\tau > t) \leq 2 \int_{\Omega_L} d\mu(\phi|B) \mathbb{P}_{\phi,\phi}(\bar{\tau} > t \wedge s) + \sup_{\phi \in B} \mathbb{P}_{\phi,\phi}(\sigma \leq s). \quad (3.7)$$

We are going to bound from above the first term on the right hand side of (3.7). We will use a Markov process with killing (see [11]). The *Dirichlet form* associated with the generator \bar{G} is the positive-semidefinite bilinear form

$$\bar{\mathcal{G}}(f, g) \equiv - \langle \bar{G}f, g \rangle_{L^2(\Omega_L, \bar{\mu})},$$

where $\langle \cdot, \cdot \rangle_{L^2(\Omega_L, \bar{\mu})}$ is the scalar product in $L^2(\Omega_L, \bar{\mu})$. Because $\bar{\mu}(A) > 0$ it is possible to define the positive semidefinite bilinear form

$$\bar{\mathcal{G}}_A(f, g) \equiv \bar{\mathcal{G}}(\hat{f}, \hat{g}), \tag{3.8}$$

with form domain

$$\text{Dom}(\bar{\mathcal{G}}_A) \equiv \left\{ f \in L^2(A, \bar{\mu}) : \hat{f} \in \text{Dom}(\bar{\mathcal{G}}) \right\}.$$

Here

$$\hat{f}(\phi) \equiv \begin{cases} f(\phi) & \text{if } \phi \in A; \\ 0 & \text{otherwise.} \end{cases}$$

Standard functional analysis methods shows that there exists a unique positive-semidefinite self-adjoint (in $L^2(A, \bar{\mu})$) operator \bar{G}_A such that:

$$\bar{\mathcal{G}}_A(f, g) \equiv - \langle \bar{G}_A f, g \rangle_{L^2(A, \bar{\mu})}.$$

This operator has a probabilistic interpretation, it is the generator of a process which evolves according to \bar{G} as long as it stays within A , but is killed when it tries to jump outside A (see [11]). The semigroup $e^{t\bar{G}_A}$ generated by \bar{G}_A is *sub-stochastic* and we have:

$$(e^{t\bar{G}_A} f)(\phi) = \mathbb{E}_{\phi, \phi} [\mathbf{1}(\bar{\tau} > t) f(\bar{\Phi}(t))]$$

for every $f \in L^2(A, \bar{\mu})$. In particular taking $f \equiv \mathbf{1}$ we obtain:

$$\mathbb{P}_{\phi, \phi}(\bar{\tau} > t) = (e^{t\bar{G}_A} \mathbf{1})(\phi).$$

Thus:

$$\begin{aligned} \int_{\Omega_L} d\mu(\phi|B) \mathbb{P}_{\phi, \phi}(\bar{\tau} > t) &= \left\langle (e^{t\bar{G}_A} \mathbf{1}), \frac{d\mu(\cdot|B)}{d\bar{\mu}} \right\rangle_{L^2(A, \bar{\mu})} \leq \\ &\leq \left\| (e^{t\bar{G}_A} \mathbf{1}) \right\|_{L^2(A, \bar{\mu})} \left\| \frac{d\mu(\cdot|B)}{d\bar{\mu}} \right\|_{L^2(A, \bar{\mu})} \leq \left\| \frac{d\mu(\cdot|B)}{d\bar{\mu}} \right\|_{L^2(A, \bar{\mu})} e^{-\lambda_1(\bar{G}_A)t}. \end{aligned} \tag{3.9}$$

Spectral theorem has been used in the last line. The spectral gap $\lambda_1(\bar{G}_A)$ is characterized by the following variational property:

$$\lambda_1(\bar{G}_A) = \inf_{\substack{f \in L^2(A, \bar{\mu}) \\ f \perp \mathbf{1}}} \frac{\bar{\mathcal{G}}_A(f, f)}{\|f\|_{L^2(A, \bar{\mu})}^2}.$$

This relation and (3.8) imply:

$$\lambda_1(\bar{G}_A) = \inf_{\substack{f \in L^2(A, \bar{\mu}) \\ f \perp \mathbf{1}}} \frac{\bar{G}(\hat{f}, \hat{f})}{\|\hat{f}\|_{L^2(\Omega_L, \bar{\mu})}^2} \geq \inf_{\substack{f \in L^2(\Omega_L, \bar{\mu}) \\ f \perp \mathbf{1}}} \frac{\bar{G}(f, f)}{\|f\|_{L^2(\Omega_L, \bar{\mu})}^2} = \lambda_1(\bar{G}).$$

By Proposition 3.2 we know that $\lambda_1(\bar{G}) \geq C_1(\beta)L^{-3}$, so (3.9) yields:

$$\int_{\Omega_L} d\mu(\phi|B)\mathbb{P}_{\phi, \phi}(\bar{\tau} > t) \leq e^{-\frac{tC_1}{L^3}} \left\| \frac{d\mu(\cdot|B)}{d\bar{\mu}} \right\|_{L^2(A, \mu)}.$$

We claim that there exists $C_2(\beta) > 0$ such that:

$$\left\| \frac{d\mu(\cdot|B)}{d\bar{\mu}} \right\|_{L^2(A, \mu)} \leq \alpha^{-1}C_2.$$

This simple technical bound is proved in the appendix (see Lemma A1.1). In conclusion:

$$\int_{\Omega_L} d\mu(\phi|B)\mathbb{P}_{\phi, \phi}(\bar{\tau} > t) \leq \alpha^{-1}C_2e^{-\frac{C_1t}{L^3}}.$$

By (3.7)

$$\int_{\Omega_L} d\mu(\phi|B)\mathbb{P}_{\phi, \phi}(\tau > t) \leq \alpha^{-1}C_2e^{-\frac{C_1(t \wedge s)}{L^3}} + \sup_{\phi \in B} \mathbb{P}_{\phi, \phi}(\sigma \leq s).$$

Taking $s = L^4$ by (3.4) we obtain:

$$\int_{\Omega_L} d\mu(\phi|B)\mathbb{P}_{\phi, \phi}(\tau > t) \leq C_2\alpha^{-1}e^{-C_1(\frac{t}{L^3} \wedge L)} + C_3\alpha^{-1}e^{-C_4\epsilon L},$$

i.e. (2.9). \square

4. The Coupling

In this section we will construct explicitly a stochastic coupling between Φ and $\bar{\Phi}$, in particular we will prove Proposition 3.1. The technique we use is an application of the so called *basic coupling*. This is a coupling between jump processes such that the processes jump together as long as possible, considering the constraint they have to jump with their own jump rates. Because the jump rates of Φ and $\bar{\Phi}$ are very close, when they are in $A = \{\phi \in \Omega_L : \|\phi\|_\infty \leq (1 - \varepsilon)L/2\}$, the two processes will evolve identically for a long time.

The first step in the construction of the coupling is to show that the jump rates of Φ and $\bar{\Phi}$ are close in A .

Lemma 4.1. *For any $\beta > \bar{\beta}$ there exists $K_1(\beta)$ and $K_2(\beta) > 0$ such that:*

$$\sup_{\substack{k=1,\dots,L \\ \phi \in A}} \left| \frac{\bar{c}_L^{L/2}(\phi, \phi \pm \delta_k)}{c_L^{L/2}(\phi, \phi \pm \delta_k)} - 1 \right| \leq K_1 e^{-m(\beta)\varepsilon K_2 L} \quad (4.1)$$

for every $L > 0$.

Proof.

To simplify we adopt the notation of the last section by writing $\bar{c} \equiv \bar{c}_L^{L/2}$, $c \equiv c_L^{L/2}$ and $\bar{\mu} \equiv \bar{\mu}_L^{L/2}$. Notice that

$$\left[\frac{\bar{c}(\phi, \phi + \delta_k)}{c(\phi, \phi + \delta_k)} \right]^2 = \frac{\bar{\mu}(\phi + \delta_k)\mu(\phi)}{\bar{\mu}(\phi)\mu(\phi + \delta_k)} = e^{[(\partial_k^+ W_L^{L/2})(\beta, \phi) - (\partial_k^+ W_L^\infty)(\beta, \phi)]} \quad (4.2)$$

for every $\phi \in A$. Here and later $(\partial_k^\pm f)(\phi) \equiv f(\phi \pm \delta_k) - f(\phi)$. If we define $p_k(\phi) \equiv (k, \phi_k)$ and $I_k \equiv \{p \in (\mathbb{Z}^2)^* : \text{dist}(p, p_k) \leq 4\}$ it simple to prove that:

$$|(\partial_k^+ W_L^{L/2})(\beta, \phi) - (\partial_k^+ W_L^\infty)(\beta, \phi)| \leq \sum_{\substack{\Lambda \cap I_k \neq \emptyset \\ \Lambda \cap (V_L^{L/2})^c \neq \emptyset}} |\Phi(\beta, \Lambda)|. \quad (4.3)$$

But if $\phi \in A$, $\Lambda \cap I_k \neq \emptyset$ and $\Lambda \cap (V_L^{L/2})^c \neq \emptyset$ then Λ should be at least of size L , *i.e.* there exists a constant $C_1 > 0$ such that $\text{diam}(\Lambda) \geq \varepsilon C_1 L$. So we can use condition (2.3) to bound the sum on the right hand side of (4.3):

$$|(\partial_k^+ W_L^{L/2})(\phi) - (\partial_k^+ W_L^\infty)(\phi)| \leq 16e^{-m(\beta)\varepsilon C_1 L}.$$

From this relation and (4.2) we have (4.1). \square

We can now prove the main result of this section.

Proof of Proposition 3.1.

We use the basic coupling. To any site $k = 1, \dots, L$ we associate two independent Poisson processes, each one with rate $c_{\max} \equiv \sup_{\phi, \psi} \{c(\phi, \psi), \bar{c}(\phi, \psi)\}$. We will denote these processes $\{N_{k,t}^+ : t \geq 0\}$ and $\{N_{k,t}^- : t \geq 0\}$ while the arrival times of each process are denoted by $\{\tau_{k,n}^+ : n \in \mathbb{N}\}$ and $\{\tau_{k,n}^- : n \in \mathbb{N}\}$ respectively. Assume that the Poisson processes associated to different sites are also mutually independent. We say that at each point in the space-time of the form $(k, \tau_{k,n}^+)$ there is a “+” mark and that at each point of the form $(k, \tau_{k,n}^-)$ there is a “-” mark.

Next we associate to each arrival time $\tau_{k,n}^\pm$ a random variable $U_{k,n}^\pm$ with uniform distribution on $[0, 1]$. All these random variables are assumed to be independent among themselves and independent from the previously introduced Poisson processes. Obviously there exists a probability space such that all these objects are defined. We have to say now how the various processes are constructed on this space. The process Φ (resp. $\bar{\Phi}$) is defined in the following manner. We know that almost surely the arrival times $\tau_{k,n}^*$, $k = 1, \dots, L$, $n \in \mathbb{N}$, $* = \pm$ are all distinct. We update the state of the process each time there is a mark at some $k = 1, \dots, L$ according to the following rule.

- If the mark that we are considering is at the point $(k, \tau_{k,n}^*)$, with $* = \pm$, and the configuration of Φ (resp. of $\bar{\Phi}$) immediately before time $\tau_{k,n}^*$ was ϕ , (resp. $\bar{\phi}$) then the configuration immediately after $\tau_{k,n}^*$ of Φ (resp. of $\bar{\Phi}$) will be $\phi \pm \delta_k$ (resp. $\bar{\phi} \pm \delta_k$) if and only if

$$c(\phi, \phi \pm \delta_k) > U_{k,n}^\pm c_{\max} \quad (\text{resp. } \bar{c}(\phi, \phi \pm \delta_k) > U_{k,n}^\pm c_{\max}).$$

Else the configuration remains the same.

It is easy to check that this construction satisfies condition *i*) of the proposition. It remains to prove (3.4).

Define $N_t = \sum_{k=1}^L (N_{k,t}^+ + N_{k,t}^-)$. This process counts the number of possible updating of the processes Φ and $\bar{\Phi}$ in the interval $[0, t]$. It is clear that $\{N_t : t \geq 0\}$ is a Poisson process with rate $\lambda \equiv 2Lc_{\max}$. For $\phi \in A$ we have:

$$\mathbb{P}_{\phi, \phi}(\sigma \leq t, \sigma \leq \bar{\tau}) = e^{-\lambda t} \sum_{n=1}^{+\infty} \frac{(\lambda t)^n}{n!} \mathbb{P}_{\phi, \phi}(\sigma \leq t, \sigma \leq \bar{\tau} | N_t = n). \quad (4.4)$$

To bound from above $\mathbb{P}_{\phi, \phi}(\sigma \leq t, \sigma \leq \bar{\tau} | N_t = n)$ observe that if Φ and $\bar{\Phi}$ are initially in the same state $\phi \in A$ and if $N_t = n$, *i.e.* there were n possible updating in $[0, t]$, then it possible that $\sigma \in [0, t]$ if and only if for some $i = 1, \dots, n$, $\psi \in \Omega_L$ and $k = 1, \dots, L$ happens that:

$$\bar{c}(\psi, \psi \pm \delta_k) > U_{k,i}^\pm c_{\max} \quad \text{and} \quad c(\psi, \psi \pm \delta_k) \leq U_{k,i}^\pm c_{\max}$$

or

$$\bar{c}(\psi, \psi \pm \delta_k) \leq U_{k,i}^\pm c_{\max} \quad \text{and} \quad c(\psi, \psi \pm \delta_k) > U_{k,i}^\pm c_{\max}.$$

The probability of this event, for fixed $i = 1, \dots, n$, $\psi \in \Omega_L$ and $k = 1, \dots, L$, is:

$$\left| \frac{\bar{c}(\psi, \psi \pm \delta_k)}{c_{\max}} - \frac{c(\psi, \psi \pm \delta_k)}{c_{\max}} \right|.$$

Moreover because $\sigma \leq \bar{\tau}$ we have:

$$\mathbb{P}_{\phi, \phi}(\sigma \leq t, \sigma \leq \bar{\tau} | N_t = n) \leq 1 - \left(1 - \sup_{\substack{k=1, \dots, L \\ \psi \in A}} \left| \frac{\bar{c}(\psi, \psi \pm \delta_k)}{c_{\max}} - \frac{c(\psi, \psi \pm \delta_k)}{c_{\max}} \right| \right)^n.$$

By Lemma 4.1 we have

$$\sup_{\substack{k=1, \dots, L \\ \psi \in A}} \left| \frac{\bar{c}(\psi, \psi \pm \delta_k)}{c_{\max}} - \frac{c(\psi, \psi \pm \delta_k)}{c_{\max}} \right| \leq C_1 e^{-m(\beta)C_2 \varepsilon L},$$

which gives

$$\mathbb{P}_{\phi, \phi}(\sigma \leq t, \sigma \leq \bar{\tau} | N_t = n) \leq 1 - \left[1 - C_1 e^{-m(\beta)C_2 \varepsilon L} \right]^n.$$

This estimate together with (4.4) gives (3.4). \square

5. Spectral Gap for \bar{G}_L^M

In this section we will prove Proposition 3.2. The strategy of the proof is the following. By a simple change of variables the *Glauber* dynamics associated with \bar{G}_L^M becomes a *Kawasaki* type dynamics, while the measure $\bar{\mu}_L^M$ becomes, in the new variables, a product measure perturbed by an infinite range interaction. This interaction term is small if β is large. Without this perturbation term the result is very simple to prove. The presence of this extra term requires a little extra work.

Consider the random variables defined by

$$\begin{aligned} \eta_1(\phi) &\equiv \phi_1 \\ \eta_2(\phi) &\equiv \phi_2 - \phi_1 \\ &\dots \\ \eta_L(\phi) &\equiv \phi_L - \phi_{L-1}. \end{aligned} \tag{5.1}$$

Obviously the map $\Omega_L \ni \phi \mapsto \Omega_L \in \eta$ is bijective with inverse map

$$T : \Omega_L \ni (\eta_1, \dots, \eta_L) \mapsto (\eta_1, \eta_1 + \eta_2, \dots, \eta_1 + \dots + \eta_L) \in \Omega_L.$$

The distribution of η is easily calculated as:

$$\bar{\mu}_L^M(\phi \in \Omega_L : \eta_1(\phi) = \eta_1, \dots, \eta_L(\phi) = \eta_L) = \mathbf{1}(|\eta_1| \leq M) \frac{e^{-\beta \sum_{i=2}^L |\eta_i| - W_L^\infty(\beta, T(\eta))}}{\bar{Z}_L^M(\beta)}.$$

Observe that $\eta(\phi)$ defined in (5.1) is the vector of the discrete derivatives of the configuration $\phi \in \Omega_L$. The study of the SOS interface can be carried out using the variables ϕ or η indifferently. We will use the last one in the sequel. Let $\bar{\nu}_L^M$ be the distribution of η , *i.e.*

$$\bar{\nu}_L^M(\eta) \equiv \mathbf{1}(|\eta_1| \leq M) \frac{e^{-\beta \sum_{i=2}^L |\eta_i| - W_L^\infty(\beta, T(\eta))}}{\bar{Z}_L^M(\beta)}. \quad (5.2)$$

The expected value operator with respect to $\bar{\nu}_L^M$ will be denoted with $\bar{\mathbf{E}}_L^M(\cdot)$, while the *covariance* form will be denoted by $\bar{\mathbf{E}}_L^M(\cdot, \cdot)$ *i.e.*

$$\bar{\mathbf{E}}_L^M(f, g) \equiv \bar{\mathbf{E}}_L^M \left[\left(f - \bar{\mathbf{E}}_L^M(f) \right) \left(g - \bar{\mathbf{E}}_L^M(g) \right) \right],$$

where $f, g \in L^2(\bar{\nu}_L^M)$. On the same Hilbert space is defined the quadratic form:

$$\bar{\mathcal{E}}_L^M(f, f) \equiv \bar{\mathbf{E}}_L^M \left[\mathbf{1}(\eta_1 < M) (\partial_{2,1} f)^2 \right] + \sum_{k=2}^{L-1} \bar{\mathbf{E}}_L^M \left[(\partial_{k+1,k} f)^2 \right] + \bar{\mathbf{E}}_L^M \left[(\partial_L^+ f)^2 \right],$$

where for any $h, k = 1, \dots, L$ $(\partial_h^+ f)(\eta) = f(\eta + \delta_h) - f(\eta)$ and $(\partial_{h,k} f)(\eta) = f(\eta + \delta_h - \delta_k) - f(\eta)$. We will use this form to estimate the spectral gap of \bar{G}_L^M :

Lemma 5.1. *There exists two constants $K_1(\beta)$ and $K_2(\beta)$ such that*

$$K_1 \inf_{f \in L^2(\bar{\nu}_L^M)} \frac{\bar{\mathcal{E}}_L^M(f, f)}{\bar{\mathbf{E}}_L^M(f, f)} \leq \lambda_1(G_L^M) \leq K_2 \inf_{f \in L^2(\bar{\nu}_L^M)} \frac{\bar{\mathcal{E}}_L^M(f, f)}{\bar{\mathbf{E}}_L^M(f, f)}. \quad (5.3)$$

Proof.

Using the fact that \bar{G}_L^M is self adjoint in $L^2(\bar{\mu}_L^M)$ it simple to check that

$$\bar{\mathcal{G}}_L^M(f, f) \equiv - \langle \bar{G}_L^M f, f \rangle_{L^2(\bar{\mu}_L^M)} = \frac{1}{2} \sum_{k=1}^L \sum_{\phi \in \Omega_L} \bar{\mu}_L^M(\phi) \bar{c}_L^M(\phi, \phi + \delta_k) \left[(\partial_k^+ f)(\phi) \right]^2.$$

If we recall the definition of the jump rates (3.3) a simple calculation shows that there exists $C_1(\beta)$ and $C_2(\beta)$ such that

$$C_1 \bar{\mu}_L^M(\phi) \mathbf{1}(\phi_1 < M) \leq \bar{\mu}_L^M(\phi) \bar{c}_L^M(\phi, \phi + \delta_1) \leq C_2 \bar{\mu}_L^M(\phi) \mathbf{1}(\phi_1 < M)$$

and

$$C_1 \bar{\mu}_L^M(\phi) \leq \bar{\mu}_L^M(\phi) \bar{c}_L^M(\phi, \phi + \delta_k) \leq C_2 \bar{\mu}_L^M(\phi) \quad k = 2, \dots, L,$$

for every $\phi \in \Omega_L$ and $L, M > 0$. This means that $\bar{\mathcal{G}}_L^M(f, f)$ can be bounded from below (from above) by $C_1/2$ (by $C_2/2$) times

$$\sum_{\phi \in \Omega_L} \bar{\mu}_L^M(\phi) \left\{ [\mathbf{1}(\phi_1 < M)(\partial_1^+ f)(\phi)]^2 + \sum_{k=1}^L [(\partial_k^+ f)(\phi)]^2 \right\}.$$

Now we use the change of variable $\phi = T\eta$ to obtain:

$$\sum_{\phi \in \Omega_L} \bar{\mu}_L^M(\phi) \left\{ [\mathbf{1}(\phi_1 < M)(\partial_1^+ f)(\phi)]^2 + \sum_{k=1}^L [(\partial_k^+ f)(\phi)]^2 \right\} = \bar{\mathcal{E}}_L^M(f^*, f^*)$$

where $f^*(\eta) \equiv f(T\eta)$. This implies

$$\frac{C_1}{2} \inf_{f^* \in L^2(\bar{\nu}_L^M)} \frac{\bar{\mathcal{E}}_L^M(f^*, f^*)}{\bar{\mathbf{E}}_L^M(f^*, f^*)} \leq \inf_{f \in L^2(\bar{\mu}_L^M)} \frac{\bar{\mathcal{G}}_L^M(f, f)}{\bar{\mu}_L^M(f, f)} \leq \frac{C_2}{2} \inf_{f^* \in L^2(\bar{\nu}_L^M)} \frac{\bar{\mathcal{E}}_L^M(f^*, f^*)}{\bar{\mathbf{E}}_L^M(f^*, f^*)}, \quad (5.4)$$

where $\bar{\mu}_L^M(f, f) \equiv \int d\bar{\mu}_L^M(\phi) f^2(\phi) - [\int d\bar{\mu}_L^M(\phi) f(\phi)]^2 = \bar{\mathbf{E}}_L^M(f^*, f^*)$. Observing that

$$\inf_{g \in L^2(\bar{\nu}_L^M)} \frac{\bar{\mathcal{E}}_L^M(g, g)}{\bar{\mathbf{E}}_L^M(g, g)} = \inf_{\substack{g \in L^2(\bar{\nu}_L^M) \\ g \geq 1}} \frac{\bar{\mathcal{E}}_L^M(g, g)}{\bar{\mathbf{E}}_L^M(g^2)}, \quad \inf_{g \in L^2(\bar{\mu}_L^M)} \frac{\bar{\mathcal{G}}_L^M(g, g)}{\bar{\mu}_L^M(g, g)} = \inf_{\substack{g \in L^2(\bar{\mu}_L^M) \\ g \geq 1}} \frac{\bar{\mathcal{G}}_L^M(g, g)}{\int d\bar{\mu}_L^M(\phi) g^2(\phi)},$$

by the variational characterization of the spectral gap and (5.4) we have (5.3). \square

We can use this lemma to prove Proposition 3.2. In fact by (5.3) it is easy to show that (3.5) is equivalent to the existence of $C_1(\beta) > 0$ such that the Poincaré inequality

$$\bar{\mathbf{E}}_L^M(f, f) \leq C_3(\beta)(L \vee M^2)L \bar{\mathcal{E}}_L^M(f, f), \quad (5.5)$$

holds for every $f \in L^2(\bar{\nu}_L^M)$. The key step of the proof of this inequality is contained in the following result.

Proposition 5.2. *There exists $\bar{\beta} > 0$ such that for every $\beta \geq \bar{\beta}$ it is possible to find $K_1(\beta) > 0$ such that*

$$\bar{\mathbf{E}}_L^M(f, f|\eta_1) \leq K_1 \sum_{k=2}^L \bar{\mathbf{E}}_L^M [(\partial_k^+ f)^2|\eta_1], \quad (5.6)$$

for any $f \in L^2(\bar{\nu}_L^M)$, $L > 0$ and $\eta_1 \in [-M, M] \cap \mathbb{Z}$.

This proposition shows the perturbative approach of the proof. In fact if $W_L^\infty = 0$ the measure $\bar{\nu}_L^M(\cdot|\eta_1)$ is a product measure. In this case, Proposition 5.2 says that there exists a uniformly

positive spectral gap for a random walk in \mathbb{Z}^{L-1} in which each component of the walk is independent from the others. It is well known that this gap exists if each component exhibits by itself a uniformly positive spectral gap, and the existence of this one site spectral gap is easily proved. Because for $\beta \rightarrow +\infty$ the perturbation W_L^∞ goes to 0, the result should be true also for large values of β .

Before proving the key result Proposition 5.2 we want to show how, from this result (5.5) follows.
Proof of Proposition 3.2.

Fix $f \in L^2(\bar{\nu}_L^M)$, a simple calculation yields:

$$\bar{\mathbf{E}}_L^M(f, f) = \bar{\mathbf{E}}_L^M \left[\bar{\mathbf{E}}_L^M(f, f|\eta_1) \right] + \bar{\mathbf{E}}_L^M \left[\bar{\mathbf{E}}_L^M(f|\eta_1), \bar{\mathbf{E}}_L^M(f|\eta_1) \right]. \quad (5.7)$$

The first term on the right hand side of this equation can be bounded using Proposition 5.2. For the second term notice that η_1 is uniformly distributed in $[-M, M] \cap \mathbb{Z}$. It is well known (see [1] for example) that this implies that there exists $C_1(\beta) > 0$ such that the one-site Poincaré inequality

$$\bar{\mathbf{E}}_L^M(g, g) \leq C_1 M^2 \bar{\mathbf{E}}_L^M \left[\mathbf{1}(\eta_1 < M) (\partial_1^+ g)^2 \right]$$

holds for every $g = g(\eta_1) \in L^2(\bar{\nu}_L^M)$ and $M > 0$. In particular this formula is true for $g(\eta_1) = \bar{\mathbf{E}}_L^M(f|\eta_1)$. Because the measure $\bar{\nu}_L^M(\cdot|\eta_1)$ does not depend on η_1 if $-M \leq \eta_1 < M$, is simple to check that

$$\partial_1^+ \bar{\mathbf{E}}_L^M(f|\eta_1) = \bar{\mathbf{E}}_L^M(\partial_1^+ f|\eta_1).$$

In conclusion we obtain

$$\bar{\mathbf{E}}_L^M \left[\bar{\mathbf{E}}_L^M(f|\eta_1), \bar{\mathbf{E}}_L^M(f|\eta_1) \right] \leq C_1 M^2 \bar{\mathbf{E}}_L^M \left[\mathbf{1}(\eta_1 < M) (\partial_1^+ f)^2 \right].$$

We can use this and (5.6) in the left hand side of (5.7) to get:

$$\bar{\mathbf{E}}_L^M(f, f) \leq C_2(\beta) \left\{ \sum_{k=2}^L \bar{\mathbf{E}}_L^M \left[(\partial_k^+ f)^2 \right] + M^2 \bar{\mathbf{E}}_L^M \left[\mathbf{1}(\eta_1 < M) (\partial_1^+ f)^2 \right] \right\}. \quad (5.8)$$

Now notice that

$$(\partial_k^+ f)(\eta) = \sum_{i=k}^{L-1} (\partial_{i+1, i} f)(\eta + \delta_{i+1}) + (\partial_L^+ f)(\eta)$$

and that the change of coordinates $\eta \mapsto \eta + \delta_j$ has bounded Jacobian. Thus (5.8) gives:

$$\bar{\mathbf{E}}_L^M(f, f) \leq C_3(\beta)(M^2 \vee L)L \left\{ \bar{\mathbf{E}}_L^M \left[\mathbf{1}(\eta_1 < M) (\partial_{2,1} f)^2 \right] + \sum_{k=2}^L \bar{\mathbf{E}}_L^M \left[(\partial_{k+1, k} f)^2 \right] + \bar{\mathbf{E}}_L^M \left[(\partial_L^+ f)^2 \right] \right\},$$

which is the same as (5.5). \square

The remaining part of this section is devoted to the proof of Proposition 5.2. It is convenient to introduce some extra notation. Recall that η_1 is independent from η_2, \dots, η_L and that $W_L^\infty(\beta, T(\eta))$ does not depend on η_1 . This implies that

$$\bar{\nu}_L^M(\eta|\eta_1) = \frac{(2M+1)e^{-\beta \sum_{i=2}^L |\eta_i| - W_L^\infty(\beta, T(\eta))}}{\bar{Z}_L^M(\beta)},$$

for every $|\eta_1| \leq M$. If we define $\eta'_1 \equiv \eta_2$, $\eta'_2 \equiv \eta_3, \dots, \eta'_{L-1} \equiv \eta_L$ and $\hat{W}_{L-1}(\eta') \equiv W_L^\infty(T(\eta))$, the last expression becomes:

$$\bar{\nu}_L^M(\eta|\eta_1) = \frac{(2M+1)e^{-\beta \sum_{i=1}^{L-1} |\eta'_i| - \hat{W}_{L-1}(\beta, \eta')}}{\bar{Z}_L^M(\beta)}.$$

Define on Ω_L the probability measure

$$\bar{\nu}_L(\eta) \equiv \frac{e^{-\beta \sum_{i=1}^L |\eta_i| - \hat{W}_L(\beta, \eta)}}{\bar{Z}_L(\beta)}, \quad \bar{Z}_L(\beta) \equiv \sum_{\eta} e^{-\beta \sum_{i=1}^L |\eta_i| - \hat{W}_L(\beta, \eta)}$$

and denote with $\bar{\mathbf{E}}_L(\cdot)$ the expected value with respect to this measure. We restate Proposition 5.2 as

Proposition 5.3. *There exists $\bar{\beta} > 0$ such that for every $\beta \geq \bar{\beta}$ it is possible to find $K_1(\beta) > 0$ such that*

$$\bar{\mathbf{E}}_L(f, f) \leq K_1(\beta) \sum_{k=1}^L \bar{\mathbf{E}}_L[(\partial_k^+ f)^2], \quad (5.9)$$

for any $f \in L^2(\bar{\nu}_L)$ and $L > 0$.

We will prove this result using the *martingale approach* outlined in [5].

Define the subsets α_j of \mathbb{N} in the following way

$$\alpha_j \equiv \begin{cases} \{j, \dots, L\} & \text{if } 1 \leq j \leq L, \\ \emptyset & \text{if } j = L+1. \end{cases}$$

For every $j = 1, \dots, L$ the restriction of η to the set α_j is denoted by $\eta_{\alpha_j} \equiv (\eta_j, \dots, \eta_L)$. If we define $f_j \equiv \bar{\mathbf{E}}_L(f|\eta_{\alpha_j})$ it is simple to check that:

$$\bar{\mathbf{E}}_L(f, f) = \sum_{i=1}^L \bar{\mathbf{E}}_L[\bar{\mathbf{E}}_L(f_j, f_j|\eta_{\alpha_{j+1}})]. \quad (5.10)$$

On the right hand side of this formula there is a sum of expected value of conditional variances. Notice that the random variable f_j , by definition, depends only on η_{α_j} . So if $\eta_{\alpha_{j+1}}$ is fixed, it depends only on η_j (we will say that f is *local* in j). This means that each of the variances on the right hand side of (5.10) is the variance of a local function.

The method we will use to prove (5.9) consists of two steps. The first step is to show that the marginal in η_j of the measure $\bar{\nu}_L(\cdot|\eta_{\alpha_{j+1}})$ exhibits a positive spectral gap uniformly in $L > 0$.

Lemma 5.4. *There exists $\bar{\beta} > 0$ such that for every $\beta > 0$ there exists $K_1(\beta) > 0$ so that*

$$\bar{\mathbf{E}}_L(f, f|\eta_{\alpha_j}) \leq K_1(\beta) \bar{\mathbf{E}}_L [(\partial_j^+ f)^2|\eta_{\alpha_j}], \quad (5.11)$$

for every $L > 0$ and $f \in L^2(\bar{\nu}_L(\cdot|\eta_{\alpha_j}))$ local in j .

Because of this lemma (5.10) becomes

$$\bar{\mathbf{E}}_L(f, f) \leq K_1(\beta) \sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2]. \quad (5.12)$$

The second step is to show that the right hand side of (5.12) can be bounded from above by the correct quadratic form:

Lemma 5.5. *There exists $\bar{\beta}$ such that*

$$\sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2] \leq 4 \sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2], \quad (5.13)$$

for every $\beta \geq \bar{\beta}$, $L > 0$ and $f \in L^2(\bar{\nu})$.

If we use Lemma 5.5 in (5.12) we obtain immediately (5.9).

In order to prove Lemma 5.4 and Lemma 5.5 we need a preliminary result.

Lemma 5.6. *There exists $\bar{\beta} > 0$ such that for every $\beta > \bar{\beta}$*

$$\exp \left[-\beta \operatorname{sign}(\eta_j) - 8e^{-m(\beta)} \right] \leq \frac{\bar{\nu}_L(\eta_j + \delta_j|\eta_{\alpha_{j+1}})}{\bar{\nu}_L(\eta_j|\eta_{\alpha_{j+1}})} \leq \exp \left[-\beta \operatorname{sign}(\eta_j) + 8e^{-m(\beta)} \right] \quad (5.14)$$

and

$$\exp \left[-16e^{-m(\beta)(i-j)} \right] \leq \frac{\bar{\nu}_L(\eta_j|\eta_{\alpha_{j+1}} + \delta_i)}{\bar{\nu}_L(\eta_j|\eta_{\alpha_{j+1}})} \leq \exp \left[16e^{-m(\beta)(i-j)} \right], \quad (5.15)$$

for every $j = 1, \dots, L-1$, $\eta \in \Omega_L$.

Proof.

The proof is divided in several steps for purpose of clarity. To keep notation simple we will write $\Delta(\eta)$ instead of $\Delta(T\eta)$ and

$$\mathcal{S}(\eta) \equiv \{\Lambda \subset V_L^\infty : \Lambda \cap \Delta(\eta) \neq \emptyset\}.$$

Recall that

$$\hat{W}_L(\beta, \eta) = \sum_{\Lambda \in \mathcal{S}(\eta)} \Phi(\beta, \Lambda).$$

Define for $z \in \mathbb{R}$ the line $r(z) \equiv \{(x, y) \in \mathbb{R}^2 : x = z\}$ and for fixed $j = 1, \dots, L-1$

$$\mathcal{S}_j(\eta) \equiv \mathcal{S}(\eta) \setminus \{\Lambda \in \mathcal{S}(\eta) : \Lambda > r(j)\}.$$

Step 1. Define

$$\hat{W}_j(\eta) \equiv \sum_{\Lambda \in \mathcal{S}_j(\eta)} \Phi(\beta, \Lambda).$$

Then

$$\bar{\nu}_L(\eta_j | \eta_{\alpha_{j+1}}) = \frac{e^{-\beta|\eta_j|} \sum_{\eta_1, \dots, \eta_{j-1}} e^{-\beta \sum_{k=1}^{j-1} |\eta_k| - \hat{W}_j(\eta)}}{\sum_{\eta_1, \dots, \eta_j} e^{-\beta \sum_{k=1}^j |\eta_k| - \hat{W}_j(\eta)}}. \quad (5.16)$$

Proof of Step 1.

Define

$$\hat{W}_j^c(\eta) \equiv \hat{W}_L(\eta) - \hat{W}_j(\eta) = \sum_{\Lambda \in \mathcal{S}(\eta) \setminus \mathcal{S}_j(\eta)} \Phi(\beta, \Lambda),$$

we claim that \hat{W}_j^c does not depend on η_1, \dots, η_j . If we assume this we obtain:

$$\begin{aligned} \bar{\nu}_L(\eta_j | \eta_{\alpha_{j+1}}) &= \frac{e^{-\beta|\eta_j|} \sum_{\eta_1, \dots, \eta_{j-1}} e^{-\beta \sum_{k=1}^{j-1} |\eta_k| - \hat{W}_L(\eta)}}{\sum_{\eta_1, \dots, \eta_j} e^{-\beta \sum_{k=1}^j |\eta_k| - \hat{W}_j(\eta)}} = \\ &= \frac{e^{-\beta|\eta_j|} \sum_{\eta_1, \dots, \eta_{j-1}} e^{-\beta \sum_{k=1}^{j-1} |\eta_k| - \hat{W}_j(\eta)}}{\sum_{\eta_1, \dots, \eta_j} e^{-\beta \sum_{k=1}^j |\eta_k| - \hat{W}_j(\eta)}}, \end{aligned}$$

i.e. (5.16).

In order to prove that $\hat{W}_j^c(\eta)$ does not depend on η_1, \dots, η_j it suffices to show that

$$\hat{W}_j^c(\eta + h\delta_k) = \hat{W}_j^c(\eta)$$

for every $h \in \mathbb{Z}$ and $k = 1, \dots, j$. Define

$$T_h : \Lambda \ni \mathcal{S}(\eta) \setminus \mathcal{S}_j(\eta) \mapsto \Lambda + h \in \mathcal{S}(\eta + h\delta_k) \setminus \mathcal{S}_j(\eta + h\delta_k).$$

This map is bijective for every $k = 1, \dots, j$. Furthermore because of the translation invariance (2.4) of $\Phi(\beta, \cdot)$ we have:

$$\begin{aligned} \hat{W}_j^c(\eta) &= \sum_{\Lambda \in \mathcal{S}(\eta) \setminus \mathcal{S}_j(\eta)} \Phi(\beta, \Lambda) = \sum_{\Lambda \in \mathcal{S}(\eta) \setminus \mathcal{S}_j(\eta)} \Phi(\beta, T_h \Lambda) = \\ &= \sum_{\Lambda \in \mathcal{S}(\eta + h\delta_k) \setminus \mathcal{S}_j(\eta + h\delta_k)} \Phi(\beta, \Lambda) = \hat{W}_j^c(\eta + h\delta_k). \end{aligned}$$

□

From Step 1 we obtain

$$\frac{\bar{\nu}_L(\eta_j + \delta_j | \eta_{\alpha_{j+1}})}{\bar{\nu}_L(\eta_j | \eta_{\alpha_{j+1}})} = e^{-\beta(|\eta_j+1|-|\eta_j|)} \frac{\sum_{\eta_1, \dots, \eta_{j-1}} e^{-\beta \sum_{k=1}^{j-1} |\eta_k| - \hat{W}_j(\eta + \delta_j)}}{\sum_{\eta_1, \dots, \eta_{j-1}} e^{-\beta \sum_{k=1}^{j-1} |\eta_k| - \hat{W}_j(\eta)}}$$

and:

$$\frac{\bar{\nu}_L(\eta_j | \eta_{\alpha_{j+1}} + \delta_i)}{\bar{\nu}_L(\eta_j | \eta_{\alpha_{j+1}})} = \frac{\sum_{\eta_2, \dots, \eta_{j-1}} e^{-\beta \sum_{i=2}^{j-1} |\eta_i| - \hat{W}_j(\eta + \delta_i)}}{\sum_{\eta_2, \dots, \eta_{j-1}} e^{-\beta \sum_{i=2}^{j-1} |\eta_i| - \hat{W}_j(\eta)}} \frac{\sum_{\eta_2, \dots, \eta_j} e^{-\beta \sum_{i=2}^j |\eta_i| - \hat{W}_j(\eta)}}{\sum_{\eta_2, \dots, \eta_j} e^{-\beta \sum_{i=2}^j |\eta_i| - \hat{W}_j(\eta + \delta_i)}},$$

for every $i > j$. By these inequalities we obtain that to prove (5.14) and (5.15) we have only to show that for every $i \geq j$

$$\sup_{\eta} |\hat{W}_j(\eta + \delta_i) - \hat{W}_j(\eta)| \leq 8e^{-m(\beta)(i+1-j)}. \quad (5.17)$$

Step 2. Define

$$\Delta_i(\eta) \equiv \Delta(\eta) \cap \{(x, y) \in \mathbb{R}^2 : x > i\}$$

and

$$\mathcal{S}_{i,j}(\eta) \equiv \{\Lambda \subset V_L^\infty : \Lambda \cap \Delta_i(\eta) \neq \emptyset, \Lambda \cap r(j) \neq \emptyset\}.$$

Then:

$$\hat{W}_j(\eta + \delta_i) - \hat{W}_j(\eta) = \sum_{\Lambda \in \mathcal{S}_{i,j}(\eta + \delta_i)} \Phi(\beta, \Lambda) - \sum_{\Lambda \in \mathcal{S}_{i,j}(\eta)} \Phi(\beta, \Lambda). \quad (5.18)$$

Proof of Step 2.

Let $\Lambda \in \mathcal{S}_j(\eta)$ be such that it intersects $\Delta(\eta)$ on the left of i . Then it also intersects $\mathcal{S}_j(\eta + \delta_i)$ in the same points. On the contrary if $\Lambda \in \mathcal{S}_j(\eta)$ intersects $\Delta(\eta + \delta_i)$ on the left of i then it intersects $\mathcal{S}_j(\eta)$. In conclusion

$$\left\{ \Lambda \in \mathcal{S}_j(\eta) : \begin{array}{l} \Lambda \text{ intersects } \Delta(\eta) \\ \text{on the left of } i \end{array} \right\} = \left\{ \Lambda \in \mathcal{S}_j(\eta + \delta_i) : \begin{array}{l} \Lambda \text{ intersects } \Delta(\eta + \delta_i) \\ \text{on the left of } i \end{array} \right\}.$$

By this relation we obtain that we can clear from the difference

$$\hat{W}_j(\eta + \delta_i) - \hat{W}_j(\eta) = \sum_{\Lambda \in \mathcal{S}_j(\eta)} \Phi(\beta, \Lambda) - \sum_{\Lambda \in \mathcal{S}_j(\eta + \delta_i)} \Phi(\beta, \Lambda),$$

all the terms $\Phi(\beta, \Lambda)$ such that Λ intersects $\Delta(\eta)$, or $\Delta(\eta + \delta_i)$, on the left of i . It follows that the sums are actually only on the Λ which neither intersects $\Delta(\eta)$ on the left of i nor intersects

$\Delta(\eta + \delta_i)$ on the left of i . Because these Λ have to intersect $\Delta(\eta)$, the intersection is on the right of i . This proves (5.18). \square

For any $S \subset (\mathbb{Z}^2)^*$ we will say that $p \in S$ is *+unstable* if:

$$p \notin (S + \mathbf{e}_y),$$

where $\mathbf{e}_y = (0, 1) \in \mathbb{R}^2$. Similarly we will say that $p \in S$ is *-unstable* if:

$$p \notin (S - \mathbf{e}_y).$$

The classes of points +unstable and -unstable of the set S will be denoted respectively with $\mathcal{I}^+(S)$ and $\mathcal{I}^-(S)$.

Step 3. *Define*

$$\begin{aligned} \mathcal{S}_{i,j}^+(\eta) &\equiv \mathcal{S}_{i,j}(\eta) \cap \mathcal{I}^+(\Delta_i(\eta)) \\ \mathcal{S}_{i,j}^-(\eta) &\equiv \mathcal{S}_{i,j}(\eta) \cap \mathcal{I}^-(\Delta_i(\eta)). \end{aligned}$$

Then

$$\hat{W}_j(\eta + \delta_i) - \hat{W}_j(\eta) = \sum_{\Lambda \in \mathcal{S}_{i,j}^-(\eta + \delta_i)} \Phi(\beta, \Lambda) - \sum_{\Lambda \in \mathcal{S}_{i,j}^+(\eta)} \Phi(\beta, \Lambda). \quad (5.19)$$

Proof.

It simple to check that:

$$\Lambda \in \mathcal{S}_{i,j}(\eta + \delta_i), \quad \Lambda \notin \mathcal{I}^-(\Delta_i(\eta + \delta_i)) \quad \iff \quad \Lambda \in \mathcal{S}_{i,j}(\eta), \quad \Lambda \notin \mathcal{I}^+(\Delta_i(\eta)).$$

This implies $\mathcal{S}_{i,j}(\eta + \delta_i) \setminus \mathcal{I}^-(\Delta_i(\eta + \delta_i)) = \mathcal{S}_{i,j}(\eta) \setminus \mathcal{I}^+(\Delta_i(\eta))$, which proves (5.19). \square

Step 4. *For any $\eta \in \Omega_L$ and $k = 1/2, 1 + 1/2, 2 + 1/2, \dots$*

$$|\mathcal{I}^\pm(\Delta(\eta)) \cap r(k)| \leq 2.$$

Proof.

If $\Delta = \Delta(\eta)$ it is clear that $\Delta = \overline{\Delta} \cup \underline{\Delta}$ where

$$\overline{\Delta} \equiv \{p \in \Delta : p \text{ is above } \Gamma(T\eta)\} \quad \underline{\Delta} \equiv \{p \in \Delta : p \text{ is below } \Gamma(T\eta)\}.$$

Notice that in general $\mathcal{I}^\pm(A \cup B) \subset \mathcal{I}^\pm(A) \cup \mathcal{I}^\pm(B)$. Thus to prove (5.19) we have only to show that:

$$|\mathcal{I}^\pm(\overline{\Delta})| = 1 \quad \text{and} \quad |\mathcal{I}^\pm(\underline{\Delta})| = 1.$$

This fact can be easily checked by using geometric considerations. \square

We are finally in a position to prove (5.17). Notice that if $\Lambda \in \mathcal{S}_{i,j}^+$ then:

- i) Λ contains $p \in \mathcal{I}^+(\Delta_i(\eta))$;
- ii) Λ intersects $r(j)$, thus because $\Lambda \subset (\mathbb{Z}^2)^*$, intersects $r(j-1/2)$.

It follows that:

$$\begin{aligned} \sum_{\Lambda \in \mathcal{S}_{i,j}^+(\eta)} \Phi(\beta, \Lambda) &\leq \sum_{k=i}^{+\infty} \sum_{p \in \mathcal{I}^+(\Delta_i(\eta)) \cap r(k+1/2)} \sum_{\substack{\Lambda \ni p \\ \Lambda \cap r(j-1/2) \neq \emptyset}} \Phi(\beta, \Lambda) \leq \\ &\leq 2 \sum_{k=i}^{+\infty} \sum_{\substack{\Lambda \ni p \\ \text{diam}(\Lambda) \geq k-j+1}} \Phi(\beta, \Lambda) \leq 2 \sum_{k=i}^{+\infty} e^{-m(\beta)(k-j+1)} \leq 4e^{-m(\beta)(k-j+1)}. \end{aligned}$$

Where β is large enough and (2.3) has been used. From this estimate we obtain (5.17) that, as we noticed before, implies (5.14) and (5.15). \square

Lemma 5.6 can be used to prove the one-site spectral gap Lemma 5.4. In fact (5.14) shows that for β large enough (recall that $m(\beta) \rightarrow +\infty$ for $\beta \rightarrow +\infty$) the measure $\bar{\nu}_L(\cdot|\eta_{\alpha_j})$ exhibits an “inward drift” (see [11]).

Proof of Lemma 5.6.

It follows immediately from [3] and [10]. \square

Now we turn to the proof of Lemma 5.5. We need a technical result

Lemma 5.7. *For every $i = 2, \dots, L$ we have:*

$$\partial_i^+ f_i = \bar{\mathbf{E}}_L(\partial_i f|\eta_{\alpha_i}) + \sum_{j=1}^{i-1} \bar{\mathbf{E}}_L(f_j^+, V_{i,j}|\eta_{\alpha_i}), \quad (5.20)$$

where $f_j^+(\eta) \equiv f_j(\eta + \delta_i)$ and $V_{i,j}(\eta) \equiv \frac{\bar{\nu}_L(\eta_j|\eta_{\alpha_{j+1}+\delta_i})}{\bar{\nu}_L(\eta_j|\eta_{\alpha_{j+1}})} - 1$.

Proof.

For every $k = 1, \dots, L$ define

$$F_k(\eta) \equiv \bar{\nu}_L(\eta_k|\eta_{\alpha_{k+1}}).$$

F_k is the marginal in η_k of $\bar{\nu}_L(\cdot|\eta_{\alpha_{k+1}})$. Notice that

$$f_i = \bar{\mathbf{E}}_L(f|\eta_{\alpha_i}) = \bar{\mathbf{E}}_L(\bar{\mathbf{E}}_L(\cdots \bar{\mathbf{E}}_L(f|\eta_{\alpha_2}) \cdots |\eta_{\alpha_{i-1}})|\eta_{\alpha_i})$$

for any $i = 2, \dots, L$. Thus

$$f_i = \sum_{\eta_1, \dots, \eta_{i-1}} f \prod_{k=1}^{i-1} F_k. \quad (5.21)$$

To compute $\partial_i^+ f_i$ we have to calculate the (discrete) derivative of a product. Notice that:

$$(\partial_i^+ gh) = (\partial_i^+ g)h^+ + g(\partial_i^+ h), \quad (5.22)$$

where $g^+(\eta) \equiv g(\eta + \delta_i)$. Using (5.22) we obtain:

$$\partial_i^+ \left(f \prod_{k=1}^{i-1} F_k \right) = (\partial_i^+ f) \prod_{k=1}^{i-1} F_k + \sum_{j=1}^{i-1} f^+ (\partial_i^+ F_j) \left(\prod_{l=1}^{j-1} F_l^+ \right) \left(\prod_{k=j+1}^{i-1} F_k \right).$$

This relation and (5.21) give:

$$\begin{aligned} \partial_i^+ f_i &= \sum_{\eta_1, \dots, \eta_{i-1}} (\partial_i^+ f) \prod_{k=1}^{i-1} F_k + \sum_{\eta_1, \dots, \eta_{i-1}} f^+ \sum_{j=1}^{i-1} (\partial_i^+ F_j) \left(\prod_{l=1}^{j-1} F_l^+ \right) \left(\prod_{k=j+1}^{i-1} F_k \right) = \\ &= \bar{\mathbf{E}}_L(\partial_i^+ f | \eta_{\alpha_i}) + \sum_{j=1}^{i-1} \sum_{\eta_j, \dots, \eta_{i-1}} (\partial_i^+ F_j) \left(\prod_{k=j+1}^{i-1} F_k \right) \sum_{\eta_1, \dots, \eta_{j-1}} f^+ \left(\prod_{l=1}^{j-1} F_l^+ \right) = \\ &= \bar{\mathbf{E}}_L(\partial_i^+ f | \eta_{\alpha_i}) + \sum_{j=1}^{i-1} \sum_{\eta_j, \dots, \eta_{i-1}} f_j^+ (\partial_i^+ F_j) \left(\prod_{k=j+1}^{i-1} F_k \right) = \\ &= \bar{\mathbf{E}}_L(\partial_i^+ f | \eta_{\alpha_i}) + \sum_{j=1}^{i-1} \sum_{\eta_j, \dots, \eta_{i-1}} f_j^+ \left(\frac{F_j^+}{F_j} - 1 \right) \left(\prod_{k=j}^{i-1} F_k \right) = \\ &= \bar{\mathbf{E}}_L(\partial_i^+ f | \eta_{\alpha_i}) + \bar{\mathbf{E}}_L(f^+ V_{i,j} | \eta_{\alpha_i}). \end{aligned}$$

From this (5.20) follows because $\bar{\mathbf{E}}_L(V_{i,j} | \eta_{\alpha_i}) = 0$. \square

Proof of Lemma 5.5.

We borrow the basic idea of the proof from [9]. We will show that for β large enough

$$\sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f_i)^2] \leq 2 \sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2] + \frac{1}{2} \sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f_i)^2], \quad (5.23)$$

for every $f \in L^2(\bar{\nu}_L)$.

Fix $f \in L^2(\bar{\nu}_L)$ and $i > 1$. By Lemma 5.7

$$(\partial_i^+ f_i)^2 \leq 2 \bar{\mathbf{E}}_L [(\partial_i^+ f)^2 | \eta_{\alpha_i}] + 2 \left[\sum_{j=1}^{i-1} \bar{\mathbf{E}}_L(f_j^+, V_{i,j} | \eta_{\alpha_i}) \right]^2. \quad (5.24)$$

We have to estimate the second term on the right hand side of this relation. By Schwartz inequality and Lemma 5.6 we obtain

$$\bar{\mathbf{E}}_L(f_j^+, V_{i,j} | \eta_{\alpha_i}) \leq 2\varepsilon^{i-j} \bar{\mathbf{E}}_L(f_j^+, f_j^+ | \eta_{\alpha_i})^{\frac{1}{2}},$$

where $\varepsilon = \varepsilon(\beta) \rightarrow 0$ for $\beta \rightarrow +\infty$. Thus if β is large enough

$$\begin{aligned} \left[\sum_{j=1}^{i-1} \overline{\mathbf{E}}_L(f_j^+, V_{i,j} | \eta_{\alpha_i}) \right]^2 &\leq 4 \left[\sum_{j=1}^{i-1} \varepsilon^{i-j} \overline{\mathbf{E}}_L(f_j^+, f_j^+ | \eta_{\alpha_i})^{\frac{1}{2}} \right]^2 \leq \\ &\leq 4 \left(\sum_{j=1}^{i-1} \varepsilon^{i-j} \right) \sum_{j=1}^{i-1} \varepsilon^{i-j} \overline{\mathbf{E}}_L(f_j^+, f_j^+ | \eta_{\alpha_i}) \leq 8\varepsilon \sum_{j=1}^{i-1} \varepsilon^{i-j} \overline{\mathbf{E}}_L(f_j^+, f_j^+ | \eta_{\alpha_i}). \end{aligned} \quad (5.25)$$

It simple to check that

$$\overline{\mathbf{E}}_L(f_j^+, f_j^+ | \eta_{\alpha_i}) = \sum_{s=j}^{i-1} \overline{\mathbf{E}}_L \left[\overline{\mathbf{E}}_L(f_s^+, f_s^+ | \eta_{\alpha_{s+1}}) | \eta_{\alpha_i} \right].$$

So by (5.11) we know that, for β large enough, there exists $C_1 > 0$ so that:

$$\overline{\mathbf{E}}_L(f_j^+, f_j^+ | \eta_{\alpha_i}) \leq C_1 \sum_{s=j}^{i-1} \overline{\mathbf{E}}_L \left[\overline{\mathbf{E}}_L \left((\partial_s^+ f_s^+)^2 | \eta_{\alpha_{s+1}} \right) | \eta_{\alpha_i} \right] = C_1 \sum_{s=j}^{i-1} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right].$$

By using this bound in (5.25) we get

$$\left[\sum_{j=1}^{i-1} \overline{\mathbf{E}}_L(f_j^+, V_{i,j} | \eta_{\alpha_i}) \right]^2 \leq \varepsilon C_2 \sum_{j=1}^{i-1} \varepsilon^{i-j} \sum_{s=j}^{i-1} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right]. \quad (5.26)$$

Exchanging the sums on the right hand side of (5.26) we have

$$\sum_{j=1}^{i-1} \varepsilon^{i-j} \sum_{s=j}^{i-1} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right] = \sum_{s=1}^{i-1} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right] \sum_{j=1}^s \varepsilon^{i-j} \leq 2 \sum_{s=1}^{i-1} \varepsilon^{i-s} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right].$$

This implies

$$\left[\sum_{j=1}^{i-1} \overline{\mathbf{E}}_L(f_j^+, V_{i,j} | \eta_{\alpha_i}) \right]^2 \leq \varepsilon C_3 \sum_{s=1}^{i-1} \varepsilon^{i-s} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right].$$

and by (5.24):

$$(\partial_i^+ f_i)^2 \leq 2 \overline{\mathbf{E}}_L \left[(\partial_i^+ f)^2 | \eta_{\alpha_i} \right] + \varepsilon C_4 \sum_{s=1}^{i-1} \varepsilon^{i-s} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s^+)^2 | \eta_{\alpha_i} \right].$$

Taking expected value on both sides of this relation and recalling that the change of variable $\eta \mapsto \eta - \delta_i$ has a bounded Jacobian (see Lemma 5.6) we obtain

$$\overline{\mathbf{E}}_L \left[(\partial_i^+ f_i)^2 \right] \leq 2 \overline{\mathbf{E}}_L \left[(\partial_i^+ f)^2 \right] + \varepsilon C_4 \sum_{s=1}^{i-1} \varepsilon^{i-s} \overline{\mathbf{E}}_L \left[(\partial_s^+ f_s)^2 \right].$$

We sum this relation for $i = 2, \dots, L$. An elementary computation gives

$$\begin{aligned} \sum_{i=2}^L \bar{\mathbf{E}}_L [(\partial_i^+ f_i)^2] &\leq 2 \sum_{i=2}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2] + C_4 \varepsilon \sum_{i=2}^L \sum_{s=1}^{i-1} \varepsilon^{i-s} \bar{\mathbf{E}}_L [(\partial_s^+ f_s)^2] = \\ &= 2 \sum_{i=2}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2] + C_4 \varepsilon \sum_{s=1}^{L-1} \sum_{i=s+1}^L \varepsilon^{i-s} \bar{\mathbf{E}}_L [(\partial_s^+ f_s)^2] \leq \\ &\leq 2 \sum_{i=2}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2] + 2C_4 \varepsilon^2 \sum_{s=1}^{L-1} \bar{\mathbf{E}}_L [(\partial_s^+ f_s)^2]. \end{aligned}$$

Recalling that $\partial_i^+ f_1 = \partial_i^+ f = \bar{\mathbf{E}}_L(\partial_i^+ f | \eta_{\alpha_1})$, this implies:

$$\sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f_i)^2] \leq 2 \sum_{i=1}^L \bar{\mathbf{E}}_L [(\partial_i^+ f)^2] + \varepsilon^2 C_5 \sum_{s=1}^L \bar{\mathbf{E}}_L [(\partial_s^+ f_s)^2].$$

To conclude the proof of (5.23) we choose $\bar{\beta} > 0$ such that $\beta \geq \bar{\beta}$ implies $C_5 \varepsilon^2(\beta) < 1/2$. \square

A. Appendix

Lemma A1.1. Define $A \equiv \{\phi \in \Omega_L : \|\phi\|_\infty \leq (1 - \varepsilon)L/2\}$ and $B \equiv \{\phi \in \Omega_L : \|\phi\|_\infty \leq \alpha L\}$ where $\varepsilon \in (0, 1/100)$ and $\alpha \in (0, 1/4)$. Then there exists $\bar{\beta} > 0$ such that for every $\beta > \bar{\beta}$ there exists $K_1(\beta) > 0$ so that

$$\sup_{\phi \in A} \frac{\mu_L^{L/2}(\phi|B)}{\bar{\mu}_L^{L/2}(\phi)} \leq \alpha^{-1} K_1. \quad (\text{A1.1})$$

Proof.

An elementary calculation shows that:

$$\frac{\mu_L^{L/2}(\phi|B)}{\bar{\mu}_L^{L/2}(\phi)} \leq e^{\sup_{\phi \in B} |W_L^\infty(\phi) - W_L^{L/2}(\phi)|} \bar{\mu}_L^{L/2}(B)^{-1}, \quad (\text{A1.2})$$

for every $\phi \in B$. Observe that:

$$W_L^\infty(\beta, \phi) - W_L^{L/2}(\beta, \phi) = \sum_{\Lambda \cap \Delta(\phi) \neq \emptyset} \Phi(\beta, \Lambda) - \sum_{\substack{\Lambda \cap \Delta(\phi) \neq \emptyset \\ \Lambda \subset V_L^{L/2}}} \Phi(\beta, \Lambda) = \sum_{\substack{\Lambda \cap \Delta(\phi) \neq \emptyset \\ \Lambda \cap (V_L^{L/2})^c \neq \emptyset}} \Phi(\beta, \Lambda).$$

Thus if $\phi \in B$, $\Lambda \cap \Delta(\phi) \neq \emptyset$ and $\Lambda \cap (V_L^{L/2})^c \neq \emptyset$, necessarily $\text{diam}(\Lambda) \geq (L/2)[(1/2) - \alpha]$. By (2.3) we obtain

$$\begin{aligned} |W_L^\infty(\beta, \phi) - W_L^{L/2}(\beta, \phi)| &\leq \sum_{\substack{\Lambda \cap \Delta(\phi) \neq \emptyset \\ \Lambda \cap (V_L^{L/2})^c \neq \emptyset}} |\Phi(\beta, \Lambda)| \leq \\ &\leq \sum_{p \in V_L^{L/2}} \sum_{\substack{\Lambda \ni p \\ \text{diam}(\Lambda) \geq (L/2)(1/2 - \alpha)}} |\Phi(\beta, \Lambda)| \leq L^2 e^{-m(\beta)(L/2)(1/2 - \alpha)} \leq C_1(\beta). \end{aligned}$$

This bound and (A1.2) give

$$\frac{\mu_L^{L/2}(\phi|B)}{\bar{\mu}_L^{L/2}(\phi)} \leq e^{C_1} \bar{\mu}_L^{L/2}(B)^{-1}.$$

To complete the proof we have to bound $\bar{\mu}_L^{L/2}(B)$ from below. We refer to [2] to prove that there exists $C_2(\beta) > 0$ such that $\bar{\mu}_L^{L/2}(B) \geq \alpha C_2(\beta)$. \square

B. References

- [1] P. Diaconis and L. Saloff-Coste: Comparison theorems for reversible Markov chains. *Ann. Appl. Probab.* **3** (1993), no. 3, 696–730.
- [2] R. Dobrushin, R. Kotecký and S. Shlosman: *Wulff Construction. A Global Shape from Local Interaction*, Translation of Mathematical Monographs, **104** (1992). AMS.
- [3] G. F. Lawler and A. D. Sokal: Bounds on the L^2 Spectrum for Markov Chains and Markov Processes: a Generalization of Cheeger’s Inequality. *Trans. Amer. Math. Soc.* **309** (1988), no. 2, 557–580.
- [4] T. M. Liggett: *Interacting particle systems*. Grundlehren der Mathematischen Wissenschaften **276** (1985). Springer-Verlag.
- [5] S. Lu and H.-T. Yau: Spectral gap and logarithmic Sobolev inequality for Kawasaki and Glauber dynamics. *Comm. Math. Phys.* **156** (1993), no. 2, 399–433.
- [6] F. Martinelli: On the two dimensional dynamical Ising model in the phase coexistence region. *J. Statist. Phys.* **76** (1994), no. 5-6, 1179–1246.
- [7] F. Martinelli: Lectures on Glauber dynamics for discrete spin models in *Lectures on probability theory and statistics (Saint-Flour, 1997)*. Lecture Notes in Math. **1717** (1999), 93–191. Springer-Verlag.
- [8] F. Martinelli and E. Olivieri: Approach to equilibrium of Glauber dynamics in the one phase region I: the attractive case. *Comm. Math. Phys.* **161** (1994), no. 3, 447–486.
- [9] F. Martinelli and E. Olivieri: Approach to equilibrium of Glauber dynamics in the one phase region II: the general case. *Comm. Math. Phys.* **161** (1994), no. 3, 487–514.

- [10] G. Posta: Spectral Gap for an Unrestricted Kawasaki Type Dynamics. *ESAIM Probability & Statistics* **1** (1997), 145–181.
- [11] A. D. Sokal and L. E. Thomas: Absence of mass gap for a class of stochastic contour models. *J. Statist. Phys.* **51** (1988), no. 5-6, 907–947.
- [12] D. W. Stroock and B. Zegarlinski: The logarithmic Sobolev inequality for discrete spin on a lattice. *Comm. Math. Phys.* **149** (1992), no. 1, 175–193.