

MONOTONICITY OF THE VARIANCE UNDER TRUNCATION AND
 VARIATIONS OF JENSEN'S INEQUALITY¹

BY Y. S. CHOW AND W. J. STUDDEN

Purdue University

Let X be a random variable and $Y = XI_{[X < b]}$ for $-\infty < b < \infty$. Then it is easy to construct examples for which $\sigma^2 X < \sigma^2 Y$; e.g. take $X = b \pm \epsilon$ with probability $\frac{1}{2}$ where $\epsilon \ll b$. However if $Y = \min(X, b)$ then we always have $\sigma^2 X \geq \sigma^2 Y$. The essential difference is that we are now replacing any value $X > b$ by b instead of 0. Similarly if $Y = \max(a, X)$ or $\max(a, \min(X, b))$ then $\sigma^2 X \geq \sigma^2 Y$. The main result in this note is Theorem 2 and Corollary 4 which contains a proof of a conditional version of the above result which was used in [2]. We will prove the above facts and some of their generalizations which provide intermediate terms in Jensen's inequality; see Corollary 1. The results and the methods of proof given below are actually special cases or slight modifications of more general inequalities involving duals of cones of generalized convex functions. See for example Karlin and Novikoff [3], Karlin and Studden [4], Ziegler [5], Barlow, Marshall and Proschan [1] and references therein. The methods used below are quite elementary and produce the desired results without recourse to dual cones.

THEOREM 1. *Let h and g be two Baire functions on the real line and X a random variable such that (a) g is nondecreasing, (b) $Eg(X)$ and $Eh(X)$ exist and $Eg(X) = Eh(X)$, (c) the function $h - g$ has one sign change from negative to positive, i.e., there exists t_0 such that $(h(t) - g(t))(t - t_0) \geq 0$ for every t . If $E\varphi(h(X))$ and $E\varphi(g(X))$ exist, then*

$$(1) \quad E\varphi(h(X)) \geq E\varphi(g(X))$$

for all continuous, convex φ on $(-\infty, \infty)$.

PROOF. By the convexity of φ ,

$$(2) \quad \varphi(h(t)) - \varphi(g(t)) \geq \varphi'(g(t))(h(t) - g(t)),$$

where φ' denotes the right (or left) derivative of φ . If $t > t_0$, then $h(t) - g(t) \geq 0$ and $\varphi'(g(t)) \geq \varphi'(g(t_0))$, so that

$$(3) \quad \varphi(h(t)) - \varphi(g(t)) \geq \varphi'(g(t_0))(h(t) - g(t)).$$

The above equation also holds for $t < t_0$, since $h(t) - g(t) \leq 0$ and $\varphi'(g(t)) \leq \varphi'(g(t_0))$. Therefore (3) holds for all t . (1) follows immediately from (3).

COROLLARY 1. *Suppose that f is nondecreasing, φ is continuous, convex and for a random variable X , EX and $Ef(X)$ exist, and $t - f(t) + Ef(X) - EX$ has one*

Received 20 February 1968; revised 25 November 1968.

¹ This research was supported by the National Science Foundation Grant GP 06073.

sign change from negative to positive. Then

$$(4) \quad E\varphi(X - EX) \geq E\varphi(f(X) - Ef(X)),$$

$$(5) \quad E\varphi(X) \geq E\varphi(f(X) - Ef(X) + EX) \geq \varphi(EX),$$

provided that all the expectations involved in (4) and (5) exist.

All of the inequalities in (4) and (5) follow from (1) by taking the obvious choices of g and h in Theorem 1. The last inequality in (5) also follows from Jensen's inequality; however, if we apply (1) by taking $h(t) = f(t) - Ef(X) + EX$ and $g(t) \equiv EX$ then the two extremes $E\varphi(X)$ and $\varphi(EX)$ result in Jensen's inequality. The interesting aspect of (5) is that it produces intermediate terms between the two extremes. Note that $\varphi(EX)$ is attained for $f(t) \equiv \text{constant}$ and $E\varphi(X)$ results if $f(t) \equiv t + \text{constant}$. Useful particular cases of (4) and (5) are provided by:

COROLLARY 2. Equations (4) and (5) hold for

$$(6) \quad \begin{aligned} f(t) = f_{a,b}(t) &= b, & t \geq b \\ &= t, & a < t < b, \\ &= a, & t \leq a, \end{aligned}$$

where $-\infty \leq a \leq b \leq \infty$, ($a = b \neq \pm\infty$) or

$$(7) \quad f(t) = f_c(t) = ct, \quad 0 \leq c \leq 1.$$

Moreover, $E\varphi(f_{a,b}(X) - Ef_{a,b}(X) + EX)$ is nonincreasing in a , nondecreasing in b , and continuous in both variables; while $E\varphi(f_c(X) - Ef_c(X) + EX)$ is nondecreasing and continuous in c .

COROLLARY 3. If $E|X|^\alpha < \infty$ for some $\alpha \geq 1$, then

$$(8) \quad E|X - EX|^\alpha \geq E|f_{a,b}(X) - Ef_{a,b}(X)|^\alpha,$$

where $f_{a,b}$ is given by (6).

As mentioned earlier the motivation for this note was in proving a conditional version of (8). This is the content of the following theorem. Various other extensions can be established by the same methods.

THEOREM 2. Let \mathcal{G} be a σ -field of measurable sets and a, b be \mathcal{G} -measurable random variables. Define $X_{a,b} = \max(a, \min(X, b))$. If $E|X| < \infty$, $E|X_{a,b}| < \infty$ and $E\varphi(X)$ exists for a continuous, convex function φ on $(-\infty, \infty)$, then almost everywhere

$$(9) \quad E(\varphi(X) | \mathcal{G}) \geq E(\varphi(X_{a,b} - E(X_{a,b} | \mathcal{G}) + E(X | \mathcal{G})) | \mathcal{G}),$$

$$(10) \quad E(\varphi(X - E(X | \mathcal{G})) | \mathcal{G}) \geq E(\varphi(X_{a,b} - E(X_{a,b} | \mathcal{G})) | \mathcal{G}).$$

PROOF. By (3),

$$\begin{aligned} \varphi(X - E(X | \mathcal{G})) - \varphi(X_{a,b} - E(X_{a,b} | \mathcal{G})) \\ \geq \varphi'(Y)(X - E(X | \mathcal{G}) - X_{a,b} + E(X_{a,b} | \mathcal{G})), \end{aligned}$$

where

$$\begin{aligned} Y &= a + E(X | \mathcal{G}) - E(X_{a,b} | \mathcal{G}), & \text{if } E(X | \mathcal{G}) \leq E(X_{a,b} | \mathcal{G}), \\ &= b + E(X | \mathcal{G}) - E(X_{a,b} | \mathcal{G}), & \text{if } E(X | \mathcal{G}) > E(X_{a,b} | \mathcal{G}). \end{aligned}$$

Put $A = [|Y| < M]$ for $0 < M < \infty$. Then $A \varepsilon \mathcal{G}$ and on A , (10) holds; letting $M \rightarrow \infty$, we have (10). The proof of (9) is similar.

COROLLARY 4. *Under the assumptions of Theorem 2,*

$$E((X - E(X | \mathcal{G}))^2 | \mathcal{G}) \geq E((X_{a,b} - E(X_{a,b} | \mathcal{G}))^2 | \mathcal{G}), \quad \text{a.e.}$$

REFERENCES

- [1] BARLOW, R. E., MARSHALL, A. W. and PROSCHAN, F. (1967). Some inequalities for star-shaped and convex functions. Boeing Scientific Research Laboratories. Document D1-82-0643.
- [2] CHOW, Y. S. (1966). On the moments of some one-sided stopping rules. *Ann. Math. Statist.* **37** 382-387.
- [3] KARLIN, S. and NOVIKOFF, A. (1963). Generalized convex inequalities. *Pacific J. Math.* **13** 1251-1279.
- [4] KARLIN, S. and STUDDEN, W. J. (1966). *Tchebycheff Systems: with Applications in Analysis and Statistics*. Interscience, New York.
- [5] ZIEGLER, ZVI (1966). Generalized convexity cones. *Pacific J. Math.* **16** 561-580.