

ON EVALUATING A COEFFICIENT OF PARTIAL CORRELATION

BY GRACE STRECKER

It is to be shown here that when the multiple correlation coefficient $R_{n; 12 \dots (n-1)}$ is found by the method of Horst¹ the partial correlation coefficient $R_{n(n-1); 12 \dots (n-2)}$ can be found in terms of the β 's. If we are interested only in evaluating a partial correlation between two variables, we may also employ the method which will be given here.

Without loss of generality the dependent variables may be chosen to be the n th and $(n - 1)$ st. The coefficient of partial correlation as given by Rietz² may be expressed in the following form:

$$(1) \quad R_{n(n-1); 12 \dots (n-2)} = \sqrt{\frac{\frac{R_{(n-1)(n-1)} - R^2}{R_{(n-1)(n-1)nn} R_{nn}}}{\frac{R_{(n-1)(n-1)}}{R_{(n-1)(n-1)nn}}}}$$

$R_{(n-1)(n-1)}$ may be treated as a new determinant R' . Regarding its elements as the coefficients of a set of normal equations ($n - 1$ in all) whose constant terms are zero, we may follow through the Doolittle elimination process. For the case where $n = 4$ we have the table given below.

In comparing this outline with the one illustrating the Doolittle elimination process for R when $n = 4$ we see that

$$\begin{aligned} \gamma'_{11} &= \gamma_{11} = \frac{A_{11}}{R^2}, \\ \gamma'_{22} &= \gamma_{22} = \frac{rA_{1122}}{R^2 A_{11}}, \\ \gamma'_{33} &= \alpha'_{33} - \sum_2^3 \beta'_{i3} = \alpha_{44} - \sum_2^3 \beta_{i4}. \end{aligned}$$

Therefore, we have

$$\begin{aligned} R' &= \frac{A_{11}}{R^2} \cdot \frac{rA_{1122}}{R^2 A_{11}} \cdot \left(\alpha_{44} - \sum_2^3 \beta_{i4} \right) \\ &= \prod_1^2 \gamma_{ii} \left(\alpha_{44} - \sum_2^3 \beta_{i4} \right). \end{aligned}$$

¹ Horst Paul, *A Short Method for Solving for a Coefficient of Multiple Correlation*, *Annals of Mathematical Statistics*, Vol. III, No. 1, Feb. 1932, pp. 40-44.

² Rietz, H. L., *Mathematical Statistics*, p. 101.



Reciprocal	1	2	3	α	β	γ	δ
$-\frac{R^2}{A_{11}}$	$\frac{A_{11}}{R^2}$ -1	$\frac{A_{12}}{R^2}$ $-\frac{A_{12}}{A_{11}}$	$\frac{A_{14}}{R^2}$ $-\frac{A_{14}}{A_{11}}$	α'_1		γ'_1	δ'_1
		$\frac{A_{22}}{R^2}$ $-\frac{A_{12}^2}{R^2 A_{11}}$	$\frac{A_{24}}{R^2}$ $-\frac{A_{12} A_{14}}{R^2 A_{11}}$	α'_2	β'_{22}		
$-\frac{R^2 A_{11}}{A A_{1122}}$		$\frac{A A_{1122}}{R^2 A_{11}}$ -1	$\frac{A A_{1124}}{R^2 A_{11}}$ $-\frac{A_{1124}}{A_{1122}}$			γ'_2	δ'_2
			$\frac{A_{44}}{R^2}$ $-\frac{A_{14}^2}{R^2 A_{11}}$ $-\frac{A A_{1124}^2}{R^2 A_{11} A_{1122}}$	α'_3	β'_{23} β'_{33}		
			$\alpha'_3 - \sum_2^3 \beta'_{i3}$ -1			γ'_3	δ'_3

In the general case:

$$\gamma'_{11} = \gamma_{11},$$

$$\gamma'_{22} = \gamma_{22},$$

$$\vdots$$

$$\gamma'_{(n-2)(n-2)} = \gamma_{(n-2)(n-2)},$$

$$\gamma'_{(n-1)(n-1)} = \alpha_{nn} - \sum_2^{n-1} \beta_{in}.$$

Hence

$$R_{(n-1)(n-1)} = R' = \prod_1^{n-2} \gamma_{ii} \left(\alpha_{nn} - \sum_2^{n-1} \beta_{in} \right).$$

Since $R = \prod_1^n \gamma_{ii}$, then $R_{(n-1)(n-1)nn} = \prod_1^{n-2} \gamma_{ii}$, from which we see that

$$\frac{R_{(n-1)(n-1)}}{R_{(n-1)(n-1)nn}} = \frac{\prod_1^{n-2} \gamma_{ii} \left(\alpha_{nn} - \sum_2^{n-1} \beta_{in} \right)}{\prod_1^{n-2} \gamma_{ii}} = \alpha_{nn} - \sum_2^{n-1} \beta_{in}.$$

But since $\alpha_{nn} = 1$, then

$$\frac{R_{(n-1)(n-1)}}{R_{(n-1)(n-1)nn}} = 1 - \sum_2^{n-1} \beta_{in}.$$

It has been shown that

$$\frac{R}{R_{nn}} = 1 - \sum_2^n \beta_{in}.$$

Substituting the above values for $\frac{R_{(n-1)(n-1)}}{R_{(n-1)(n-1)nn}}$ and $\frac{R}{R_{nn}}$ in equation (1), we have

$$R_{n(n-1); 12 \dots (n-2)} = \sqrt{\frac{1 - \sum_2^{n-1} \beta_{in} - \left(1 - \sum_2^n \beta_{in} \right)}{1 - \sum_2^{n-1} \beta_{in}}},$$

or

$$R_{n(n-1); 12 \dots (n-2)} = \sqrt{\frac{\beta_{nn}}{1 - \sum_2^{n-1} \beta_{in}}}.$$

Hence it is seen that when the β 's given by Horst (page 42) are calculated, it is an easy matter to solve for the partial correlation $R_{n(n-1); 12 \dots (n-2)}$.

ST. LOUIS UNIVERSITY.