

EXACT CONVERGENCE RATE IN THE LIMIT THEOREMS OF ERDŐS-RÉNYI AND SHEPP

BY PAUL DEHEUVELS, LUC DEVROYE, AND JAMES LYNCH

*Université Paris VI, McGill University, and the
 Pennsylvania State University*

The original Erdős-Rényi theorem states that $U_n/(\alpha k) \rightarrow 1$ almost surely for a large class of distributions, where $U_n = \sup_{0 \leq i \leq n-k} (S_{i+k} - S_i)$, $S_i = X_1 + \dots + X_i$ is a partial sum of i.i.d. random variables, $k = \kappa(n) = [c \log n]$, $c > 0$, and $\alpha > 0$ is a number depending only upon c and the distribution of X_1 . We prove that the lim sup and the lim inf of $(U_n - \alpha k)/\log k$ are almost surely equal to $(2t^*)^{-1}$ and $-(2t^*)^{-1}$, respectively, where t^* is another positive number depending only upon c and the distribution of X_1 . The same limits are obtained for the random variable $T_n = \sup_{1 \leq i \leq n} (S_{i+\kappa(i)} - S_i)$ studied by Shepp.

1. Introduction. We are concerned with the asymptotic behavior of

$$U_n = \sup_{0 \leq i \leq n-k} \{S_{i+k} - S_i\},$$

for $k = [c \log n]$, where $c > 0$, $S_0 = 0$, $S_i = X_1 + \dots + X_i$, and X_1, X_2, \dots are independent, identically distributed random variables having moment generating function $\phi(t)$ and satisfying the conditions

- (A) $E(X_1) = 0$;
- (B) X_1 is nondegenerate, i.e., $P(X_1 = x) < 1$ for all x ;
- (C) $t_0 = \sup\{t; \phi(t) = E(e^{tX_1}) < \infty\} > 0$.

If c is related to α via the equation

$$\exp\left(-\frac{1}{c}\right) = \inf_t \phi(t) e^{-t\alpha},$$

Erdős and Rényi (1970) proved that, for any $\alpha \in \{\phi'(t)/\phi(t), 0 < t < t_0\}$,

$$\lim_{n \rightarrow \infty} \frac{U_n}{\alpha k} = 1 \quad \text{almost surely.}$$

Earlier, Shepp (1964) had obtained a related theorem under the same conditions by showing that

$$\lim_{n \rightarrow \infty} \frac{T_n}{\alpha k} = 1 \quad \text{almost surely,}$$

for

$$T_n = \sup_{1 \leq i \leq n} \{S_{i+\kappa(i)} - S_i\},$$

where $\kappa(i) = [c \log i]$.

Received October 1983; revised March 1985.

AMS 1980 subject classification. Primary 60F15; secondary 60F10.

Key words and phrases. Erdős-Rényi laws, large deviations, moving averages, laws of large numbers, law of the iterated logarithm.

These fundamental results were followed by a flurry of refinements and extensions. Among the refinements, we cite the work of S. Csörgő (1979) and of M. Csörgő and Steinebach (1981), the latter two of whom proved that

$$\frac{U_n}{\alpha k} = 1 + o(k^{-1/2}) \quad \text{almost surely.}$$

In this paper we show that the $o(k^{-1/2})$ term can be replaced by $O(k^{-1} \log k)$ and that this replacement is the best we can achieve. We also prove that the almost sure behavior of T_n is identical to that of U_n .

Before stating our results in detail, we need to specify the range of values of c and α that will be covered by our theorems. This will be done in Section 2. Section 3 presents a large deviation estimate applicable to our problem. Section 4 contains the proof of our main theorems.

2. Properties of the moment generating function. Let $X = X_1$ be a random variable satisfying conditions (ABC) and define

$$m(t) = \frac{\phi'(t)}{\phi(t)} = \frac{E(Xe^{tX})}{E(e^{tX})}.$$

Then $m(0) = 0$ and $m(\cdot)$ is strictly increasing on $[0, t_0)$ and continuously differentiable on $(0, t_0)$. Define further

$$A = \lim_{t \uparrow t_0} m(t) \quad \text{and} \quad c_0 = 1 \int_0^{t_0} tm'(t) dt.$$

Throughout, we will consider only the interval $[0, t_0)$ for t , where $0 < t_0 \leq \infty$. Let $c = c(\alpha)$ and $\rho = \rho(\alpha)$ be defined by

$$\rho = \exp\left(-\frac{1}{c}\right) = \inf_t \phi(t)e^{-t\alpha}.$$

THEOREM 1. (1) For any $t \in (0, t_0)$, $m(t) \in (0, A)$. Conversely, for any $\alpha \in (0, A)$, there exists a unique $t^* = t^*(\alpha) \in (0, t_0)$ such that $m(t^*) = \alpha$;

(2) For any $\alpha \in (0, A)$,

$$\rho = \exp\left(-\frac{1}{c}\right) = \phi(t^*)e^{-t^*\alpha} \quad \text{and} \quad c \in (c_0, \infty);$$

(3) For any $c \in (c_0, \infty)$, there exists a unique $\alpha \in (0, A)$ such that $c = c(\alpha)$.

PROOF. First, on $(0, t_0)$, ϕ is continuously infinitely differentiable. Next, $\phi(0) = 1$ and ϕ is nondecreasing on $[0, t_0)$. This follows from Gurland's inequality [see Gurland (1967)] and the inequalities, for $s \geq t$,

$$E(e^{sX}) = E(e^{tX}(e^{tX})^{(s-t)/t}) \geq E(e^{tX})E(e^{(s-t)X}) \geq E(e^{tX}).$$

Furthermore, ϕ is convex because $\phi''(t) = E(X^2e^{tX}) \geq 0$.

The function m is continuous and strictly increasing on $[0, t_0)$. This follows from the Cauchy-Schwarz inequality $E^2(Xe^{tX}) \leq E(X^2e^{tX})E(e^{tX})$, which im-

plies that $m'(t) = (\phi'(t)\phi(t) - \phi^2(t))/\phi^2(t) \geq 0$. This last inequality has to be strict by (B).

We have $m(0) = \lim_{t \downarrow 0} m(t) = \phi'(0)/\phi(0) = E(X) = 0$. On the other hand, $A = \lim_{t \uparrow t_0} m(t) \leq \text{ess sup } X$. Here, equality occurs when $\text{ess sup } X < \infty$, or when $t_0 = \infty$, $\text{ess sup } X = \infty$, and we have an inequality in the other cases [see Petrov (1965), p. 288].

Consider now the equation $m(t) = \alpha$, and its solution $t^* = t^*(\alpha)$. For all $0 \leq \alpha < A$, there is a unique solution in the range $0 \leq t < t_0$. Conversely, as t takes all values in $[0, t_0)$, $m(t)$ takes all values in $[0, A)$.

Next, $\log \phi(t) - t\alpha$ has first derivative $m(t) - \alpha$ and strictly positive second derivative $m'(t)$ on $[0, t_0)$. Thus it has a unique minimum on $[0, t_0)$ as the solution of the equation $m(t) = \alpha$. This proves (1), and allows us to write

$$\log \rho(\alpha) = \log \phi(t^*) - \alpha t^* = -\frac{1}{c}.$$

Since $m(t^*(\alpha)) = \alpha$, it follows that $t^{*'}(\alpha)m'(t^*(\alpha)) = 1$, and that

$$(\log \rho(\alpha))' = -t^{*'}(\alpha), \quad 0 \leq \alpha < A.$$

Noting that $\rho(0) = \inf_t \phi(t) = \phi(0) = 1$, it follows that

$$\frac{1}{c} = -\int_0^\alpha (\log \rho(\theta))' d\theta = \int_0^\alpha t^{*'}(\theta) d\theta = \int_0^{t^*(\alpha)} tm'(t) dt.$$

Clearly, c is a continuous function of $\alpha \in (0, A)$, strictly increasing in α , with $\lim_{\alpha \downarrow 0} c = \infty$, and $\lim_{\alpha \uparrow A} c = c_0 = 1/\int_0^{t_0} tm'(t) dt$. Thus, for every value $\alpha \in (0, A)$, there exists a unique value $c \in (c_0, \infty)$ and vice versa. This completes the proof of Theorem 1.

REMARK 1. In the sequel we shall make use of the fact that $\sigma^2(t) = m'(t) > 0$ on $(0, t_0)$. The function σ is continuous on $[0, t_0)$, such that

$$\sigma^2(0) = \lim_{t \downarrow 0} \sigma^2(t) = E(X^2) \leq \infty.$$

REMARK 2. We shall need also the fact that $(1/t)\log \phi(t)$ is strictly increasing in t . This last result is a consequence of Jensen's inequality, for $s > t$,

$$E^{1/t}(e^{tX}) \leq E^{1/s}(e^{sX}),$$

with equality if and only X is constant, which is not allowed by (B).

THEOREM 2. $c_0 = 0$ in all cases except the following two:

(i) $A < \infty$, $t_0 < \infty$. (This covers a class of distributions with $\text{ess sup } X_1 = \infty$). In that case, $c_0 = 1/(At_0 - \log \phi(t_0))$;

(ii) $A < \infty$, $t_0 = \infty$. (This occurs if and only if $\text{ess sup } X_1 < \infty$). In that case, we have $A = \text{ess sup } X_1$, $P(X_1 = A) > 0$, and $c_0 = -1/\log P(X_1 = A)$.

PROOF. It is easy to show that if $\text{ess sup } X < \infty$, then $t_0 = \infty$ and $A < \infty$. Conversely, by Petrov (1965), if $t_0 = \infty$ and $\text{ess sup } X = \infty$, then $A = \infty$.

It follows that $\text{ess sup } X < \infty$ if and only if $A < \infty$ and $t_0 = \infty$. In that case, we have $A = \text{ess sup } X$.

Let us now characterize the distributions for which $c_0 = 1/\int_0^A t^*(\alpha) d\alpha = 0$. Since t^* is an increasing function of α , we see that $c_0 = 0$ if $A = \infty$. Let us assume that $A < \infty$. There are then two cases:

(i) $t_0 < \infty$. In that case, $\int_0^A t^*(\alpha) d\alpha \leq At_0 < \infty$ and hence $c_0 \neq 0$. We have here

$$\frac{1}{c_0} = - \lim_{\alpha \uparrow A} \log \rho(\alpha) = \lim_{\alpha \uparrow A} (\alpha t^* - \log \phi(t^*)) = At_0 - \log \phi(t_0).$$

(ii) $t_0 = \infty$. By the remarks above, we must have then $A = \text{ess sup } X < \infty$. In that case, we have $c_0 < \infty$. Furthermore,

$$\begin{aligned} \int_0^A t^*(\alpha) d\alpha &= \int_0^\infty tm'(t) dt \\ &= \int_0^\infty (A - m(t)) dt = \int_0^\infty \frac{E((A - X)e^{tX})}{E(e^{tX})} dt = - \int_0^\infty \frac{E(Ye^{tY})}{E(e^{tY})} dt, \end{aligned}$$

where $Y = X - A$.

Put $\zeta(t) = E(e^{tY})$. We have

$$\begin{aligned} \int_0^A t(\alpha) d\alpha &= - \int_0^\infty \frac{\zeta'(t)}{\zeta(t)} dt = - \lim_{t \uparrow \infty} \log \zeta(t) \\ &= - \log P(Y = 0) = - \log P(X = A). \end{aligned}$$

Here, we have used the fact that by the dominated convergence theorem, $E(e^{tY}) \rightarrow P(Y = 0)$ as $t \uparrow \infty$. This proves Theorem 2.

REMARK 3. A number of authors have apparently ignored the fact that there exist distributions for which $A \neq \text{ess sup } X_1$ and yet fulfill condition (i) of Theorem 2. By taking a density decreasing as $e^{-x}x^{-3}$, we get $A < \text{ess sup } X_1 = \infty$, and $t_0 < \infty$, as sought.

3. Application of Petrov's large-deviation theorem. In this section, we use the hypotheses and notation of Sections 1 and 2.

THEOREM 3 (Petrov, 1965).

$$P(S_n \geq n\alpha) \sim \frac{\psi(t^*)}{\sqrt{n}} \exp\left(-\frac{n}{c}\right) = \frac{\psi(t^*)}{\sqrt{n}} \exp(n(\log \phi(t^*) - t^*\alpha)),$$

uniformly for $\alpha \in [\epsilon, \min(A - \epsilon, 1/\epsilon)]$, where $\epsilon > 0$ is arbitrary, and $\psi(t^*) > 0$ is a finite number depending upon t^* and the distribution of X_1 only.

For nonlattice distributions, one can take $\psi(t^*) = (t^*\sigma(t^*)\sqrt{2\pi})^{-1}$, while for lattice distributions with span H , one can take

$$\psi(t^*) = \frac{H}{1 - e^{-Ht^*}} \cdot \frac{1}{\sigma(t^*)\sqrt{2\pi}}.$$

REMARK 4. Cramér (1938) proved a similar result for more restricted classes of random variables and Bahadur and Ranga Rao (1960) obtained another result that comes close to Theorem 3 [see Nagaev (1979) for a general discussion of large deviation results].

We will repeatedly use the following corollaries of Theorem 3.

COROLLARY 1. *Let $\alpha \in (0, A)$ and let y_n be a sequence of numbers satisfying $ny_n^2 \rightarrow 0$ as $n \rightarrow \infty$. Then, uniformly over all sequences z_n with $|z_n| \leq |y_n|$, we have*

$$P(S_n \geq n(\alpha + z_n)) \sim \frac{\psi(t^*)}{\sqrt{n}} \exp\left(-\frac{n}{c}\right) \exp(-nz_n t^*).$$

PROOF. The proof is based upon Theorem 3 jointly with the following observations taken from Section 2: $t^* = t^*(\alpha)$ is a continuous function of α , and thus $\psi(t^*)$ is a continuous function of α too. The derivative of $-(1/c) = \log \rho(\alpha)$ with respect to α is $-t^*$.

COROLLARY 2. *For all $\varepsilon \in \mathbb{R}$, $\alpha \in (0, A)$, we have*

$$\frac{\psi(t^*) + o(1)}{k^{1/2+(\pm 1/2+\varepsilon)}} \leq nP\left(S_k \geq \alpha k + \left(\pm \frac{1}{2} + \varepsilon\right) \frac{\log k}{t^*}\right) \leq \frac{e^{1/c}\psi(t^*) + o(1)}{k^{1/2+(\pm 1/2+\varepsilon)}}.$$

PROOF. It follows directly from Corollary 1 and the observation that $e^{(k+1)/c} \geq n \geq e^{k/c}$.

4. The main theorems. In the remainder of this section, we will need the following increasing sequence of integers:

$$n_j = \inf\{n; [c \log n] = j\}.$$

It is clear that $k = \kappa(n) = j$ for $n_j \leq n < n_{j+1}$.

LEMMA 1.

- (i) $\limsup_{n \rightarrow \infty} (U_n - \alpha k) / \log k \leq \frac{1}{2t^*}$ almost surely;
- (ii) $\limsup_{n \rightarrow \infty} (T_n - \alpha k) / \log k \leq \frac{1}{2t^*}$ almost surely.

PROOF. (i) For $n_j \leq n < n_{j+1}$, we know that $k = j$, and $U_n \leq U_{n_{j+1}-1}$. Thus, for $\varepsilon > 0$,

$$P\left(U_n \geq \alpha k + \left(\frac{1}{2} + \varepsilon\right) \frac{\log k}{t^*} \text{ i.o.}\right) \leq P\left(U_{n_{j+1}-1} \geq \alpha j + \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*} \text{ i.o. (in } j)\right).$$

By Corollary 2, since $j = [c \log(n_{j+1} - 1)]$,

$$P\left(U_{n_{j+1}-1} \geq \alpha j + \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) \leq n_{j+1} P\left(S_j \geq \alpha j + \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) = O(j^{-1-\varepsilon}),$$

which is summable in j . The result follows by Borel–Cantelli.

(ii) Likewise, for $n_j \leq n < n_{j+1}$, we have $T_n \leq T_{n_{j+1}-1} = \sup_{0 \leq i \leq j} \Lambda_i$, where $\Lambda_i = \sup_{n_i \leq l < n_{i+1}} \{S_{l+\kappa(l)} - S_l\}$. It follows that, for any $\varepsilon > 0$,

$$P\left(\Lambda_j \geq \alpha j + \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*} \text{ i.o.}\right) = 0 \Rightarrow P\left(T_n \geq \alpha k + \left(\frac{1}{2} + \varepsilon\right) \frac{\log k}{t^*} \text{ i.o.}\right) = 0.$$

By Corollary 2,

$$\begin{aligned} P\left(\Lambda_j \geq \alpha j + \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) &\leq (n_{j+1} - n_j)P\left(S_j \geq \alpha j + \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) \\ &= O(j^{-1-\varepsilon}), \end{aligned}$$

which is summable in j . The result follows by Borel–Cantelli.

LEMMA 2.

- (i) For any $\varepsilon > 0$, $P\left((U_n - \alpha k)/\log k \leq -\frac{1}{2t^*} + \varepsilon\right) \rightarrow 1$;
- (ii) For any $\varepsilon > 0$, $P\left((T_n - \alpha k)/\log k \leq -\frac{1}{2t^*} + \varepsilon\right) \rightarrow 1$.

PROOF. (i) By Corollary 2, we have, for any $\varepsilon > 0$,

$$\begin{aligned} P\left((U_n - \alpha k)/\log k \geq \frac{-1}{2t^*} + \varepsilon\right) \\ \leq nP\left(S_k \geq \alpha k + \left(-\frac{1}{2} + \varepsilon t^*\right) \frac{\log k}{t^*}\right) \rightarrow 0, \text{ hence result.} \end{aligned}$$

(ii) For $n_j \leq n < n_{j+1}$, we have, using ε/t^* in place of ε ,

$$P\left((T_n - \alpha k)/\log k \geq \frac{-1}{2t^*} + \frac{\varepsilon}{t^*}\right) \leq P\left((T_{n_{j+1}-1} - \alpha j)/\log j \geq \frac{-1}{2t^*} + \frac{\varepsilon}{t^*}\right).$$

Proceeding as in the proof of Lemma 1, let $\Theta_j = T_{n_{j+1}-1} = \sup_{0 \leq i \leq j} \Lambda_i$. By Lemma 1(ii), for any $\varepsilon > 0$, $P(\Theta_j \geq \alpha j + (1/2 + \varepsilon)(\log j)/t^*) \rightarrow 0$. It follows by change of index from j to $j - [j^{\varepsilon/2}]$ that

$$P\left(\Theta_{j-[j^{\varepsilon/2}]} \geq \alpha j + \left(-\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) \rightarrow 0.$$

Finally by Corollary 2, we have

$$\begin{aligned} P\left(\sup_{j-[j^{\varepsilon/2}] < i \leq j} \Lambda_i \geq \alpha j + \left(-\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) \\ \leq \sum_{j-[j^{\varepsilon/2}] < i \leq j} P\left(\Lambda_i \geq \alpha i + \left(-\frac{1}{2} + \varepsilon\right) \frac{\log i}{t^*}\right) = O(j^{-\varepsilon/2}) \rightarrow 0. \end{aligned}$$

This suffices for the proof.

COROLLARY 3.

- (i) $\liminf_{n \rightarrow \infty} (U_n - \alpha k) / \log k \leq -\frac{1}{2t^*}$ almost surely;
- (ii) $\liminf_{n \rightarrow \infty} (T_n - \alpha k) / \log k \leq -\frac{1}{2t^*}$ almost surely.

PROOF. This statement uses the observations that for any sequence of events A_n with $P(A_n) \rightarrow 1$, we must have $P(A_n \text{ i.o.}) = 1$. It follows directly from Lemma 2.

Having established the easy halves of our main results, we now turn to the more complicated parts. We shall make use of the following lemmas.

LEMMA 3 (Chung and Erdős, 1952). For arbitrary sequences of events A_1, \dots, A_n , we have

$$P\left(\bigcup_{i=1}^n A_i\right) \geq \frac{\left(\sum_{i=1}^n P(A_i)\right)^2}{\sum_{i=1}^n P(A_i) + \sum_{i \neq j} P(A_i \cap A_j)}.$$

LEMMA 4. Let $1 \leq i \leq k$ and let $S_i = X_1 + \dots + X_i$, $S'_{k-i} = X_{i+1} + \dots + X_k$, and $S''_i = X_{k+1} + \dots + X_{k+i}$. Then for any x and q and for any $t \in (0, t^*)$, we have

$$P(S_i + S'_{k-i} \geq x, S'_{k-i} + S''_i \geq x) \leq (\phi(t^*))^{k-i} e^{-t^*q} + P(S_k \geq x)(\phi(t))^i e^{-t(x-q)}.$$

PROOF OF LEMMA 4. Note first that $P(S_n \geq \alpha n + u) \leq E(e^{t(S_n - n\alpha - u)}) = (\phi(t)e^{-t\alpha})^n e^{-tu}$. From there we get, for any $0 < t \leq t^*$, by Jensen's inequality,

$$P(S_n \geq s) \leq (\phi(t))^n e^{-ts} \quad \text{and} \quad P(S_n \geq s) \leq (\phi(t^*))^n e^{-t^*s}.$$

Next, we have

$$\begin{aligned} &P(S_i + S'_{k-i} \geq x, S'_{k-i} + S''_i \geq x) \\ &\leq P(S'_{k-i} \geq q) + P(S''_i \geq x - q)P(S_i + S'_{k-i} \geq x) \\ &\leq (\phi(t^*))^{k-i} e^{-t^*q} + (\phi(t))^i e^{-t(x-q)}P(S_k \geq x) \end{aligned}$$

as sought.

LEMMA 5. For any $\epsilon > 0$, we have with $\epsilon' = \epsilon/t^*$:

- (i) $P\left((U_n - \alpha k) / \log k \geq \frac{-1}{2t^*} - \epsilon'\right) \rightarrow 1$;
- (ii) $P\left((T_n - \alpha k) / \log k \geq \frac{-1}{2t^*} - \epsilon'\right) \rightarrow 1$.

PROOF. (i) We will use Lemma 3 for events

$$A_i = \left\{ S_{i+k} - S_i \geq \alpha k + \left(-\frac{1}{2} - \varepsilon \right) \frac{\log k}{t^*} \right\}, \quad 0 \leq i \leq n - k.$$

Noting that A_i and A_j are independent when $|i - j| > k$ and that $P(A_i) = P(A_0)$, we have

$$\begin{aligned} & \sum_{i=0}^{n-k} P(A_i) + \sum_{i \neq j} P(A_i \cap A_j) \\ &= \left(\sum_{i=0}^{n-k} P(A_i)^2 \right) + \sum_{1 \leq |i-j| \leq k} (P(A_i \cap A_j) - P(A_i)P(A_j)) \\ & \quad + \sum_{i=0}^{n-k} (P(A_i) - P^2(A_i)). \end{aligned}$$

It follows that we need only show that, for $\varepsilon > 0$,

$$(*) \quad nP(A_0) \rightarrow \infty,$$

and

$$(**) \quad n \sum_{i=1}^k P(A_0 \cap A_i) = o\left((nP(A_0))^2\right).$$

By Corollary 2, $(\psi(t^*) + o(1))k^\varepsilon \leq nP(A_0) \leq (e^{1/c}\psi(t^*) + o(1))k^\varepsilon$. Hence $(*)$ is satisfied and $(**)$ amounts to

$$(**) \quad n \sum_{i=1}^k P(A_0 \cap A_i) = o(k^{2\varepsilon}).$$

Put in the inequality of Lemma 4, $x = \alpha k + (-1/2 - \varepsilon)(\log k)/t^*$, and $q = \alpha k - (i/t^*)\log \phi(t^*) + (2/t^*)\log i$. Let $t \in (0, t^*)$ be fixed. We have by Lemma 4 that

$$\begin{aligned} P(A_0 \cap A_i) &= P(S_i + S'_{k-i} \geq x, S'_{k-i} + S''_i \geq x) \\ &\leq (\phi(t^*))^{k-i} e^{-t^*q} + P(A_0)(\phi(t))^i e^{-t(x-q)} \\ &= i^{-2} e^{-k/c} + P(A_0) k^{(t/t^*)(\varepsilon+1/2)} i^{2t/t^*} e^{-\theta i} \\ &\leq i^{-2} e^{-k/c} + P(A_0) k^{\varepsilon+5/2} e^{-\theta i}, \end{aligned}$$

where $\theta = t((1/t^*)\log \phi(t^*) - (1/t)\log \phi(t)) > 0$ by Remark 2.

Let $l = [k^{\varepsilon/2}]$. We have

$$\begin{aligned} n \sum_{i=1}^k P(A_0 \cap A_i) &\leq 2nP(A_0) + n \sum_{i=l+1}^{k-l} P(A_0 \cap A_i) \\ &\leq 2nP(A_0) + e^{1/c} \sum_{i=l+1}^{\infty} i^{-2} + nP(A_0) k^{\varepsilon+5/2} \frac{e^{-\theta l}}{1 - e^{-\theta}} \\ &= k^{\varepsilon/2} nP(A_0) O(1) = O(k^{3\varepsilon/2}), \end{aligned}$$

which suffices for $(**)$, since $nP(A_0) = O(k^\varepsilon)$.

(ii) Let again $\Lambda_j = \sup_{n_j \leq i < n_{j+1}} \{S_{i+j} - S_i\}$, and note for further use that $T_n \geq \Lambda_j$ for $n \geq n_{j+1}$. Let $N = n_{j+1} - n_j \leq e^{(j+2)/c}$, $N \geq e^{j/c}(e^{1/c} - 1) - 1 \sim e^{j/c}(e^{1/c} - 1)$. We obtain by the same arguments as above, with j replacing k in the definition of A_i ,

$$P\left(\Lambda_j \geq \alpha j + \left(-\frac{1}{2} - \varepsilon\right) \frac{\log j}{t^*}\right) = P\left(\bigcup_{i=1}^N A_i\right) \rightarrow 1,$$

whenever

$$(*) \quad NP(A_0) \rightarrow \infty,$$

and

$$(**) \quad N \sum_{i=1}^j P(A_0 \cap A_i) = o\left((NP(A_0))^2\right).$$

We have just proved that $(*)$ and $(**)$ hold if $\varepsilon > 0$. This implies that, as $j \rightarrow \infty$

$$P\left(T_n \geq \alpha j + \left(-\frac{1}{2} - \varepsilon\right) \frac{\log j}{t^*}\right) \geq P\left(\Lambda_j \geq \alpha j + \left(-\frac{1}{2} - \varepsilon\right) \frac{\log j}{t^*}\right) \rightarrow 1,$$

uniformly in $n_{j+1} \leq n < n_{j+2}$. This, in turn, implies that

$$P\left(T_n \geq \alpha k + \left(-\frac{1}{2} - 2\varepsilon\right) \frac{\log k}{t^*}\right) \rightarrow 1,$$

which completes the proof of Lemma 5, $\varepsilon > 0$ being arbitrary.

The first of the two main theorems of this paper follows:

THEOREM 4. *For any $\alpha \in (0, A)$ or, equivalently, for any $c = c(\alpha) \in (c_0, \infty)$, we have*

- (i) $(U_n - \alpha k) / \log k \rightarrow -\frac{1}{2t^*}$ in probability;
- (ii) $(T_n - \alpha k) / \log k \rightarrow -\frac{1}{2t^*}$ in probability.

PROOF. Combine Lemmas 2 and 5.

We proceed with the sequence of lemmas directed toward the second of our main theorems.

LEMMA 6.

- (i) $\limsup_{n \rightarrow \infty} (U_n - \alpha k) / \log k \geq \frac{1}{2t^*}$ almost surely;
- (ii) $\limsup_{n \rightarrow \infty} (T_n - \alpha k) / \log k \geq \frac{1}{2t^*}$ almost surely.

PROOF. Let $R_j = \sup_{n_j \leq n < n_{j+1} - j} \{S_{n+j} - S_n\}$. Since $k = \kappa(n) = j$ when $n_j \leq n < n_{j+1}$, it is straightforward that $R_j \leq \min\{T_m, U_m\}$, for $n_j \leq m < n_{j+1}$. It follows that

$$P\left(T_m \geq \alpha k + \left(\frac{1}{2} - \varepsilon\right) \frac{\log k}{t^*} \text{ i.o.}\right) = P\left(U_m \geq \alpha k + \left(\frac{1}{2} - \varepsilon\right) \frac{\log k}{t^*} \text{ i.o.}\right) = 1$$

for any $\varepsilon > 0$, if

$$P\left(R_j \geq \alpha j + \left(\frac{1}{2} - \varepsilon\right) \frac{\log j}{t^*} \text{ i.o.}\right) = 1$$

for any $\varepsilon > 0$. Since the R_j 's form a sequence of independent random variables, the latter probability is one if and only if

$$\sum_j P\left(R_j \geq \alpha j + \left(\frac{1}{2} - \varepsilon\right) \frac{\log j}{t^*}\right) = \sum_j P_j = \infty.$$

Next, we note that $P_j = P(\bigcup_{i=1}^N A_i)$, where $A_i = \{S_{i+j} - S_i \geq x\}$, $x = \alpha j + (1/2 - \varepsilon)(\log j)/t^*$, and $N = n_{j+1} - n_j - j \leq e^{(j+2)/c}$, $N \geq e^{j/c}(e^{1/c} - 1) - j - 2 \sim e^{j/c}(e^{1/c} - 1)$.

By Lemma 3, it follows that

$$P_j = P\left(\bigcup_{i=1}^N A_i\right) \geq \frac{(NP(A_0))^2}{NP(A_0) + (NP(A_0))^2 + 2N \sum_{l=1}^j P(A_0 \cap A_l)}.$$

By Corollary 2 and our bounds on N , we note that, for some appropriate constants $c_1 > 0$ and $c_2 > 0$, we have

$$(c_1 + o(1))j^{-1+\varepsilon} \leq NP(A_0) \leq (c_2 + o(1))j^{-1+\varepsilon}.$$

Summarizing and simplifying, we obtain

$$P_j \geq (1 + o(1)) \frac{c_1^2 j^{-2(1-\varepsilon)}}{c_2 j^{-(1-\varepsilon)} + 2N \sum_{l=1}^j P(A_0 \cap A_l)}.$$

We will show further on that there exists a constant $c_3 > 0$ such that

$$N \sum_{l=\lceil j^{\varepsilon/2} \rceil}^j P(A_0 \cap A_l) \leq (c_3 + o(1))j^{-1}.$$

But this is all we need, because

$$\begin{aligned} N \sum_{l=1}^j P(A_0 \cap A_l) &\leq (c_3 + o(1))j^{-1} + NP(A_0)j^{\varepsilon/2} \\ &\leq (c_3 + o(1))j^{-1} + (c_2 + o(1))j^{-1+(3\varepsilon/2)} \\ &\sim c_2 j^{-1+3\varepsilon/2}. \end{aligned}$$

From there, we conclude that

$$P_j = P\left(\bigcup_{i=1}^N A_i\right) \geq (1 + o(1)) \frac{c_1^2}{2c_2} j^{-1+(\varepsilon/2)},$$

which is not summable in j and we are done.

To bound $P(A_0 \cup A_l)$, we use Lemma 4 with $x = \alpha j + (1/2 - \varepsilon)(\log j)/t^*$, $k = j$, $i = l$, $q = \alpha j - (l/t^*)\log \phi(t^*) + (1 + 2/\varepsilon)(\log l)/t^*$, the constants t , t^* , and θ are as in the proof of Lemma 5. This gives

$$P(A_0 \cap A_l) \leq e^{-j/c} l^{-(1+(2/\varepsilon))} + P(A_0) e^{-\theta l j^{(t/t^*)(\varepsilon-1/2)}} l^{(t/t^*)(1+2/\varepsilon)}.$$

If we sum over all $l \geq [j^{\varepsilon/2}]$ and multiply by $N \leq e^{(j+2)/c}$, we see that

$$\begin{aligned} N \sum_{l=[j^{\varepsilon/2}]}^j P(A_0 \cap A_l) &\leq e^{2/c} \left\{ ([j^{\varepsilon/2}]^{2/\varepsilon})^{-1} \right. \\ &\quad \left. + (c_2 + o(1)) j^{-1+\varepsilon} \frac{\exp(-\theta j^{\varepsilon/2})}{1 - e^{-\theta}} j^{(t/t^*)(1/2+\varepsilon+2/\varepsilon)} \right\} \\ &\leq (c_3 + o(1)) j^{-1}, \end{aligned}$$

for some positive constant c_3 .

This concludes the proof of Lemma 6.

In view of Lemma 1, Corollary 3, and Lemma 6, only one piece of the puzzle is missing, i.e.,

LEMMA 7.

- (i) $\liminf_{n \rightarrow \infty} (U_n - \alpha k) / \log k \geq \frac{-1}{2t^*}$ almost surely;
- (ii) $\liminf_{n \rightarrow \infty} (T_n - \alpha k) / \log k \geq \frac{-1}{2t^*}$ almost surely.

PROOF. For $n_j \leq n < n_{j+1}$, we know that $U_n \geq U_{n_j}$ and $T_n \geq T_{n_j}$. Thus, by the Borel-Cantelli lemma, we are done if we can show that, for all $\varepsilon > 0$,

$$\begin{aligned} \sum_{j=1}^{\infty} P\left(U_{n_j} < \alpha j - \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) &< \infty \\ \text{and } \sum_{j=1}^{\infty} P\left(T_{n_j} < \alpha j - \left(\frac{1}{2} + \varepsilon\right) \frac{\log j}{t^*}\right) &< \infty. \end{aligned}$$

Consider the set J_j of all integers of the form $r[j^{\varepsilon/2}]$, $r = 0, 1, 2, \dots$. For integer l , let us also define the quantity, for each fixed j ,

$$Q_l = \sup_{2lj \leq i < (2l+1)j; i \in J_j} \{S_{i+j} - S_i\}.$$

It is noteworthy that Q_0, Q_1, Q_2, \dots are independent random variables for each j and that

$$U_{n_j} \geq \sup_{0 \leq l \leq L} Q_l,$$

where L is the largest integer such that $(2L + 1)j - 1 \leq n_j - j$, i.e.,

$$L = \left\lceil \frac{1}{2} \left(\frac{n_j - j + 1}{j} - 1 \right) \right\rceil \geq \frac{n_j - 2j}{2j} - 1 = \frac{n_j - 4j}{2j} \sim \frac{n_j}{2j}.$$

By all this, we have

$$\begin{aligned} P'_j &= P\left(U_{n_j} < \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*} \right) \leq \prod_{l=0}^L P\left(Q_l < \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*} \right) \\ &\leq \exp\left(- (1 + o(1)) \frac{n_j}{2j} P\left(Q_0 \geq \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*} \right) \right), \end{aligned}$$

where we used the independence of the Q_l 's and the inequality $1 - u \leq e^{-u}$. Let N_l be the number of indices i in the intersection $J_j \cap \{2lj, \dots, (2l + 1)j - 1\}$. This number N_l satisfies, uniformly in l , $N_l \sim j^{1-\varepsilon/2}$ as $j \rightarrow \infty$ (and this is why we need only consider Q_0). Put, in the sequel, $N = N_0$.

By a simple Bonferroni inequality

$$\begin{aligned} P\left(Q_0 \geq \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*} \right) &\geq NP\left(S_j \geq \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*} \right) \\ &\quad - 2N \sum_{r=1}^{N-1} P\left(S_j \geq \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*}, \right. \\ &\quad \left. S_{j+r\lceil j^{\varepsilon/2} \rceil} - S_{r\lceil j^{\varepsilon/2} \rceil} \geq \alpha j - \left(\frac{1}{2} + \varepsilon \right) \frac{\log j}{t^*} \right). \end{aligned}$$

By Corollary 2, the first term in this lower bound is larger than

$$(1 + o(1))j^{1-\varepsilon/2}e^{-j/c}\psi(t^*)j^\varepsilon \sim \psi(t^*)j^{1+\varepsilon/2}e^{-j/c}.$$

We will show that the second term in the lower bound is $o(\text{first term})$, so that P'_j is not greater than

$$\exp\left(- (1 + o(1)) \frac{n_j}{2} j^{\varepsilon/2} \psi(t^*) e^{-j/c} \right) \leq \exp\left(- (1 + o(1)) \frac{1}{2} \psi(t^*) j^{\varepsilon/2} \right),$$

which is summable in j by the integral test. This proves (i).

Let $x = \alpha j - (1/2 + \varepsilon)(\log j)/t^*$ and $m = r\lceil j^{\varepsilon/2} \rceil$. Then we need only show that

$$(*) \quad \sum_{r=1}^{N-1} P(S_j \geq x, S_{j+m} - S_m \geq x) = O(j^\varepsilon e^{-j/c}).$$

To do so, we will once again use Lemma 4, with t, t^* , and θ defined as in the proof of Lemma 5, and with the formal replacements

$$k = j, \quad i = m = r\lceil j^{\varepsilon/2} \rceil, \quad q = \alpha j - r\lceil j^{\varepsilon/2} \rceil \frac{1}{t^*} \log \phi(t^*) + \frac{2}{t^*} \log(r\lceil j^{\varepsilon/2} \rceil).$$

The r th term of (*) is bounded from above by

$$e^{-j/c} (r [j^{\epsilon/2}])^{-2} + P(S_j \geq x) j^{(t/t^*)(1/2+\epsilon)} (r [j^{\epsilon/2}])^{(2t/t^*)} e^{-\theta r [j^{\epsilon/2}]},$$

which, taking into account that $r [j^{\epsilon/2}] \leq j$, $\epsilon < \frac{1}{2}$ (without loss of generality) and $t/t^* < 1$, is by Corollary 2 not greater than

$$e^{-j/c} r^{-2} j^{-\epsilon} (1 + o(1)) + (e^{1/c} \psi(t^*) + o(1)) e^{-j/c} j^{3+\epsilon} e^{-\theta r [j^{\epsilon/2}]},$$

where the “ $o(1)$ ” terms are uniform in $r \geq 1$.

Summing over r gives the bound

$$(**) (1 + o(1)) e^{-j/c} \left(\frac{\pi^2}{6j^\epsilon} + \frac{e^{1/c} \psi^\epsilon(t^*) j^{3+\epsilon} e^{-\theta N [j^{\epsilon/2}]}}{1 - \exp(-\theta [j^{\epsilon/2}])} \right) = o(e^{-j/c}) = o(j^\epsilon e^{-j/c}),$$

as requested. This completes the proof of (i).

The proof of (ii) is based on the same arguments as the proof of (i), but with slight modifications. We first replace Q_l by Q'_l , defined by

$$Q'_l = \sup_{2lj \leq i < (2l+1)j; i \in J} \{S_{i+\kappa(i)} - S_i\}.$$

We have evidently, for any $0 \leq M \leq L$,

$$T_{n_j} \geq \sup_{M \leq l \leq L} Q'_l.$$

Next, we choose $M = o(L)$ such that, for any $i \in \{2Mj, \dots, (2L + 1)j\}$, we have

$$j - \frac{\log j}{t^*} o(1) \leq \kappa(i) \leq j,$$

where the “ $o(1)$ ” term is uniform in i as $j \rightarrow \infty$. We can take here $M = [L/\log \log j]$. We get then, by the same arguments as above

$$P_j'' = P\left(T_{n_j} < \alpha j - \left(\frac{1}{2} + \epsilon\right) \frac{\log j}{t^*}\right) \leq \exp\left(-\sum_{l=M}^L P\left(Q'_l \geq \alpha j - \left(\frac{1}{2} + \epsilon\right) \frac{\log j}{t^*}\right)\right).$$

By Bonferroni, we have

$$P\left(Q'_l \geq \alpha j - \left(\frac{1}{2} + \epsilon\right) \frac{\log j}{t^*}\right) \geq \sum_{i \in I_l} P(S_{\kappa(i)} \geq x) - \sum_{r \neq s \in I_l} P(S_{r+\kappa(r)} - S_r \geq x, S_{s+\kappa(s)} - S_s \geq x),$$

where $I_l = J_j \cap \{2lj, \dots, (2l + 1)j - 1\}$.

By Corollaries 1 and 2 we have evidently, for j large enough,

$$P(S_{\kappa(i)} \geq x) \geq (\psi(t^*) + o(1)) e^{-j/c} j^{3\epsilon/4},$$

uniformly in $i \in I_l$, so that

$$P(S_{\kappa(i)} \geq x) \geq \frac{e^{1/c} \psi(t^*) + o(1)}{e^{j/c} j^{-\epsilon/2}} \geq (1 + o(1)) P\left(S_j \geq \alpha j - \left(\frac{1}{2} + \frac{\epsilon}{2}\right) \frac{\log j}{t^*}\right).$$

Likewise, by Lemma 4 and the argument we have just used to prove (i), we deduce from (* *) that

$$\sum_{r \neq s \in I_l} P(S_{r+\kappa(r)} - S_r \geq x, S_{s+\kappa(s)} - S_s \geq x) = o\left(\sum_{i \in I_l} P(S_{\kappa(i)} \geq x)\right).$$

It follows that

$$P'_j \leq \exp\left(- (L - M)(1 + o(1))P\left(Q_0 \geq \alpha j - \left(\frac{1}{2} + \frac{\varepsilon}{2}\right)\frac{\log j}{t^*}\right)\right).$$

Since $L - M = L(1 + o(1)) = (n_j/(2j))(1 + o(1))$, the expression above yields the same upper bound as was obtained for P'_j with $\varepsilon/2$ replacing ε . But we have proved that $\sum_j P'_j < \infty$ for all $\varepsilon > 0$. This concludes the proof of Lemma 7.

The theorem that gives the exact convergence rate for the Erdős–Rényi and Shepp limit theorems has now been proved:

THEOREM 5. *For any $\alpha \in (0, A)$, or equivalently, for any $c = c(\alpha) \in (c_0, \infty)$, we have*

- (i) $\limsup_{n \rightarrow \infty} (U_n - \alpha k)/\log k = \frac{1}{2t^*}$ almost surely;
- (ii) $\liminf_{n \rightarrow \infty} (U_n - \alpha k)/\log k = -\frac{1}{2t^*}$ almost surely.

In statements (i)–(ii), U_n can be replaced by T_n .

PROOF. Combine Corollary 3 with Lemmas 1, 6, and 7.

REMARK 5. It can be seen that the methods we have used can be extended to the case where $k = \kappa(n)$ is a nondecreasing sequence such that $\kappa(n) - c \log n = o(\log \log n)$.

REMARK 6. What happens when $c \in (0, c_0)$, corresponding to cases (i)–(ii) of Theorem 2, will be discussed in forthcoming papers (Deheuvels, 1985 and Deheuvels–Devroye, 1983 and 1985). Related results concerning Erdős–Rényi laws for moving quantiles are to be found in Deheuvels and Steinebach (1984).

Acknowledgment. We are grateful to the referee for his careful reading and helpful suggestions.

REFERENCES

- BAHADUR, R. R. and RANGA RAO, R. (1960). On deviations of the sample mean. *Ann. Math. Statist.* **31** 1015–1027.
- CHUNG, K. L. and ERDŐS, P. (1952). On the application of the Borel–Cantelli lemma. *Trans. Amer. Math. Soc.* **72** 179–186.
- CRAMER, H. (1938). Sur un nouveau théorème limite de la théorie des probabilités. *Actualités Scientifiques et Industrielles* **736** 5–23, Hermann, Paris.

- CSÖRGŐ, M. and STEINEBACH, J. (1981). Improved Erdős-Rényi and strong approximation laws for increments of partial sums. *Ann. Probab.* **9** 988-996.
- CSÖRGŐ, S. (1979). Erdős-Rényi laws. *Ann. Statist.* **7** 772-787.
- DEHEUVELS, P. (1985). On the Erdős-Rényi theorem for random fields and sequences and its relationships with the theory of runs and spacings. *Z. Wahrsch. verw. Gebiete* **70**, 91-115.
- DEHEUVELS, P. and DEVROYE, L. 1983. Limit laws related to the Erdős-Rényi theorem. Technical Report 83-6, LSTA, Université Paris VI.
- DEHEUVELS, P. and DEVROYE, L. (1985). Limit laws of the Erdős-Rényi type. Submitted to *Ann. Probab.*
- DEHEUVELS, P. and STEINEBACH, J. (1984). On an improved Erdős-Rényi strong law for moving quantiles. Technical Report 84-22, LSTA, Université Paris VI.
- ERDŐS, P. and RÉNYI, A. (1970). On a new law of large numbers. *J. Analyse Math.* **23** 103-111.
- GURLAND, J. (1967). An inequality satisfied by the expectation of the reciprocal of a random variable. *Amer. Statist.* **21** 24-25.
- NAGAEV, S. V. (1979). Large deviations of sums of independent random variables. *Ann. Probab.* **7** 745-789.
- PETROV, V. V. (1965). On the probabilities of large deviations of sums of independent random variables. *Theory Probab. App.* **10** 287-298.
- SHEPP, L. A. (1964). A limit theorem concerning moving averages. *Ann Math. Statist.* **35** 424-428.

PAUL DEHEUVELS
 INSTITUT DE STATISTIQUE
 LSTA UNIVERSITÉ PARIS VI
 4 PLACE JUSSIEU
 75230 PARIS CEDEX 05
 FRANCE

LUC DEVROYE
 SCHOOL OF COMPUTER SCIENCE
 MCGILL UNIVERSITY
 805 SHERBROOKE STREET WEST
 MONTREAL, PQ
 CANADA H3A 2K6

JAMES LYNCH
 DEPARTMENT OF STATISTICS
 THE PENNSYLVANIA STATE UNIVERSITY
 UNIVERSITY PARK
 PENNSYLVANIA 16802