

AN OCCUPANCY PROBLEM

K. S. KHATTAB AND A. M. KHIDR

ABSTRACT. Under a certain restriction, n numbered balls are distributed into m urns. A match occurs if the ball numbered i lies in the urn with the number i . The number of matches and its asymptotic probability distribution are found.

1. Introduction. Let the n numbered balls be distributed into m numbered equal urns under the restriction (R); if the ball numbered i lies in the urn numbered j , then no ball with a number higher than i is permitted to lie in an urn with a smaller number than j , $1 \leq i \leq n$, $1 \leq j \leq m$.

It is clear that the number of distributions $A(n, m)$ of n balls into the m urns under the restriction (R) is equal to the number of increasing words formed of n letters out of m distinct letters and thus (see [3, p. 10]), $A(n, m) = \binom{n+m-1}{n}$. We say a match has occurred in the j th urn if the ball numbered j lies in the urn with number j , $1 \leq j \leq \min(n, m)$. Let $W(n, m, r)$ be the number of distributions in which there are exactly r matches $0 \leq r \leq \min(n, m)$. We will prove the following theorems.

THEOREM 1.

- (a) For $n \neq m$, $W(n, m, r) = W(n-1, m, r) + W(n, m-1, r)$,
- (b) For $n = m$, $W(n, n, r) = W(n-1, n, r-1) + W(n, n-1, r)$,
- (c) For $r \neq 0$, $W(n, m, r) = W(m, n, r)$

It is found that the value of $W(n, m, r)$ which satisfies Theorem 1 is given by Theorem 2,

THEOREM 2.

$$W(n, m, r) = \begin{cases} \binom{n+m-1}{m-r} \frac{n-m+2r}{n+r} & m \leq n, \\ \binom{n+m-1}{n-r} \frac{m-n+2r}{m+r} & m > n. \end{cases}$$

As an application in probability theory we obtain the following result.

Received by the editors on March 30, 1983 and in revised form on November 2, 1983.

THEOREM 3. Let $W = W(n, m, r) / \binom{n+m-1}{n}$ and $Z = W / \sqrt{n}$, then $\lim P(Z > z) = e^{-\alpha z - z^2}$, where the limit on n and m are such that $\alpha = \lim_{n, m \rightarrow \infty} (n - m) / \sqrt{n}$.

Similar problems were solved in [1] and [2] leading to theorems corresponding to theorems 1, 2, 3.

2. Proof of Theorem 1. (a) If the n th ball is placed into the m th urn then we have to distribute the $n-1$ remaining balls into the m urns under the restriction (R) to give r matches, the number of these possibilities equals $W(n - 1, m, r)$.

If on the other hand the n th ball is not in the m th urn, then the n balls are to be distributed under (R), into the $m - 1$ urns to get r matches, the number of these possibilities equals $W(n, m - 1, r)$, hence.

$$(2.1) \quad W(n, m, r) = W(n - 1, m, r) + W(n, m - 1, r).$$

(b) The first summand is as in the last summand in part (a), the second summand has a different shape, since if the $(n - 1)$ -st ball is in the n th urn this gives a match; therefore the $n - 1$ remaining balls have to be distributed into the n urns (including the n th urn) under (R) to get $r - 1$ matches.

(c) We prove, $W(n, m, r) = W(m, n, r)$, $r \neq 0$, by double induction on m, n . Assume that i) $W(n - 1, m, r) = W(m, n - 1, r)$ and ii) $W(n, m - 1, r) = W(m - 1, n, r)$.

Adding corresponding summands in (i) and (ii) and by (a) of theorem 1, we get

$$(2.2) \quad W(n, m, r) = W(m, n, r), \quad r \neq 0$$

The following tables give the number of cases in which there are exactly r matches for certain r .

Urns														
$r=1$	1	2	3	4	$r=2$	1	2	3	4	$r=3$	1	2	3	4
1	1	1	1	1		0	0	0	0		0	0	0	0
2	1	2	3	4		0	1	1	1		0	0	0	0
3	1	3	5	9		0	1	4	5		0	0	1	1
4	1	4	9	14		0	1	5	14		0	0	1	6

3. Proof of Theorem 2. By double induction on $m, n, m \leq n$, assume,

$$W(n - 1, n, r) = \binom{n + m - 2}{m - r} \frac{n - m + 2r - 1}{n + r - 1},$$

$$W(n, m - 1, r) = \binom{n + m - 2}{m - r - 1} \frac{n - m + 2r + 1}{n + r}.$$

By a simple calculation and (a) of Theorem 1, we get

$$(3.1) \quad W(n-1, m, r) + W(n, m-1, r) = \binom{n+m-1}{n-r} \frac{n-m+2r}{n+r} = W(n, m, r).$$

For $n < m$ the result is obtained by interchanging n by m and using (c) of Theorem 1. The case $n = m$ can be proved similarly.

4. Proof of Theorem 3. In what follows θ denotes on unspecified bounded function which may be different from one line to another, such that $0 < \theta < 1$, over the range of argument specified. Using $\text{Ln}(1 - Z) = Z + (1/2)Z^2 + \theta(Z) Z^3$, then

$$(4.1) \quad \begin{aligned} P(W > k) &= \binom{n+m-1}{m-k} / \binom{n+m-1}{n} \\ &= \left(\frac{m}{n}\right)^{k-1} \frac{\left(1-\frac{1}{m}\right)\left(1-\frac{2}{m}\right)\cdots\left(1-\frac{k-1}{m}\right)}{\left(1+\frac{1}{n}\right)\left(1+\frac{2}{n}\right)\cdots\left(1+\frac{k-1}{n}\right)}, \quad m \leq n. \end{aligned}$$

Hence,

$$(4.2) \quad \begin{aligned} \ln P(W > k) &= \ln\left(\frac{m}{n}\right)^{k-1} + \sum_{i=1}^{k-1} \ln\left(1-\frac{i}{m}\right) - \sum_{j=1}^{k-1} \ln\left(1+\frac{j}{n}\right) \\ &= \ln\left(\frac{m}{n}\right)^{k-1} - \left(\frac{1}{m} + \frac{1}{n}\right) \frac{k(k+1)}{2} \\ &\quad - \frac{1}{2} \left(\frac{1}{m^2} - \frac{1}{n^2}\right) \frac{k(k+1)(2k+1)}{6} \\ &\quad - \theta(k) \frac{1}{3} \left(\frac{1}{m^3} + \frac{1}{n^3}\right) \frac{k^2(k+1)^2}{4}. \end{aligned}$$

Replace k by \sqrt{n} , z and m by $n - \alpha \sqrt{n}$ such that $\alpha \rightarrow 0$ as $n \rightarrow m$, and taking the limit as $n, m \rightarrow \infty$ all terms in $P(W > k)$ tend to 0 except the first and the second term, hence

$$(4.3) \quad \begin{aligned} \lim_{n \rightarrow \infty} P(Z > z) &\simeq \lim_{n \rightarrow \infty} \left(1 - \frac{\alpha}{n}\right)^{\sqrt{nz}} e^{-\left(\frac{2n-\alpha\sqrt{n}}{n(n-\alpha\sqrt{n})}\right) \frac{\sqrt{n}z(\sqrt{nz}-1)}{2}} \\ &\simeq e^{(-z^2+\alpha z)}. \end{aligned}$$

The case $n < m$ gives the same result where $\alpha \rightarrow 0$ as $m \rightarrow n$.

REFERENCES

1. A.M. Khidr, Indian J. Pure Applied. Math. 12 (12) (1981), 1402-1407.
2. L. Takas, On the comparison of a theoretical and an empirical distribution function, J. Applied. Prob. 8 (1971), 321-330.
3. I. Tomescu, and S. Rudeanu, Introduction to Combinatorics, Colletts, England, 1975.

