



Ann. Funct. Anal. 4 (2013), no. 1, 11–17

ANNALS OF FUNCTIONAL ANALYSIS

ISSN: 2008-8752 (electronic)

URL: www.emis.de/journals/AFA/

CONFORMAL NETS AND KK-THEORY

SEBASTIANO CARPI¹, ROBERTO CONTI^{2*} AND ROBIN HILLIER³

Communicated by N. P. Landsman

ABSTRACT. Given a completely rational conformal net \mathcal{A} on S^1 , its fusion ring acts faithfully on the K-group $K_0(\mathfrak{K}_{\mathcal{A}})$ of a certain universal C^* -algebra $\mathfrak{K}_{\mathcal{A}}$ associated to \mathcal{A} , as shown in a previous paper. We prove here that this action can actually be identified with a Kasparov product, thus paving the way for a fruitful interplay between conformal field theory and KK-theory.

1. INTRODUCTION

In the operator algebraic approach a chiral conformal field theory is described by a conformal net \mathcal{A} on S^1 , namely an inclusion preserving map $I \mapsto \mathcal{A}(I)$ from the set of proper (nonempty, nondense, open) intervals of the unit circle S^1 into the family of von Neumann algebras (actually type III₁ factors) acting on a fixed separable (complex) Hilbert space \mathcal{H} called the *vacuum Hilbert space* of the theory, which is conformally covariant and satisfies some other natural conditions, see e.g. [16] and the references therein. The superselection structure of the theory is then captured by the representation theory of the net \mathcal{A} and can also be described in terms of localized (DHR) endomorphisms.

In a recent paper [4] together with Weiner we have started an investigation of K-theoretical aspects of conformal nets. In the present note we take one further step along these lines and provide an interpretation of some of the results obtained there in the framework of Kasparov KK-theory. At a first glance, this might appear merely as an academic curiosity. However, in view of the striking success of KK-theory on the one hand (e.g. [9] for an overview) and the noncommutative

Date: Received: 24 July 2012; Accepted: 13 August 2012.

* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 81Txx; Secondary 46Lxx, 58B34, 19K35.

Key words and phrases. Operator algebra, conformal field theory, conformal net, superselection sector, fusion ring, K-theory, Kasparov product.

geometrization program for (super-) conformal nets and their representations [19, 6, 5] on the other hand, we feel that there are good reasons to give a closer look at this subject, as it could possibly reveal a lot of potential for further investigations.

We recall the results in [4] relevant for the present analysis, referring the reader to that paper for more details, notation, and references. Let \mathcal{A} be a completely rational local conformal net on S^1 . We can then define the so-called locally normal universal C^* -algebra $C_{\text{ln}}^*(\mathcal{A})$ which is canonically associated to the superselection theory of the net. It may be expressed as a quotient of the universal algebra $C^*(\mathcal{A})$ previously defined by Fredenhagen in [12] or, equivalently, as the image of $C^*(\mathcal{A})$ in the so-called locally normal universal representation $(\pi_{\text{ln}}, \mathcal{H}_{\text{ln}})$. The surprising fact is that $C_{\text{ln}}^*(\mathcal{A})$ turns out to be isomorphic to the direct sum $B(\mathcal{H})^{\oplus n}$ of n copies of the algebra $B(\mathcal{H})$ of all bounded operators on the vacuum Hilbert space of \mathcal{A} . Here, $n < \infty$ is the number of sectors of \mathcal{A} . Consequently, it has trivial K-theory and several other properties which do not make it really attractive as a C^* -algebra.

We therefore consider its separable C^* -subalgebra $\mathfrak{K}_{\mathcal{A}}$ generated by the finite projections. It is isomorphic to $\mathcal{K}^{\oplus n} \subset B(\mathcal{H})^{\oplus n}$, where $\mathcal{K} := K(\mathcal{H})$ is the algebra of compact operators on \mathcal{H} . Clearly $\mathfrak{K}_{\mathcal{A}}$ is an ideal of $C_{\text{ln}}^*(\mathcal{A})$ and in fact it is the largest among the norm-separable ideals. We have

$$K_0(\mathfrak{K}_{\mathcal{A}}) = \mathbb{Z}^n, \quad K_1(\mathfrak{K}_{\mathcal{A}}) = 0.$$

Furthermore, we get a faithful semiring action of the fusion semiring $\mathcal{R}_{\mathcal{A}}$ of \mathcal{A} on $K_0(\mathfrak{K}_{\mathcal{A}})$, via restriction of endomorphisms of $C_{\text{ln}}^*(\mathcal{A})$ to $\mathfrak{K}_{\mathcal{A}}$ and subsequent pushforward to $K_0(\mathfrak{K}_{\mathcal{A}})$. Namely, as shown in [4, Th.4.4], there is an injective semiring homomorphism

$$\eta : \mathcal{R}_{\mathcal{A}} \rightarrow \text{End}(K_0(\mathfrak{K}_{\mathcal{A}}))$$

satisfying $\eta_{[\rho]} = (\hat{\rho}|_{\mathfrak{K}_{\mathcal{A}}})_*$ for every localized covariant endomorphism ρ of $C^*(\mathcal{A})$, where $[\rho] \in \mathcal{R}_{\mathcal{A}}$ is the sector (i.e., the unitary equivalence class) represented by ρ , and $\hat{\rho}$ is a normal $*$ -homomorphism of $C_{\text{ln}}^*(\mathcal{A})$ naturally associated to ρ , mapping $\mathfrak{K}_{\mathcal{A}}$ into itself. Hereafter, we shall use “[ρ]” exclusively to denote the sector determined by ρ and not its KK-class.

We shall give an interpretation of this action in terms of KK-theory. This fact can be seen as an illustration in the conformal nets setting of a comment of Izumi [13, page 118], based on previous ideas of Kajiwara and Watatani [15, 21], on the relation between sector theory and KK-theory. There they work out the most natural way of associating a KK-class to a $*$ -homomorphism between two simple C^* -algebras. In our context, however, it is a priori unclear what are the correct C^* -algebras to be considered and whether we can get rid of their assumptions. In [10, 11], Evans and Gannon discuss KK-theoretical aspects of CFT in relation to modular invariants and twisted K-theory. Following some of their ideas, we shall provide a KK-theoretical interpretation of modular invariants in our setting towards the end of the present article.

Another instance of the emergence of KK-theoretical concepts in AQFT has been recently pointed out by Conti and Morsella [7], where the DHR sectors of

scaling limit nets (as defined by Buchholz-Verch, in order to cast the renormalization group analysis into operator algebraic terms) are described by maps of the original global quasi-local C^* -algebra on four-dimensional Minkowski space-time that are suitable modifications of the asymptotic morphisms of Connes and Higson.

We close this introduction by mentioning that other relationships of seemingly quite different nature between KK-theory and quantum field theory have been investigated by other authors, see e.g. [3].

2. BASICS OF KK-THEORY

It is well-known that the push-forward of $*$ -homomorphisms gives rise to elements in KK-theory. We want to see how this works in our setting in relation to the semiring action η recalled above. For the convenience of the reader and in order to fix the notation let us introduce here KK-theory very briefly, while referring to [1] and [14] for details. All C^* -algebras we consider below will be separable and stably unital. In particular, the various approaches to KK-theory become all equivalent [1, Sec.17&18] and $K_0(A)$ is the Grothendieck group of the projection semigroup of the C^* -algebra $\mathcal{K} \otimes A$. Moreover, we will restrict ourselves to the ungraded algebra case.

A *Kasparov* (A, B) -module is a tuple (\mathcal{E}, ϕ, F) , where \mathcal{E} is a countably generated graded Hilbert B -module, $\phi : A \rightarrow B(\mathcal{E})$ is a graded $*$ -homomorphism, and $F \in B(\mathcal{E})$ has degree one, such that

$$(F - F^*)\phi(a), \quad (F^2 - \mathbf{1})\phi(a), \quad [F, \phi(a)]$$

lie all in $K(\mathcal{E})$, the compact operators on \mathcal{E} , for all $a \in A$. With the usual concept of homotopy, one defines $KK(A, B)$ as the set of homotopy equivalence classes of Kasparov (A, B) -modules. Moreover, if we have two Kasparov (A, B) -modules $(\mathcal{E}_i, \phi_i, F_i)$ and a unitary in $B(\mathcal{E}_1, \mathcal{E}_2)$ intertwining the ϕ_i and the F_i , then the two Kasparov modules are homotopically equivalent. There is a direct sum for Kasparov (A, B) -modules, which passes to the quotient $KK(A, B)$; cf. [1, Sec.17] for all these statements.

For our immediate purposes, the most relevant facts about KK-theory can be summarized as follows:

Theorem 2.1. *Let A, B, C be C^* -algebras as above.*

- (1) *$KK(A, B)$ is an abelian group with respect to the above addition, and KK is a bifunctor from the category of C^* -algebras to abelian groups [1, 17.3&17.8].*
- (2) *There is a canonical identification of $KK(\mathbb{C}, A)$ with $K_0(A)$ (as additive groups), [1, 17.5.5].*
- (3) *Every $*$ -homomorphism $\phi : A \rightarrow B$ naturally defines a $KK(A, B)$ -element $\{\phi\}$ as the homotopy class of $(B, \phi, 0)$, where we have identified $B(B)$ with the multiplier algebra $\mathcal{M}(B)$.*
- (4) *If two $*$ -homomorphisms $\phi, \psi : A \rightarrow B$, are unitarily equivalent in $\mathcal{M}(B)$, then the induced $KK(A, B)$ -elements coincide, [14, Sec.1.3].*

(5) *There exists a bilinear map \times , the so-called Kasparov product*

$$KK(A, B) \times KK(B, C) \rightarrow KK(A, C),$$

which is associative, [1, Sec.18].

(6) *If $\psi : A \rightarrow B$ and $\phi : B \rightarrow C$ are $*$ -homomorphisms then*

$$\{\psi\} \times \{\phi\} = \{\phi \circ \psi\}.$$

If id_A is the identity automorphism of A then $\{\text{id}_A\}$ is the neutral element in $KK(A, A)$ for the Kasparov product. Hence $KK(A, A)$ is a unital ring.

(7) *The adjoint of the pairing $KK(\mathbb{C}, A) \times KK(A, B) \rightarrow KK(\mathbb{C}, B)$ defines, via the identification in (2), a map $\gamma : KK(A, B) \rightarrow \text{Hom}(K_0(A), K_0(B))$, namely*

$$\gamma(y)(x) = x \times y, \quad x \in K_0(A), y \in KK(A, B).$$

For $y = \{\phi\}$ with $\phi : A \rightarrow B$ a $$ -homomorphism, we have*

$$\gamma(\{\phi\}) = \phi_*,$$

the push-forward of ϕ in K -theory.

We only mention that property (6) may be obtained as a consequence of the functoriality properties of the KK -product [1, 18.7.1] and the relation $\{\phi\} = \phi^* \{\text{id}_C\}$. The canonical identification in (2) is given by

$$[p_+] - [p_-] \in K_0(A) \mapsto \left[\mathcal{H}_A \oplus \mathcal{H}_A, \phi_{p_+} \oplus \phi_{p_-}, \begin{pmatrix} 0 & \mathbf{1} \\ \mathbf{1} & 0 \end{pmatrix} \right] \in KK(\mathbb{C}, A),$$

where \mathcal{H}_A is the Hilbert module $A \otimes l^2(\mathbb{Z})$, the grading on $\mathcal{H}_A \oplus \mathcal{H}_A$ is $\mathbf{1} \oplus -\mathbf{1}$, $p_{\pm} \in A \otimes \mathcal{K}$ are projections, and $\phi_{p_{\pm}}$ is the map: $t \in \mathbb{C} \mapsto tp_{\pm} \in \mathcal{M}(A \otimes \mathcal{K})$. It has, in particular, the property that push-forwards in $K_0(A)$ and $KK(\mathbb{C}, A)$ mutually correspond to each other. All this can be easiest seen as outlined in [8, Sec.1] following Cuntz' quasi-homomorphism picture; alternatively one may use [1, 17.5.5]. Finally, property (7) is a consequence of (2) and the functoriality property [1, 17.8.2].

3. MAIN RESULT

Let us return to our setting, using the notation from the above theorem and Section 4 in [4].

Theorem 3.1. *Let \mathcal{A} be a completely rational net on S^1 , and let $\eta : \mathcal{R}_{\mathcal{A}} \rightarrow \text{End}(K_0(\mathfrak{K}_{\mathcal{A}}))$ be the injective semiring homomorphism recalled above. Then there is an injective unital semiring homomorphism $j : \mathcal{R}_{\mathcal{A}} \rightarrow KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$ such that*

$$x \times j([\rho]) = \eta_{[\rho]}(x), \quad [\rho] \in \mathcal{R}_{\mathcal{A}}, \quad x \in K_0(\mathfrak{K}_{\mathcal{A}}) = KK(\mathbb{C}, \mathfrak{K}_{\mathcal{A}}). \quad (3.1)$$

Proof. Using Theorem 2.1(1), define

$$j([\rho]) := \{\hat{\rho}|_{\mathfrak{K}_{\mathcal{A}}}\} \in KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}}), \quad [\rho] \in \mathcal{R}_{\mathcal{A}}.$$

This map is well-defined since if ρ and ρ' are equivalent endomorphisms in $C^*(\mathcal{A})$ then $\hat{\rho}$ and $\hat{\rho}'$ are equivalent endomorphisms of $C_{\text{ln}}^*(\mathcal{A})$. As the latter contains $\mathfrak{K}_{\mathcal{A}}$ as an ideal and it is weakly closed in $B(\mathcal{H}_{\text{ln}})$, it can be easily identified with the

multiplier algebra $\mathcal{M}(\mathfrak{K}_{\mathcal{A}})$. Then by Theorem 2.1(2), $\hat{\rho}|_{\mathfrak{K}_{\mathcal{A}}}$ and $\hat{\rho}'|_{\mathfrak{K}_{\mathcal{A}}}$ give rise to Kasparov $(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$ -modules, which according to (3) are equivalent and so define the same KK-element $\{\hat{\rho}|_{\mathfrak{K}_{\mathcal{A}}}\}$.

Furthermore, j is multiplicative, namely $j([\rho_1][\rho_2]) = j([\rho_1]) \times j([\rho_2])$ and $j([\text{id}])$ is the unit in $KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$. Indeed, using the commutativity of the composition of DHR sectors (due to the existence of a unitary braiding) and point (6) in Theorem 2.1,

$$\begin{aligned} j([\rho_1][\rho_2]) &= j([\rho_1 \circ \rho_2]) = j([\rho_2 \circ \rho_1]) = \{\widehat{\rho_2 \circ \rho_1}|_{\mathfrak{K}_{\mathcal{A}}}\} \\ &= \{\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}} \circ \hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\} = \{\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\} \times \{\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}\} = j([\rho_1]) \times j([\rho_2]) . \end{aligned}$$

Next, we check additivity of j , namely $j([\rho_1] \oplus [\rho_2]) = j([\rho_1]) + j([\rho_2])$. If s_1 and s_2 are a pair of isometries generating a copy of O_2 inside $C^*(\mathcal{A})$ then a representative of $[\rho_1] \oplus [\rho_2] \in \mathcal{R}_{\mathcal{A}}$ is the endomorphism $\rho = s_1\rho_1(\cdot)s_1^* + s_2\rho_2(\cdot)s_2^*$. Therefore, using the fact that $\hat{\rho}(x) = \hat{s}_1\hat{\rho}_1(x)\hat{s}_1^* + \hat{s}_2\hat{\rho}_2(x)\hat{s}_2^*$, $x \in C_{\text{in}}^*(\mathcal{A})$, where $\hat{s}_i = \pi_{\text{in}}(s_i)$, $i = 1, 2$, one has

$$j([\rho_1] \oplus [\rho_2]) = \{(s_1\rho_1(\cdot)s_1^* + s_2\rho_2(\cdot)s_2^*)|_{\mathfrak{K}_{\mathcal{A}}}\} = \{\hat{s}_1\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_1^* + \hat{s}_2\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_2^*\} .$$

The subsequent Lemma 3.2 and its proof (with $A = C_{\text{in}}^*(\mathcal{A})$ and $I = \mathfrak{K}_{\mathcal{A}}$) imply

$$T(\mathfrak{K}_{\mathcal{A}} \oplus \mathfrak{K}_{\mathcal{A}}) = \mathfrak{K}_{\mathcal{A}}, \quad T(\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}} \oplus \hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}})T^* = \hat{s}_1\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_1^* + \hat{s}_2\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_2^*,$$

so that, according to Theorem 2.1(3) and the introduction of Kasparov modules, the classes of $(\mathfrak{K}_{\mathcal{A}}, \hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}, 0) \oplus (\mathfrak{K}_{\mathcal{A}}, \hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}, 0)$ and $(\mathfrak{K}_{\mathcal{A}}, \hat{s}_1\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_1^* + \hat{s}_2\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_2^*, 0)$ in $KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$ coincide, thus

$$\{\hat{s}_1\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_1^* + \hat{s}_2\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}\hat{s}_2^*\} = \{\hat{\rho}_1|_{\mathfrak{K}_{\mathcal{A}}}\} + \{\hat{\rho}_2|_{\mathfrak{K}_{\mathcal{A}}}\} = j([\rho_1]) + j([\rho_2]).$$

Finally, identity (3.1) follows at once from point (7) in Theorem 2.1. Since η is injective according to [4, Theorem 4.4], j has to be injective, too. \square

Lemma 3.2. *Let A be a unital C^* -algebra containing a copy of the Cuntz algebra O_2 and let I be a closed two-sided ideal in A . Then $I \oplus I \simeq I$ as right Hilbert I -modules.*

Proof. Let $s_1, s_2 \in A$ be a pair of isometries such that $s_1s_1^* + s_2s_2^* = 1$. Define the linear operator $T : I \oplus I \rightarrow I$ by $T(a, b) := s_1a + s_2b$, $a, b \in I$. It is easy to see that T is an isometric and surjective right I -module map, so that it is indeed unitary, [18, 3.2]. \square

4. SOME REMARKS

First of all, Theorem 3.1 says that the left action of $\mathcal{R}_{\mathcal{A}}$ on $K_0(\mathfrak{K}_{\mathcal{A}})$ ‘‘factorizes’’ through a right action of $KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$ and can be expressed in terms of a Kasparov product. This construction appears to depend on various special properties of the algebra $\mathfrak{K}_{\mathcal{A}}$ which has been defined only for completely rational conformal nets. It is not evident how to generalize the definition of $\mathfrak{K}_{\mathcal{A}}$ in order to find the

analogues of [4, Th.4.4] and Theorem 3.1 for a more general class of conformal nets.

Second, write $\tilde{\mathcal{R}}_{\mathcal{A}}$ for the Grothendieck ring generated by the semiring $\mathcal{R}_{\mathcal{A}}$ and \tilde{j} for the unique ring homomorphism extending j to $\tilde{\mathcal{R}}_{\mathcal{A}}$. It is clearly injective, owing to the preceding theorem. We would like to stress then that $\tilde{j}(\tilde{\mathcal{R}}_{\mathcal{A}}) \subset KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$ is always a *proper subring* if the number of sectors n is greater than 1, i.e., for non-holomorphic theories. This can be easily seen because $\tilde{\mathcal{R}}_{\mathcal{A}} \simeq K_0(\mathfrak{K}_{\mathcal{A}})$ is commutative while $KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$ is non-commutative. To prove the latter statement, we simply apply the universal coefficient theorem [2, V.1.5.8], which holds for the algebra $\mathfrak{K}_{\mathcal{A}}$ since it lies in the so-called bootstrap class [2, V.1.5.4], and which says

$$KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}}) \simeq \text{End}(K_0(\mathfrak{K}_{\mathcal{A}})) \simeq M_n(\mathbb{Z}).$$

Our last comment concerns *modular invariants*. In [10, Sec.7] it has been pointed out that modular invariants can be regarded as certain KK-classes. As we shall see now, an interpretation of modular invariants as KK-classes can be directly formulated in our setting. For a completely rational conformal net the braiding is always non-degenerate [17, Cor.37]. Then by Rehren's construction there are invertible complex vector space endomorphisms $S, T \in \text{End}(\mathbb{C} \otimes_{\mathbb{Z}} \tilde{\mathcal{R}}_{\mathcal{A}})$ giving rise to a representation of the modular group $\text{SL}(2, \mathbb{Z})$ on the fusion algebra $\mathbb{C} \otimes_{\mathbb{Z}} \tilde{\mathcal{R}}_{\mathcal{A}}$, cf. [20]. A modular invariant is a \mathbb{Z} -module endomorphisms $Z \in \text{End}_{\mathbb{Z}}(\tilde{\mathcal{R}}_{\mathcal{A}})$ such that:

- $\text{id} \otimes_{\mathbb{Z}} Z \in \text{End}(\mathbb{C} \otimes_{\mathbb{Z}} \tilde{\mathcal{R}}_{\mathcal{A}})$ commutes with S and T (modular invariance);
- $Z(\mathcal{R}_{\mathcal{A}}) \subset \mathcal{R}_{\mathcal{A}}$ (positivity);
- the matrix element Z_{11} in the distinguished basis of sectors is 1 (uniqueness of the vacuum).

Hence, as a consequence of the additive group isomorphism $\tilde{\mathcal{R}}_{\mathcal{A}} \simeq K_0(\mathfrak{K}_{\mathcal{A}})$, every modular invariant Z can be identified with an element in $\text{End}(K_0(\mathfrak{K}_{\mathcal{A}})) \simeq KK(\mathfrak{K}_{\mathcal{A}}, \mathfrak{K}_{\mathcal{A}})$.

Acknowledgement. We would like to thank Roberto Longo for stimulating comments on the relationship between the the representation theory of conformal nets and KK-theory. We also thank Martin Grensing for a very useful discussion on Kasparov products and for drawing to our attention some references.

REFERENCES

1. B. Blackadar, *K-theory for operator algebras*, 2nd edition, Mathematical Sciences Research Institute Publications 5, Cambridge University Press, Cambridge, 1998.
2. B. Blackadar, *Operator algebras. Theory of C^* -algebras and von Neumann algebras*, Encyclopaedia of Mathematical Sciences, 122, Operator Algebras and Non-commutative Geometry, III, Springer-Verlag, Berlin, 2006.
3. J. Brodzki, V. Mathai, J. Rosenberg and R. Szabo, *D-branes, RR-fields and duality on noncommutative manifolds*, Comm. Math. Phys. **277** (2008), 643–706.
4. S. Carpi, R. Conti, R. Hillier and M. Weiner, *Representations of conformal nets, universal C^* -algebras and K-theory*, Commun. Math. Phys. DOI: 10.1007/s00220-012-1561-5.

5. S. Carpi, R. Hillier, Y. Kawahigashi and R. Longo, *Spectral triples and the super-Virasoro algebra*, Comm. Math. Phys. **259** (2010), 1–27.
6. S. Carpi, Y. Kawahigashi and R. Longo, *Structure and classification of superconformal nets*, Ann. Henri Poincaré **9** (2008), 1069–1121.
7. R. Conti and G. Morsella, *Asymptotic morphisms and superselection theory in the scaling limit*, J. Noncommut. Geom. (to appear), arXiv: 1205.6380.
8. J. Cuntz, *Generalized homomorphisms between C^* -algebras and KK-theory*, Dynamics and processes, 31–45, LNM 1031, Springer, Berlin, 1983.
9. J. Cuntz, R. Meyer and J.M. Rosenberg, *Topological and bivariant KK-theory*, Oberwolfach Seminars, 36, Birkhäuser Verlag, Basel, 2007.
10. D.E. Evans and T. Gannon, *Modular invariants and twisted K-theory*, Commun. Number Theory Phys. **277** (2009), 209–296.
11. D.E. Evans and T. Gannon, *Modular invariants and twisted K-theory II. Dynkin diagram symmetries*, arXiv:1012.1634.
12. K. Fredenhagen, *Generalizations of the theory of superselection sectors*, The algebraic theory of superselection sectors. Introduction and recent results (Palermo, 1989), 379–387, World Sci. Publ., River Edge, NJ, 1990.
13. M. Izumi, *Inclusions of simple C^* -algebras*, J. Reine Angew. Math. **547** (2002), 97–138.
14. K.K. Jensen and K. Thomsen. *Elements of KK-theory*, Birkhäuser, Boston, MA, 1991.
15. T. Kajiwara and Y. Watatani, *Jones index theory by Hilbert C^* -bimodules and K-theory*, Trans. Amer. Math. Soc. **352** (2000), 3429–3472.
16. Y. Kawahigashi and R. Longo, *Classification of local conformal nets. Case $c < 1$* , Ann. of Math. **160** (2004), 493–522.
17. Y. Kawahigashi, R. Longo and M. Müger, *Multi-interval subfactors and modularity of representations in conformal field theory*, Comm. Math. Phys. **219** (2001), 631–669.
18. E.C. Lance. *Hilbert C^* -modules. A toolkit for operator algebraists*, London Mathematical Society Lecture Note Series, 210. Cambridge University Press, Cambridge, 1995.
19. R. Longo. *Notes for a quantum index theorem*. Comm. Math. Phys. **222** (2001), 45–96.
20. K.H. Rehren, *Braid group statistics and their superselection rules*, The algebraic theory of superselection sectors. Introduction and recent results (Palermo, 1989), 333–355, World Sci. Publ., River Edge, NJ, 1990.
21. Y. Watatani, *Index for C^* -subalgebras*. Mem. Amer. Math. Soc. 83, 1990.

¹ DIPARTIMENTO DI ECONOMIA, UNIVERSITÀ DI CHIETI-PESCARA “G. D’ANNUNZIO” VIALE PINDARO, 42, I-65127 PESCARA, ITALY.

E-mail address: s.carpi@unich.it

² SBAI, SEZIONE DI MATEMATICA, SAPIENZA UNIVERSITÀ DI ROMA, VIA A. SCARPA, 16, I-00161 ROMA, ITALY.

E-mail address: roberto.conti@sbai.uniroma1.it

³ DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA “TOR VERGATA”, VIA DELLA RICERCA SCIENTIFICA, 1, I-00133 ROMA, ITALY.

E-mail address: hillier@mat.uniroma2.it