



Ann. Funct. Anal. 3 (2012), no. 1, 19–31

ANNALS OF FUNCTIONAL ANALYSIS

ISSN: 2008-8752 (electronic)

URL: [www.emis.de/journals/AFA/](http://www.emis.de/journals/AFA/)

## $p$ -POWER QUASICONCAVITY OF A REARRANGEMENT INVARIANT FUNCTION SPACE

CHONGSUNG LEE<sup>1\*</sup> AND KYUGEUN CHO<sup>2</sup>

Communicated by C. P. Niculescu

ABSTRACT. We define the  $p$ -power quasiconcave function and show relationships between  $p$ -power quasiconcave fundamental function and r.i. spaces like Lorentz space and Marcinkiewicz space.

### 1. INTRODUCTION

Let  $E$  be a rearrangement invariant function space (or r.i. space, in short), which consists of measurable functions defined on a measure space  $(\Omega, \Sigma, \mu)$ . The general theory on r.i. space can be found in [4]. In this paper, we focus on r.i. space defined on a positive real line with Lebesgue measure since our objectivity is to investigate relationships between the fundamental function  $\varphi_E = \|\chi_{[0,t]}\|_E$  and some geometric properties of a space  $E$ . We say a positive function  $\varphi(t)$  on the positive real line is called quasiconcave if  $\varphi(t)$  is positive and nondecreasing and  $\varphi(t)/t$  is nonincreasing. In [6], it has been revealed that the necessary and sufficient condition of a positive function  $\varphi(t)$  is a fundamental function of r.i. space  $\varphi(t)$  that is quasiconcave and  $\varphi(0) = 0$ . The followings are the known results of a positive concave function  $\varphi(t)$  on  $[0, \infty)$  and their proof can be found in [3] and [5].

**Property 1.1.** A positive function  $\varphi(t)$  is equivalent to a positive concave function if and only if

$$\varphi(t) \leq C \max\left(1, \frac{t}{s}\right) \varphi(s) \quad \text{for all } s, t > 0.$$

---

*Date:* Received: 7 October 2011; Accepted: 30 December 2011.

\* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 46B03; Secondary 46B04, 46B42.

*Key words and phrases.*  $p$ -power concavity, upper  $p$ -estimate, rearrangement invariant space.

In particular, when  $C = 1$ , there exists a concave function  $\tilde{\varphi}(t)$  such that

$$\frac{1}{2}\tilde{\varphi}(t) \leq \varphi(t) \leq \tilde{\varphi}(t).$$

Indeed, we can find a concave function which is equivalent to a given quasiconcave function.

**Property 1.2.** If  $\varphi(t)$  is positive and everywhere finite on  $(0, \infty)$ , which satisfies  $\varphi(t_1 \cdot t_2) \leq \varphi(t_1) \cdot \varphi(t_2)$ , then we get

$$\lim_{t \rightarrow \infty} \frac{\log \varphi(t)}{\log t} = \inf_{1 < t} \frac{\log \varphi(t)}{\log t} = \bar{\alpha}$$

and

$$\lim_{t \rightarrow 0} \frac{\log \varphi(t)}{\log t} = \inf_{t < 1} \frac{\log \varphi(t)}{\log t} = \underline{\alpha}$$

Furthermore, we have  $-\infty < \underline{\alpha} \leq \bar{\alpha} < \infty$ .

**Definition 1.3.** For a given positive, everywhere finite function  $\varphi(t)$  on  $(0, \infty)$ , we define the function  $D_\varphi(s)$ , which is called the dilation function of  $\varphi(t)$ , by

$$D_\varphi(s) = \sup_{0 < t < \infty} \frac{\varphi(st)}{\varphi(t)}.$$

When  $\varphi(t)$  is quasiconcave, it is easy to show that  $D_\varphi(s)$  is everywhere finite and satisfies the submultiplicativity condition of Property 1.2. Therefore, we can define the following two indices, which were introduced by Zippin for the fundamental function  $\varphi_E(t)$  of r.i. space E (see [8]).

**Definition 1.4.** Let  $\varphi(t)$  be a positive quasiconcave function. We then define two indices,  $\bar{r}(\varphi)$  and  $\underline{r}(\varphi)$ , which will be called the upper and lower indices of  $\varphi(t)$ , by

$$\bar{r}(\varphi) = \lim_{t \rightarrow \infty} \frac{\log D_\varphi(t)}{\log t} = \inf_{1 < t} \frac{\log D_\varphi(t)}{\log t}$$

and

$$\underline{r}(\varphi) = \lim_{t \rightarrow 0} \frac{\log D_\varphi(t)}{\log t} = \sup_{0 < t < 1} \frac{\log D_\varphi(t)}{\log t}.$$

The following simple facts are also mentioned in [8] for the case of a fundamental function  $\varphi_E(t)$  and [3].

**Property 1.5.** Let  $\varphi(t)$  be a quasiconcave function. We then have

- i)  $0 \leq \underline{r}(\varphi) \leq \bar{r}(\varphi) \leq 1$
- ii)  $\underline{r}(\varphi) + \bar{r}(\varphi) = 1$
- iii) If  $\Psi(t)$  is equivalent to  $\varphi(t)$ , then  $\bar{r}(\Psi) = \bar{r}(\varphi)$  and  $\underline{r}(\Psi) = \underline{r}(\varphi)$ .

**Property 1.6.** Let  $\varphi(t)$  be nondecreasing and  $D_\varphi(s_1) \leq s_1$  for some  $s_1 > 1$ . Then there exists a concave function  $\Psi(t)$  which is equivalent to  $\varphi(t)$ .

2.  $p$ -POWER QUASICONCAVE FUNCTION

**Definition 2.1.** Let  $\varphi(t)$  be a positive quasiconcave function on  $[0, \infty)$ . We say that  $\varphi(t)$  is  $p$ -power quasiconcave with a constant  $C$  if it satisfies

$$\sum_{i=1}^n \varphi^p(a_i) \leq C \varphi^p \left( \sum_{i=1}^n a_i \right), \quad \text{for all } a_i \text{ in } [0, \infty). \quad (2.1)$$

In particular, if  $\varphi(t)$  is concave and satisfies (2.1), we say that  $\varphi(t)$  is  $p$ -power concave with a constant  $C$ .

It is clear that the concave function  $x^{1/p}$  is  $p$ -power concave. In the following, we consider some conditions, which are useful in showing the existence of nontrivial  $p$ -power quasiconcave functions.

- Lemma 2.2.**
- i) If  $\varphi(t)$  is a  $p$ -power quasiconcave function with a constant  $C$ , then  $\varphi(t)$  is also  $q$ -power quasiconcave with a constant  $C^{q/p}$  when  $p \leq q$ .
  - ii) Let  $\varphi(t)$  satisfy (2.1) and  $\Psi(t)$  be a positive nondecreasing function. Then  $\varphi(t)\Psi(t)$  also satisfies (2.1).
  - iii) Let  $\varphi(t)$  be a positive  $p$ -power quasiconcave function on  $[0, \infty)$ . If  $\Psi(t)$  is equivalent to  $\varphi(t)$ , then  $\Psi(t)$  also satisfies (2.1).

*Proof.* i) It is clear since  $\|\cdot\|_p \geq \|\cdot\|_q$  for  $p \leq q$ . Indeed, we have

$$\left\{ \sum_{i=1}^n \varphi^q(a_i) \right\}^{1/q} \leq \left\{ \sum_{i=1}^n \varphi^p(a_i) \right\}^{1/p} \leq C^{1/p} \varphi \left( \sum_{i=1}^n a_i \right).$$

Hence,

$$\sum_{i=1}^n \varphi^q(a_i) \leq C^{q/p} \varphi^q \left( \sum_{i=1}^n a_i \right).$$

ii)

$$\begin{aligned} \sum_{i=1}^n \varphi^p(a_i) \Psi^p(a_i) &\leq \left( \sum_{i=1}^n \varphi^p(a_i) \right) \Psi^p \left( \sum_{i=1}^n a_i \right) \\ &\leq C \varphi^p \left( \sum_{i=1}^n a_i \right) \Psi^p \left( \sum_{i=1}^n a_i \right) \end{aligned}$$

iii) Suppose that  $C_1 \varphi(t) \leq \Psi(t) \leq C_2 \varphi(t)$ . Then,

$$\begin{aligned} \sum_{i=1}^n \Psi^p(a_i) &\leq \sum_{i=1}^n C_2^p \varphi^p(a_i) \leq C_2^p \cdot C \varphi^p \left( \sum_{i=1}^n a_i \right) \\ &\leq C \cdot \left( \frac{C_2}{C_1} \right)^p \Psi^p \left( \sum_{i=1}^n a_i \right). \end{aligned}$$

□

**Theorem 2.3.** *Let  $\varphi(t)$  be a positive quasiconcave function with  $\bar{r}(\varphi) < 1$  and  $w(t)$  be a quasiconcave function. If  $\Psi(t) = \varphi(t)w^\alpha(t)$  for  $0 \leq \alpha < 1 - \bar{r}(\varphi)$ , then  $\Psi(t)$  is quasiconcave. Furthermore, if  $\varphi(t)$  is  $p$ -power quasiconcave, there exists a concave function  $\Phi(t)$  which is also  $p$ -power quasiconcave and equivalent to  $\Psi(t)$ .*

*Proof.* By Property 1.6, it is enough to show that  $\Psi(t)$  is nondecreasing and  $D_\Psi(s_1) \leq s_1$ , for some  $s_1 > 1$ . Since  $\varphi(t)$  and  $w(t)$  are quasiconcave and  $\alpha$  is nonnegative,  $\Psi(t)$  is nondecreasing. In order to show that  $D_\Psi(s_1) \leq s_1$ , for some  $s_1 > 1$ , take  $\epsilon$  such that  $\alpha + \epsilon \leq 1 - \bar{r}(\varphi)$ . With this  $\epsilon$ , choose  $s_1$  sufficiently large such that  $D_\varphi(s_1) \leq s_1^{\bar{r}(\varphi) + \epsilon}$  from the definition of upper index of  $\varphi$ . Then, for all  $t > 0$ ,

$$\begin{aligned} \frac{\Psi(s_1 t)}{\Psi(t)} &= \frac{\varphi(s_1 t)\{w(s_1 t)\}^\alpha}{\varphi(t)\{w(t)\}^\alpha} \\ &= \frac{\varphi(s_1 t)\{w(s_1 t)/s_1 t\}^\alpha (s_1 t)^\alpha}{\varphi(t)\{w(t)/t\}^\alpha t^\alpha} \\ &\leq \frac{\varphi(s_1 t)}{\varphi(t)} s_1^\alpha, \\ &\quad \text{since } w(t)/t \text{ is nonincreasing and } \alpha \text{ is nonnegative,} \\ &\leq D_\varphi(s_1) s_1^\alpha \\ &\leq s_1^{\alpha + \bar{r}(\varphi) + \epsilon} \leq s_1 \end{aligned}$$

since  $\bar{r}(\varphi) + \alpha + \epsilon \leq 1$  and  $s_1 \geq 1$ . This implies that  $D_\Psi(s_1) \leq s_1$  for some  $s_1 > 1$ . Hence,  $\Psi(t)$  is quasiconcave and  $p$ -power quasiconcave by Lemma 2. 2. ii. The existence of a concave function which is equivalent to  $\Psi(t)$  is also obtained by Property 1.1  $\square$

The above theorem tells us that for a given  $p$ -power quasiconcave function  $\varphi(t)$ , we can construct another  $p$ -power quasiconcave function which is nonequivalent to  $\varphi(t)$ . The next example is a  $p$ -power concave function which is not equivalent to  $x^{1/p}$ . The following concave function has  $1/p$  as its lower and upper indices. Thus, the converse of Property 1.5-iii) is not true.

**Example 2.4.** Let  $p > 1$  be fixed. Define

$$\varphi(t) = \begin{cases} 1 & 0 \leq t \leq 1 \\ 1 + \log t & 1 \leq t \end{cases}$$

Define  $\Psi(t) = t^{1/p}\varphi(t)$ . We now compute the dilation function  $D_\Psi(s)$  of  $\Psi(t)$ . When  $s > 1$ , a simple calculation shows that

$$\frac{\Psi(st)}{\Psi(t)} = \begin{cases} s^{1/p} & 0 < t \leq 1/s \\ s^{1/p}(1 + \log st) & 1/s < t \leq 1 \\ s^{1/p}(1 + \log st)/(1 + \log t) & 1 < t. \end{cases}$$

Therefore,

$$\begin{aligned} D_\Psi &= \sup_{0 < t} \{\Psi(st)/\Psi(t)\} \\ &= \sup_{1/s < t \leq 1} s^{1/p}(1 + \log st) = s^{1/p}(1 + \log s). \end{aligned}$$

From this, we get

$$\begin{aligned} \bar{r}(\Psi) &= \lim_{s \rightarrow \infty} [\log \{s^{1/p}(1 + \log s)\} / \log s] \\ &= 1/p + \lim_{s \rightarrow \infty} \log\{1 + \log s\} / \log s = 1/p \end{aligned}$$

Since  $1/p$  is strictly less than 1 and  $\Psi(t)$  is nondecreasing, we take  $\alpha$  such that  $\bar{r}(\Psi) = \frac{1}{p} < \alpha \leq 1$ . We then have  $D_\Psi(s) \leq s^\alpha \leq s$  for  $s > 1$  hence there is a concave function  $\bar{\Psi}(t)$  by Property 1.6. When  $s \leq 1$ , we get

$$\frac{\Psi(st)}{\Psi(t)} = \begin{cases} s^{1/p} & t \leq 1 \\ s^{1/p}(1 + \log t) & 1 < t \leq 1/s \\ s^{1/p}(1 + \log st)/(1 + \log t) & 1/s < t. \end{cases}$$

Thus, by computation,

$$\begin{aligned} D_\Psi(s) &= \sup_{0 < t} \Psi(st)/\Psi(t) \\ &= \sup_{0 < t \leq 1} \Psi(st)/\Psi(t) = s^{1/p}. \end{aligned}$$

Therefore,  $\underline{r}(\Psi) = 1/p$ . Finally,  $\bar{\Psi}(t)$  is not equivalent to  $t^{1/p}$  since  $\Psi(t)$  is not equivalent to  $t^{1/p}$ . Also,  $\bar{\Psi}(t)$  is  $p$ -power concave by Lemma 2.2.

**Theorem 2.5.** *Let  $\varphi(t)$  be a  $p$ -power quasiconcave function with a constant  $C$  and a lower index  $\underline{r}(\varphi)$ . Then,  $1/\underline{r}(\varphi) \leq p$ .*

*Proof.* By taking  $t$  for each  $a_i$  in (2.1), we get

$$D_\varphi(1/n) \leq C^{1/p} (1/n)^{1/p}.$$

Dividing both sides by  $\log(1/n) < 0$ , we get

$$\frac{D_\varphi(1/n)}{\log(1/n)} \geq \frac{(1/p)\log C}{\log(1/n)} + \frac{(1/p)\log(1/n)}{\log(1/n)}.$$

Letting  $n$  go to  $\infty$ , we have  $\underline{r}(\varphi) \geq 1/p$ . □

From this result, we know that if  $\varphi(t)$  is  $p$ -power quasiconcave, then we have  $p \geq 1$  by Property 1.5. We now consider conditions under which a given quasiconcave function  $\varphi(t)$  is  $p$ -power quasiconcave.

**Lemma 2.6.** *Let  $\varphi(t)$  be a positive quasiconcave function with  $\underline{r}(\varphi) > 0$ . Then there exists a constant  $1 \leq C$  and a positive quasiconcave differentiable function  $\Psi(t)$  with  $\Psi(0) = 0$  such that*

$$\varphi(t) \leq \Psi(t) \leq C\varphi(t)$$

and

$$\frac{1}{C} \frac{\Psi(t)}{t} \leq \frac{d\Psi}{dt}(t) \leq \frac{\Psi(t)}{t}.$$

*Proof.* Define  $\Psi(t) = \int_0^t \frac{\varphi(x)}{x} dx$  and let  $\tilde{\varphi}(x) = \frac{x}{\varphi(x)}$ .

$\Psi(t)$  is nondecreasing and we show that  $\frac{\Psi(t)}{t}$  is nonincreasing. We have, for  $t_1 \leq t_2$ ,

$$\begin{aligned} \frac{\Psi(t_1)}{t_1} &= \frac{1}{t_1} \int_0^{t_1} \frac{\varphi(x)}{x} dx \\ &\geq \frac{1}{t_1} \int_0^{t_1} \frac{\varphi(t_1)}{t_1} dx = \frac{\varphi(t_1)}{t_1}. \end{aligned}$$

Thus,

$$\begin{aligned} \Psi(t_2) &= \int_0^{t_2} \frac{\varphi(x)}{x} dx \\ &= \int_0^{t_1} \frac{\varphi(x)}{x} dx + \int_{t_1}^{t_2} \frac{\varphi(x)}{x} dx \\ &\leq \Psi(t_1) + \frac{\varphi(t_1)}{t_1} (t_2 - t_1) \\ &\leq \Psi(t_1) + \frac{\Psi(t_1)}{t_1} (t_2 - t_1) \\ &= \frac{t_2}{t_1} \Psi(t_1). \end{aligned}$$

Therefore, we have shown that  $\Psi(t)$  is quasicave. We now show that  $\Psi(t)$  is equivalent to  $\varphi(t)$ . Since  $\underline{r}(\varphi) > 0$ , we know that  $\bar{r}(\tilde{\varphi}) < 1$  by Property 1.5. We take  $\alpha$  such that  $\bar{r}(\tilde{\varphi}) < \alpha < 1$ . By definition of  $\bar{r}(\tilde{\varphi})$ , there exists  $M > 1$  such that

$$D_{\tilde{\varphi}}(s) \leq s^\alpha \quad \text{for } s > M.$$

Now, let  $t$  be fixed and let  $\beta = tM$ . If  $x$  is in  $(0, t)$  we have  $M < \frac{\beta}{x}$ . Thus we get

$$\frac{\tilde{\varphi}(\beta)}{\tilde{\varphi}(x)} \leq D_{\tilde{\varphi}}\left(\frac{\beta}{x}\right) \leq \left(\frac{\beta}{x}\right)^\alpha$$

and

$$\frac{\tilde{\varphi}(t)}{\tilde{\varphi}(\beta)} \leq 1.$$

We then have

$$\begin{aligned}
 \Psi(t) &= \int_0^t \frac{1}{\tilde{\varphi}(x)} dx = \frac{1}{\tilde{\varphi}(t)} \cdot \frac{\tilde{\varphi}(t)}{\tilde{\varphi}(\beta)} \int_0^t \frac{\tilde{\varphi}(\beta)}{\tilde{\varphi}(x)} dx \\
 &\leq \frac{1}{\tilde{\varphi}(t)} \int_0^{\frac{\beta}{M}} \left(\frac{\beta}{x}\right)^\alpha dx \\
 &= \frac{\beta^\alpha}{\tilde{\varphi}(t)} \frac{1}{1-\alpha} \left(\frac{\beta}{M}\right)^{1-\alpha} \\
 &= \frac{\beta}{1-\alpha} \cdot \frac{1}{M^{1-\alpha}} \cdot \frac{1}{\tilde{\varphi}(t)} \\
 &= \frac{M^\alpha}{1-\alpha} \frac{t}{\tilde{\varphi}(t)} = \frac{M^\alpha}{1-\alpha} \varphi(t).
 \end{aligned}$$

Now take  $C = \frac{M^\alpha}{1-\alpha}$  then we have the right inequality.

For the left inequality, the nonincreasing property of  $\frac{\varphi(t)}{t}$  gives

$$\varphi(t) = \frac{t}{\tilde{\varphi}(t)} = \int_0^t \frac{dx}{\tilde{\varphi}(t)} \leq \int_0^t \frac{dx}{\tilde{\varphi}(x)} = \Psi(t).$$

By definition of  $\Psi(t)$ , we have  $\frac{d\Psi(t)}{dt} = \frac{\varphi(t)}{t}$ . Thus,

$$\frac{1}{C} \Psi(t) \leq \varphi(t) = t \cdot \frac{d\Psi(t)}{dt} \leq \Psi(t)$$

and we get the result by dividing the above inequality by  $t$ . □

**Theorem 2.7.** *Let  $\varphi(t)$  be a quasiconcave function with  $\underline{r}(\varphi) > 0$ . Then there exists a finite  $p$  such that  $\varphi(t)$  becomes a  $p$ -power concave function.*

*Proof.* By Lemma 2.2, it is enough to show that  $\Psi(t) = \int_0^t \frac{\varphi(x)}{x} dx$  satisfies (2.1) since  $\Psi(t)$  is equivalent to  $\varphi(t)$  by Lemma 2.6. First, we want to show that  $\frac{\Psi^p(t)}{t}$  is nondecreasing for some finite  $p$ . Note that  $\Psi(t)$  is differentiable. We then have

$$\begin{aligned}
 \frac{d}{dt} \left( \frac{\Psi^p(t)}{t} \right) &= \frac{p\Psi^{p-1}(t)\Psi'(t) \cdot t - \Psi^p(t)}{t^2} \\
 &= \frac{\Psi^{p-1}(t)}{t^2} \{p\Psi'(t) \cdot t - \Psi(t)\} \\
 &\geq \frac{\Psi^{p-1}(t)}{t^2} \left\{ \frac{p\Psi(t)}{C} - \Psi(t) \right\} \quad \text{by Lemma 2.6} \\
 &= \frac{\Psi^p(t)}{t^2} \left( \frac{p}{C} - 1 \right).
 \end{aligned}$$

Hence, for  $1 \leq C < p < \infty$ ,  $\frac{d}{dt} \left( \frac{\Psi^p(t)}{t} \right) \geq 0$  and so  $\frac{\Psi^p(t)}{t}$  is nondecreasing. We now show that  $\Psi(t)$  satisfies (2.1).

$$\begin{aligned} \Psi^p \left( \sum_{i=1}^n a_i \right) &= \sum_{i=1}^n a_i \left\{ \Psi^p \left( \sum_{i=1}^n a_i \right) / \sum_{i=1}^n a_i \right\} \\ &\geq \sum_{i=1}^n a_i \{ \Psi^p(a_i) / a_i \} \\ &= \sum_{i=1}^n \Psi^p(a_i). \end{aligned}$$

□

**Theorem 2.8.** *Let  $\varphi(t)$  be the fundamental function of r.i. space  $E$ .*

*i) Suppose that  $\varphi(t)$  is  $p_1$ -power quasiconcave. If  $E$  satisfies an upper  $q$ -estimate, then  $q \leq p_1$ .*

*ii) Suppose that  $\tilde{\varphi}(t) = \frac{t}{\varphi(t)}$  is equivalent to  $\Psi(t)$  which is  $p_2$ -power quasiconcave. If  $E$  satisfies a lower  $q$  estimate, then  $q \geq \tilde{p}_2$  where  $1/p_2 + 1/\tilde{p}_2 = 1$ .*

*Proof.* i) Since  $\varphi(t)$  is  $p_1$ -power quasiconcave and the fundamental function of r.i.-space, there exist constant  $C_1$  and  $C_2$  such that

$$\frac{1}{C_1} n^{1/p_1} \leq \varphi(n) \leq C_2 n^{1/q}$$

for all integers  $n$ . Thus we have  $q \leq p_1$ .

ii) Since  $\tilde{\varphi}(t)$  is equivalent to the  $p_2$ -power concave function, there exists  $C_3$  such that

$$\tilde{\varphi}\left(\frac{1}{n}\right) = \left(\frac{1}{n}\right) \varphi\left(\frac{1}{n}\right) \leq C_3 \left(\frac{1}{n}\right)^{1/p_2}.$$

By the lower  $q$  estimate property, we have  $C_4$  such that

$$\varphi\left(\frac{1}{n}\right) \leq C_4 \left(\frac{1}{n}\right)^{1/q}.$$

Thus,

$$\frac{1}{C_3} \left(\frac{1}{n}\right)^{1-(1/p_2)} \leq \varphi\left(\frac{1}{n}\right) \leq C_4 \left(\frac{1}{n}\right)^{1/q}$$

for all integers. Thus, we have  $1/q \leq 1 - (1/p_2) = 1/\tilde{p}_2$ .

□

### 3. MAIN RESULT

We now apply the  $p$ -power quasiconcave property to some r.i. space like Lorentz space and Marcinkiewicz space. Although there are several versions of these spaces, we take Sharpley's version with minor modifications [7]. Let  $f$  be a real valued function defined on  $[0, \infty)$  with Lebesgue measure  $\mu$ . The distribution function of  $f$ , denoted by  $\lambda_f(t)$ , is defined by  $\mu(\{x : |f(x)| > t\})$ . We define  $f^*(t)$



is the non-increasing, right continuous function which is equimeasurable with  $f$  and  $f^{**}(t) = \frac{1}{t} \int_0^t f^*(x)dx$ .

For an explicit formula of  $f^*(t)$ , we have  $f^*(t) = \inf\{y \geq 0 : \lambda_f(y) \leq t\}$  [1].

**Definition 3.1.** Let  $\varphi(t)$  be a quasiconcave function on  $[0, \infty)$  with  $0 < \underline{r}(\psi) \leq \bar{r}(\psi) < 1$  and  $1 \leq q < \infty$ .

The Lorentz space  $\Lambda_{\psi,q}$  is the set of all measurable functions  $f$  such that  $f^*$  exists and

$$\|f\|_{\Lambda_{\psi,q}} = \left\{ \int_0^\infty (f^{**}(t)\psi(t))^q \frac{dt}{t} \right\}^{1/q} < \infty.$$

The Marcinkiewicz space  $M_\psi$  is the set of all measurable functions  $f$  such that

$$\|f\|_{M_\psi} = \sup_{t>0} \psi(t)f^{**}(t) < \infty.$$

By definition, we can easily show that  $\|f\|_{\Lambda_{\varphi,q}}$  and  $\|f\|_{M_\varphi}$  are equivalent to  $\|f\|_{\Lambda_{\psi,q}}$  and  $\|f\|_{M_\psi}$  respectively when  $\varphi$  and  $\psi$  are equivalent. It is well known that  $\int_0^t f^*(x)dx = \sup_{\mu(E)=t} \int_A |f|d\mu$  (See page 64 in [3]). Thus we have an alternate form of  $\|f\|_{M_\varphi}$  :

$$\|f\|_{M_\psi} = \sup_{\mu(E)>0} \frac{\psi(\mu(E))}{\mu(E)} \int_A |f|d\mu.$$

For the space  $\Lambda_{\psi,q}$ , we have also useful form of its norm

$$\|f\|_{\Lambda_{\psi,q}}^* = \left\{ \int_0^\infty [f^*(t)\psi(t)]^q \frac{dt}{t} \right\}^{1/q},$$

which is equivalent to  $\|f\|_{\Lambda_{\psi,q}}$  (see [7, Theorem 2.3]).

We now introduce some functional which is convenient to compute the norm in  $\Lambda_{\Psi,q}$ . We modify the proof in Sharpley's version of Lorentz space (see [3, 7]).

**Lemma 3.2.** *i) Let  $f$  be an element in  $\Lambda_{\psi,q}$ . Then, the functional*

$$\|f\|_{\Lambda_{\psi,q}}^0 = \left\{ \int_0^\infty ((f^*(t))^q d\psi^q) \right\}^{1/q}$$

*is equivalent to  $\|f\|_{\Lambda_{\psi,q}}$ .*

*ii) The fundamental function  $\varphi_{\Lambda_{\psi,q}}$  in  $\Lambda_{\psi,q}$  is equivalent to  $\psi(t)$ .*

*iii) If  $f \in \Lambda_{\psi,q}$ , we have*

$$\|f\|_{\Lambda_{\psi,q}}^0 = \left\{ q \int_0^\infty y^{q-1} [\psi^q(\lambda_f(y))] dy \right\}^{1/q}.$$

*Proof.* i) By Lemma 2.6,  $\psi(t)$  has an equivalent quasiconcave differentiable function  $\Psi(t) = \int_0^t \frac{\psi}{x} dx$  and we can renorm  $\Lambda_{\psi,q}$  by replacing  $\psi(t)$  with  $\Psi(t)$ . Thus we may assume, for some constant  $C$ ,

$$\frac{1}{C} \frac{\psi(t)}{t} \leq \frac{d\psi}{dt}(t) \leq \frac{\psi(t)}{t}.$$

Then, by the improper Stieltjes integral, we have

$$\begin{aligned} \left\{ \|f\|_{\Lambda_{\psi,q}}^0 \right\}^q &= \psi(+0)(f^*(+0))^q + \int_{+0}^{\infty} (f^*(t))^q d\psi^q, \\ &\text{since } 0 < r(\psi), \text{ it is easy to get } \lim_{s \rightarrow 0} \psi(s) = 0, \\ &= \int_0^{\infty} (f^*(t))^q q\psi^{q-1}(t) \frac{d\psi(t)}{dt} t \frac{dt}{t}. \end{aligned}$$

Since  $\frac{q}{C}\psi^q(t) \leq q\psi^{q-1} \cdot \frac{d\psi(t)}{dt} \cdot t \leq q\psi^q(t)$ , we have

$$\frac{q}{C} \|f\|_{\Lambda_{\psi,q}}^* \leq \|f\|_{\Lambda_{\psi,q}}^0 \leq q \|f\|_{\Lambda_{\psi,q}}^*.$$

Thus  $\|f\|_{\psi_{\Lambda,q}}^0$  is equivalent to  $\|f\|_{\Lambda_{\psi,q}}$ .

- ii) Since  $\|\chi_{[0,t]}\|_{\Lambda_{\psi,q}}^0 = \left\{ \int_0^t d\psi^q \right\}^{1/q} = \psi(t)$ , the fundamental function  $\varphi_{\Lambda_{\psi,q}}$  is equivalent to  $\psi(t)$ .
- iii) Since simple functions are dense in  $\Lambda_{\psi,q}$  (see [2, 7]), we show the result for a simple function  $f = \sum_{i=1}^n a_i \chi_{A_i}$ , where  $\{A_i\}$  are pairwise disjoint measurable sets. Without loss of generality, we may assume  $a_1 > \dots > a_n$ . Define  $d_i = \sum_{j=1}^i \mu(A_j)$  and  $d_0 = 0$ . Then, we have

$$\lambda_f(y) = \begin{cases} d_i & a_{i+1} \leq y < a_i \\ 0 & a_1 \leq y \end{cases}$$

Hence,

$$\begin{aligned} \|f\|_{\Lambda_{\psi,q}}^0 &= \left\{ \int_0^{\infty} (f^*(t))^q d\psi^q \right\}^{1/q} \\ &= \left\{ \sum_{i=1}^n a_i^q [\psi^q(d_i) - \psi^q(d_{i-1})] \right\}^{1/q} \end{aligned}$$

On the other hand,

$$\begin{aligned} &\left( q \int_0^{\infty} y^{q-1} \{\psi[\lambda_f(y)]\}^q dy \right)^{1/q} \\ &= \left\{ \psi^q(d_n) a_n^q + \psi^q(d_{n-1})(a_{n-1}^q - a_n^q) + \dots + \psi^q(d_1)(a_1^q - a_2^q) \right\}^{1/q} \\ &= \left( \sum_{i=1}^n a_i^q [\psi^q(d_i) - \psi^q(d_{i-1})] \right)^{1/q} \\ &= \|f\|_{\Lambda_{\psi,q}}^0 \end{aligned}$$

□

**Theorem 3.3.** *If  $\psi(t)$  is  $p$ -power quasiconcave with constant  $C$ , the space  $\Lambda_{\psi,q}$  satisfies a lower- $p$ -estimate for  $1 \leq q < p$ .*

*Proof.* Since a lower estimate is metric invariant, we use the equivalent functional  $\|f\|_{\Lambda_{\psi,q}}^0$  by Lemma 3.2. Let  $\{f_i\}$  be elements in  $\Lambda_{\psi,q}$  with pairwise disjoint support. Note that  $\sum_{i=1}^n \lambda_{f_i}(y) = \lambda \sum_{i=1}^n f_i(y)$ .

By Lemma 3.2, we have

$$\begin{aligned}
 \left\{ \sum_{i=1}^n \left( \|f_i\|_{\Lambda_{\psi,q}}^0 \right)^p \right\}^{1/p} &= \left( \sum_{i=1}^n \left\{ \int_0^\infty \psi^q(\lambda_{f_i}(y)) dy^q \right\}^{p/q} \right)^{1/p} \\
 &= \left( \sum_{i=1}^n \left\{ \int_0^\infty [\psi^p(\lambda_{f_i}(y))]^{q/p} dy^q \right\}^{p/q} \right)^{1/p} \\
 &\leq \left( \int_0^\infty \left\{ \sum_{i=1}^n \psi^p(\lambda_{f_i}(y)) \right\}^{q/p} dy^q \right)^{1/q} \\
 &\quad \text{since } \sum_{i=1}^n \|f_i\|_{L_r} \leq \left\| \sum_{i=1}^n f_i \right\|_{L_r} \text{ when } r < 1 \\
 &\leq \left( \int_0^\infty \left\{ C\psi^p \left[ \sum_{i=1}^n \lambda_{f_i}(y) \right] \right\}^{q/p} dy^q \right)^{1/q} \\
 &\quad \text{since } \psi \text{ is } p\text{-power quasiconcave} \\
 &= C^{1/p} \left( \int_0^\infty \left\{ \psi \left[ \sum_{i=1}^n \lambda_{f_i}(y) \right] \right\}^q dy^q \right)^{1/q} \\
 &= C^{1/p} \left( \int_0^\infty \left\{ \psi(\lambda_{\sum_{i=1}^n f_i}(y)) \right\}^q dy^q \right)^{1/q} \\
 &= C^{1/p} \left\| \sum_{i=1}^n f_i \right\|_{\Lambda_{\psi,q}}^0
 \end{aligned}$$

□

**Theorem 3.4.** *Let  $\psi(t)$  be quasiconcave such that  $\tilde{\psi}(t) = \frac{t}{\psi(t)}$  is  $q$ -power quasiconcave with constant  $C$ . Then  $M_\psi$  satisfies an upper  $p$ -estimate where  $\frac{1}{p} + \frac{1}{q} = 1$ .*

*Proof.* Let  $\{f_i\}_{i=1}^n$  be a set of measurable functions with disjoint supports  $\{E_i\}_{i=1}^n$  respectively. Let  $E$  be any measurable set in  $[0, \infty)$ . Define  $F_i = A \cap E_i$ . Since

each  $E_i$ 's are disjoint, we know  $\sum_{i=1}^n \mu(F_i) \leq \mu(E)$ . Thus

$$\begin{aligned}
\frac{\psi(\mu(E))}{\mu(E)} \int_E \sum_{i=1}^m f_i &= \frac{\psi(\mu(E))}{\mu(E)} \sum_{i=1}^m \int_E f_i \\
&\leq \frac{\psi(\mu(E))}{\mu(E)} \sum \frac{\mu(F_i)}{\psi(\mu(F_i))} \|f_i\|_{M_\psi} \\
&\leq \frac{\psi(\mu(E))}{\mu(E)} \left\{ \sum_{i=1}^n \left( \frac{\mu(F_i)}{\psi(\mu(F_i))} \right)^q \right\}^{1/q} \left\{ \sum_{i=1}^m \|f_i\|_{M_\psi}^p \right\}^{1/p} \\
&\leq \frac{\psi(\mu(E))}{\mu(E)} \left\{ \sum_{i=1}^n \tilde{\psi} \left( \sum_{i=1}^m \mu(F_i) \right)^q \right\}^{1/q} \left\{ \sum_{i=1}^m \|f_i\|_{M_\psi}^p \right\}^{1/p} \\
&= \frac{\psi(\mu(E))}{\mu(E)} \left\{ \sum_{i=1}^n \tilde{\psi} \left( \sum_{i=1}^m \mu(F_i) \right) \right\} \left\{ \sum_{i=1}^m \|f_i\|_{M_\psi}^p \right\}^{1/p} \\
&= C \frac{\psi(\mu(E))}{\mu(E)} \frac{\sum_{i=1}^n \mu(F_i)}{\psi(\sum_{i=1}^n \mu(F_i))} \left\{ \sum_{i=1}^m \|f_i\|_{M_\psi}^p \right\}^{1/p} \\
&\leq C \left\{ \sum_{i=1}^m \|f_i\|_{M_\psi}^p \right\}^{1/p},
\end{aligned}$$

since  $\frac{\psi(t)}{t}$  is nonincreasing and  $\sum_{i=1}^n \mu(F_i) \leq \mu(E)$ .

We thus have

$$\left\| \sum_{i=1}^n f_i \right\|_{M_{\psi_i}} \leq C \left\{ \sum_{i=1}^n \|f_i\|_{M_\psi}^p \right\}^{1/p}.$$

□

The following are easily obtained from the above theorems.

**Corollary 3.5.** *Weak  $L_p$  satisfies an upper  $p$ -estimate.*

**Corollary 3.6.** *Let  $\psi(t)$  be a quasiconcave with  $\bar{r}(\psi) < 1$ . Then there exists  $p$  such that  $1 < p < \infty$  and  $M_\psi$  satisfies an upper  $p$ -estimate.*

**Acknowledgement.** This work was supported by the Inha University Research Grant.

## REFERENCES

1. M. Cwikel, *The dual of weak  $L_p$* , Ann. Inst. Fourier, Grenoble **25** (1975), 81–126
2. R.A. Hunt, *On  $L_{p,q}$  spaces*, Enseignement Math. **12** (1966), 249–276.
3. S. Krein, J. Petunim and E. Semenov, *Interpolation of linear operators*, English transl., Transl. Math. Mono. vol 54, AMS Providence, R.I. 1980

4. J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces II, Function Spaces*, Ergeb. der Math. u. ihr. Grenzgeb, 97, Berlin, Springer, 1979
5. J. Peetree, *Concave majorants of positive functions*, Acta. Math. Acad. Sci. Hungar. **21** (1970), 327–333
6. E.M. Semenov, *Embedding theorems for Banach spaces of measurable functions* (Russian), Dokl. Akad. Nauk SSSR **156** (1964), 1292–1295.
7. R. Sharpley, *Spaces  $\Lambda_\alpha(X)$  and interpolation*, J. Functional Analysis **11** (1972), 479–513
8. M. Zippin, *Interpolation of operators of weak type between rearrangement invariant function spaces*, J. Funct. Anal. **7** (1971), 267–284

<sup>1</sup> DEPARTMENT OF MATHEMATICS EDUCATION, INHA UNIVERSITY, INCHON 402-751, KOREA.

*E-mail address:* [cslee@inha.ac.kr](mailto:cslee@inha.ac.kr)

<sup>2</sup> BANGMOK COLLEGE OF GENERAL EDUCATION, MYONG JI UNIVERSITY, YONG-IN 449-728, KOREA.

*E-mail address:* [kgjo@mju.ac.kr](mailto:kgjo@mju.ac.kr)