

## A FIXED POINT APPROACH TO THE STABILITY OF $\varphi$ -MORPHISMS ON HILBERT $C^*$ -MODULES

GH. ABBASPOUR TABADKAN<sup>1\*</sup> AND M. RAMEZANPOUR<sup>2</sup>

Communicated by J. Chmieliński

ABSTRACT. Let  $E, F$  be two Hilbert  $C^*$ -modules over  $C^*$ -algebras  $A$  and  $B$  respectively. In this paper, by the alternative fixed point theorem, we give the Hyers-Ulam-Rassias stability of the equation

$$\langle U(x), U(y) \rangle = \varphi(\langle x, y \rangle) \quad (x, y \in E),$$

where  $U : E \rightarrow F$  is a mapping and  $\varphi : A \rightarrow B$  is an additive map.

### 1. INTRODUCTION AND PRELIMINARIES

A *pre-Hilbert  $A$ -module* is a right module  $E$  over  $C^*$ -algebra  $A$ , with a map  $\langle \cdot, \cdot \rangle : E \times E \rightarrow A$  which is conjugate linear in the first, linear in its second argument and satisfies

- (i)  $\langle x, ya \rangle = \langle x, y \rangle a \quad (x, y \in E, a \in A)$ ,
- (ii)  $\langle x, y \rangle^* = \langle y, x \rangle \quad (x, y \in E)$ ,
- (iii)  $\langle x, x \rangle \geq 0 \quad (x \in E)$ ,
- (iv)  $\langle x, x \rangle = 0 \Rightarrow x = 0$ .

A *Hilbert  $A$ -module* (briefly *Hilbert module*) is a pre-Hilbert  $A$ -module that is complete in the norm defined by  $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$ . For more details about Hilbert modules see [12].

Let  $E, F$  be two Hilbert modules over  $C^*$ -algebras  $A$  and  $B$  respectively and  $\varphi : A \rightarrow B$  be a map. A mapping  $U : E \rightarrow F$  is called a  $\varphi$ -morphism if

$$\langle U(x), U(y) \rangle = \varphi(\langle x, y \rangle) \quad (x, y \in E).$$

---

*Date:* Received: 25 July 2010; Revised: 22 September 2010; Accepted: 29 September 2010.

\* Corresponding author.

2010 *Mathematics Subject Classification.* Primary 39B82; Secondary 46L08.

*Key words and phrases.* Hyers-Ulam-Rassias stability, Hilbert  $C^*$ -modules.

This kind of mappings were introduced by Bakić and Guljaš [3]. The first author together with Moslehian and Niknam [1] used this kind of mappings to introduce dynamical systems on Hilbert modules. Also Abbaspour and Skeide in [2] investigated the relation between  $\varphi$ -morphisms, where they called them generalized module mappings, and ternary homomorphisms.

The stability problem of functional equations had been first raised by Ulam [18] by the following question: For what metric groups  $G$  is it true that an  $\epsilon$ -automorphism of  $G$  is necessarily near to a strict automorphism? A partial answer to the above question has been given as follows. Suppose  $E_1$  and  $E_2$  are two real Banach spaces and  $f : E_1 \rightarrow E_2$  is a mapping. If there exist  $\delta \geq 0$  and  $p \geq 0$ ,  $p \neq 1$  such that

$$\|f(x+y) - f(x) - f(y)\| \leq \delta(\|x\|^p + \|y\|^p)$$

for all  $x, y \in E_1$ , then there is a unique additive mapping  $T : E_1 \rightarrow E_2$  such that

$$\|f(x) - T(x)\| \leq \frac{2\delta\|x\|^p}{|2 - 2^p|} \quad (x \in E_1).$$

This result is called the Hyers-Ulam-Rassias stability of the additive Cauchy equation. Indeed Hyers [10] obtained the above result for  $p = 0$ . Then Rassias [17] generalized the result of Hyers to the case where  $0 \leq p < 1$ . Gajda [9] solved the problem for  $p > 1$  and gave an example that a similar result does not hold for  $p = 1$ . For the case  $p < 0$ , recently Lee [13] has shown that  $f$  should be an additive map. Thus the Hyers-Ulam-Rassias stability of the additive Cauchy equation holds for  $p \in \mathbb{R} \setminus \{1\}$ .

Let  $X$  be a set. A function  $d : X \times X \rightarrow [0, \infty]$  is called a generalized metric on  $X$  if  $d$  satisfies

- (1)  $d(x, y) = 0$  if and only if  $x = y$ ,
- (2)  $d(x, y) = d(y, x)$ ,
- (3)  $d(x, y) \leq d(x, z) + d(z, x)$ .

Generalized metric space  $(X, d)$  is called complete if each Cauchy sequence converges in  $X$ .

In 2003, Radu [16] employed the following theorem to prove the stability of a Cauchy functional equation. Later many authors, [7, 11, 14, 15] used this strategy to give the stability of functional equations. Before stating the theorem we recall that a mapping  $J : X \rightarrow X$  is called a strictly contractive operator with the Lipschitz constant  $L$ , if

$$d(J(x), J(y)) < Ld(x, y) \quad (x, y \in X).$$

**Theorem 1.1.** ([8]) *Let  $(X, d)$  be a generalized complete metric space and  $J : X \rightarrow X$  be a strictly contractive operator with the Lipschitz constant  $L < 1$ . If there exists a nonnegative integer  $k$  such that  $d(J^{k+1}x, J^kx) < \infty$  for some  $x \in X$ , then the following are true:*

- (a) *The sequence  $\{J^n x\}$  converges to a fixed point  $x^*$  of  $J$ ,*
- (b)  *$x^*$  is the unique fixed point of  $J$  in*

$$X^* = \{y \in X \mid d(J^k x, y) < \infty\},$$

(c) if  $y \in X^*$ , then

$$d(y, x^*) \leq \frac{1}{1-L} d(Jy, y).$$

In [4], Badora and Chmieliński, investigated the stability and superstability of inner product preserving mappings on Hilbert spaces. After then Chmieliński and Moslehian [6] investigated this problem in the framework of Hilbert  $C^*$ -modules; see also [5]. We mention that each  $\varphi$ -morphism is in fact a mapping preserving inner product modulo  $\varphi$ . In this paper, by using the alternative fixed point theorem for generalized metric spaces, the stability of  $\varphi$ -morphisms on Hilbert  $C^*$ -modules is considered. Throughout the paper we assume that  $E$  and  $F$  are two Hilbert  $C^*$ -modules over  $C^*$ -algebras  $A$  and  $B$  respectively and  $\varphi : A \rightarrow B$  is an additive map.

## 2. MAIN RESULTS

**Definition 2.1.** A mapping  $U : E \rightarrow F$  is called an approximate  $\varphi$ -morphism if there exists a control function  $\tau : E^2 \rightarrow \mathbb{R}$  such that

$$\|\langle U(x), U(y) \rangle - \varphi(\langle x, y \rangle)\| \leq \tau(x, y)$$

holds for each  $x, y \in E$ .

As a consequence of Theorem 2.4 we will show that under some conditions on control function  $\tau$  each approximate  $\varphi$ -morphism is near to a  $\varphi$ -morphism.

**Example 2.2.** We know that each  $C^*$ -algebra  $A$  is a Hilbert  $C^*$ -module over itself with the inner product defined by  $\langle a, b \rangle = a^*b$ . Let  $A$  be a unital  $C^*$ -algebra,  $a \in A$ ,  $\epsilon = \|a^*a - 1\|$  and  $\varphi : A \rightarrow A$  be a  $*$ -homomorphism. If we define  $U(x) = a\varphi(x)$  then we have

$$\begin{aligned} \|\langle U(x), U(y) \rangle - \varphi(\langle x, y \rangle)\| &= \|\varphi(x^*)a^*a\varphi(y) - \varphi(x^*)\varphi(y)\| \\ &= \|\varphi(x^*)(a^*a - 1)\varphi(y)\| \\ &\leq \epsilon\|x\|\|y\| \\ &\leq \frac{\epsilon}{2}(\|x\|^2 + \|y\|^2) \end{aligned}$$

If  $a$  is a unitary element then  $U$  is a  $\varphi$ -morphism, otherwise  $U$  is an approximate  $\varphi$ -morphism with control function  $\tau(x, y) = \frac{\epsilon}{2}(\|x\|^2 + \|y\|^2)$ .

**Lemma 2.3.** If  $U : E \rightarrow F$  is a mapping such that  $\|U(x+y) - U(x) - U(y)\| \leq \tau(x, y)$  for some control function  $\tau : E^2 \rightarrow \mathbb{R}$  and there is  $0 < L < 1$  with  $\tau(2x, 2y) \leq 2L\tau(x, y)$ , then there exists a unique additive map  $\psi : E \rightarrow F$  such that  $\|U(x) - \psi(x)\| \leq \frac{1}{2-2L}\tau(x, x)$ .

*Proof.* Let  $X = \{g : E \rightarrow F : g \text{ is a mapping}\}$  and define

$$d(g, h) = \inf\{c \geq 0 : \|g(x) - h(x)\| \leq c\tau(x, x) \quad \forall x \in E\},$$

for  $g, h \in X$ . Then  $(X, d)$  is a complete generalized metric space. Now we consider the mapping  $J : X \rightarrow X$  by  $J(g)(x) = \frac{1}{2}g(2x)$ . We can write for any  $g, h \in X$ ,

$$\|g(x) - h(x)\| \leq d(g, h)\tau(x, x) \quad (x \in E),$$

therefore for  $x \in E$ ,

$$\|J(g)(x) - J(h)(x)\| = \left\| \frac{1}{2}g(2x) - \frac{1}{2}h(2x) \right\| \leq \frac{1}{2}d(g, h)\tau(2x, 2x) \leq Ld(g, h)\tau(x, x).$$

Hence  $d(J(g), J(h)) \leq Ld(g, h)$ . Since  $d(J(U), U) \leq \frac{1}{2} < \infty$ , Theorem 1.1 implies that

- (i)  $J$  has a unique fixed point  $\psi : E \rightarrow F$  in the set  $X^* = \{g \in X : d(g, U) < \infty\}$ .
- (ii)  $d(J^n(U), \psi) \rightarrow 0$  as  $n \rightarrow \infty$ . This implies that  $\lim_{n \rightarrow \infty} \frac{U(2^n x)}{2^n} = \psi(x)$  for all  $x \in E$ .
- (iii)  $d(U, \psi) \leq \frac{d(U, J(U))}{1-L} \leq \frac{1}{2-2L}$ . That is,  $\|U(x) - \psi(x)\| \leq \frac{1}{2-2L}\tau(x, x)$  for all  $x \in E$ .

Moreover, for each  $x, y \in E$  we have,

$$\begin{aligned} \|\psi(x+y) - \psi(x) - \psi(y)\| &= \lim_{n \rightarrow \infty} \left\| \frac{U(2^n(x+y))}{2^n} - \frac{U(2^n x)}{2^n} - \frac{U(2^n y)}{2^n} \right\| \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{2^n} \tau(2^n x, 2^n y) \\ &\leq \lim_{n \rightarrow \infty} L^n \tau(x, y) \\ &= 0. \end{aligned}$$

Hence  $\psi$  is an additive map. Now let  $\psi' : E \rightarrow F$  be another additive map such that

$$\|U(x) - \psi'(x)\| \leq \frac{1}{2-2L}\tau(x, x) \quad (x \in E),$$

so  $J(\psi') = \psi'$  and  $d(U, \psi') \leq \frac{1}{2-2L}$ . In other words  $\psi'$  is a fixed point of  $J$  in  $X^*$ . Thus  $\psi' = \psi$ .  $\square$

**Theorem 2.4.** *Let  $U : E \rightarrow F$  be a mapping and  $\varphi : A \rightarrow B$  be an additive map such that for some control function  $\rho : E^2 \rightarrow \mathbb{R}$ ,  $\|\langle Ux, Uy \rangle - \varphi(\langle x, y \rangle)\| \leq \rho(x, y)$  for all  $x, y \in E$ . Let*

$$\begin{aligned} \tau(x, y) &= \left( \rho(x+y, x+y) + \rho(x+y, x) + \rho(x, x+y) + \rho(x+y, y) + \rho(y, x+y) \right. \\ &\quad \left. + \rho(x, x) + \rho(y, y) + \rho(x, y) + \rho(y, x) \right)^{\frac{1}{2}} \end{aligned}$$

and suppose there is  $0 < L < 1$  such that  $\tau(2x, 2y) \leq 2L\tau(x, y)$ . Then there exists a unique  $\varphi$ -morphism  $T : E \rightarrow F$  such that  $\|U(x) - T(x)\| \leq \frac{1}{2-2L}\tau(x, x)$  for all  $x \in X$ .

*Proof.* For all  $x, y, z \in E$  we have

$$\begin{aligned}
& \|\langle U(x+y) - U(x) - U(y), U(z) \rangle\| \\
&= \|\langle U(x+y) - U(x) - U(y), U(z) \rangle - \varphi(\langle x+y, z \rangle) + \varphi(\langle x, z \rangle) \\
&\quad + \varphi(\langle y, z \rangle)\| \\
&\leq \|\langle U(x+y), U(z) \rangle - \varphi(\langle x+y, z \rangle)\| + \|\langle U(x), U(z) \rangle - \varphi(\langle x, z \rangle)\| \\
&\quad + \|\langle U(y), U(z) \rangle - \varphi(\langle y, z \rangle)\| \\
&\leq \rho(x+y, z) + \rho(x, z) + \rho(y, z).
\end{aligned}$$

Thus

$$\begin{aligned}
& \|U(x+y) - U(x) - U(y)\|^2 \\
&= \|\langle U(x+y) - U(x) - U(y), U(x+y) - U(x) - U(y) \rangle\| \\
&\leq \|\langle U(x+y) - U(x) - U(y), U(x+y) \rangle\| \\
&\quad + \|\langle U(x+y) - U(x) - U(y), U(x) \rangle\| \\
&\quad + \|\langle U(x+y) - U(x) - U(y), U(y) \rangle\| \\
&\leq \rho(x+y, x+y) + \rho(x, x+y) + \rho(y, x+y) + \rho(x+y, x) + \rho(x, x) \\
&\quad + \rho(y, x) + \rho(x+y, y) + \rho(x, y) + \rho(y, y).
\end{aligned}$$

It follows that

$$\|U(x+y) - U(x) - U(y)\| \leq \tau(x, y).$$

By Lemma 2.3, there is a unique additive map  $T : E \rightarrow F$  such that

$$\|U(x) - T(x)\| \leq \frac{1}{2-2L}\tau(x, x) \quad (x \in E).$$

Then

$$T(x) = \lim_{n \rightarrow \infty} \frac{U(2^n x)}{2^n}.$$

Now for each  $x, y \in E$  we have

$$\begin{aligned}
\|\langle Tx, Ty \rangle - \varphi(\langle x, y \rangle)\| &= \lim_{n \rightarrow \infty} \frac{1}{4^n} \|\langle U(2^n x), U(2^n y) \rangle - \varphi(\langle 2^n x, 2^n y \rangle)\| \\
&\leq \lim_{n \rightarrow \infty} \frac{1}{4^n} \rho(2^n x, 2^n y) \leq \lim_{n \rightarrow \infty} \frac{1}{4^n} \tau(2^n x, 2^n y)^2 \\
&= \lim_{n \rightarrow \infty} \left( \frac{1}{2^n} \tau(2^n x, 2^n y) \right)^2 \leq \left( \lim_{n \rightarrow \infty} L^n \tau(x, y) \right)^2 \\
&= 0.
\end{aligned}$$

This shows that  $T$  is a  $\varphi$ -morphism. Since each  $\varphi$ -morphism is an additive map Lemma 2.3 implies that  $T$  is the unique  $\varphi$ -morphism as desired.  $\square$

One can replace the condition  $\tau(2x, 2y) \leq 2L\tau(x, y)$  on the control function  $\tau$  by

$$\tau(x, y) \leq \frac{1}{2}L\tau(2x, 2y)$$

and obtain the following results.

**Lemma 2.5.** *If  $U : E \rightarrow F$  is a mapping such that  $\|U(x+y) - U(x) - U(y)\| \leq \tau(x, y)$  for some control function  $\tau : E^2 \rightarrow \mathbb{R}$  and there is  $0 < L < 1$  with  $\tau(x, y) \leq \frac{1}{2}L\tau(2x, 2y)$ , then there exists a unique additive map  $\psi : E \rightarrow F$  such that  $\|U(x) - \psi(x)\| \leq \frac{L}{2-2L}\tau(x, x)$ .*

**Theorem 2.6.** *Let  $U : E \rightarrow F$  be a mapping and  $\varphi : A \rightarrow B$  be an additive map such that for some control function  $\rho : E^2 \rightarrow \mathbb{R}$ ,  $\|\langle Ux, Uy \rangle - \varphi(\langle x, y \rangle)\| \leq \rho(x, y)$  for all  $x, y \in E$ . Let*

$$\begin{aligned} \tau(x, y) = & \left( \rho(x+y, x+y) + \rho(x+y, x) + \rho(x, x+y) + \rho(x+y, y) + \rho(y, x+y) \right. \\ & \left. + \rho(x, x) + \rho(y, y) + \rho(x, y) + \rho(y, x) \right)^{\frac{1}{2}} \end{aligned}$$

and suppose there is  $0 < L < 1$  such that  $\tau(x, y) \leq \frac{1}{2}L\tau(2x, 2y)$ . Then there exists a unique  $\varphi$ -morphism  $T : E \rightarrow F$  such that  $\|U(x) - T(x)\| \leq \frac{L}{2-2L}\tau(x, x)$  for all  $x \in X$ .

For a real number  $p$  let  $E_p$  denote either the whole space  $E$  if  $p \geq 0$  or  $E \setminus \{0\}$  if  $p < 0$ .

**Corollary 2.7.** *Let  $U : E \rightarrow F$  be a mapping and  $\varphi : A \rightarrow B$  be an additive map such that for some  $p \neq 2$ ,*

$$\|\langle Ux, Uy \rangle - \varphi(\langle x, y \rangle)\| \leq c(\|x\|^p + \|y\|^p) \quad (x, y \in E_p).$$

Then there exists a unique  $\varphi$ -morphism  $T : E \rightarrow F$  such that

$$\|U(x) - T(x)\| \leq \frac{\sqrt{6c(2^p + 2)}}{|2 - 2^{\frac{p}{2}}|} \|x\|^{\frac{p}{2}} \quad (x \in E_p).$$

*Proof.* Define  $\rho : E_p \times E_p \rightarrow \mathbb{R}$  by  $\rho(x, y) = c(\|x\|^p + \|y\|^p)$ , then apply Theorems 2.4 and 2.6 with

$$\tau(x, y) = \sqrt{6c(\|x+y\|^p + \|x\|^p + \|y\|^p)}$$

□

*Remark 2.8.* If  $E$  and  $F$  are two Hilbert  $C^*$ -modules over the same  $C^*$ -algebra  $A$  and  $\varphi : A \rightarrow A$  is the identity map, then [6, Corollary 4.2] is a consequence of the above corollary.

Applying Theorem 2.4 and 2.6 with  $\rho(x, y) = c\|x\|^p\|y\|^p$  we have the next result.

**Corollary 2.9.** *Let  $U : E \rightarrow F$  be a mapping and  $\varphi : A \rightarrow B$  be an additive map such that for some  $p \neq 1$ ,*

$$\|\langle Ux, Uy \rangle - \varphi(\langle x, y \rangle)\| \leq c\|x\|^p\|y\|^p \quad (x, y \in E_p).$$

Then there exists a unique  $\varphi$ -morphism  $T : E \rightarrow F$  such that

$$\|U(x) - T(x)\| \leq \frac{\sqrt{c(2^p + 2)}}{|2 - 2^p|} \|x\|^p \quad (x \in E_p).$$

**Acknowledgement.** The authors would like to thank the referee for his/her useful comments.

#### REFERENCES

1. Gh. Abbaspour Tabadkan, M.S. Moslehian and A. Niknam, *Dynamical systems on Hilbert  $C^*$ -modules*, Bull. Iranian Math. Soc. **31** (2005), 25–35.
2. Gh. Abbaspour Tabadkan and M. Skeide, *Generators of dynamical systems on Hilbert modules*, Commun. on Stoch. Anal. **1** (2007), no. 2, 193–207.
3. D. Bakić and B. Guljaš, *On class of module maps of Hilbert  $C^*$ -modules*, Math. Commun. **7** (2002), 177–192.
4. R. Badora and J. Chmieliński, *Decomposition of mappings approximately inner product preserving*, Nonlinear Anal. **62** (2005), 1015–1023.
5. J. Chmieliński, D. Ilišević, M.S. Moslehian and Gh. Sadeghi, *Perturbation of the Wigner equation in inner product  $C^*$ -modules*, J. Math. Phys. **49** (2008), no. 3, 033519, 8 pp.
6. J. Chmieliński and M.S. Moslehian, *Approximately  $C^*$ -inner product preserving mappings*, Bull. Korean Math. Soc. **45** (2008), no. 1, 157–167.
7. L. Cădariu and V. Radu, *Fixed points and the stability of Jensen's functional equation*, J. Inequal. Pure Appl. Math. **4** (2003), no.1, Art. 4.
8. J. Diaz and B. Margolis, *A fixed point theorem of the alternative for contractions on a generalized complete metric space*, Bull. Amer. Math. Soc. **74** (1968), 305–309.
9. Z. Gajda, *On stability of additive mappings*, Inter. J. Math. Sci. **14** (1991), 431–434.
10. D.H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Soc. U.S.A. **27** (1941), 222–224.
11. S.-M. Jung, *A fixed point approach to the stability of an equation of the square spiral*, Banach J. Math. Anal. **1** (2007), no. 2, 148–153.
12. E.C. Lance, *Hilbert  $C^*$ -Modules*, LMS Lecture note Series 210, Cambridge Univ. Press, 1995.
13. Y.-H. Lee, *On the stability of the monomial functional equation*, Bull. Korean Math. Soc. **45** (2008), no. 2, 397–403.
14. A.K. Mirmostafae, *Approximate isometris in Hilbert  $C^*$ -modules*, Math. Commun. **14** (2009), no.1, 167–176.
15. C. Park and Th.M. Rassias, *Fixed points and the stability of the Cauchy functional equation*, Aust. J. Math. Anal. Appl. **6** (2009), no. 1, Art. 14.
16. V. Radu, *The fixed point alternative and stability of functional equations*, Fixed point theory **4** (2003), 190–199.
17. Th.M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
18. S.M. Ulam, *A collection of the mathematical problems*, Interscience Publ, New Yourk, 1960.

<sup>1</sup> SCHOOL OF MATHEMATICS AND COMPUTER SCIENCES, DAMGHAN UNIVERSITY, DAMGHAN, IRAN.

*E-mail address:* [abbaspour@du.ac.ir](mailto:abbaspour@du.ac.ir)

<sup>2</sup> SCHOOL OF MATHEMATICS AND COMPUTER SCIENCES, DAMGHAN UNIVERSITY, DAMGHAN, IRAN.

*E-mail address:* [md\\_ramezanpour@yahoo.com](mailto:md_ramezanpour@yahoo.com), [ramezanpour@du.ac.ir](mailto:ramezanpour@du.ac.ir)