

**A REMARK ON TILED ORDERS OVER
A LOCAL DEDEKIND DOMAIN**

Dedicated to Professor Hisao Tominaga
on his 60th birthday

By

Hisaaki FUJITA

Let R be a noetherian domain with the quotient ring K . An R -order in the full $n \times n$ matrix ring $(K)_n$ is called *tiled* if it contains n orthogonal idempotents (cf. [3]). There are many papers on noetherian ring theory which contain tiled R -orders as examples. Concerning global dimension, tiled R -orders are studied by K.L. Fields [1], R.B. Tarsy [10], [11], V.A. Jategaonkar [2], [3], [4] and K.W. Roggenkamp [8], [9].

In [5], B.J. Müller introduced the concept of *links* between prime ideals of Fully Bounded Noetherian (FBN) rings to study localizability of semiprime ideals. Recently in [6], he initiated a detailed study of the link graph and announced some results on FBN prime rings of Krull dimension one, especially, with self-injective dimension one.

In this note, we shall attempt a study on the link graph of tiled orders over a local Dedekind domain, which are FBN prime rings of Krull dimension one and have arbitrarily large global dimension (cf. [1], [7] and Example 3.5).

After recalling some definitions and notations, in Section 1, we shall point out that the link graph coincides with the quiver of orders introduced by A. Wiedemann and K.W. Roggenkamp [12].

Confining ourselves to tiled R -orders between $(R)_n$ and its radical, in Section 2, we shall prove the following.

THEOREM. *Let R be a local Dedekind domain with the maximal ideal πR and the quotient ring K . Let A be a basic tiled R -order between $(R)_n$ and $(\pi R)_n$, $Q(A)$ the quiver of the $R/\pi R$ -algebra $A = \Lambda/(\pi R)_n$ and M_1, \dots, M_n the maximal ideals of A . Then, there is a link from M_i to M_j if and only if there is an arrow from i to j in $Q(A)$, or else i is a non-domain and j is a non-range in $Q(A)$.*

We shall give some remarks after proving the theorem. We shall add an

Appendix in which we shall announce global dimension of some special Λ , i.e., $Q(\Lambda)$ is a tree, of A_n -type, a cycle and so on.

1. Preliminaries

Let R be a local Dedekind domain with the maximal ideal πR and the quotient ring K . Let $(K)_n$ be the full $n \times n$ matrix ring over K and Λ be a *tilted* R -order in $(K)_n$ (i.e., Λ contains n orthogonal primitive idempotents). By virtue of [4, Lemma 1], we may assume $\Lambda = (\pi^{\lambda_{ij}} R) \subset (R)_n$ where λ_{ij} 's are non-negative integers and $\lambda_{ii} = 0$ for $1 \leq i \leq n$. Since Λ is a subring of $(R)_n$, it holds that

$$(*-1) \quad \lambda_{ik} + \lambda_{kj} \geq \lambda_{ij} \quad \text{for all } 1 \leq i, j, k \leq n.$$

Since R is a local ring, Λ is semiperfect. Using $(*-1)$, it is easily checked that Λ is basic if and only if Λ satisfies

$$(*-2) \quad \text{If } i \neq j, \lambda_{ij} = 0 \text{ implies } \lambda_{ji} \neq 0.$$

In what follows, $\Lambda = (\pi^{\lambda_{ij}} R)$ is a basic tilted R -order contained in $(R)_n$.

Since Λ is finitely generated over a local Dedekind domain, Λ is an FBN prime ring of Krull dimension one.

For $1 \leq k \leq n$, let $M_k = (\pi^{m_{kij}} R) \subset \Lambda$ where $m_{kij} = 1$ (if $i = j = k$) λ_{ij} (otherwise). Then M_1, \dots, M_n are the maximal ideals of Λ .

There exists a *link* from M_i to M_j (denoted by $M_i \rightsquigarrow M_j$) if $M_i \cap M_j \cong M_j M_i$ holds (cf. [5, Remark (2), p 236]).

Let I_1, I_2 be ideals of R . Since R is a local Dedekind domain, $I_1 = \pi^a R$, $I_2 = \pi^b R$ for some integers $a, b \geq 0$. Then $I_1 \supset I_2$ if and only if $a \leq b$. We define an order between ideals of R by $I_1 \leq I_2$ if and only if $a \leq b$. (The symbol " \leq " may not be confused in the context.) We shall use Max and Min among ideals of R under the above order.

Put $M_{kij} = \pi^{m_{kij}} R$ for $1 \leq i, j, k \leq n$. For $1 \leq k, h \leq n$, put

$$X_{kh} = \text{Max}\{M_{ikh}, M_{jkh}\},$$

$$Y_{kh} = \text{Min}\{M_{jk\ell} M_{i\ell h} \mid 1 \leq \ell \leq n\}.$$

Then $M_i \cap M_j = (X_{kh})$ and $M_j M_i = (Y_{kh})$.

LEMMA 1.1. *If $(k, h) \neq (j, i)$, then $X_{kh} = Y_{kh}$.*

PROOF. It holds that $M_{ikh} \leq X_{kh} \leq Y_{kh} \leq M_{jkh} M_{ikh}$. If $X_{kh} \neq Y_{kh}$, we have $M_{jkh} \neq R$. Hence $k = j$. Similarly $h = i$.

For $1 \leq i \leq n$, let e_i be the matrix in Λ with (i, i) -entry equal to 1 and all others

0, and put $P_i = e_i A$, $J_i = \text{rad}(P_i) = e_i J$, where $J = \text{rad}(A)$. $\mathbf{P}(X)$ denote a projective cover of a module X .

We now repeat the (right-handed) definition of the valued quiver of the tiled R -order given by A. Wiedemann and K.W. Roggenkamp [12]. (The links that we have been using are right-handed, but [12] is left-handed.)

A *valued quiver* $Q = (Q_0, Q_1, d, r, v)$ consists of a finite set Q_0 of vertices and a finite set Q_1 of arrows. d and r are maps from Q_1 to Q_0 such that $d(\alpha)$ is the domain and $r(\alpha)$ is the range of an arrow $\alpha \in Q_1$. v is a map from Q_1 to non-negative integers. Forgetting the valuation map v from a valued quiver, we call it a *quiver*.

For the tiled R -order $A = (\pi^{ij} R) \subset (R)_n$, the *valued quiver* $Q(A)$ of A is defined by the vertices $Q(A)_0 = \{1, \dots, n\}$, there exists an arrow $\alpha \in Q(A)_1$ with $d(\alpha) = i$, $r(\alpha) = j$ if P_i is isomorphic to a direct summand of $\mathbf{P}(J_j)$, and $v(\alpha) = \lambda_{ji}$. In [12], a procedure is given to construct a tiled R -order $A(Q)$ from a certain valued quiver Q and it is shown that $A = A(Q(A))$ [12, § 2, Theorem 1].

PROPOSITION 1.2. *The link graph between maximal ideals of A coincides with the quiver of A .*

PROOF. There is an arrow from i to j in $Q(A) \iff P_i$ is isomorphic to a direct summand of $\mathbf{P}(J_j) \iff e_j J e_i \oplus e_j J^2 e_i \iff X_{ji} \cong Y_{ji} \iff M_i \rightsquigarrow M_j$ by Lemma 1.1.

COROLLARY 1.3. *If A has finite global dimension, then all maximal ideals of A are idempotent.*

PROOF. It follows from [12, § 1, Lemma 3] that $Q(A)$ has no loops (i.e., there is no arrows from a vertex to itself). So, by the proposition, $M_i \rightsquigarrow M_i$, and hence $M_i = M_i^2$ for all $1 \leq i \leq n$.

REMARK. There is a tiled R -order of infinite global dimension with all of whose maximal ideals are idempotent. See Example 3.4.

An ideal I is *eventually idempotent* if $I^m = I^{m+1} = \dots$ for some integer $m > 0$. Of course, there are many tiled R -orders with non-idempotent maximal ideals, however, we note the following.

PROPOSITION 1.4. *Let A be the tiled R -order in $(R)_n$ ($n \geq 2$). Then all maximal ideals of A are eventually idempotent.*

PROOF. Fix $1 \leq k \leq n$. By induction on ℓ , define

$$\begin{cases} m_{kij}^1 = m_{kij}, \\ m_{kij}^{\ell+1} = \min\{m_{kih}^{\ell} + m_{khj} \mid 1 \leq h \leq n\}. \end{cases}$$

Then it is easily shown that $m_{kij}^{\ell} \leq \ell$ (if $i=j=k$) and $=\lambda_{ij}$ (otherwise). It follows that $M_k^d = M_k^{d+1} = \dots$, where $d = \min\{\lambda_{kh} + \lambda_{hk} \mid 1 \leq h \leq n, h \neq k\}$.

2. The link graph of Λ and the quiver of the factor algebra

Let $\Lambda = (\pi^{\lambda_{ij}}R)$ be a basic tiled R -order between $(R)_n$ and $(\pi R)_n$ (i.e., $(R)_n \supset \Lambda \supset (\pi R)_n$). Put $\Lambda_{ij} = \pi^{\lambda_{ij}}R$, $A = \Lambda/(\pi R)_n$ and $N = \text{rad}(A)$. Then A is a basic $R/\pi R$ -algebra. The quiver $Q(A)$ of A is defined by the set of vertices $Q(A)_0 = \{1, \dots, n\}$ and there is an arrow from i to j if


$$(*)-3) \quad e_j N e_i / e_j N^2 e_i \neq 0.$$

Let $J = (J_{ij})$ be the Jacobson radical of Λ and $B_{ij} = \text{Min}\{J_{ik} J_{kj} \mid 1 \leq k \leq n\}$. Then $(*)-3)$ is equivalent to $J_{ji}/(B_{ji} + \pi R) \neq 0$. So $Q(A)$ has no loops. It follows from $(*)-2)$ that $Q(A)$ has no oriented cycles. Let \mathcal{D} (resp. \mathcal{R}) denote a subset of $Q(A)_0$ consisting of non-domains (resp. non-ranges) in $Q(A)$.


Here, we give an example which helps the reader's understanding of the theorem.

EXAMPLE 2.1. Let $\Lambda = \begin{pmatrix} R & \pi R & \pi R \\ \pi R & R & \pi R \\ \pi R & R & R \end{pmatrix}$. Then $A \cong \begin{pmatrix} \mathbf{k} & 0 & 0 \\ 0 & \mathbf{k} & 0 \\ 0 & \mathbf{k} & \mathbf{k} \end{pmatrix}$, where $\mathbf{k} = R/\pi R$.

The quiver $Q(A)$ is given by



So $\mathcal{D} = \{1, 3\}$ and $\mathcal{R} = \{1, 2\}$. Thus the link graph is given by



- LEMMA 2.2. (1) If $d \in \mathcal{D}$, then $\Lambda_{jd} = \pi R$ for all $j \neq d$.
 (2) If $r \in \mathcal{R}$, then $\Lambda_{ri} = \pi R$ for all $i \neq r$.

PROOF. (1) For each $j \neq d$, since $e_j N e_d / e_j N^2 e_d = 0$,

(i)
$$\Lambda_{jd} / (B_{jd} + \pi R) = 0.$$

Assume that $\Lambda_{jd} = R$ for some $j \neq d$. Then by (i), $B_{jd} = R$. Since $B_{jd} = \text{Min}\{J_{j\ell} J_{\ell d} \mid$

$1 \leq \ell \leq n$, $J_{j\ell_1} J_{\ell_1 d} = R$ for some $1 \leq \ell_1 \leq n$. Clearly, $J_{j\ell_1} = J_{\ell_1 d} = R$ and $j \neq \ell_1 \neq d$. Hence $A_{\ell_1 d} = J_{\ell_1 d} = R$, and by (i), $B_{\ell_1 d} = R$. Repeating the above argument, we obtain $\ell_0 = j, \ell_1, \dots, \ell_n$ ($\neq d$) and $J_{\ell_{i-1}\ell_i} = R$ ($1 \leq i \leq n$). So $\ell_n = \ell_k$ ($= \ell$, say) for some $0 \leq h < k \leq n$. We get $\pi R \supset J_{\ell\ell_{h+1}} \cdots J_{\ell_{k-1}\ell} = R$, a contradiction. This completes the proof of (1). Similarly, (2) is proved.

PROOF OF THE THEOREM. (\implies) It follows from Lemma 1.1 that $X_{ji} \not\cong Y_{ji}$.

Case 1. $X_{ji} = \pi R$.

Then $Y_{ji} \geq \pi^2 R$. It holds that

$$(ii) \quad \begin{cases} M_{iji} = M_{jji} = \pi R & \text{if } i = j, \\ M_{iji} = A_{ji} = M_{jji} = \pi R & \text{if } j \neq i. \end{cases}$$

Hence $Y_{ji} = \text{Min} \{M_{jjk} M_{iki} \mid 1 \leq k \leq n\} = \text{Min} [\{A_{jk} A_{ki} \mid 1 \leq k \leq n, k \neq i, j\} \cup \{\pi^2 R\}] \leq \pi^2 R$. Therefore $\pi^2 R = Y_{ji} \leq A_{jk} A_{ki}$ for $1 \leq k \leq n, k \neq i, j$. Hence by (ii), $A_{jk} = \pi R$ if $k \neq j$ and $A_{ki} = \pi R$ if $k \neq i$. Thus

$$\begin{aligned} e_k N e_i / e_k N^2 e_i &\cong A_{ki} / (B_{ki} + \pi R) = \pi R / (B_{ki} + \pi R) = 0 & \text{if } k \neq i, \\ e_j N e_k / e_j N^2 e_k &\cong A_{jk} / (B_{jk} + \pi R) = \pi R / (B_{jk} + \pi R) = 0 & \text{if } k \neq j. \end{aligned}$$

Consequently, $i \in \mathcal{D}$ and $j \in \mathcal{R}$.

Case 2. $X_{ji} = R$.

Then $Y_{ji} \geq \pi R$. It holds that $i \neq j, M_{iji} = A_{ji} = M_{jji} = R$. Observe that

$$(iii) \quad \begin{aligned} B_{ji} &= \text{Min} \{J_{jk} J_{ki} \mid 1 \leq k \leq n\} \\ &= \text{Min} \{M_{jjk} M_{iki} \mid 1 \leq k \leq n\} \\ &= Y_{ji}. \end{aligned}$$

Thus $e_j N e_i / e_j N^2 e_i \cong A_{ji} / (B_{ji} + \pi R) = R / \pi R \neq 0$. Hence $i \rightarrow j \in Q(A)_1$.

(\longleftarrow) Case 1. $i \in \mathcal{D}$ and $j \in \mathcal{R}$.

It follows from Lemma 2.2 that $M_{jjk} = M_{iki} = \pi R$ for all $k = 1, \dots, n$. Hence $X_{ji} = \pi R < \pi^2 R = Y_{ji}$. Therefore $M_i \rightsquigarrow M_j$.

Case 2. There is an arrow from i to j in $Q(A)$.

It holds that $i \neq j, A_{ji} / (B_{ji} + \pi R) \neq 0$, so that $A_{ji} = R$ and $B_{ji} \geq \pi R$. Since $i \neq j, M_{iji} = A_{ji} = M_{jji}$. Hence $X_{ji} = A_{ji} = R$. By (iii), $Y_{ji} = B_{ji} \geq \pi R > R = X_{ji}$. Therefore $M_i \rightsquigarrow M_j$.

REMARKS. (1) We note the following fact that is shown in the proof; If $M_i \rightsquigarrow M_j$ and $M_i \cap M_j = (X_{kh})$, then

$$\begin{cases} X_{ji} = \pi R \iff i \in \mathcal{D} \text{ and } j \in \mathcal{R}, \\ X_{ji} = R \iff i \rightarrow j \in Q(A)_1. \end{cases}$$


(2) In our vein, we note that the link graph of maximal ideals of A is connected.

(3) A maximal ideal M_i of A is not idempotent if and only if i is an isolated vertex in $Q(A)$.


(4) $Q(A)$ has full information about A . As for the link graph, there are tiled R -orders A_1, A_2 with the same link graph, but $Q(A_1)$ is different from $Q(A_2)$ where $A_i = A_i / (\pi R)_n$ ($i=1,2$).

EXAMPLE 2.3. Let

$$A_1 = \begin{pmatrix} R & \pi R & \pi R & \pi R \\ R & R & \pi R & \pi R \\ R & R & R & \pi R \\ R & R & \pi R & R \end{pmatrix} \quad \text{and} \quad A_2 = \begin{pmatrix} R & R & R & R \\ \pi R & R & \pi R & \pi R \\ \pi R & R & R & \pi R \\ \pi R & R & \pi R & R \end{pmatrix}.$$



It follows from the theorem that the link graphs of A_1 and A_2 are given by



PROPOSITION 2.4. Let A_1, A_2 be the tiled R -orders between $(R)_n$ and $(\pi R)_n$ with the same link graph and put $A_i = A_i / (\pi R)_n$ ($i=1,2$). Then $Q(A_1)$ is connected if and only if $Q(A_2)$ is connected.

PROOF. Suppose that $Q(A_1)$ is disconnected and put $Q(A_1) = \mathcal{U} \dot{\cup} \mathcal{V}$ (disjoint union) with \mathcal{U} connected.

If $\mathcal{U}_0 = \{m\}$, then the link graph has a loop on m . Assume that $Q(A_2)$ is connected. Then there exist a non-domain (or non-range) vertex v in \mathcal{V} and an arrow $v \rightarrow m$ (or $m \rightarrow v$) in $Q(A_2)$. Hence the link graph of A_2 has no loops on m , a contradiction. Therefore $Q(A_2)$ is disconnected.

Let \mathcal{U}_0 be not a singleton and assume that $Q(A_2)$ is connected. Then (i) there exist a source s_0 in \mathcal{U} and a non-domain d in \mathcal{V} with $d \xrightarrow{a} s_0 \in Q(A_2)$, or (ii) there exist a sink s_i in \mathcal{U} and a non-range r in \mathcal{V} with $s_i \xrightarrow{a} r \in Q(A_2)$.

In the case (i), since $Q(A_2)$ is connected, there exist a sink x and a path $s_0 \rightarrow \dots \rightarrow x$ in $Q(A_2)$. Assume $x \in \mathcal{V}_0$. Then there is an arrow β in the path from s_0 to x such that $d(\beta) \in \mathcal{U}_0$ and $r(\beta) \in \mathcal{V}_0$. Then $d(\beta)$ is a sink in \mathcal{U} . Since s_0 is a range of α in $Q(A_2)$, there must exist an arrow from s to s_0 in $Q(A_2)$ for each sink s in \mathcal{U} . Hence $Q(A_2)$ has an oriented cycle $s_0 \rightarrow \dots \rightarrow d(\beta)$, a contradiction. Therefore $x \in \mathcal{U}_0$. Similarly, there exist a source y and a path $y \rightarrow \dots \rightarrow d$ in $Q(A_2)$ with $y \in \mathcal{V}_0$. Hence there is an arrow $x \rightarrow y$ in the link graph of A_2 . But, since x is not a sink in \mathcal{U} , the link graph of A_1 has no arrows from x to y , a contradiction. Similarly, we can deduce a contradiction in the case (ii). Therefore $Q(A_2)$ is disconnected.

3. Appendix: Global dimension of some special Λ

Let Λ be a basic tiled R -order between $(R)_n$ and $(\pi R)_n$ and put $A = \Lambda / (\pi R)_n$. Before attacking some special cases, we shall note the following proposition whose proof mainly depends on the infinite global dimensional criterion given by V.A. Jategaonkar [4].

PROPOSITION 3.1. *If Λ has finite global dimension, then the quiver $Q(A)$ is connected.*

PROOF. Suppose that $Q(A)$ is disconnected. Since there is a permutation matrix $u \in (R)_n$ such that

$$\Lambda \cong u \Lambda u^{-1} = \begin{pmatrix} R & * & \pi R \cdots \pi R \\ \vdots & \ddots & \vdots \\ * & R & \pi R \cdots \pi R \\ \pi R \cdots \pi R & R & * \\ \vdots & \vdots & \vdots \\ \pi R \cdots \pi R & * & R \end{pmatrix} (s,$$

we may assume that $\Lambda = u \Lambda u^{-1}$. Put $M = (R, \dots, R) = R^n$ and

$$M \supset B_1 = (R, \dots, \overset{s}{R}, \pi R, \dots, \pi R) \supset \pi M,$$

$$M \supset B_2 = (\pi R, \dots, \pi R, R, \dots, R) \supset \pi M.$$

Then B_1, B_2, M are right Λ -modules and $M/\pi M \cong B_1/\pi M \oplus B_2/\pi M$ as right $\Lambda/\pi \Lambda$ -modules. Thus it follows from [4, Lemma 1.7] that $\text{proj. dim}(M_\Lambda) = \infty$, so that $\text{gl. dim } \Lambda = \infty$.

PROPOSITION 3.2. *If $Q(A)$ is a tree, then $\text{gl. dim } \Lambda \leq 3$.*

PROOF. Put $i^- = \{j \in Q(A)_0 \mid d(\alpha) = j, r(\alpha) = i \text{ for some } \alpha \in Q(A)_1\}$, $\mathcal{S} = \{j \in Q(A)_0 \mid j \text{ is a sink in } Q(A)\}$ and $L = (R, \dots, R) = R^n$.

Case 1 (i) $i^- = \emptyset$, \mathcal{S} is a singleton.

Let $\mathcal{S} = \{j\}$. Then $P_j = L$. Since $J_i = (\pi R, \dots, \pi R) = \pi L \cong L$, $\text{proj. dim}(J_i) = 0$.

Case 1 (ii) $i^- = \emptyset$, $\mathcal{S} = \{s_1, \dots, s_t\}$ ($t \geq 2$).

For a subset \mathcal{X} of $Q(A)_0$, put $\mathcal{P}_{\mathcal{S}}(\mathcal{X}) = \{j \in Q(A)_0 \mid \text{there is a path } j \rightarrow \dots \rightarrow x \text{ in } Q(A) \text{ or } j = x \text{ for some } x \in \mathcal{X}\}$. Pick up a sink s_1 . Since $Q(A)$ is connected, there exist a sink s_2 outside $\mathcal{P}_{\mathcal{S}}(s_1)$ and $j_1 \in \mathcal{P}_{\mathcal{S}}(s_1)$ such that there is a path $j_1 \rightarrow \dots \rightarrow s_2$ in $Q(A)$ which branches at the vertex j_1 . Since $Q(A)$ has no cycles, such j_1 is unique. If $\mathcal{P}_{\mathcal{S}}(s_1, s_2) \neq Q(A)_0$, repeat the above procedure. After some repetitions, we reach $Q(A)_0 = \mathcal{P}_{\mathcal{S}}(s_1, \dots, s_t)$ with vertices j_1, \dots, j_{t-1} . Then using canonical maps, we obtain short exact sequences

$$\begin{aligned} 0 &\longrightarrow P_{j_1} \longrightarrow P_{s_1} \oplus P_{s_2} \longrightarrow M_2 \longrightarrow 0, \\ 0 &\longrightarrow P_{j_2} \longrightarrow M_2 \oplus P_{s_3} \longrightarrow M_3 \longrightarrow 0, \\ &\quad \quad \quad \vdots \\ 0 &\longrightarrow P_{j_{t-1}} \longrightarrow M_{t-1} \oplus P_{s_t} \longrightarrow M_t \longrightarrow 0. \end{aligned}$$

Then $\text{proj. dim}(M_t) = \text{proj. dim}(M_{t-1}) = \dots = \text{proj. dim}(M_2) = 1$ and $M_t = L$. Since $J_i = \pi L \cong L$, $\text{proj. dim}(J_i) = 1$.

Case 2 i^- is a singleton.

Let $i^- = \{j\}$. Then $J_i \cong P_j$. Hence $\text{proj. dim}(J_i) = 0$.

Case 3 (i) $i^- = \{j_1, \dots, j_u\}$ ($u \geq 2$), \mathcal{S} is a singleton.

Using canonical maps, we obtain short exact sequences

$$\begin{aligned} 0 &\longrightarrow L \longrightarrow P_{j_1} \oplus P_{j_2} \longrightarrow N_2 \longrightarrow 0, \\ 0 &\longrightarrow L \longrightarrow N_2 \oplus P_{j_3} \longrightarrow N_3 \longrightarrow 0, \\ &\quad \quad \quad \vdots \\ 0 &\longrightarrow L \longrightarrow N_{u-1} \oplus P_{j_u} \longrightarrow N_u \longrightarrow 0. \end{aligned}$$

Then $N_u = J_i$. Hence $\text{proj. dim}(J_i) = \text{proj. dim}(L) + 1 = 1$.

Case 3 (ii) $i^- = \{j_1, \dots, j_u\}$, $\mathcal{S} = \{s_1, \dots, s_t\}$ ($u, t \geq 2$).

As in Case 3 (i), $\text{proj. dim}(J_i) = \text{proj. dim}(L) + 1 = 2$ from Case 1 (ii).


Therefore $\text{gl. dim } \Lambda = \sup \{\text{proj. dim}(J_i) \mid 1 \leq i \leq n\} + 1 \leq 3$.

REMARK. It follows from the proof that $\text{gl. dim } \Lambda \leq 2$ iff $Q(A)$ has a unique source or a unique sink. This is a special case of [8, Theorem].

EXAMPLE 3.3. Let $Q(A)$ be of A_n -type and put m be the number of vertices at which directions of arrows are changed. Then if $m=0, 1, \geq 2$, then $\text{gl. dim } A=1, 2, 3$, respectively.

PROOF. This follows from the proof of Prop. 3.2.


EXAMPLE 3.4. Let $Q(A)$ be a cycle



If $t=1, \geq 2$, then $\text{gl. dim } A=2, \infty$, respectively. It follows from Remark (3) in § 2 that the maximal ideals of A are idempotent, while $\text{gl. dim } A=\infty$ if $t \geq 2$.

In [1] and [7], there is an example which is a tiled R -order between $(R)_n$ and $(\pi R)_n$ with enough large global dimension. If $n=2^m$, then its global dimension is m . Next one is such an example with smaller n . Calculations of Examples 3.4 and 3.5 are left to the reader.

EXAMPLE 3.5. For $m \geq 2$ and $0 \leq k \leq m-1$, let $Q(A_k)$ be



Let A_k be the tiled R -order between $(R)_n$ and $(\pi R)_n$ such that $A_k = A_k / (\pi R)_n$ where $n=2m$. Then $A_0 \subset A_1 \subset \dots \subset A_{m-2} \subset A_{m-1}$ and $\text{gl. dim } A_k = k+3$ for $0 \leq k \leq m-2$ and $\text{gl. dim } A_{m-1} = \infty$.

If $m=2(3)$ and $k=0(1)$, then $n=4(6)$ and $\text{gl. dim } A_k=3(4)$. It is verified by computation that $4(6)$ is the smallest n with global dimension $3(4)$.

References

- [1] Fields, K. L., Examples of orders over discrete valuation rings, *Math. Z.* **111** (1969), 126-130.
- [2] Jategaonkar, V. A., Global dimension of triangular orders over a discrete valuation ring, *Proc. Amer. Math. Soc.* **38** (1973), 8-14.
- [3] Jategaonkar, V. A., Global dimension of tiled orders over commutative noetherian domains, *Trans. Amer. Math. Soc.* **190** (1974), 357-374.

- [4] Jategaonkar, V. A., Global dimension of tiled orders over a discrete valuation ring, *Trans. Amer. Math. Soc.* **196** (1974), 313–330.
- [5] Müller, B. J., Localization in fully bounded noetherian rings, *Pacific J. Math.* **67** (1976), 233–245.
- [6] Müller, B. J., Links between maximal ideals in bounded noetherian prime rings of Krull dimension one, “Methods in Ring Theory” NATO ASI Series, D. Reidel Publishing Company 1984, 347–377.
- [7] Robson, J. C., Some constructions of rings of finite global dimension, *Glasgow Math. J.* **26** (1985), 1–12.
- [8] Roggenkamp, K. W., Some examples of orders of global dimension two, *Math. Z.* **154** (1977), 225–238.
- [9] Roggenkamp, K. W., Orders of global dimension two, *Math. Z.* **160** (1978), 63–67.
- [10] Tarsy, R. B., Global dimension of orders, *Trans. Amer. Math. Soc.* **151** (1970), 335–340.
- [11] Tarsy, R. B., Global dimension of triangular orders, *Proc. Amer. Math. Soc.* **28** (1971), 423–426.
- [12] Wiedemann, A. and Roggenkamp, K. W., Path orders of global dimension two, *J. Algebra* **80** (1983), 113–133.

Institute of Mathematics
University of Tsukuba
Sakura-mura Ibaraki 305, Japan