

GRAPHS AND FINITE DISTRIBUTIVE PARTIAL LATTICES

By

Juhani NIEMINEN

Abstract

The Hasse diagram graph of a finite distributive partial lattice is characterized by means of prime convexes.

Median graphs constitute a well known and widely studied class of graphs; see for example the papers [1] and [2] and the references therein. They constitute a subclass of the Hasse diagram graphs of distributive partial lattices. In this paper we give a characterization for the Hasse diagram graphs G of finite distributive partial lattices by means of prime convexes of G . This characterization generalizes that of Mulder and Schrijver for median graphs reprinted in [1, Theorem 2.2].

A meetsemilattice S is a partial lattice if for any two elements a, b having an upper bound in S also the element $a \vee b$ belongs to S . Clearly every finite meetsemilattice is a partial lattice. A partial lattice S is distributive if its every subset $[k] = \{s | s \leq k\}$ is a distributive lattice. A finite distributive partial lattice S can be embedded in the distributive lattice $I(S)$ of ideals of S , where the join of two ideals I and J is $I \vee J = \{s | s \leq i \vee j, i \in I \text{ and } j \in J\}$. By using this lattice we see that one shortest path joining two points a and b of the Hasse diagram graph S contains the point $a \wedge b$, and if $a > b$, then every point c , $a \geq c \geq b$, is on some shortest a - b path.

The graphs $G = (V, X)$ considered here are finite, connected and undirected without loops and multiple lines. The points of G constitute the set V and its lines the set X . A pointset $A \subset V$ of G is called a *convex* if A contains all points of any shortest a - b path (of any a - b geodesic) for every two points $a, b \in A$. The intersection of two convexes is also a convex and thus the least convex containing a given pointset B of G is $\bigcap \{C | C \text{ is a convex and } B \subset C\}$. This set is briefly denoted by $\langle B \rangle$. A convex $A \neq V$ is called *prime* if the set $V \setminus A$ is also a convex. The sets \emptyset and V are trivial prime convexes. A graph G has the prime convex intersection property (is a *prime convex intersection graph*) if its every

convex A is the intersection of all prime convexes containing A . By [1, Theorem 2.2], every median graph is a prime convex intersection graph. The class of prime convex intersection graphs is rather wide: for example every complete graph belongs to this class.

Let $a, b, c \in V$. A point t satisfying the distance conditions $d(a, b) = d(a, t) + d(t, b)$, $d(b, c) = d(b, t) + d(t, c)$ and $d(a, c) = d(a, t) + d(t, c)$ is a *median* of the points a, b and c . A graph is a median graph if its all three points have exactly one median.

If A is a subset of a set U , then $\bar{A} = U \setminus A$ is its complement in U .

When proving the main theorem of this note we need two auxiliary results which we prove first.

LEMMA 1. *A connected graph G is a prime convex intersection graph if and only if for any nonempty convex A and any point x , $x \notin A$, there is a prime convex P separating A and x , i. e. $A \subset P$ and $x \in \bar{P}$.*

PROOF. If G is a prime convex intersection graph, A its nonempty convex and x its point such that $x \notin A$, there is a prime convex P separating A and x , because otherwise A cannot be represented as an intersection of prime convexes of G . Conversely, if there is a prime convex separating any convex A and any point x of the lemma then G is a prime convex intersection graph. Indeed, if there is a nonempty convex A which cannot be expressed as the intersection of prime convexes, then the intersection contains a point x not belonging to A . By assumption there is a prime convex P separating A and x , and thus the intersection cannot contain the point x , and the lemma follows.

LEMMA 2. *The convex $\langle a, b \rangle$ of a prime convex intersection graph G consists of points on a - b geodesics for every pair $a, b \in V$.*

PROOF. Let a and b be a pair of points such that the convex $\langle a, b \rangle$ contains at least one point v which is not on any a - b geodesic. This implies the existence of two points x and z , x is on an a - b geodesic and z is on another a - b geodesic, such that no point x_1, \dots, x_m of an x - z geodesic $x = x_0, x_1, \dots, x_m, x_{m+1} = z$ is on any a - b geodesic. Clearly a and b can be chosen such that every convex $\langle u, w \rangle$ with $d(u, w) < d(a, b)$ is the set of all points on u - w geodesics. We may assume further that $d(a, b) \geq d(x, b)$, $d(z, b) \geq d(x, b)$, and that x and z are as near to b as possible. Let us consider the point x_1 . Because $d(a, x) < d(a, b)$, the convex $\langle a, x \rangle$ consists of points on a - x geodesics, and thus $x_1 \notin \langle a, x \rangle$. By Lemma 1, the prime convex intersection property of G implies now the existence of a prime

convex P separating $\langle a, x \rangle$ and $x_1: \langle a, x \rangle \subset P$ and $x_1 \in \bar{P}$. Because $x_1 \in \langle a, b \rangle$, we have $x_1, b \in \bar{P}$. Let $x = b_0, b_1, b_2, \dots, b_{k-1}, b_k = b$ be the points of an x - b geodesic. Because x and z are as near to b as possible, $d(z, b) \geq d(x, b)$ and $d(x_1, z) \geq d(x_1, x) = 1$, then a b_i - x_1 geodesic goes over $x, i = 1, \dots, k$. This implies that there is no prime convex separating $\langle a, x \rangle$ and x_1 , which is a contradiction. Thus the assumption is false and the convex $\langle a, b \rangle$ consists of points on a - b geodesics for every pair $a, b \in V$, and the lemma follows.

Now we can present the characterization theorem of this note.

THEOREM. *A connected graph G is isomorphic to the Hasse diagram graph of a finite distributive partial lattice if and only if the following two conditions hold:*

- (i) *G is a prime convex intersection graph;*
- (ii) *$\bigcap \{\bar{P} | P \in \mathcal{K}\} \neq \phi$ or $\mathcal{K} = \phi$ for the collection \mathcal{K} of all nontrivial prime convexes in G having the following property: if $P_1 \in \mathcal{K}$, there are $P_2, P_3, \dots, P_n \in \mathcal{K} (n \geq 3)$ such that $P_i \cap P_j \neq \phi$ and $P_1 \cap P_2 \cap \dots \cap P_n = \phi$.*

PROOF. Mulder and Schrijver proved that a connected graph G is a median graph if and only if G is a prime convex intersection graph and its prime convexes satisfy the Helly property [1, Theorem 2.2]. The condition (ii) above is nothing but a weakened Helly property for prime convexes of G .

Assume first that G is the Hasse diagram graph of a finite distributive partial lattice S .

(i) Let $x \in S$. The element corresponding x in the ideal lattice $I(S)$ of S is $(x]$. Because $I(S)$ is distributive, one $(z]$ - $(x]$ geodesic goes over the element $(z] \wedge (x] = (z \wedge x]$. Thus, if the distance $d((z], (x]) = n$ in $I(S)$, then $d(z, x) = n$ in S , because the z - $z \wedge x$ - x path always belongs to S . In particular, if C is a convex of the Hasse diagram graph of $I(S)$, then the set $\{x | (x] \in C \text{ in } I(S)\} = C_s$ is a convex in S . Moreover, if C is a prime convex in $I(S)$, then C_s is a prime convex in S . Let A be a nonempty convex of G , x a point of G with $x \notin A$ and A^* the least convex of the graph $G(I(S))$ of $I(S)$ with the property: $(z] \in A^*$ in $G(I(S))$ if $z \in A$ in G . Clearly, $(x] \notin A^*$ in $G(I(S))$. Because $I(S)$ is a distributive lattice, the graph $G(I(S))$ is a median graph and has thus the prime convex intersection property. Hence there is a prime convex C in $G(I(S))$ separating A^* and $(x]$, which implies that the prime convex C_s separates A and x in G . By Lemma 1, this proves [that G has the prime convex intersection property, and thus (i) holds for G .

(ii) Assume that the collection \mathcal{K} of the theorem is nonempty. We prove

that least element 0 of S belongs to $\cap \{\bar{P} | P \in \mathcal{K}\}$, from which the assertion follows. In fact, we prove the assertion for $n=3$; the proofs are the same for other values of n and hence they are omitted. Let $P_1, P_2, P_3 \in \mathcal{K}$ be three prime convexes of G such that $P_i \cap P_j \neq \emptyset$ and $P_1 \cap P_2 \cap P_3 = \emptyset$. The sets $P_1 \cap P_2, P_1 \cap P_3$ and $P_2 \cap P_3$ are convexes of G , and because S is finite, every one of them has a least element, and let them be $a \in P_1 \cap P_2, b \in P_1 \cap P_3$ and $c \in P_2 \cap P_3$. Assume that $0 \notin \cap \{\bar{P} | P \in \mathcal{K}\}$, which means that 0 belongs to at least one set of \mathcal{K} , say to P_1 . Because $0, a, b \in P_1$, then also $a \wedge b \wedge c \in P_1$. The relation $a, c \in P_2$ implies that $a \wedge c \in P_2$. On the other hand, $a \geq a \wedge c \geq a \wedge b \wedge c$, where $a, a \wedge b \wedge c \in P_1$, and thus $a \wedge c \in P_1$. Accordingly, $a \wedge c \in P_1 \cap P_2$, and because a is the least element in this convex, $a = a \wedge c \geq c$. Similarly we see that $b \leq c$. Because there is an upper bound c for a and b , the element $a \vee b$ exists, and as well known, an a - b geodesic goes over $a \vee b$ in the Hasse diagram graph of a finite distributive lattice. Thus $a \vee b \in P_1$. Because $c, b \in P_3$ and $c \geq a \vee b$, the element $a \vee b$ belongs to P_3 , and analogously we see that $a \vee b \in P_2$. Now, $a \vee b \in P_1 \cap P_2 \cap P_3$, which intersection should be empty, and hence the assumption $0 \notin \cap \{\bar{P} | P \in \mathcal{K}\}$ must be false. This proves the property (ii).

Assume conversely that G is a graph satisfying the properties (i) and (ii) of the theorem. We choose an arbitrary point from the set $\cap \{\bar{P} | P \in \mathcal{K}\}$ and denote it by h . Let a and b be two arbitrary points in V and let us consider the intersection $\langle h, a \rangle \cap \langle h, b \rangle \cap \langle a, b \rangle$. Because the convexes $\langle h, a \rangle, \langle h, b \rangle$ and $\langle a, b \rangle$ are the intersections of corresponding prime convexes, we can substitute the intersection $\langle h, a \rangle \cap \langle h, b \rangle \cap \langle a, b \rangle$ by the expression

$$(\cap \{P_i | P_i \text{ is a prime convex and } \langle h, a \rangle \subset P_i\}) \cap (\cap \{U_j | U_j \text{ is a prime convex and } \langle h, b \rangle \subset U_j\}) \cap (\cap \{W_k | W_k \text{ is a prime convex and } \langle a, b \rangle \subset W_k\}).$$

Now, $P_i \cap W_k, P_i \cap U_j, U_j \cap W_k \neq \emptyset$, and if $\langle h, a \rangle \cap \langle h, b \rangle \cap \langle a, b \rangle = \emptyset$, then $h \notin \cap \{\bar{P} | P \in \mathcal{K}\}$, which is a contradiction. Thus $\langle h, a \rangle \cap \langle h, b \rangle \cap \langle a, b \rangle \neq \emptyset$. Moreover, this intersection contains exactly one element. This can be seen as follows: Every prime convex P of G (or its complement \bar{P}) contains at least two of the points a, b, h . If the intersection $\langle h, a \rangle \cap \langle h, a \rangle \cap \langle a, b \rangle$ contains two disjoint points x and y , then every P (or \bar{P}) contains both x and y , and the convex x cannot be separated from the point y , which contradicts (i) by Lemma 1. Thus $\langle h, a \rangle \cap \langle h, b \rangle \cap \langle a, b \rangle = \{d\}$. According to Lemma 2, a convex $\langle x, z \rangle$ consists of points on x - z geodesics. Thus the relation $\{d\} = \langle h, a \rangle \cap \langle h, b \rangle \cap \langle a, b \rangle$ shows that every triple h, a, b , where a and b are arbitrary points of G , has a unique median.

We order now the points of V as follows:

$$a \leq b \iff a \text{ is on a } b\text{-}h \text{ geodesic} \iff a \in \langle h, b \rangle.$$

This definition suggests us to define the meet $a \wedge b$ as the unique median d of the points a, b and h . Assume that c is a point such that $c \in \langle h, a \rangle \cap \langle h, a \rangle$ and $c \notin \langle h, d \rangle$. The intersection $\langle h, d \rangle \cap \langle c, b \rangle$ is empty, because if x belongs to this intersection, then the $d\text{-}x\text{-}c\text{-}h$ path is a $d\text{-}h$ geodesic and $c \in \langle d, h \rangle$, which is a contradiction. There is a prime convex P separating the convexes $\langle h, d \rangle$ and $\langle c, b \rangle$: $\langle h, d \rangle \subset \bar{P}$ and $\langle c, b \rangle \subset P$. Indeed, as seen above, the points h, d and c have a median u which is on a $d\text{-}h$ geodesic and thus belongs to the convex $\langle d, h \rangle$. By the prime convex intersection property of G and Lemma 1, there is a prime convex P separating $\langle c, b \rangle$ and u ($\langle c, b \rangle \subset P$ and $u \in \bar{P}$). If now h or d belongs to P , then also u belongs to P because u is on a $c\text{-}h$ geodesic as well as on a $c\text{-}d$ geodesic. Thus $h, d \in \bar{P}$, whence also $\langle h, d \rangle \subset \bar{P}$. If $a \in \bar{P}$, then $c \in \bar{P}$ because it is on an $a\text{-}h$ geodesic, and thus a must belong to P . Because d is on an $a\text{-}b$ geodesic, the relation $a, b \in P$ implies a contradiction, and hence $c \in \langle h, d \rangle$. This proves that d is a maximum lower bound of a and b , and thus the order defined on V is a meetsemilattice order. Accordingly, V is a meetsemilattice with h as the least element. Because V is finite, it is a partial lattice. The Hasse diagram graph of V is isomorphic to G : When a line belongs to an $x\text{-}h$ geodesic, there is nothing to prove, and hence we assume that the line (a, b) of G does not belong to any $x\text{-}h$ geodesic. This is possible only if $d(a, h) = d(b, h)$. But then a, b and h have no median, which is absurd, and the isomorphism follows.

It remains to show that every set $(k] = \{v \mid v \in V \text{ and } v \leq k\}$ is a distributive lattice. By the order definition above, $\langle h, k \rangle = (k]$. Every convex A of a prime convex intersection graph induces a prime convex intersection graph. By Mulder and Schrijver [1, Theorem 2.2], a prime convex intersection graph $\langle h, k \rangle$ is a median graph (and then the Hasse diagram graph of a distributive lattice with h as the least element and k as the greatest element by [1, Theorem 3.1]) if its prime convexes needed to separate its convexes satisfy the Helly property. The prime convexes needed to separate the convexes of $\langle h, k \rangle$ are obtained from the prime convexes of \mathcal{K} by intersecting them with $\langle h, k \rangle$. Let now P_1, P_2, \dots, P_m be prime convexes of \mathcal{K} such that $P_i \cap P_j \cap \langle h, k \rangle \neq \emptyset$. We denote the corresponding prime convexes of $\langle h, k \rangle$ by $P_i^0 = P_i \cap \langle h, k \rangle$. By Lemma 2, the convex $\langle h, k \rangle$ consists of points on $h\text{-}k$ geodesics in G . If $h, k \notin P_i^0$, then P_i^0 is not prime because its every point is on some $h\text{-}k$ geodesic. Hence either h or k belongs to P_i^0 . The relation $h \in P_i^0$ contradicts the property $h \in \bigcap \{\bar{P} \mid P \in \mathcal{K}\}$, and thus $k \in P_i^0$, and this relation holds for every $i, i = 1, \dots, m$. Then $k \in P_1^0 \cap P_2^0 \cap \dots \cap P_m^0$, and the Helly property of the prime convexes needed to separate the convexes of $\langle h,$

k follows. This proves the distributivity of $\langle h, k \rangle = (k]$, and thus G is the Hasse diagram graph of a finite distributive partial lattice.

The author likes to express his sincere thanks to the referee for his valuable suggestions and comments.

References

- [1] Bandelt, H. J. and Hedlikova, J., Median algebras, *Discrete Math.* **45** (1983), 1-30.
- [2] Bandelt, H. J. and Mulder, H. M., Infinite median graphs, $(0, 2)$ -graphs and hypercubes, *J. Graph Theory* **7** (1983), 487-497.

Faculty of Technology
University of Oulu
90570 Oulu
Finland