

A NOTE ON THE PROJECTIVE NORMALITY OF SPECIAL LINE BUNDLES ON ABELIAN VARIETIES

By

Akira OHBUCHI

Dedicated to Professor Yukihiro Kodama on his 60th birthday

Introduction.

Let L be an ample line bundle on an abelian variety A of dimension g defined over an algebraically closed field k . It is well known that $L^{\otimes 2}$ is base point free and $L^{\otimes 3}$ is very ample and projectively normal. Moreover we know that

$$\Gamma(A, L^{\otimes a}) \otimes \Gamma(A, L^{\otimes b}) \longrightarrow \Gamma(A, L^{\otimes a+b})$$

is surjective if $a \geq 2$ and $b \geq 3$ (Koizumi [3], Sekiguchi [8], [9]). But in the case of $a=b=2$, this map is not surjective in general. In this paper we determine the condition of projective normality of $L^{\otimes 2}$ for some ample line bundle L . Our result is as follows.

THEOREM. *If L is a symmetric ample line bundle of separable type, $l(A, L)$ is odd and assume that $\text{char}(k) \neq 2$, then $L^{\otimes 2}$ is projectively normal if and only if $Bs|L| \cap A[2] = \emptyset$.*

In §1 we prove the above theorem for abelian varieties defined over \mathbf{C} . In §2 we give the Mumford's theory of a theta group (Mumford [4], [5]). In §3 we prove the above theorem in general by the theory in §2.

Notations.

$\text{char}(k)$: The characteristic of a field k

f^* : The pull back defined by a morphism f

\underline{L} : The invertible sheaf associated to a line bundle L

\mathcal{O}_A : The invertible sheaf of a variety A

(L^g) : The self intersection number

$|L|$: The set of all effective Cartier divisors which define a line bundle L

$Bs|L|$: The set defined by $\bigcap_{D \in |L|} D$

$\Gamma(A, L)$: The set of global sections of a line bundle L

$l(A, L)$: The dimension of $\Gamma(A, L)$ as a vector space

Received May 12, 1987.

T_x : The translation morphism on an abelian variety A defined by $T_x(y) = x + y$ where x and y are elements of A

$K(L)$: The subgroup of an abelian variety A defined by $\{x \in A; T_x^*L \cong L\}$ where L is a line bundle on A

$A[n]$: The set of all points of order n of an abelian variety A

\mathbf{Z} : The ring of integers

\mathbf{R} : The field of real numbers

\mathbf{C} : The field of complex numbers

§ 1. The \mathbf{C} case.

First we recall a definition of projective normality.

DEFINITION. Let M be an ample line bundle on an abelian variety A . We call that M is projectively normal if

$$\Gamma(A, M)^{\otimes n} \longrightarrow \Gamma(A, M^{\otimes n})$$

is surjective for every $n \geq 1$.

Next we define a theta function defined on \mathbf{C}^g .

DEFINITION. Let m', m'' be elements of \mathbf{R}^g and let τ be an element of a Siegel upper half space H_g . We define $\theta \begin{bmatrix} m' \\ m'' \end{bmatrix}(\tau, z)$ by

$$\theta \begin{bmatrix} m' \\ m'' \end{bmatrix}(\tau, z) = \sum_{\zeta \in \mathbf{Z}} e((1/2)^t(\zeta + m')\tau(\zeta + m') + {}^t(\zeta + m')(z + m''))$$

where $e(x)$ means $e^{2\pi\sqrt{-1}x}$ and z is contained in \mathbf{C}^g .

Let d_1, \dots, d_g be positive integers with $d_1 | \dots | d_g$. We define an integral matrix e by

$$\begin{bmatrix} d_1 & & & 0 \\ & \ddots & & \\ & & \ddots & \\ 0 & & & d_g \end{bmatrix}.$$

For an element τ of H_g we define an abelian variety A by $\mathbf{C}^g / \langle \tau, e \rangle$ where $\langle \tau, e \rangle$ is a lattice subgroup of \mathbf{C}^g defined by $\tau\mathbf{Z}^g + e\mathbf{Z}^g$. Let \mathcal{A} be a Riemann form on $\langle \tau, e \rangle$ defined by

$$\mathcal{A}(\tau x + ey, \tau x' + ey') = {}^t x e y' - {}^t x' e y$$

where x, x', y, y' are elements of \mathbf{Z}^g . It is well known that this \mathcal{A} defines an

algebraic equivalence class of line bundles on A . Now we take a line bundle L on A satisfying that L is symmetric and the global sections of L are generated by $\theta \begin{bmatrix} \eta \\ 0 \end{bmatrix}(\tau, z)$ where η runs over a complete system of representative of $(1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$.

LEMMA 1. *The basis of $\Gamma(A, L^{\otimes 2^n})$ is given by $\theta \begin{bmatrix} \eta \\ 0 \end{bmatrix}(2^n\tau, 2^n z)$ where η runs over a complete system of representative of $(1/2^n d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/2^n d_g)\mathbf{Z}/\mathbf{Z}$ ($n=1, 2, \dots$). Moreover $\Gamma(A, L^{\otimes s})$ is generated by $\theta \begin{bmatrix} \xi \\ \sigma \end{bmatrix}(\tau, 2z)$ where ξ runs over a complete system of representative of $(1/2d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/2d_g)\mathbf{Z}/\mathbf{Z}$ and σ runs over a complete system of representative of $((1/2)\mathbf{Z}/\mathbf{Z})^g$.*

PROOF. This is well known fact (cf. Igusa [2], p. 72, Theorem 4, and p. 84, Theorem 6).

LEMMA 2 (Multiplication formula). *If $\eta', \eta'', \xi', \xi''$ are contained in \mathbf{R}^g and τ is contained in H_g , then*

$$\begin{aligned} \theta \begin{bmatrix} \eta' \\ \eta'' \end{bmatrix}(\tau, z) \theta \begin{bmatrix} \xi' \\ \xi'' \end{bmatrix}(\tau, z) &= (1/2^g) \sum_{a'' \in ((1/2)\mathbf{Z}/\mathbf{Z})^g} e(-2^t \eta' a'') \\ &\quad \cdot \theta \begin{bmatrix} \eta' + \xi' \\ (\eta'' + \xi'')/2 + a'' \end{bmatrix}(\tau/2, (z_1 + z_2)/2) \\ &\quad \cdot \theta \begin{bmatrix} \eta' - \xi' \\ (\eta'' - \xi'')/2 + a'' \end{bmatrix}(\tau/2, (z_1 - z_2)/2) \end{aligned}$$

where z_1 and z_2 are contained in \mathbf{C}^g .

PROOF. This is also well known fact (cf. Igusa [2], p. 139, Theorem 2).

LEMMA 3. *If η, η' are elements of $(1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$, d_g is odd and $\varepsilon, \varepsilon'$ are contained in \mathbf{Z}^g , then*

$$\begin{aligned} \sum_{\sigma \in ((\mathbf{Z}/2\mathbf{Z})^g)} (-1)^{t_{\sigma\varepsilon\varepsilon'}} \theta \begin{bmatrix} \eta + (\sigma/2) \\ 0 \end{bmatrix}(2\tau, 2z) \theta \begin{bmatrix} \eta' + (\sigma + \varepsilon)/2 \\ 0 \end{bmatrix}(2\tau, 2z) \\ = \theta \begin{bmatrix} \eta + \eta' + (\varepsilon/2) \\ e\varepsilon'/2 \end{bmatrix}(\tau, 2z) \theta \begin{bmatrix} \eta - \eta' + (\varepsilon/2) \\ e\varepsilon'/2 \end{bmatrix}(\tau, 0). \end{aligned}$$

PROOF. By lemma 2, we obtain

$$\begin{aligned} & \theta \begin{bmatrix} \eta + (\sigma/2) \\ 0 \end{bmatrix} (2\tau, 2z) \theta \begin{bmatrix} \eta' + (\sigma + \varepsilon)/2 \\ 0 \end{bmatrix} (2\tau, 2z) \\ &= (1/2^g) \sum_{a'' \in ((1/2)\mathbf{Z}/\mathbf{Z})^g} e^{-2^t(\eta + (\sigma/2))a''} \\ & \quad \cdot \theta \begin{bmatrix} \eta + \eta' + (\varepsilon/2) \\ a'' \end{bmatrix} (\tau, 2z) \theta \begin{bmatrix} \eta - \eta' + (\varepsilon/2) \\ a'' \end{bmatrix} (\tau, 0). \end{aligned}$$

Hence

$$\begin{aligned} & \sum_{\sigma \in (\mathbf{Z}/2\mathbf{Z})^g} (-1)^{t\sigma\varepsilon\varepsilon'} \theta \begin{bmatrix} \eta + (\sigma/2) \\ 0 \end{bmatrix} (2\tau, 2z) \theta \begin{bmatrix} \eta' + (\sigma + \varepsilon)/2 \\ 0 \end{bmatrix} (2\tau, 2z) \\ &= (1/2^g) \sum_{a'' \in ((1/2)\mathbf{Z}/\mathbf{Z})^g} e^{-2^t\eta a''} \sum_{\sigma \in (\mathbf{Z}/2\mathbf{Z})^g} (-1)^{t\sigma(\varepsilon\varepsilon' - 2a'')} \\ & \quad \cdot \theta \begin{bmatrix} \eta + \eta' + (\varepsilon/2) \\ a'' \end{bmatrix} (\tau, 2z) \theta \begin{bmatrix} \eta - \eta' + (\varepsilon/2) \\ a'' \end{bmatrix} (\tau, 0) \\ &= e^{(t\eta\varepsilon\varepsilon')} \theta \begin{bmatrix} \eta + \eta' + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 2z) \theta \begin{bmatrix} \eta - \eta' + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 0) \\ &= \theta \begin{bmatrix} \eta + \eta' + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 2z) \theta \begin{bmatrix} \eta - \eta' + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 0). \end{aligned}$$

Therefore we obtain this lemma.

LEMMA 4. *If M is an ample line bundle on an abelian variety A , then*

$$\Gamma(A, M^{\otimes a}) \otimes \Gamma(A, M^{\otimes b}) \longrightarrow \Gamma(A, M^{\otimes a+b})$$

is surjective for $a \geq 2$ and $b \geq 3$.

PROOF. See Koizumi [3] or Sekiguchi [8], [9].

LEMMA 5. *Under the notation of lemma 3, if there exists some $\eta_0 \in (1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$ with $\theta \begin{bmatrix} \eta_0 + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 0) \neq 0$, then $\theta \begin{bmatrix} \eta + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 2z)$ is in the image of $\Gamma(A, L^{\otimes 2})^{\otimes 2} \rightarrow \Gamma(A, L^{\otimes 4})$ for every $\eta \in (1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$.*

PROOF. Let η_1 be an element of $(1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$. In this case, we obtain that

$$\theta \begin{bmatrix} 2\eta_1 + \eta_0 + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 2z) \theta \begin{bmatrix} \eta_0 + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 0)$$

is contained in the image of $\Gamma(A, L^{\otimes 2})^{\otimes 2} \rightarrow \Gamma(A, L^{\otimes 4})$ by lemma 3. Hence $\theta \begin{bmatrix} 2\eta_1 + \eta_0 + (\varepsilon/2) \\ \varepsilon\varepsilon'/2 \end{bmatrix} (\tau, 2z)$ is contained in the image of $\Gamma(A, L^{\otimes 2})^{\otimes 2} \rightarrow \Gamma(A, L^{\otimes 4})$. As

d_g is odd, therefore the set $\{2\eta_1 + \eta_0; \eta_1 \in (1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}\}$ is equal to $(1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$.

Therefore we obtain this lemma.

LEMMA 6. *Under the assumption of lemma 3, the following conditions are equivalent:*

- a) *For every $\varepsilon, \varepsilon' \in \mathbf{Z}^g$, there exists some $\eta \in (1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$ with $\theta \begin{bmatrix} \eta + (\varepsilon/2) \\ e\varepsilon'/2 \end{bmatrix}(\tau, 0) \neq 0$;*
- b) *$Bs|L| \cap A[2] = \emptyset$.*

PROOF. As

$$\theta \begin{bmatrix} m' + \xi' \\ m'' + \xi'' \end{bmatrix}(\tau, z) = e((1/2)^t \xi' \tau \xi' + {}^t \xi'(z + \xi'' + m'')) \theta \begin{bmatrix} m' \\ m'' \end{bmatrix}(\tau, z + \tau \xi' + \xi'')$$

(cf. Igusa [2], p. 50, ($\theta.3$)), therefore the condition $\theta \begin{bmatrix} \eta + (\varepsilon/2) \\ e\varepsilon'/2 \end{bmatrix}(\tau, 0) \neq 0$ is equivalent to $\theta \begin{bmatrix} \eta \\ 0 \end{bmatrix}(\tau, (\tau\varepsilon + e\varepsilon')/2) \neq 0$. Hence this lemma is clear because $A = \mathbf{C}^g / \langle \tau, e \rangle$.

THEOREM. *If $l(A, L)$ is odd, then $L^{\otimes 2}$ is projectively normal if and only if $Bs|L| \cap A[2] = \emptyset$.*

PROOF. By lemma 1, a basis of $\Gamma(A, L^{\otimes 4})$ consists of $\theta \begin{bmatrix} \eta \\ \sigma \end{bmatrix}(\tau, 2z)$ where η runs over a complete system of representative of $(1/2d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/2d_g)\mathbf{Z}/\mathbf{Z}$ and σ runs over a complete system of representative of $((1/2)\mathbf{Z}/\mathbf{Z})^g$. Hence $\Gamma(A, L^{\otimes 2})^{\otimes 2} \rightarrow \Gamma(A, L^{\otimes 4})$ is surjective if and only if for every $\varepsilon, \varepsilon' \in \mathbf{Z}^g$, there exists some $\eta_0 \in (1/d_1)\mathbf{Z}/\mathbf{Z} \oplus \cdots \oplus (1/d_g)\mathbf{Z}/\mathbf{Z}$ such that $\theta \begin{bmatrix} \eta_0 + (\varepsilon/2) \\ e\varepsilon'/2 \end{bmatrix}(\tau, 0) \neq 0$ by lemma 5. Hence we obtain this theorem by lemma 4 and lemma 6.

§ 2. Review of a theta group.

In this section we recall the Mumford's theory for a theta group (cf. Mumford [4], [5]). Let A be an abelian variety of dimension g defined over an algebraically closed field k with $\text{char}(k) \neq 2$. We fix these notations.

DEFINITION. *Let M be an ample line bundle on A . We call that M is of separable type if $\text{char}(k) \nmid l(A, M)$.*

DEFINITION. Let M be an ample line bundle on A and be of separable type. We define a theta group $G(M)$ by

$$\{(x, \phi); x \in K(M) \text{ and } \phi: M \xrightarrow{\sim} T_x^*M\}.$$

This $G(M)$ is a group by the following way:

$$(x, \phi) \cdot (y, \rho) = (x + y, T_x^* \phi \cdot \rho).$$

It is well known that $K(M)$ is isomorphic to the following abelian group via Weil pairing:

$$K(M) = \mathbf{Z}/d_1\mathbf{Z} \oplus \cdots \oplus \mathbf{Z}/d_g\mathbf{Z} \oplus \overbrace{(\mathbf{Z}/d_1\mathbf{Z} \oplus \cdots \oplus \mathbf{Z}/d_g\mathbf{Z})}$$

where $d_1 | \cdots | d_g$ and \hat{G} means $\text{Hom}(G, k^*)$ for a group G . Here we denote by $k^* = k - \{0\}$. In this situation, we set $\delta_M = (d_1, \dots, d_g)$ and put $H(\delta_M) = \mathbf{Z}/d_1\mathbf{Z} \oplus \cdots \oplus \mathbf{Z}/d_g\mathbf{Z}$. We define a Heisenberg group $G(\delta_M)$.

DEFINITION. In the above notations, we define a Heisenberg group $G(\delta_M)$ by

$$G(\delta_M) = k^* \times H(\delta_M) \times H(\hat{\delta}_M).$$

This $G(\delta_M)$ is a group by the following way:

$$(t, (x, m)) \cdot (t', (x', m')) = (tt'm'(x), (x + x', m + m'))$$

where $x, x' \in H(\delta_M)$, $m, m' \in H(\hat{\delta}_M)$ and $t, t' \in k^*$. The following theorem is fundamental.

THEOREM. In the above notations, the following two horizontal sequences are exact and isomorphic:

$$\begin{array}{ccccccc} 0 & \longrightarrow & k^* & \longrightarrow & G(M) & \longrightarrow & K(M) \longrightarrow 0 \\ & & \parallel & \curvearrowright & \downarrow & \curvearrowright & \downarrow \\ 0 & \longrightarrow & k^* & \longrightarrow & G(\delta_M) & \longrightarrow & H(\delta_M) \times H(\hat{\delta}_M) \longrightarrow 0. \end{array}$$

PROOF. See Mumford [4], p. 294, Corollary of Th. 1.

DEFINITION. Let $z = (x, \phi)$ be an element of $G(M)$. We define a map U_z as follows:

$$U_z: \Gamma(A, M) \xrightarrow{\Gamma(\phi)} \Gamma(A, T_x^*M) \xrightarrow{T_{-x}^*} \Gamma(A, M).$$

It is clear that $\Gamma(A, M)$ is a $G(M)$ -module by this way. Next we define a vector space $V(\delta_M)$ and its $G(\delta_M)$ -module structure.

DEFINITION. The vector space $V(\delta_M)$ is defined as follows:

$$V(\delta_M) = \text{the set of all maps from } H(\delta_M) \text{ to } k.$$

Let $(t, (x, m))$ be an element of $G(\delta_M)$. We define an automorphism $U_{(t, (x, m))}$ of $V(\delta_M)$ as follows:

$$U_{(t, (x, m))}(f)(y) = tm(y)f(x+y)$$

where $f \in V(\delta_M)$ and $y \in H(\delta_M)$.

The following theorem is also fundamental.

THEOREM. If $\alpha: G(M) \xrightarrow{\sim} G(\delta_M)$ is an isomorphism given in the above theorem, then we obtain an isomorphism

$$\Gamma(A, M) \xrightarrow{\sim} V(\delta_M)$$

as $G(M) \cong G(\delta_M)$ -modules.

PROOF. See Mumford [4], p. 295, proposition 3, and p. 297, theorem 2.

DEFINITION. Let x be an element of $H(\delta_M)$. We define $\delta_x \in V(\delta_M)$ by $\delta_x(y) = 1$ if $y = x$ and $\delta_x(y) = 0$ if $y \neq x$.

It is clear that $U_{(t, (x, m))}(\delta_u) = tm(u-x)\delta_{u-x}$.

§ 3. The general case.

Let L be an ample line bundle on an abelian variety A of dimension g . Throughout of this section, we assume that L is of separable type and $l(A, L)$ is odd and L is symmetric. We fix an isomorphism $G(L^{\otimes 4}) \cong G(4\delta_L)$ and identify two vector spaces $\Gamma(A, L^{\otimes 4})$ and $V(4\delta_L)$ by the isomorphism in § 2.

LEMMA 1. Let f be an element of $V(4\delta_L) \cong \Gamma(A, L^{\otimes 4})$ defined by $f = \sum_{u \in H(4\delta_L) \text{ and } 2u=0} \delta_u$. Then f is in the image of $2_A^*: \Gamma(A, L) \rightarrow \Gamma(A, L^{\otimes 4})$ for some isomorphism $2_A^*L \cong L^{\otimes 4}$ where $2_A(x) = 2x$ ($x \in A$).

PROOF. This lemma is trivial.

By the above lemma, we obtain $\theta \in \Gamma(A, L)$ with $2_A^*\theta = f$. We fix these notations through this section.

DEFINITION. Let x be an element of $H(4\delta_L)$ and σ be an element of $H(\hat{\delta}_L)$. We define an element of $\theta \begin{bmatrix} x \\ \sigma \end{bmatrix}$ of $V(4\delta_L)$ by

$$\theta \begin{bmatrix} x \\ \sigma \end{bmatrix} = U_z(2_A^* \theta)$$

where z is an element of $G(L^{\otimes 4})$ corresponding to $(1, (x, \sigma))$ which is an element of $G(4\delta_L)$.

LEMMA 2. Let x, u be elements of $H(4\delta_L)$ and σ, u^* be elements of $H(\hat{\delta}_L)$. If $2u=0$ and $2u^*=0$, then $\theta \begin{bmatrix} x+u \\ \sigma+u^* \end{bmatrix} = u^*(x)\theta \begin{bmatrix} x \\ \sigma \end{bmatrix}$.

PROOF. By the definition, we obtain that

$$\theta \begin{bmatrix} x \\ \sigma \end{bmatrix} = \sum_{2\zeta=0} \sigma(\zeta-x)\delta_{\zeta-x}.$$

Therefore

$$\begin{aligned} \theta \begin{bmatrix} x+u \\ \sigma+u^* \end{bmatrix} &= \sum_{2\zeta=0} (\sigma+u^*)(\zeta-x-u)\delta_{\zeta-x-u} \\ &= \sum_{2\zeta=0} (\sigma+u^*)(\zeta-x)\delta_{\zeta-x} \\ &= \sum_{2\zeta=0} u^*(\zeta-x)\sigma(\zeta-x)\delta_{\zeta-x}. \end{aligned}$$

As $2u^*=0$, hence $u^*(\zeta)=1$ for every $\zeta \in H(4\delta_L)$ with $2\zeta=0$. Therefore

$$\begin{aligned} \theta \begin{bmatrix} x+u \\ \sigma+u^* \end{bmatrix} &= u^*(-x) \sum_{2\zeta=0} \sigma(\zeta-x)\delta_{\zeta-x} \\ &= u^*(x)\theta \begin{bmatrix} x \\ \sigma \end{bmatrix}. \end{aligned}$$

So we obtain this lemma.

LEMMA 3. The vector space $\Gamma(A, L^{\otimes 4})$ is spanned by the elements $\theta \begin{bmatrix} x \\ \sigma \end{bmatrix}$ where $x \in H(4\delta_L)$ and $\sigma \in (\hat{\mathbf{Z}}/4\mathbf{Z})^g$ which is regarded the subgroup of order of $H(4\hat{\delta}_L)$.

PROOF. We regard that $H(\delta_L)$ and $(\mathbf{Z}/4\mathbf{Z})^g$ are the subgroups of $H(4\delta_L)$ in the canonical way. For every $\xi \in H(\delta_L)$, $\tau \in (\mathbf{Z}/4\mathbf{Z})^g$ and $\sigma \in (\hat{\mathbf{Z}}/4\mathbf{Z})^g$, we obtain

$$\begin{aligned} \theta \begin{bmatrix} \xi+\tau \\ \sigma \end{bmatrix} &= \sum_{2\zeta=0} \sigma(\zeta-\xi-\tau)\delta_{\zeta-\xi-\tau} \\ &= \sum_{2\zeta=0} \sigma(\zeta-\tau)\delta_{\zeta-\xi-\tau} \end{aligned}$$

because as $l(A, L)$ is odd, $\sigma(\xi)=1$. Therefore

$$\begin{aligned} \sum_{\sigma \in (\mathbb{Z}/4\mathbb{Z})^g} \sigma(\tau) \theta \begin{bmatrix} \xi + \tau \\ \sigma \end{bmatrix} &= \sum_{\sigma \in (\mathbb{Z}/4\mathbb{Z})^g} \sum_{2\zeta=0} \sigma(\tau) \sigma(\zeta - \tau) \delta_{\zeta - \xi - \tau} \\ &= \sum_{2\zeta=0} \left(\sum_{\sigma \in (\mathbb{Z}/4\mathbb{Z})^g} \sigma(\zeta) \right) \delta_{\zeta - \xi - \tau} \\ &= 4^g \delta_{\zeta - \xi - \tau}. \end{aligned}$$

Therefore we obtain this lemma.

Let x be a closed point of A and M be an ample line bundle on A of separable type. We define $M(x)$ by

$$M(x) = \underline{M}_x \otimes_{\mathcal{O}_{A,x}} k(x)$$

where \underline{M}_x and $\mathcal{O}_{A,x}$ are the stalk of M and \mathcal{O}_A at x respectively, and $k(x)$ is a residue field of $\mathcal{O}_{A,x}$. It is clear that $M(x) \cong k$. We choose an isomorphism $\lambda_0: M(0) \xrightarrow{\cong} k$. We fix an isomorphism $G(M) \cong G(\delta_M)$. For every $w \in K(M)$, we take $(w, \phi_w) \in G(M)$ which is corresponding to an element of $G(\delta_M)$ with a type $(1, (x, m))$ by the above isomorphism.

DEFINITION. We defined $\lambda_w: M(w) \rightarrow k$ by

$$\lambda_w: M(w) = (T_x^* M)(0) \xleftarrow{\phi_w(0)} M(0) \xrightarrow{\lambda_0} k$$

where $w \in K(M)$ and $\phi_w(0)$ is given by ϕ_w .

DEFINITION. Under the above notations, we define q_w^M by

$$q_w^M: \Gamma(A, M) \xrightarrow{\text{canonical map}} M(w) \xrightarrow{\lambda_w} k.$$

REMARK. For any $z = (x, \phi) \in G(M)$ and any $s \in \Gamma(A, M)$, the conditions $q_w^M(U_z(s)) = 0$ and $q_{w+x}^M(s) = 0$ are equivalent.

REMARK. The condition that $q_w^M(s) = 0$ for every $s \in \Gamma(A, M)$ implies that w is contained in $B_S | M |$.

REMARK. If M is a symmetric ample line bundle on A , then the conditions $q_w^{M^{\otimes 4}}(2_A^* s) = 0$ and $q_{2w}^M(s) = 0$ are equivalent for any $s \in \Gamma(A, M)$.

DEFINITION. We define $q_{L^{\otimes 4}}(x)$ by

$$q_{L^{\otimes 4}}(x) = q_0^{L^{\otimes 4}}(\delta_x)$$

where $x \in K(L^{\otimes 4})$ and $\delta_x \in V(4\delta_L) \cong \Gamma(A, L^{\otimes 4})$. Moreover we define $q \begin{bmatrix} x \\ \sigma \end{bmatrix}$ by

$$q \begin{bmatrix} x \\ \sigma \end{bmatrix} = q_0^{L^{\otimes 4}} \left(\theta \begin{bmatrix} x \\ \sigma \end{bmatrix} \right)$$

where $x \in H(4\delta_L)$ and $\sigma \in (\mathbf{Z}/4\mathbf{Z})^g$.

Now the isomorphism $G(L^{\otimes 4}) \cong G(4\delta_L)$ induces an isomorphism $G(L^{\otimes 2}) \cong G(2\delta_L)$; these isomorphisms define the symmetric theta structure for $(L^{\otimes 2}, L^{\otimes 4})$ (cf. Mumford [4], p. 317). We identify the two vector spaces $\Gamma(A, L^{\otimes 2})$ and $V(2\delta_L)$ by means of the isomorphism $G(L^{\otimes 2}) \cong G(2\delta_L)$.

LEMMA 4 (*Multiplication formula*). *If δ_x and $\delta_{x'}$ are elements of $\Gamma(A, L^{\otimes 2})$, then*

$$\delta_x \cdot \delta_{x'} = \sum_{\zeta \in H(4\delta_L) \text{ and } 2\zeta=0} q_{L^{\otimes 4}}(x - x' + \zeta) \delta_{x+x'+\zeta}$$

where \cdot is a canonical map $\Gamma(A, L^{\otimes 2})^{\otimes 2} \rightarrow \Gamma(A, L^{\otimes 4})$ and $\underline{x}, \underline{x}' \in H(4\delta_L)$ satisfying $2\underline{x} = x$, $2\underline{x}' = x'$. Here we regard $H(2\delta_L)$ as a subgroup of $H(4\delta_L)$ in the canonical way.

PROOF. See Mumford [4], p. 330.

Let x, x' be elements of $H(\delta_L)$, and ξ, ξ' be elements of $(\mathbf{Z}/2\mathbf{Z})^g$. We take $\underline{x}, \underline{x}' \in H(\delta_L)$ and $\underline{\xi}, \underline{\xi}' \in (\mathbf{Z}/4\mathbf{Z})^g$ satisfying $2\underline{x} = x$, $2\underline{x}' = x'$, $2\underline{\xi} = \xi$ and $2\underline{\xi}' = \xi'$.

LEMMA 5. *Under the above notations,*

$$\delta_{x+\xi} \cdot \delta_{x'+\xi'} = (1/4^g) \sum_{\sigma \in (\mathbf{Z}/4\mathbf{Z})^g} \sigma(\xi) q \begin{bmatrix} -\underline{x} + \underline{x}' - \underline{\xi} + \underline{\xi}' \\ \sigma \end{bmatrix} \theta \begin{bmatrix} -\underline{x} - \underline{x}' - \underline{\xi} - \underline{\xi}' \\ \sigma \end{bmatrix}$$

PROOF. For $\delta_{x+\xi}$ and $\delta_{x'+\xi'} \in \Gamma(A, L^{\otimes 2})$, we obtain that

$$\begin{aligned} \delta_{x+\xi} \cdot \delta_{x'+\xi'} &= \sum_{\zeta \in H(4\delta_L) \text{ and } 2\zeta=0} q_{L^{\otimes 4}}(x - x' + \underline{\xi} - \underline{\xi}' + \zeta) \delta_{\underline{x} + \underline{x}' + \underline{\xi} + \underline{\xi}' + \zeta} \\ &= \sum_{2\zeta=0} q_{L^{\otimes 4}}(x - x' + \underline{\xi} - \underline{\xi}' + \zeta) (1/4^g) \sum_{\sigma \in (\mathbf{Z}/4\mathbf{Z})^g} \sigma(-\underline{\xi} - \underline{\xi}' - \zeta) \\ &\quad \cdot \theta \begin{bmatrix} -\underline{x} - \underline{x}' - \underline{\xi} - \underline{\xi}' - \zeta \\ \sigma \end{bmatrix} \\ &= (1/4^g) \sum_{\sigma} \sum_{2\zeta=0} \sigma(-\underline{\xi} - \underline{\xi}' - \zeta) q_{L^{\otimes 4}}(x - x' + \underline{\xi} - \underline{\xi}' + \zeta) \\ &\quad \cdot \theta \begin{bmatrix} -\underline{x} - \underline{x}' - \underline{\xi} - \underline{\xi}' - \zeta \\ \sigma \end{bmatrix}. \end{aligned}$$

On the other hand, $\theta \begin{bmatrix} x+u \\ \sigma+u^* \end{bmatrix} = u^*(x) \theta \begin{bmatrix} x \\ \sigma \end{bmatrix}$ for $2u=0$ and $2u^*=0$. Moreover in above situation, $\sigma(\underline{x}) = \sigma(\underline{x}') = 1$. Hence

$$\begin{aligned} \delta_{x+\xi} \cdot \delta_{x'+\xi'} &= (1/4^g) \sum_{\sigma} \sigma(-2\xi) \theta \left[\begin{matrix} -x-x'-\xi-\xi' \\ \sigma \end{matrix} \right] \sum_{\zeta} \sigma(x-x'+\xi-\xi'+\zeta) \\ &\quad \cdot q_{L^{\otimes 4}}(x-x'+\xi-\xi'+\zeta) \\ &= (1/4^g) \sum_{\sigma} \sigma(\xi) \theta \left[\begin{matrix} -x-x'-\xi-\xi' \\ \sigma \end{matrix} \right] \\ &\quad \cdot q_0^{L^{\otimes 4}}(\sum_{\zeta} \sigma(x-x'+\xi-\xi'+\zeta) \delta_{x-x'+\xi-\xi'+\zeta}) \\ &= (1/4^g) \sum_{\sigma} \sigma(\xi) \theta \left[\begin{matrix} -x-x'-\xi-\xi' \\ \sigma \end{matrix} \right] q \left[\begin{matrix} -x+x'-\xi-\xi' \\ \sigma \end{matrix} \right] \end{aligned}$$

Therefore we obtain this lemma.

THEOREM. *Under the above notations, $L^{\otimes 2}$ is projectively normal if and only if $Bs|L| \cap A[2] = \phi$.*

PROOF. Replacing the lemmas for the theorem in § 1 by the above lemmas, the proof of the theorem in § 1 still works in general case.

COROLLARY. *If M is an ample line bundle and is of separable type on an abelian variety A , then $Bs|M| = \phi$ and $l(A, M) = \text{odd}$ imply that $M^{\otimes 2}$ is projectively normal.*

To conclude this section, we give an easy criterion for the base point freeness of a line bundle M on an abelian variety A . We assume that M is of separable type. Let $\alpha: G(M) \rightarrow G(\delta_M)$ be an isomorphism. As α induces $\bar{\alpha}: K(M) \rightarrow H(\delta_M) \oplus H(\hat{\delta}_M)$, we put $H(M)$ by $\bar{\alpha}^{-1}(H(\delta_M))$. Let B be an abelian variety defined by $A/H(M)$ and $\pi: A \rightarrow B$ the canonical morphism. In this situation, the line bundle M is given by $M \cong \pi^*N$ where N is a principal line bundle on B . In this notations, we obtain the following proposition.

PROPOSITION. $Bs|M| = \pi^{-1}(\bigcap_{x \in \pi(K(M))} T_x^* \theta)$ where $\theta \in |N|$.

PROOF. As there exists a canonical isomorphism

$$\Gamma(A, M) \cong \bigoplus_{x \in \pi(K(M))} \Gamma(B, T_x^*N)$$

therefore this proposition is clear.

The following proposition is also clear.

PROPOSITION. *Let M be as in above. If $Bs|M|$ is finite and $(M^g) > (g!)^2$, then $Bs|M| = \phi$.*

PROOF. If $Bs|M| \neq \emptyset$, then there is a point $q \in Bs|M|$. By the definition of $K(M)$, $q+K(M)$ is also contained in $Bs|M|$. As

$$\text{the cardinality of } Bs|M| \leq (M^g),$$

hence

$$\text{the order of } K(M) = ((M^g)/g!)^2 \leq (M^g).$$

Therefore we obtain $(M^g) \leq (g!)^2$.

References

- [1] Geemen, B., Schottky-Jung relations and vector bundles on hyperelliptic curves. Thesis University of Utrecht (1985).
- [2] Igusa, J., Theta functions. Springer-Verlag (1972).
- [3] Koizumi, S., Theta relations and projective normality of abelian varieties. Amer. J. Math. **98**, 865-889 (1976).
- [4] Mumford, D., On the equation defining abelian varieties I. Invent. Math. **1**, 289-354 (1966).
- [5] Mumford, D., Abelian varieties. Oxford University Press (1970).
- [6] Ohbuchi, A., Some remarks on ample line bundles on abelian varieties. Manuscripta Math. **57**, 225-238 (1987).
- [7] Saint-Donat, B., Projective models of K3 surfaces. Amer. J. Math. **96**, 602-632 (1974).
- [8] Sekiguchi, T., On projective normality of abelian varieties. J. Math. Soc. Japan **28**, 307-322 (1976).
- [9] Sekiguchi, T., On projective normality of abelian varieties II. J. Math. Soc. Japan **29**, 709-727 (1977).

Department of Mathematics
 Faculty of Education
 Yamaguchi University
 16771-1, Oh-aza Yoshida
 Yamaguchi-shi, Yamaguchi 753,
 Japan