

NEW CRITERIA FOR MEROMORPHIC p -VALENT STARLIKE FUNCTIONS

By

M.K. AOUF and H.M. HOSSEN

Abstract. Let $B_n(\alpha)$ be the class of functions of the form

$$f(z) = \frac{a_{-p}}{z^p} + \sum_{k=0}^{\infty} a_k z^k \quad (a_{-p} \neq 0, p \in N = \{1, 2, \dots\})$$

which are regular in the punctured disc $U^* = \{z : 0 < |z| < 1\}$ and satisfying

$$\operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{D^n f(z)} - (p+1) \right\} < -\alpha \quad (n \in N_0 = \{0, 1, \dots\}, |z| < 1, 0 \leq \alpha < p),$$

where

$$D^n f(z) = \frac{a_{-p}}{z^p} + \sum_{m=1}^{\infty} (p+m)^n a_{m-1} z^{m-1}.$$

It is proved that $B_{n+1}(\alpha) \subset B_n(\alpha)$. Since $B_0(\alpha)$ is the class of meromorphically p -valent starlike functions of order α , all functions in $B_n(\alpha)$ are p -valent starlike. Further a property preserving integrals is considered.

1. Introduction.

Let Σ_p denote the class of functions of the form

$$f(z) = \frac{a_{-p}}{z^p} + \sum_{k=0}^{\infty} a_k z^k \quad (a_{-p} \neq 0, p \in N = \{1, 2, \dots\}) \quad (1.1)$$

which are regular in the punctured disc $U^* = \{z : 0 < |z| < 1\}$. Define

$$D^0 f(z) = f(z), \quad (1.2)$$

KEY WORDS—Regular, p -valent, starlike, meromorphic.

AMS (1991) Subject Classification, 30C45–30C50.

Received November 6, 1993.

$$\begin{aligned}
 D^1 f(z) &= \frac{a_{-p}}{z^p} + (p+1)a_0 + (p+2)a_1 z + (p+3)a_2 z^2 + \dots \\
 &= \frac{(z^{p+1} f(z))'}{z^p}.
 \end{aligned} \tag{1.3}$$

$$D^2 f(z) = D(D^1 f(z)), \tag{1.4}$$

and for $n=1, 2, \dots$,

$$\begin{aligned}
 D^n f(z) &= D(D^{n-1} f(z)) = \frac{a_{-p}}{z^p} + \sum_{m=1}^{\infty} (p+m)^n a_{m-1} z^{m-1} \\
 &= \frac{(z^{p+1} D^{n-1} f(z))'}{z^p}.
 \end{aligned} \tag{1.5}$$

In this paper, we shall show that a function $f(z)$ in Σ_p , which satisfies one of the conditions

$$\operatorname{Re} \left\{ \frac{D^{n+1} f(z)}{D^n f(z)} - (p+1) \right\} < -\alpha, \quad (z \in U = \{z: |z| < 1\}), \tag{1.6}$$

for some $\alpha (0 \leq \alpha < p)$ and $n \in N_0 = \{0, 1, 2, \dots\}$, is meromorphically p -valent starlike in U^* . More precisely, it is proved that, for the classes $B_n(\alpha)$ of functions in Σ_p satisfying (1.6).

$$B_{n+1}(\alpha) \subset B_n(\alpha) \tag{1.7}$$

holds. Since $B_0(\alpha)$ equals $\Sigma_p^*(\alpha)$ (the class of meromorphically p -valent starlike functions of order α), the starlikeness of members of $B_n(\alpha)$ is a consequence of (1.7). Further for $c > 0$, let

$$F(z) = \frac{c}{z^{c+p}} \int_0^z t^{c+p-1} f(t) dt. \tag{1.8}$$

It is shown that $F(z) \in B_n(\alpha)$ whenever $f(z) \in B_n(\alpha)$. Some known results of Bajpai [1], Goel and Sohi [2] and Uralegaddi and Somanatha [5] are extended. In [4] Ruscheweyh obtained the new criteria for univalent functions.

2. Properties of the class $B_n(\alpha)$.

In proving our main results (Theorem 1 and Theorem 2 below), we shall need the following lemma due to Jack [3].

LEMMA. *Let $w(z)$ be non-constant regular in $U = \{z: |z| < 1\}$, $w(0) = 0$. If $|w(z)|$ attains its maximum value on the circle $|z| = r < 1$ at z_0 , we have $z_0 w'(z_0) = k w(z_0)$ where k is a real number, $k \geq 1$.*

THEOREM 1. $B_{n+1}(\alpha) \subset B_n(\alpha)$ for each integer $n \in N_0$.

PROOF. Let $f(z) \in B_{n+1}(\alpha)$. Then

$$\operatorname{Re} \left\{ \frac{D^{n+2}f(z)}{D^{n+1}f(z)} - (p+1) \right\} < -\alpha, \quad |z| < 1. \tag{2.1}$$

We have to show that (2.1) implies the inequality

$$\operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{D^n f(z)} - (p+1) \right\} < -\alpha. \tag{2.2}$$

Define a regular function $w(z)$ in U by

$$\frac{D^{n+1}f(z)}{D^n f(z)} - (p+1) = -\frac{p+(2\alpha-p)w(z)}{1+w(z)}. \tag{2.3}$$

Clearly $w(0)=0$. Equation (2.3) may be written as

$$\frac{D^{n+1}f(z)}{D^n f(z)} = \frac{1+(2p+1-2\alpha)w(z)}{1+w(z)}. \tag{2.4}$$

Differentiating (2.4) logarithmically and using the identity (easy to verify)

$$z(D^n f(z))' = D^{n+1}f(z) - (p+1)D^n f(z), \tag{2.5}$$

we obtain

$$\frac{\frac{D^{n+2}f(z)}{D^{n+1}f(z)} - (p+1) + \alpha}{p-\alpha} = \frac{2zw'(z)}{(1+w(z))(1+(2p+1-2\alpha)w(z))} - \frac{1-w(z)}{1+w(z)}. \tag{2.6}$$

We claim that $|w(z)| < 1$ in U . For otherwise (by Jack's lemma) there exists a point z_0 in U such that

$$z_0 w'(z_0) = kw(z_0) \tag{2.7}$$

where $|w(z_0)| = 1$ and $k \geq 1$. From (2.6) and (2.7), we obtain

$$\frac{\frac{D^{n+2}f(z_0)}{D^{n+1}f(z_0)} - (p+1) + \alpha}{p-\alpha} = \frac{2kw(z_0)}{(1+w(z_0))(1+(2p+1-2\alpha)w(z_0))} - \frac{1-w(z_0)}{1+w(z_0)}. \tag{2.8}$$

Thus

$$\operatorname{Re} \left\{ \frac{\frac{D^{n+2}f(z_0)}{D^{n+1}f(z_0)} - (p+1) + \alpha}{p-\alpha} \right\} \geq \frac{1}{2(1+p-\alpha)} > 0, \tag{2.9}$$

which contradicts (2.1). Hence $|w(z)| < 1$ in U and from (2.3) it follows that $f(z) \in B_n(\alpha)$.

THEOREM 2. Let $f(z) \in \Sigma_p$ satisfy the condition

$$\operatorname{Re} \left\{ \frac{D^{n+1}f(z)}{D^n f(z)} - (p+1) \right\} < -\alpha + \frac{p-\alpha}{2(p-\alpha+c)} \quad (z \in U) \tag{2.10}$$

for a given $n \in N_0$ and $c > 0$. Then

$$F(z) = \frac{c}{z^{c+p}} \int_0^z t^{c+p-1} f(t) dt$$

belongs to $B_n(\alpha)$.

PROOF. From the definition of $F(z)$, we have

$$z(D^n F(z))' = cD^n f(z) - (c+p)D^n F(z) \quad (2.11)$$

and also

$$z(D^n F(z))' = D^{n+1}F(z) - (p+1)D^n F(z). \quad (2.12)$$

Using (2.11) and (2.12) the condition (2.10) may be written as

$$\operatorname{Re} \left\{ \frac{\frac{D^{n+2}F(z)}{D^{n+1}F(z)} + (c-1)}{1 + (c-1)\frac{D^n F(z)}{D^{n+1}F(z)}} - (p+1) \right\} < -\alpha + \frac{p-\alpha}{2(p-\alpha+c)}. \quad (2.13)$$

We have to prove that (2.13) implies the inequality

$$\operatorname{Re} \left\{ \frac{D^{n+1}F(z)}{D^n F(z)} - (p+1) \right\} < -\alpha. \quad (2.14)$$

Define $w(z)$ in U by

$$\frac{D^{n+1}F(z)}{D^n F(z)} - (p+1) = -\frac{p+(2\alpha-p)w(z)}{1+w(z)}. \quad (2.15)$$

Clearly $w(z)$ is regular and $w(0)=0$. The equation (2.15) may be written as

$$\frac{D^{n+1}F(z)}{D^n F(z)} = \frac{1+(2p+1-2\alpha)w(z)}{1+w(z)}. \quad (2.16)$$

Differentiating (2.16) logarithmically and using (2.5), we obtain

$$\frac{D^{n+2}F(z)}{D^{n+1}F(z)} - \frac{D^{n+1}F(z)}{D^n F(z)} = \frac{2(p-\alpha)zw'(z)}{(1+w(z))(1+(2p+1-2\alpha)w(z))}. \quad (2.17)$$

The above equation may be written as

$$\begin{aligned} \frac{\frac{D^{n+2}F(z)}{D^{n+1}F(z)} + (c-1)}{1 + (c-1)\frac{D^n F(z)}{D^{n+1}F(z)}} - (p+1) &= \frac{D^{n+1}F(z)}{D^n F(z)} - (p+1) \\ &+ \left[\frac{2(p-\alpha)zw'(z)}{(1+w(z))(1+(2p+1-2\alpha)w(z))} \right] \cdot \left[\frac{1}{1 + (c-1)\frac{D^n F(z)}{D^{n+1}F(z)}} \right]. \end{aligned}$$

which, by using (2.15) and (2.16), reduces to

$$\frac{\frac{D^{n+2}F(z)}{D^{n+1}F(z)} + (c-1)}{1 + (c-1)\frac{D^n F(z)}{D^{n+1}F(z)}} - (p+1) = -\left[\alpha + (p-\alpha)\frac{1-w(z)}{1+w(z)} \right] + \frac{2(p-\alpha)zw'(z)}{(1+w(z))[c+(c+2(p-\alpha))w(z)]}.$$

The remaining part of the proof is similar to that of Theorem 1.

REMARKS. (i) Putting $p=1$, $a_{-1}=1$, $n=0$ and $\alpha=0$ in Theorem 2, we get the result of Goel and Sohi [2, Corollary 1].

(ii) For $p=1$, $a_{-1}=1$, $n=0$, $\alpha=0$ and $c=1$ the above theorem extends a result of Bajpai [1, Theorem 1].

THEOREM 3. $f(z) \in B_n(\alpha)$ if and only if

$$F(z) = \frac{1}{z^{1+p}} \int_0^z t^p f(t) dt \in B_{n+1}(\alpha).$$

PROOF. From the definition of $F(z)$ we have

$$D^n(zF'(z)) + (1+p)D^n F(z) = D^n f(z).$$

That is,

$$z(D^n F(z))' + (1+p)D^n F(z) = D^n f(z). \tag{2.18}$$

By using the identity (2.5), (2.18) reduces to $D^n f(z) = D^{n+1}F(z)$. Hence $D^{n+1}f(z) = D^{n+2}F(z)$.

Therefore

$$\frac{D^{n+1}f(z)}{D^n f(z)} = \frac{D^{n+2}F(z)}{D^{n+1}F(z)}$$

and the result follows.

REMARK. Putting $p=1$ in the above theorems, we get the results obtained by Uralegaddi and Somanatha [5].

References

[1] S.K. Bajpai, A note on a class of meromorphic univalent functions, Rev. Roum. Math. Pures Appl. 22 (1977), 295-297.
 [2] R.M. Goel and N.S. Sohi, On a class of meromorphic functions, Glas. Mat. 17 (1981), 19-28.
 [3] I.S. Jack, Functions starlike and convex of order α , J. London Math. Soc. (2) 3 (1971), 469-474.

- [4] S. Ruscheweyh, New criteria for univalent functions, *Proc. Amer. Math. Soc.* **49** (1975), 109-115.
- [5] B. A. Uralegaddi and C. Somanatha, New criteria for meromorphic starlike univalent functions, *Bull. Austral. Math. Soc.* **43** (1991), 137-140.

Department of Mathematics
Faculty of Science
University of Mansoura
Mansoura, Egypt