

A FAMILY OF BRAIDED COSEMISIMPLE HOPF ALGEBRAS OF FINITE DIMENSION

By

Satoshi SUZUKI

0. Introduction

Recently some finite dimensional cosemisimple Hopf algebras were constructed [Mas2] [F] [G]. We aim to give a plain and systematic description of cosemisimple Hopf algebras of low dimension. For this purpose we construct them as quotient bialgebras of a sufficiently large bialgebra. This way has the advantage of defining homomorphisms and determining braidings.

In this paper we define and study a family of finite dimensional cosemisimple Hopf algebras

$$\mathcal{F} = \{A_{NL}^{(++)}, A_{NL}^{(+-)}, A_{NL}^{(-+)}, A_{NL}^{(--)} \mid N \geq 1, L \geq 2\},$$

which consists of quotients of a bialgebra B over an algebraically closed field k with $chk \neq 2$.

This family contains the “non-trivial” cosemisimple Hopf algebras of dimension 8, 12 if $chk \neq 3$.

In Section 1 we review basic definitions and results.

In Section 2 quadratic bialgebras B , $B^{(+)}$ and $B^{(-)}$ are constructed. We use B to construct the family \mathcal{F} , and $B^{(\pm)}$ to obtain braidings on the members of a subfamily of \mathcal{F} . These bialgebras B , $B^{(\pm)}$ are cosemisimple, and we determine all braidings on them.

In Section 3 we define the family \mathcal{F} as a set of quotient bialgebras of the bialgebra B . We write $A_{NL}^{(+1,-1)} = A_{NL}^{(+-)}$, etc. Let $\nu, \lambda = \pm 1$. Our main results are as follows.

i) $A_{NL}^{(\nu\lambda)}$ is a non-cocommutative involutory cosemisimple Hopf algebra of dimension $4NL$, which is non-commutative unless $(L, \lambda) = (2, +1)$. $A_{NL}^{(\nu\lambda)}$ is furthermore semisimple if $(\dim A_{NL}^{(\nu\lambda)}) \cdot 1 \neq 0$.

- ii) Any non-commutative subHopf algebra of $A_{NL}^{(\nu\lambda)}$ generated by a simple subcoalgebra is a member of the family.
- iii) All braidings on $A_{NL}^{(\nu\lambda)}$ are determined.
- iv) We determine when $A_{N_1L_1}^{(\nu_1\lambda_1)}$ and $A_{N_2L_2}^{(\nu_2\lambda_2)}$ are isomorphic.

1. Preliminaries [D]

We follow Sweedler's book [S] and Montgomery's book [M] for terminology of Hopf algebras.

In this section we review basic definitions and results. They are due to Doi [D].

Let B be a bialgebra over a field k , $\tau : B \otimes B \rightarrow k$ a k -linear map which is invertible with respect to the convolution product. (B, τ) is called a *braided* bialgebra if the following three conditions hold:

- (1) $\Sigma\tau(x_1, y_1)x_2y_2 = \Sigma y_1x_1\tau(x_2, y_2)$
- (2) $\tau(xy, z) = \Sigma\tau(x, z_1)\tau(y, z_2)$
- (3) $\tau(x, yz) = \Sigma\tau(x_1, z)\tau(x_2, y)$

for $x, y, z \in B$.

Then the following conditions are automatically satisfied:

$$\begin{aligned} \tau(x, 1) &= \varepsilon(x) = \tau(1, x), \\ \Sigma\tau(x_1, y_1)\tau(x_2, z_1)\tau(y_2, z_2) &= \Sigma\tau(y_1, z_1)\tau(x_1, z_2)\tau(x_2, y_2) \quad \text{for } x, y, z \in B. \end{aligned}$$

We call this τ a *braiding* on B .

PROPOSITION 1.1 ([H, Proposition 1.2]). *Let (B, τ) be a braided bialgebra generated by a subcoalgebra C , (I) the bi-ideal generated by a coideal I of B . Then τ induces a braiding on the bialgebra $B/(I)$ iff $\tau = 0$ on $C \otimes I + I \otimes C$.*

If (B, τ) is a braided bialgebra, ${}^t\tau^{-1}$ is another braiding on B , where ${}^t\tau^{-1}(x, y) = \tau^{-1}(y, x)$, and the braiding τ is said to be *symmetric* if ${}^t\tau^{-1} = \tau$.

Let C be a coalgebra over k , $\sigma : C \otimes C \rightarrow k$ an invertible k -linear map. For any bialgebra B , a linear map $f : C \rightarrow B$ is called a σ -map if

$$\Sigma\sigma(x_1, y_1)f(x_2)f(y_2) = \Sigma f(y_1)f(x_1)\sigma(x_2, y_2), \quad x, y \in C.$$

Let $T(C)$ be the tensor (bi-)algebra and I_σ is the (bi-)ideal generated by

$$(4) \quad \Sigma\sigma(x_1, y_1)x_2y_2 - \Sigma y_1x_1\sigma(x_2, y_2), \quad x, y, z \in C.$$

We can form the bialgebra $M(C, \sigma) = T(C)/I_\sigma$, which is called is the *quadratic bialgebra associated with* (C, σ) .

REMARK 1.2. i) The map $i : C \hookrightarrow T(C) \rightarrow M(C, \sigma)$ is an injective coalgebra σ -map.

ii) If B is a bialgebra and $f : C \rightarrow B$ is a σ -(coalgebra) map, then there is a unique (*bi*-) algebra map $\hat{f} : M(C, \sigma) \rightarrow B$ such that $\hat{f} \circ i = f$.

iii) $M(C, \sigma)$ has a natural algebra-gradation $\{C^n\}_{n \geq 0}$.

iv) $M(C, \sigma)^{op} = M(C, \sigma^{-1}) = M(C, {}^t\sigma)$, $M(C, \sigma) = M(C, {}^t\sigma^{-1})$.

Let (C, σ) be as above. The map σ is called a *Yang-Baxter form* (or *YB-form*) if for all $x, y, z \in C$,

$$(5) \quad \Sigma\sigma(x_1, y_1)\sigma(x_2, z_1)\sigma(y_2, z_2) = \Sigma\sigma(y_1, z_1)\sigma(x_1, z_2)\sigma(x_2, y_2).$$

We call (C, σ) a *YB-coalgebra* if σ is a YB-form.

REMARK 1.3. If σ is a YB-form on C , so is ${}^t\sigma^{-1}$.

A YB-form σ is said to be *symmetric* if ${}^t\sigma^{-1} = \sigma$.

PROPOSITION 1.4 ([D, Theorem 2.6]). *If (C, σ) is a YB-coalgebra, σ uniquely extends to a braiding $\tilde{\sigma}$ on $M(C, \sigma)$.*

We note that if (C, σ) is a YB-coalgebra then $M(C, \sigma)$ has another braiding ${}^t\tilde{\sigma}^{-1}$.

COROLLARY 1.5. *$\tilde{\sigma}$ is symmetric iff σ is symmetric.*

For a bialgebra B , a Hopf algebra H and a bialgebra map $\iota : B \rightarrow H$, we call (H, ι) (or simply H) a *Hopf closure* of B if the following universality holds: for any Hopf algebra A and any bialgebra map $f : B \rightarrow A$, there is a unique Hopf algebra map $\tilde{f} : H \rightarrow A$ such that $\tilde{f} \circ \iota = f$. See [Man] [H] [D].

PROPOSITION 1.6 ([T2] [D, Theorem 3.6] [H]). *Let $M(C, \sigma)$ be the quadratic bialgebra associated with (C, σ) , $d(\neq 0)$ a grouplike element of $M(C, \sigma)$. If there is a map $j : C \rightarrow M(C, \sigma)$ such that*

$$\Sigma i(x_1)j(x_2) = \varepsilon(x)d = \Sigma j(x_1)i(x_2) \quad \text{for all } x \in C,$$

then d is central and the (well-defined) localization $M(C, \sigma)[d^{-1}]$ becomes a Hopf algebra. Moreover it is a Hopf closure of $M(C, \sigma)$, and it follows that $M(C, \sigma)[d^{-1}] = M(C, \sigma)[G^{-1}]$, where G is the set of grouplike elements of $M(C, \sigma)$. If (C, σ) is a YB-coalgebra, $M(C, \sigma)[d^{-1}]$ has a braiding.

2. YB-coalgebras and quadratic bialgebras

From now on we work over an algebraically closed field k whose characteristic, chk , is not 2. Indices of Kronecker's δ_{ij} , X_{ij} , etc. are considered modulo 2.

In this section we define some YB-coalgebras and examine quadratic bialgebras associated with them.

Set $C = M_2(k)^*$, the dual coalgebra of the 2×2 -matrix algebra $M_2(k)$, and let $\{X_{ij}\}_{1 \leq i, j \leq 2}$ be the comatrix basis of C , namely it spans C and satisfies

$$\Delta(X_{ij}) = \sum_{k=1}^2 X_{ik} \otimes X_{kj}, \quad \varepsilon(X_{ij}) = \delta_{ij}.$$

For any coalgebra D and $Y_{ij} \in D$, $1 \leq i, j \leq 2$, if the linear map $C \rightarrow D$, $X_{ij} \mapsto Y_{ij}$, is an injective coalgebra map, we denote the image by

$$\text{span}_k(Y_{ij}) = \text{span}_k \begin{pmatrix} Y_{11} & Y_{12} \\ Y_{21} & Y_{22} \end{pmatrix}.$$

Let $\lambda = \pm 1$. Now for any $\alpha \in k^\times = k - \{0\}$, we define linear maps $\sigma_{(\alpha)}$, $\tau_{(\alpha)}^{(\pm 1)} = \tau_{(\alpha)}^{(\pm)}$: $C \otimes C \rightarrow k$ as follows (see [D, Example 2.8] for $\tau^{(\lambda)}$):

$\sigma_{(\alpha)}$	X_{11}	X_{12}	X_{21}	X_{22}	$\tau_{(\alpha)}^{(\lambda)}$	X_{11}	X_{12}	X_{21}	X_{22}
X_{11}	0	0	0	0	X_{11}	α	0	0	1
X_{12}	0	α	1	0	X_{12}	0	0	0	0
X_{21}	0	1	α	0	X_{21}	0	0	0	0
X_{22}	0	0	0	0	X_{22}	λ	0	0	α

PROPOSITION 2.1. $\sigma_{(\alpha)}$, $\tau_{(\alpha)}^{(\lambda)}$ ($\alpha \in k^\times$) are YB-forms on C .

PROOF. We show that $\sigma_{(\alpha)} = \sigma$ is a YB-form.

We can write $\sigma(X_{i,j+1}, X_{l,m+1}) = \delta_{ij} \delta_{lm} \alpha^{\delta_{il}}$.

For X_{ij} , X_{lm} and X_{uv} , observe that

$$\begin{aligned} & \Sigma_{a,b,c} \sigma(X_{ia}, X_{lb}) \sigma(X_{aj}, X_{uc}) \sigma(X_{bm}, X_{cv}) \\ &= \sigma(X_{i,i+1}, X_{l,l+1}) \sigma(X_{i+1,j}, X_{u,u+1}) \sigma(X_{l+1,m}, X_{u+1,v}) \\ &= \delta_{ij} \delta_{lm} \delta_{uv} \alpha^{\delta_{il}} \alpha^{\delta_{i+1,u}} \alpha^{\delta_{lu}}, \end{aligned}$$

and

$$\begin{aligned} & \Sigma_{a,b,c} \sigma(X_{lb}, X_{uc}) \sigma(X_{ia}, X_{cv}) \sigma(X_{aj}, X_{bm}) \\ &= \sigma(X_{l,l+1}, X_{u,u+1}) \sigma(X_{i,i+1}, X_{u+1,v}) \sigma(X_{i+1,j}, X_{l+1,m}) \\ &= \delta_{uv} \delta_{ij} \delta_{lm} \alpha^{\delta_{lu}} \alpha^{\delta_{i,u+1}} \alpha^{\delta_{il}}. \end{aligned}$$

Thus Condition (5) is satisfied.

The inverse is given by

$$\sigma_{(\alpha)}^{-1} = \sigma_{(\alpha^{-1})}.$$

Therefore $\sigma_{(\alpha)}$ is a YB-form for $\alpha \in k^\times$.

It is easy to check that $\tau_{(\alpha)}^{(\lambda)}$ is also a YB-form on C . \square

Therefore $(C, \sigma_{(\alpha)})$ and $(C, \tau_{(\alpha)}^{(\lambda)})$ are YB-coalgebras for all $\alpha \in k^\times$.

REMARK 2.2. $\{\sigma_{(\alpha)}, \tau_{(\beta)}^{(+)} \mid \alpha, \beta \in k^\times\}$, $\{\tau_{(\alpha)}^{(+)}, \tau_{(\beta)}^{(-)} \mid \alpha, \beta \in k^\times\}$ form subgroups of the unit group of $M_2(k)^{\otimes 2}$.

Next we examine the defining relations of the quadratic bialgebras associated with them.

PROPOSITION 2.3.

i) The ideal I_σ , where $\sigma = \sigma_{(\alpha)}$, is generated by the following:

$$\begin{aligned} & \{X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2, X_{j,j+1} X_{ii} - \alpha X_{i+1,i+1} X_{j+1,j}\} \quad \text{if } \alpha^2 = 1, \\ & \{X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2, X_{ij} X_{lm} (i+j+l+m \equiv 1)\} \quad \text{if } \alpha^2 \neq 1. \end{aligned}$$

ii) The ideal $I_{\tau^{(\lambda)}}$, where $\tau^{(\lambda)} = \tau_{(\alpha)}^{(\lambda)}$, is generated by the following:

$$\begin{aligned} & \{X_{11} X_{22} - X_{22} X_{11}, X_{12} X_{21} - \lambda X_{21} X_{12}, X_{i2} X_{i1} - \alpha X_{i1} X_{i2}, X_{2j} X_{1j} - \lambda \alpha X_{1j} X_{2j}\} \\ & \hspace{20em} \text{if } \alpha^2 = \lambda, \\ & \{X_{11} X_{22} - X_{22} X_{11}, X_{12} X_{21} - \lambda X_{21} X_{12}, X_{ij} X_{lm} (i+j+l+m \equiv 1)\} \\ & \hspace{20em} \text{if } \alpha^2 \neq \lambda. \end{aligned}$$

PROOF. i) For X_{ij} , X_{lm} , observe that

$$\begin{aligned} \Sigma \sigma(X_{ia}, X_{lb}) X_{aj} X_{bm} &= \sigma(X_{i,i+1}, X_{l,l+1}) X_{i+1,j} X_{l+1,m} \\ &= \alpha^{\delta_{il}} X_{i+1,j} X_{l+1,m}, \end{aligned}$$

$$\begin{aligned}\Sigma X_{lb}X_{ia}\sigma(X_{aj}, X_{bm}) &= X_{l,m+1}X_{i,j+1}\sigma(X_{j+1,j}, X_{m+1,m}) \\ &= X_{l,m+1}X_{i,j+1}\alpha^{\delta_{jm}}.\end{aligned}$$

Thus the subset

$$\{\alpha^{\delta_{il}}X_{ij}X_{lm} - X_{l+1,m+1}X_{i+1,j+1}\alpha^{\delta_{jm}} \mid 1 \leq i, j, l, m \leq 2\}$$

generates the ideal I_σ . The above polynomials are written as follows:

$$\begin{cases} \alpha X_{ij}^2 - X_{i+1,j+1}^2 & \text{if } i = l, j = m, \\ X_{ij}X_{lj} - X_{l+1,j+1}X_{i+1,j+1} & \text{if } i \neq l, j = m, \\ \alpha X_{ij}X_{im} - X_{i+1,m+1}X_{i+1,j+1} & \text{if } i = l, j \neq m, \\ X_{ij}X_{lm} - X_{l+1,m+1}X_{i+1,j+1} & \text{if } i \neq l, j \neq m \text{ (i.e., } l \equiv i+1, m \equiv j+1 \text{)}. \end{cases}$$

ii) This is similarly shown as i). □

REMARK 2.4. i) For the bialgebra $M(C, \sigma_{(-1)})$, see the quantum conformal group in [Man].

ii) $M(C, \tau_{(\pm 1)}^{(+)})$ are the quantum matrix bialgebras $M_{\pm 1}(2)$.

iii) $M(C, \tau_{(\sqrt{-1})}^{(-)})$ is Takeuchi's two-parameter bialgebra $M_{\alpha, \beta}(2)$ for $\alpha = \sqrt{-1}$, $\beta = -\sqrt{-1}$ ([T1], [D]).

Define $B = M(C, \sigma_{(\alpha)})$ for $\alpha^2 \neq 1$ and $B^{(\lambda)} = M(C, \tau_{(\alpha)}^{(\lambda)})$ for $\alpha^2 \neq \lambda$. We write $B^{(\pm 1)} = B^{(\pm)}$. These definitions, ignoring choice of α , are reasonable by Proposition 2.3.

On the other hand, we see by Proposition 1.1 that braidings $\tilde{\sigma}_{(\pm 1)}$, $\tilde{\tau}_{(\pm\sqrt{\lambda})}^{(\lambda)}$ are induced on B , $B^{(\lambda)}$, respectively, via the canonical surjections

$$M(C, \sigma_{(\pm 1)}) \rightarrow B, \quad M(C, \tau_{(\pm\sqrt{\lambda})}^{(\lambda)}) \rightarrow B^{(\lambda)}.$$

Note that $\{X_{ij}X_{lm} \mid i+j+l+m \equiv 1\}$ spans a coideal of $T(C)$.

Therefore we have the following claim:

CLAIM 2.5.

- i) $\sigma_{(\alpha)} : C \otimes C \rightarrow k$ extends to a braiding $\tilde{\sigma}_{(\alpha)}$ on B for every $\alpha \in k^\times$.
- ii) $\tau_{(\alpha)}^{(\lambda)} : C \otimes C \rightarrow k$ extends to a braiding $\tilde{\tau}_{(\alpha)}^{(\lambda)}$ on $B^{(\lambda)}$ for every $\alpha \in k^\times$.

We examine the coalgebra structure of B .

PROPOSITION 2.6.

i) B has the following set as a basis

$$\{X_{11}^{n-r} \overbrace{X_{22}X_{11}X_{22} \dots}^r, X_{12}^{n-r} \overbrace{X_{21}X_{12}X_{21} \dots}^r \mid n \geq 0, 0 \leq r \leq n\}.$$

ii) The grouplike elements but 1 in B are given by

$$X_{11}^{2s} \pm X_{12}^{2s} \quad (s \geq 1).$$

Then are central non-zero divisors.

iii) The simple subcoalgebras of B which are not spanned by grouplike elements are of dimension 4. They are given by

$$C_{st} = \text{span}_k \left(\begin{array}{cc} X_{11}^{2s} \overbrace{X_{11}X_{22}X_{11} \dots}^t & X_{12}^{2s} \overbrace{X_{12}X_{21}X_{12} \dots}^t \\ X_{12}^{2s} \overbrace{X_{21}X_{12}X_{21} \dots}^t & X_{11}^{2s} \overbrace{X_{22}X_{11}X_{22} \dots}^t \end{array} \right) \quad (s \geq 0, t \geq 1).$$

iv) B is cosemisimple. The n th component C^n ($n \geq 1$) of B is decomposed as the sum of simple subcoalgebras as follows:

$$C^n = \begin{cases} \sum_{n=2s+t} C_{st}, & \text{if } n \text{ is odd;} \\ \sum_{n=2s+t} C_{st} + k(X_{11}^n \pm X_{12}^n), & \text{if } n \text{ is even.} \end{cases}$$

PROOF. i) It is verified in the same manner as Theorem 3.1.i) below.

ii), iii), iv) It is easy to see that $X_{11}^{2s} \pm X_{12}^{2s}$ is grouplike for $s \geq 1$. By i) and the defining relations of B , it is a central non-zero divisor. C is isomorphic to C_{st} as coalgebras by

$$\begin{aligned} X_{11} &\mapsto X_{11}^{2s} \overbrace{X_{11}X_{22}X_{11} \dots}^t, \\ X_{12} &\mapsto X_{12}^{2s} X_{12}X_{21}X_{12} \dots, \\ X_{21} &\mapsto X_{12}^{2s} X_{21}X_{12}X_{21} \dots, \\ X_{22} &\mapsto X_{11}^{2s} X_{22}X_{11}X_{22} \dots \end{aligned}$$

By i) we have that

$$\begin{aligned} B &= k \cdot 1 + \sum k(X_{11}^{2s} \pm X_{12}^{2s}) + \sum C_{st} \\ &= k \cdot 1 \oplus \{\oplus_{s \geq 1} k(X_{11}^{2s} \pm X_{12}^{2s})\} \oplus \{\oplus_{s \geq 0, t \geq 1} C_{st}\}. \end{aligned}$$

Thus ii), iii), iv) are done. □

PROPOSITION 2.7.

i) $B^{(\lambda)}$ has the following set as a basis

$$\{X_{11}^u X_{22}^v, X_{12}^u X_{21}^v \mid u + v \geq 0\}.$$

ii) The grouplike elements but 1 in $B^{(\lambda)}$ are given by

$$X_{11}^u X_{22}^u \pm \sqrt{\lambda^u} X_{12}^u X_{21}^u \quad (u \geq 1).$$

They are non-zero divisors.

iii) The simple subcoalgebras of $B^{(\lambda)}$ which are not spanned by grouplike elements are all of dimension 4. They are given by

$$D_{uv} = \text{span}_k \begin{pmatrix} X_{11}^u X_{22}^v & X_{12}^u X_{21}^v \\ X_{21}^u X_{12}^v & X_{22}^u X_{11}^v \end{pmatrix}, \quad (u \leq v).$$

iv) $B^{(\lambda)}$ is cosemisimple. The n th component C^n ($n \geq 1$) of $B^{(\lambda)}$ is decomposed as the sum of simple subcoalgebras as follows:

$$C^n = \begin{cases} \sum_{n=u+v, u \leq v} D_{uv}, & \text{if } n \text{ is odd;} \\ \sum_{n=u+v, u \leq v} D_{uv} + k(X_{11}^{n/2} X_{22}^{n/2} \pm \sqrt{\lambda^{n/2}} X_{12}^{n/2} X_{21}^{n/2}), & \text{if } n \text{ is even.} \end{cases}$$

We omit the proof.

COROLLARY 2.8. Let $\langle C_{st} \rangle$ denote the sub-bialgebra generated by the simple subcoalgebra $C_{st} \subset B$. Then as bialgebras,

$$B \cong \langle C_{st} \rangle \simeq \begin{cases} B, & \text{if } t \text{ is odd;} \\ B^{(+)}, & \text{if } t \text{ is even.} \end{cases}$$

We omit the proof. See the proof of Theorem 3.5 below.

Define linear maps $\sigma_{\alpha\beta} = \beta\sigma_{(\alpha\beta^{-1})}$, $\tau_{\alpha\beta}^{(\lambda)} = \beta\tau_{(\alpha\beta^{-1})}^{(\lambda)}$ for $\alpha, \beta \in k^\times$, $\lambda = \pm 1$. They are also YB-forms on C . The YB-form $\sigma_{\alpha\beta}$ extends to a braiding $\tilde{\sigma}_{\alpha\beta}$ on B , and $\tau_{\alpha\beta}^{(\lambda)}$ extends to a braiding $\tilde{\tau}_{\alpha\beta}^{(\lambda)}$ on $B^{(\lambda)}$.

PROPOSITION 2.9. i) $\sigma_{\alpha\beta}$ is symmetric iff $\alpha^2 = 1 = \beta^2$. $\tau_{\alpha\beta}^{(\lambda)}$ is symmetric iff $\alpha^2 = 1, \beta^2 = \lambda$.

ii) The set of braidings on B is $\{\tilde{\sigma}_{\alpha\beta} \mid \alpha, \beta \in k^\times\}$, and that on $B^{(\lambda)}$ is $\{\tilde{\tau}_{\alpha\beta}^{(\lambda)} \mid \alpha, \beta \in k^\times\}$.

PROOF. i) We note that ${}^t\sigma_{\alpha\beta} = \sigma_{\alpha\beta}$, ${}^t\tau_{\alpha\beta}^{(\lambda)} = \tau_{\alpha, \lambda\beta}^{(\lambda)}$. The statement follows from these.

ii) We show the statement with B . The statement with $B^{(\lambda)}$ is similarly verified.

We have obtained braidings $\tilde{\sigma}_{\alpha\beta}(\alpha, \beta \in k^\times)$ on B .

Let σ be a braiding. Note that the second component C^2 of B has a basis

$$\{X_{11}^2, X_{12}^2, X_{11}X_{22}, X_{22}X_{11}, X_{12}X_{21}, X_{21}X_{12}\}.$$

So for X_{ij}, X_{lm} , it follows that

$$\begin{aligned} \Sigma\sigma(X_{ia}, X_{lb})X_{aj}X_{bm} &= \sigma(X_{ij}, X_{lm})X_{jj}X_{mm} + \sigma(X_{i,j+1}, X_{l,m+1})X_{j+1,j}X_{m+1,m}, \\ \Sigma X_{lb}X_{ia}\sigma(X_{aj}, X_{bm}) &= X_{ll}X_{ii}\sigma(X_{ij}, X_{lm}) + X_{l,l+1}X_{i,i+1}\sigma(X_{i+1,j}, X_{l+1,m}). \end{aligned}$$

These must be equal, so we obtain the following by Proposition 2.6.i):

$$\begin{aligned} \sigma(X_{ij}, X_{lm})X_{jj}X_{mm} &= X_{ll}X_{ii}\sigma(X_{ij}, X_{lm}), \\ \sigma(X_{i,j+1}, X_{l,m+1})X_{j+1,j}X_{m+1,m} &= X_{l,l+1}X_{i,i+1}\sigma(X_{i+1,j}, X_{l+1,m}). \end{aligned}$$

The above equations imply that $\sigma|_{C \otimes C}$ is given as follows with some $\alpha, \beta, \gamma \in k$:

σ	X_{11}	X_{12}	X_{21}	X_{22}
X_{11}	γ	0	0	0
X_{12}	0	α	β	0
X_{21}	0	β	α	0
X_{22}	0	0	0	γ

Moreover it follows by Condition (2) that

$$\begin{aligned} 0 &= \sigma(0, X_{12}) = \sigma(X_{11}X_{12}, X_{12}) \\ &= \sigma(X_{11}, X_{11})\sigma(X_{12}, X_{12}) + \sigma(X_{11}, X_{12})\sigma(X_{12}, X_{22}) = \gamma\alpha, \end{aligned}$$

and

$$\begin{aligned} 0 &= \sigma(0, X_{12}) = \sigma(X_{11}X_{21}, X_{12}) \\ &= \sigma(X_{11}, X_{11})\sigma(X_{21}, X_{12}) + \sigma(X_{11}, X_{12})\sigma(X_{21}, X_{22}) = \gamma\beta. \end{aligned}$$

We have that $\gamma = 0$, $\alpha, \beta \in k^\times$ since σ is invertible.

Therefore $\sigma|_{C \otimes C} = \sigma_{\alpha\beta}$, so $\sigma = \tilde{\sigma}_{\alpha\beta}$. □

We describe a Hopf closure of the bialgebra B .

Set $d_{\pm} = X_{11}^2 \pm X_{12}^2$. These are central grouplike elements. For example, observe that

$$\begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \end{pmatrix} \begin{pmatrix} X_{11} & X_{21} \\ X_{12} & X_{22} \end{pmatrix} = d_+ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} X_{11} & X_{21} \\ X_{12} & X_{22} \end{pmatrix} \begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \end{pmatrix},$$

and

$$\begin{aligned} \begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \end{pmatrix} \begin{pmatrix} X_{11} & -X_{21} \\ -X_{12} & X_{22} \end{pmatrix} &= d_- \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} X_{11} & -X_{21} \\ -X_{12} & X_{22} \end{pmatrix} \begin{pmatrix} X_{11} & X_{12} \\ X_{21} & X_{22} \end{pmatrix}. \end{aligned}$$

Using Proposition 1.6 and Proposition 2.6, we have the following.

PROPOSITION 2.10. *The Hopf closure H of B is given by*

$$H = B[d_+^{-1}] = B[d_-^{-1}] = B[G(B)^{-1}],$$

where $G(B)$ is the set of grouplike elements in B . This Hopf algebra is braided and cosemisimple, and includes B as a sub-bialgebra. Furthermore, H is involutory. In fact, the antipode S is determined by

$$S(X_{ij}) = X_{ji}d_+^{-1} = (-1)^{i+j}X_{ji}d_-^{-1}.$$

3. Quotients of the bialgebra B

In this section we define and study a family of finite dimensional cosemisimple bi(Hopf) algebras which are quotients of the bialgebra B over an algebraically closed field k with $chk \neq 2$.

It will be shown that the family contains the “non-trivial” cosemisimple Hopf algebras of dimension 8 ([Mas2]) and of dimension 12 ([F]) if $chk \neq 3$. See also Gelaki’s Hopf algebras of dimension $4p$, where $p(\geq 3)$ is prime ([G]).

We construct the family. It is easy to see by Proposition 2.6 that for $L \geq 2$, $N \geq 1$ and $\lambda, \nu = \pm 1$, the following subsets

$$\begin{aligned} &\{\overbrace{X_{22}X_{11}X_{22}\cdots}^L - \overbrace{X_{11}X_{22}X_{11}\cdots}^L, \overbrace{X_{21}X_{12}X_{21}\cdots}^L - \lambda \overbrace{X_{12}X_{21}X_{12}\cdots}^L\}, \\ &\{1 - (X_{11}^{2N} + \nu X_{12}^{2N})\} \end{aligned}$$

span coideals of B . Let J_L^λ and I_N^ν be the ideals generated by these coideals respectively, which are bi-ideals.

We can form the bialgebra

$$A_{NL}^{(v\lambda)} = B/J_L^\lambda + I_N^v.$$

We write $A_{NL}^{(+,-)} = A_{NL}^{(+1,-1)}$, etc. Let π be the following surjective bialgebra map:

$$\pi : B \rightarrow A_{NL}^{(v\lambda)}, \quad X_{ij} \mapsto \bar{X}_{ij} = x_{ij}.$$

THEOREM 3.1.

i) $A_{NL}^{(v\lambda)}$ has the following set as a basis

$$\{x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t, x_{12}^s \overbrace{x_{21}x_{12}x_{21} \cdots}^t \mid 1 \leq s \leq 2N, 0 \leq t \leq L-1\}.$$

Thus $\dim A_{NL}^{(v\lambda)} = 4NL$.

ii) Let $G(A_{NL}^{(v\lambda)}) = G$ be the set of grouplike elements of $A_{NL}^{(v\lambda)}$. Then

$$G = \{x_{11}^{2s} \pm x_{12}^{2s}, x_{11}^{2s} \overbrace{x_{11}x_{22}x_{11} \cdots}^L \pm \sqrt{\lambda} x_{12}^{2s} \overbrace{x_{12}x_{21}x_{12} \cdots}^L \mid 1 \leq s \leq N\}.$$

iii) The simple subcoalgebras of $A_{NL}^{(v\lambda)}$ which are not spanned by grouplike elements are given by

$$C_{st} = \text{span}_k \left(\begin{array}{cc} x_{11}^{2s} \overbrace{x_{11}x_{22}x_{11} \cdots}^t & x_{12}^{2s} \overbrace{x_{12}x_{21}x_{12} \cdots}^t \\ x_{12}^{2s} \overbrace{x_{21}x_{12}x_{21} \cdots}^t & x_{11}^{2s} \overbrace{x_{22}x_{11}x_{22} \cdots}^t \end{array} \right)$$

for $0 \leq s \leq N-1, 1 \leq t \leq L-1$.

iv) $|G(A_{NL}^{(v\lambda)})| = 4N$, and there are exactly $N(L-1)$ simple subcoalgebras of dimension 4.

v) $A_{NL}^{(v\lambda)}$ is non-cocommutative and cosemisimple. It is non-commutative unless $(L, \lambda) = (2, +1)$.

vi) $A_{NL}^{(v\lambda)}$ is an involutory Hopf algebra.

vii) Let $\Lambda = \sum x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t (1 \leq s \leq 2N, 0 \leq t \leq L-1)$. Then Λ is a non-zero two-sided integral.

viii) $A_{NL}^{(v\lambda)}$ is semisimple if $chk \nmid NL$.

PROOF. i) Let B' be the algebra $k\langle X, Y \rangle / \{X^2 - Y^2\}$ and $\lambda, v = \pm 1$. Let V be the k -vector space with a basis $\{\langle s, t \rangle \in V \mid s \geq 1, 0 \leq t \leq L-1\}$.

We define the following ideals of B' :

$$J_L^{\lambda'} = (\overbrace{YXYX \cdots}^L - \lambda \overbrace{XYXY \cdots}^L),$$

$$I_N^{\nu'} = (1 - \nu X^{2N}).$$

We prove i) step-by-step.

(Step 1) We define a right B' -module structure on V .

Define the actions of X and Y as follows:

$$X : \langle s, t \rangle \mapsto \begin{cases} \langle s, t+1 \rangle, & \text{if } t \text{ is odd, } t \leq L-2, \\ \lambda \langle s+1, L-1 \rangle, & t = L-1, \\ \langle s+1, 0 \rangle, & \text{if } t \text{ is even, } t = 0, \\ \langle s+2, t-1 \rangle, & t \geq 2, \end{cases}$$

$$Y : \langle s, t \rangle \mapsto \begin{cases} \langle s+2, t-1 \rangle, & \text{if } t \text{ is odd,} \\ \langle s, t+1 \rangle, & \text{if } t \text{ is even, } t \leq L-2, \\ \lambda \langle s+1, L-1 \rangle, & t = L-1. \end{cases}$$

It is easy to see $X^2 \equiv Y^2$ in $\text{End}_k(V)$.

Thus we have a right B' -module structure on V .

(Step 2) We claim the subspace W spanned by

$$\{\langle q(2N) + s, t \rangle - \nu^q \langle s, t \rangle \mid 1 \leq s \leq 2N, q \geq 1, 0 \leq t \leq L-1\}$$

is a submodule of V .

For example, when $t = L-1$ is odd and $s = 2N$, observe the following:

$$\begin{aligned} X : \langle q(2N) + 2N, L-1 \rangle &\mapsto \lambda \langle q(2N) + 2N + 1, L-1 \rangle \\ &= \lambda \langle (q+1)(2N) + 1, L-1 \rangle \\ &\equiv \lambda \nu^{q+1} \langle 1, L-1 \rangle \pmod{W}, \end{aligned}$$

and

$$\begin{aligned} X : \nu^q \langle 2N, L-1 \rangle &\mapsto \nu^q \lambda \langle 2N + 1, L-1 \rangle \\ &= \nu^q \lambda \langle 1 \cdot (2N) + 1, L-1 \rangle \\ &\equiv \nu^q \lambda \nu \langle 1, L-1 \rangle \pmod{W}. \end{aligned}$$

(Step 3) The action of B' induces the $B'/J_L^{\lambda'}$ -module structure on V .

We check it case-by-case.

When L is even, for each $0 \leq 2u \leq L - 2$, observe the following:

$$\begin{aligned}
 \overbrace{YX \cdots X}^L : \cdot \langle s, 2u \rangle &\xrightarrow{(YX)^{L/2-u-1}} \langle s, L - 2 \rangle \\
 &\xrightarrow{YX} \lambda \langle s + 1, L - 1 \rangle \\
 &\xrightarrow{(YX)^u} \lambda \langle s + 1 + 4u, L - 1 - 2u \rangle, \\
 \cdot \langle s, 2u + 1 \rangle &\xrightarrow{(YX)^u} \langle s + 4u, 1 \rangle \xrightarrow{YX} \langle s + 4u + 3, 0 \rangle \\
 &\xrightarrow{(YX)^{L/2-u-1}} \langle s + 4u + 3, L - 2u - 2 \rangle.
 \end{aligned}$$

$$\begin{aligned}
 \overbrace{XY \cdots Y}^L : \cdot \langle s, 2u \rangle &\xrightarrow{(XY)^u} \langle s + 4u, 0 \rangle \xrightarrow{XY} \langle s + 4u + 1, 1 \rangle \\
 &\xrightarrow{(XY)^{L/2-u-1}} \langle s + 4u + 1, L - 2u - 1 \rangle, \\
 \cdot \langle s, 2u + 1 \rangle &\xrightarrow{(XY)^{L/2-u-1}} \langle s, L - 1 \rangle \\
 &\xrightarrow{XY} \lambda \langle s + 3, L - 2 \rangle \\
 &\xrightarrow{(XY)^u} \lambda \langle s + 3 + 4u, L - 2 - 2u \rangle.
 \end{aligned}$$

Thus it follows that $\overbrace{YX \cdots X}^L \equiv \lambda \overbrace{XY \cdots Y}^L$ in $\text{End}_k(V)$.

When L is odd (so $L \geq 3$), for each $2 \leq 2u \leq L - 1$, observe the following:

$$\begin{aligned}
 \overbrace{YX \cdots Y}^L : \cdot \langle s, 0 \rangle &\xrightarrow{(YX)^{(L-1)/2}} \langle s, L - 1 \rangle \xrightarrow{Y} \lambda \langle s + 1, L - 1 \rangle, \\
 \cdot \langle s, 2u \rangle &\xrightarrow{(YX)^{(L-1)/2-u}} \langle s, L - 1 \rangle \xrightarrow{Y} \lambda \langle s + 1, L - 1 \rangle \\
 &\xrightarrow{(XY)^u} \lambda \langle s + 1 + 4u, L - 1 - 2u \rangle. \\
 \cdot \langle s, 2u - 1 \rangle &\xrightarrow{(YX)^{u-1}} \langle s + 4u - 4, 1 \rangle \xrightarrow{YX} \langle s + 4u - 1, 0 \rangle \\
 &\xrightarrow{(YX)^{(L-1)/2-u}} \langle s + 4u - 1, L - 1 - 2u \rangle \xrightarrow{Y} \langle s + 4u - 1, L - 2u \rangle.
 \end{aligned}$$

$$\begin{aligned}
& \overbrace{XY \cdots X}^L : \cdot \langle s, 0 \rangle \mapsto \langle s+1, L-1 \rangle, \\
& \cdot \langle s, 2u \rangle \xrightarrow{(XY)^u} \langle s+4u, 0 \rangle \xrightarrow{X} \langle s+4u+1, 0 \rangle \\
& \quad \xrightarrow{(YX)^{(L-1)/2-u}} \langle s+4u+1, L-2u-1 \rangle, \\
& \cdot \langle s, 2u-1 \rangle \xrightarrow{(XY)^{(L-1)/2-u}} \langle s, L-2 \rangle \xrightarrow{XY} \lambda \langle s+1, L-1 \rangle \\
& \quad \xrightarrow{(XY)^{u-1}} \lambda \langle s+4u-3, L-2u+1 \rangle \xrightarrow{X} \lambda \langle s+4u-1, L-2u \rangle.
\end{aligned}$$

Thus we have that $\overbrace{YX \cdots Y}^L \equiv \lambda \overbrace{XY \cdots X}^L$ in $\text{End}_k(V)$.

In either case V becomes a right $B'/J_L^{\lambda'}$ -module by the action.

(Step 4) V/W is a $B'/J_L^{\lambda'} + I_N^{\nu'}$ -module of dimension $2NL$.

Since V/W has the set $\{\langle s, t \rangle \mid 1 \leq s \leq 2N, 0 \leq t \leq L-1\}$ as a basis, V/W has dimension $2NL$.

The action of X^2 is given by $X^2 : \langle s, t \rangle \mapsto \langle s+2, t \rangle$.

Thus for $1 \leq s \leq 2N$, $0 \leq t \leq L-1$, it follows that

$$X^{2N} : \langle s, t \rangle \mapsto \langle s+2N, t \rangle = \langle 1 \cdot (2N) + s, t \rangle \equiv \nu \langle s, t \rangle \pmod{W}.$$

So we have that $1 \equiv \nu X^{2N}$ in $\text{End}_k(V/W)$.

Thus it is done.

(Step 5) We construct a right $A_{NL}^{\nu\lambda}$ -module $M = (V/W) \oplus (V/W)$.

There are two algebra maps

$$\begin{aligned}
& \pi'_0 : B \rightarrow B'/J_L^{\nu'} + I_N^{\nu'}, \\
& X_{11} \mapsto \bar{X} = x, \quad X_{22} \mapsto \bar{Y} = y, \\
& X_{i,i+1} \mapsto 0,
\end{aligned}$$

and

$$\begin{aligned}
& \pi'_1 : B \rightarrow B'/J_L^{\lambda'} + I_N^{\nu'}, \\
& X_{12} \mapsto \bar{X} = x, \quad X_{21} \mapsto \bar{Y} = y, \\
& X_{ii} \mapsto 0.
\end{aligned}$$

They induce algebra maps

$$\begin{aligned}
& \pi_0 : A_{NL}^{(\nu\lambda)} \rightarrow B'/J_L^{\nu'} + I_N^{\nu'}, \\
& x_{11} \mapsto x, \quad x_{22} \mapsto y, \quad x_{i,i+1} \mapsto 0, \\
& \pi_1 : A_{NL}^{(\nu\lambda)} \rightarrow B'/J_L^{\lambda'} + I_N^{\nu'}, \\
& x_{12} \mapsto x, \quad x_{21} \mapsto y, \quad x_{ii} \mapsto 0.
\end{aligned}$$

Using these, we obtain the right $A_{NL}^{(\nu\lambda)}$ -module $V/W = V_0$ through π_0 with a basis

$$\{\langle s, t \rangle_0 = \langle s, t \rangle \mid 1 \leq s \leq 2N, 0 \leq t \leq L-1\},$$

and the right $A_{NL}^{(\nu\lambda)}$ -module $V/W = V_1$ through π_1 with a basis

$$\{\langle s, t \rangle_1 = \langle s, t \rangle \mid 1 \leq s \leq 2N, 0 \leq t \leq L-1\}.$$

Let M be the right $A_{NL}^{(\nu\lambda)}$ -module $V_0 \oplus V_1$. We note that M has dimension $4NL$.

(Step 6) It follows that $M \simeq A_{NL}^{(\nu\lambda)}$ as right $A_{NL}^{(\nu\lambda)}$ -modules.

Define an $A_{NL}^{(\nu\lambda)}$ -module map $\phi : A_{NL}^{(\nu\lambda)} \rightarrow M$ and a k -linear map $\psi : M \rightarrow A_{NL}^{(\nu\lambda)}$ as follows:

$$\phi : A_{NL}^{(\nu\lambda)} \rightarrow M, \quad a \mapsto \{\langle 2N, 0 \rangle_0 + \nu \langle 2N, 0 \rangle_1\} \cdot a,$$

$$\psi : M \rightarrow A_{NL}^{(\nu\lambda)}, \quad \langle s, t \rangle_0 \mapsto x_{11}^s \overbrace{x_{22} x_{11} x_{22} \cdots}^t,$$

$$\langle s, t \rangle_1 \mapsto x_{12}^s \overbrace{x_{21} x_{12} x_{21} \cdots}^t.$$

It is easy to see that ψ is surjective and that $\phi \circ \psi$ is the identity map on M . Therefore we have that $M \simeq A_{NL}^{(\nu\lambda)}$ as $A_{NL}^{(\nu\lambda)}$ -modules, in particular $\dim A_{NL}^{(\nu\lambda)} = \dim M = 4NL$.

This completes the proof of i).

ii) \sim v) These are easily verified by i). Since $A_{NL}^{(\nu\lambda)}$ is generated by $\{x_{ij}\}$, it is commutative iff $(L, \lambda) = (2, +1)$.

vi) There is an algebra map $B \rightarrow B^{op}$, $X_{ij} \mapsto X_{ji} \cdot (X_{11}^{2(2N-1)} + X_{12}^{2(2N-1)})$, and this induces an algebra map S ,

$$\begin{array}{ccc} B & \longrightarrow & B^{op} \\ \downarrow \pi & & \downarrow \\ A_{NL}^{(\nu\lambda)} & \xrightarrow{S} & (A_{NL}^{(\nu\lambda)})^{op}. \end{array}$$

The anti-algebra map S is an antipode of $A_{NL}^{(\nu\lambda)}$, which is given by

$$\begin{aligned} S : x_{ij} &\mapsto x_{ji} (x_{11}^{2(2N-1)} + x_{12}^{2(2N-1)}) \\ &= x_{ji} (x_{11}^2 + x_{12}^2)^{-1}. \end{aligned}$$

So $A_{NL}^{(\nu\lambda)}$ is an involutory Hopf algebra.

vii) The element $\Lambda = \sum x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t$ ($1 \leq s \leq 2N, 0 \leq t \leq L-1$) is non-zero by i).

Recall that Λ is called a left (resp. right) integral if $a\Lambda$ (resp. Λa) = $\varepsilon(a)\Lambda$ for all $a \in A_{NL}^{(\nu\lambda)}$.

It is enough to check on the subset $\{x_{ij}\}$. Observe the following.

$$\begin{aligned}
x_{12}\Lambda &= x_{21}\Lambda = 0 \\
&= \varepsilon(x_{12})\Lambda = \varepsilon(x_{21})\Lambda. \\
x_{11}\Lambda &= \sum x_{11}^{s+1} \overbrace{x_{22}x_{11}x_{22} \cdots}^t = \sum x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t = \Lambda \\
&= \varepsilon(x_{11})\Lambda. \\
x_{22}\Lambda &= \sum x_{22}x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t \\
&= \sum_{s:\text{even}} x_{11}^s x_{22} \overbrace{x_{22}x_{11}x_{22} \cdots}^t + \sum_{s:\text{odd}} x_{11}^{s-1} x_{22}x_{11} \overbrace{x_{22}x_{11}x_{22} \cdots}^t \\
&= \sum_{s:\text{even}, t=0} x_{11}^s x_{22} + \sum_{s:\text{even}, t=1} x_{11}^{s+2} + \sum_{s:\text{even}, t \geq 2} x_{11}^{s+3} \overbrace{x_{22}x_{11}x_{22} \cdots}^{t-2} \\
&\quad + \sum_{s:\text{odd}, t \leq L-3} x_{11}^{s-1} \overbrace{x_{22}x_{11}x_{22} \cdots}^{t+2} + \sum_{s:\text{odd}, t=L-2} x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^{L-1} \\
&\quad + \sum_{s:\text{odd}, t=L-1} x_{11}^{s+2} \overbrace{x_{22}x_{11}x_{22} \cdots}^{L-2} \\
&= \sum_{s:\text{even}} x_{11}^s x_{22} + \sum_{s:\text{even}} x_{11}^s + \sum_{s:\text{odd}, 0 \leq t \leq L-3} x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t \\
&\quad + \sum_{s:\text{even}, 2 \leq t \leq L-1} x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^t + \sum_{s:\text{odd}} x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^{L-1} \\
&\quad + \sum_{s:\text{odd}} x_{11}^s \overbrace{x_{22}x_{11}x_{22} \cdots}^{L-2} \\
&= \Lambda \\
&= \varepsilon(x_{22})\Lambda.
\end{aligned}$$

Thus Λ is a left integral. It is similarly shown that Λ is a right integral. Therefore Λ is a non-zero two-sided integral.

viii) It follows that $\varepsilon(\Lambda) = 2NL \neq 0$ iff $chk \not\perp NL$. □

REMARK 3.2. For the multiplication relations of $A_{NL}^{(v\lambda)}$, we note the following.

- x_{ij}^2 is central.
- $x_{ii}^{2N+1} = x_{ii}$, and $x_{i,i+1}^{2N+1} = vx_{i,i+1}$.
- $x_{11}^{4N} + x_{12}^{4N} = 1$.
- $(x_{11}^{2s} + \mu x_{12}^{2s})^{-1} = x_{11}^{2(2N-s)} + \mu x_{12}^{2(2N-s)}$ for $1 \leq s \leq N, \mu = \pm 1$.

Set $h_{\pm} = x_{11}^2 \pm x_{12}^2$ and $g = \overbrace{x_{11}x_{22}x_{11}\cdots}^L + \sqrt{\lambda} \overbrace{x_{12}x_{21}x_{12}\cdots}^L$ for a fixed $\sqrt{\lambda}$. C_m denotes the cyclic group of order m .

PROPOSITION 3.3. i) The subgroup $\langle h_+, h_- \rangle$ of G is central in $A_{NL}^{(v\lambda)}$, and the order is $2N$. As groups

$$\langle h_+, h_- \rangle \simeq \begin{cases} C_N \times C_2, & \text{if } (N, v) = (\text{even}, +1); \\ C_{2N}, & \text{otherwise.} \end{cases}$$

ii) $G \subset Z(A_{NL}^{(v\lambda)})$, the center of $A_{NL}^{(v\lambda)}$, iff $g \in Z(A_{NL}^{(v\lambda)})$ iff $(L, \lambda) = (\text{even}, +1)$.

PROOF. i) The order of $\langle h_+, h_- \rangle$ is $2N$ by Theorem 3.1.

If $(N, v) =$

$$\begin{cases} (\text{even}, +1), & \text{then } \langle h_+, h_- \rangle = \langle h_+ \rangle \times \langle x_{11}^{2N} - x_{12}^{2N} \rangle, \\ (\text{even}, -1), & \text{then } \langle h_+, h_- \rangle = \langle h_+ \rangle = \langle h_- \rangle, \\ (\text{odd}, +1), & \text{then } \langle h_+, h_- \rangle = \langle h_- \rangle, \\ (\text{odd}, -1), & \text{then } \langle h_+, h_- \rangle = \langle h_+ \rangle. \end{cases}$$

ii) Note that $G = \langle h_+, h_- \rangle \cup \langle h_+, h_- \rangle g$. So it follows that $G \subset Z(A_{NL}^{(v\lambda)})$ iff $g \in Z(A_{NL}^{(v\lambda)})$.

It is easy to see that

$$g \text{ is central} \Leftrightarrow \begin{cases} x_{ii} \cdot \overbrace{x_{11}x_{22}\cdots}^L = \overbrace{x_{11}x_{22}\cdots}^L \cdot x_{ii}, \\ x_{i,i+1} \cdot \overbrace{x_{12}x_{21}\cdots}^L = \overbrace{x_{12}x_{21}\cdots}^L \cdot x_{i,i+1}, \end{cases} \quad \text{for } i = 1, 2. \quad \square$$

REMARK 3.4.

- i) The dimension of a simple subcoalgebra of $A_{NL}^{(v\lambda)}$ is either 1 or $2^2 = 4$.
- ii) The simple subcoalgebra C_{01} generates $A_{NL}^{(v\lambda)}$ as an algebra.

iii) For the YB-coalgebra $(C, \sigma_{\alpha\beta})$, $C \simeq C_{01} \subset A_{NL}^{(\nu\lambda)}$, $X_{ij} \mapsto x_{ij}$, is a coalgebra $\sigma_{\alpha\beta}$ -map.

We identify C and C_{01} .

iv) $A_{12}^{(+)}$ ($\simeq A_{12}^{(-)}$, see Prop.3.12 below) is the “non-trivial” semisimple Hopf algebra of dimension 8 ([Mas2]). The ideal decomposition is given as follows:

$$\begin{aligned} A_{12}^{(+)} &= k(x_{11} + x_{22} + x_{11}^2 + x_{11}x_{22}) \oplus k(x_{11} - x_{22} - x_{11}^2 + x_{11}x_{22}) \\ &\quad \oplus k(x_{11} - x_{22} + x_{11}^2 - x_{11}x_{22}) \oplus k(x_{11} + x_{22} - x_{11}^2 - x_{11}x_{22}) \\ &\quad \oplus \text{span}_k\{x_{12}, x_{21}, x_{12}^2, x_{12}x_{21}\}. \end{aligned}$$

v) Since the subHopf algebra $K = k\langle h_+, h_- \rangle$ is normal, $A_{NL}^{(\nu\lambda)}K^+$ is a Hopf ideal, where $K^+ = \text{Ker } \varepsilon_K$. So $A_{NL}^{(\nu\lambda)}/A_{NL}^{(\nu\lambda)}K^+ = \bar{A}$ is a Hopf algebra of dimension $2L$. It is easy to see that the elements $\bar{x}_{11} = a$, $\bar{x}_{22} = b \in \bar{A}$ are grouplike and generate \bar{A} as an algebra. This means that \bar{A} is a group-algebra. Moreover let $ab = c$, then the order of c is L . Then,

$$\begin{aligned} \bar{A} &= k\langle a, b \mid a^2 = 1 = b^2, \overbrace{baba \cdots}^L = \overbrace{abab \cdots}^L \rangle \\ &= k\langle a, c \mid a^2 = 1, c^L = 1, \overbrace{aca^{-1}}^L = c^{-1} \rangle \\ &= kD_L, \quad \text{where } D_L \text{ is the dihedral group of order } 2L. \end{aligned}$$

Thus we obtain a short exact sequence by means of [Mas1, Definition 1.3]

$$1 \rightarrow K \hookrightarrow A_{NL}^{(\nu\lambda)} \rightarrow kD_L \rightarrow 1.$$

vi) As bialgebras

$$\begin{aligned} B/J_2^\lambda &= B/(X_{11}X_{22} - X_{22}X_{11}, X_{12}X_{21} - \lambda X_{21}X_{12}) \\ &= k\langle X_{ij} \rangle / (X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2, X_{ij}X_{lm} \ (i+j+l+m \equiv 1), \\ &\quad X_{11}X_{22} - X_{22}X_{11}, X_{12}X_{21} - \lambda X_{21}X_{12}) \\ &= B^{(\lambda)} / (X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2). \end{aligned}$$

Thus $A_{N2}^{(\nu\lambda)}$ is furthermore a quotient bialgebra of $B^{(\lambda)}$:

$$A_{N2}^{(\nu\lambda)} = B^{(\lambda)} / (X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2, 1 - (X_{11}^{2N} + \nu X_{12}^{2N})).$$

We note that $\{X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2\}$ spans a coideal of $B^{(\lambda)}$ and that $\{1 - (X_{11}^{2N} + \nu X_{12}^{2N})\}$ spans a coideal modulo the coideal $\text{span}_k\{X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2\}$.

Recall that C_{st} denotes a simple subcoalgebra of dimension 4 of $A_{NL}^{(\nu\lambda)}$ for $0 \leq s \leq N-1$, $1 \leq t \leq L-1$. Let $\langle C_{st} \rangle$ denote the subHopf algebra generated by C_{st} . It is easy to see that $\langle C_{st} \rangle$ is commutative iff either t is even or $(L, \lambda) = (2t, +1)$. So it follows that t is odd if $\langle C_{st} \rangle$ is non-commutative.

We show that $\langle C_{st} \rangle$ is a member of the family $\{A_{NL}^{(\nu\lambda)}\}$ if t is odd.

Set

$$\begin{aligned} \text{GCD}(L, t) &= m_L, & \text{GCD}(N, 2s+t) &= m_N, \\ L/m_L &= L_0, & N/m_N &= N_0, & t/m_L &= t_0, & (2s+t)/m_N &= (s, t)_0, \\ & & & & (2 \leq L_0 \leq L, 1 \leq N_0 \leq N). \end{aligned}$$

THEOREM 3.5. *Assume that t is odd and $C_{st} \subset A_{NL}^{(\nu\lambda)}$. Then*

$$\langle C_{st} \rangle \simeq A_{N_0 L_0}^{(\nu\lambda)} \quad \text{as Hopf algebras.}$$

PROOF. Let t be odd, and fix $0 \leq s \leq N-1$ and $1 \leq t \leq L-1$. We note that integers $2s+t$, t_0 , $(s, t)_0$, m_L and m_N are also odd.

Set

$$\begin{aligned} z_{11} &= x_{11}^{2s} \overbrace{x_{11} x_{22} \cdots x_{11}}^t, & z_{12} &= x_{12}^{2s} \overbrace{x_{12} x_{21} \cdots x_{12}}^t, \\ z_{21} &= x_{12}^{2s} \overbrace{x_{21} x_{12} \cdots x_{21}}^t, & z_{22} &= x_{11}^{2s} \overbrace{x_{22} x_{11} \cdots x_{22}}^t. \end{aligned}$$

The map $\omega : A_{N_0 L_0}^{(\nu\lambda)} \rightarrow \langle C_{st} \rangle$, $x_{ij} \mapsto z_{ij}$, is a (well-defined) surjective Hopf algebra map. This is easily verified.

We show that the map ω is injective.

Recall and set that

$$\begin{aligned} G_0 &= G(A_{N_0 L_0}^{(\nu\lambda)}) \\ &= \{x_{11}^{2u} \pm x_{12}^{2u} x_{11}^{2u} \cdot \overbrace{x_{11} x_{22} x_{11} \cdots}^{L_0} \pm \sqrt{\lambda} x_{12}^{2u} \cdot \overbrace{x_{12} x_{21} x_{12} \cdots}^{L_0} \mid 1 \leq u \leq N_0\}, \\ (C_{uv})_0 &= C_{uv} \subset A_{N_0 L_0}^{(\nu\lambda)}. \end{aligned}$$

Then it follows that

$$A_{N_0 L_0}^{(\nu\lambda)} = kG_0 \oplus \Sigma(C_{uv})_0.$$

Thus it is enough to show that ω is injective on kG_0 and on $\Sigma(C_{uv})_0$.

It is easy to see that ω is injective on kG_0 .

So we show that ω is injective on $\Sigma(C_{uv})_0$.

First we examine $\omega((C_{uv})_0)$ for $0 \leq u \leq N_0 - 1$, $1 \leq v \leq L_0 - 1$.

Let $tv = qL + r$, for some q , $0 \leq r \leq L - 1$. It is easy to see that $r \neq 0$, so it follows that $1 \leq r$, $L - r \leq L - 1$.

For $x_{11}^{2u} \overbrace{x_{11}x_{22}x_{11} \cdots}^v \in (C_{uv})_0$, observe that

$$\begin{aligned}
& \omega(x_{11}^{2u} \overbrace{x_{11}x_{22}x_{11} \cdots}^v) \\
&= z_{11}^{2u} \overbrace{z_{11}z_{22}z_{11} \cdots}^v \\
&= (x_{11}^{2s} \overbrace{x_{11}x_{22} \cdots x_{11}}^t)^{2u} \cdot \overbrace{(x_{11}^{2s} \cdot x_{11}x_{22} \cdots x_{11})(x_{11}^{2s} \cdot x_{22}x_{11} \cdots x_{22}) \cdots}^v \\
&= x_{11}^{2(2s+t)u} x_{11}^{2sv} \overbrace{x_{11}x_{22}x_{11} \cdots}^{tv} \\
&= x_{11}^{2(2s+t)u} x_{11}^{2sv} \\
&\quad \times \begin{cases} x_{11}^{qL} \cdot \overbrace{x_{11}x_{22}x_{11} \cdots}^r, & \text{if } q \text{ is even,} \\ x_{11}^{(q-1)L} \cdot \overbrace{x_{11}x_{22}x_{11} \cdots}^{L+r}, & \text{if } q \text{ is odd} \end{cases} \\
&= \begin{cases} x_{11}^{2\{(2s+t)u+sv+(q/2)L\}} \cdot \overbrace{x_{11}x_{22}x_{11} \cdots}^r, & \text{if } q \text{ is even,} \\ x_{11}^{2\{(2s+t)u+sv+((q-1)/2)L+r\}} \cdot \overbrace{x_{22}x_{11}x_{22} \cdots}^{L-r}, & \text{if } q \text{ is odd} \end{cases} \\
&\neq 0.
\end{aligned}$$

Let

$$(a, b) = \begin{cases} ((2s+t)u + sv + \frac{q}{2}L \bmod N, r), & \text{if } q \text{ is even,} \\ ((2s+t)u + sv + \frac{(q-1)}{2}L + r \bmod N, L - r), & \text{if } q \text{ is odd,} \\ (0 \leq a \leq N - 1, 1 \leq b \leq L - 1). \end{cases}$$

So we have that

$$0 \neq \omega(x_{11}^{2u} \overbrace{x_{11}x_{22}x_{11} \cdots}^v) \in \omega((C_{uv})_0) \cap C_{ab}.$$

Since C_{ab} is a simple subcoalgebra, it follows that

$$\omega((C_{uv})_0) = C_{ab} \subset A_{NL}^{(v\lambda)}.$$

Thus ω is injective on $(C_{uv})_0$.

Next assume that there are $0 \leq u, u' \leq N_0 - 1, 1 \leq v, v' \leq L_0 - 1$ such that $\omega((C_{uv})_0) = \omega((C_{u'v'})_0)$.

Let $tv' = q'L + r', 1 \leq r' \leq L - 1$.

It is easy to see that $q \equiv q' \pmod{2}$ implies $u = u'$ and $v = v'$.

So let q be even and q' odd. This implies that $q + q' + 1$ is even and that $L = r + r'$.

We have that $t(v + v') = (q + q' + 1)L$, so it follows that $L_0 | v + v'$.

It follows that $L_0 = v + v'$, since $1 \leq v, v' \leq L_0 - 1$.

So we have $t = (q + q' + 1)m_L$, and this means that t is even. A contradiction.

Thus $\omega((C_{uv})_0) = \omega((C_{u'v'})_0)$ iff $u = u', v = v'$, so ω is injective on $\Sigma(C_{uv})_0$. Therefore we have the injectivity of ω .

This completes the proof of the theorem. \square

It is easy to see that the following lemma holds.

LEMMA 3.6. *Assume that A_1 and A_2 are bialgebras over an algebraically closed field. If the bialgebra $A_1 \otimes A_2$ is generated by a simple subcoalgebra as an algebra, then so is $A_i, i = 1, 2$. Moreover if any simple subcoalgebra of $A_1 \otimes A_2$ has dimension 1 or n^2 , then either A_1 or A_2 is pointed.*

COROLLARY 3.7.

i) *Assume that $A_{NL}^{(v\lambda)}$ is non-commutative, i.e. $(L, \lambda) \neq (2, +1)$, and $C_{st} \subset A_{NL}^{(v\lambda)}$.*

Then

$$\langle C_{st} \rangle = A_{NL}^{(v\lambda)} \quad \text{iff } t \text{ is odd, } (L, t) = 1 \text{ and } (N, 2s + t) = 1.$$

ii) *Assume simply that t is odd and $C_{st} \subset A_{NL}^{(v\lambda)}$. Then*

$$\langle C_{st} \rangle = A_{NL}^{(v\lambda)} \quad \text{iff } (L, t) = 1, (N, 2s + t) = 1.$$

iii) *Let N be $2^n m$, and m odd. Then*

$$A_{NL}^{(v\lambda)} \simeq A_{2^n, L}^{(v\lambda)} \otimes kC_m \quad \text{as Hopf algebras.}$$

iv) *If $A_{2^n, L}^{(v\lambda)}$ is non-commutative, then it is indecomposable as the tensor product of its subHopf algebras.*

PROOF. i), ii) These follow from the dimensionality.

iii) Let N be $2^n m$ and m odd. We may assume that $m \geq 3$. Now let $s = (m-1)/2$, $t = 1$, then it follows that $2s + t = m$, $N_0 = 2^n$, $L_0 = L$, and $\langle C_{st} \rangle \simeq A_{2^n, L}^{(v\lambda)}$.

Let $f = x_{11}^{2 \cdot 2^n} + vx_{12}^{2 \cdot 2^n}$. Then f is a central grouplike element with order m , and $C_{st} \cdot f = C_{s't}$, where $s' = 2^n + (m-1)/2 \leq N-1$.

For such s , s' and t , it follows that

$$\begin{aligned} (2s' + t, N) &= \left(2 \left\{ 2^n + \frac{m-1}{2} \right\} + 1, 2^n m \right) \\ &= (2^{n+1} + m, 2^n m) \\ &= 1. \end{aligned}$$

Thus the simple subcoalgebra $C_{st} \cdot f = C_{s't}$ generates $A_{NL}^{(v\lambda)}$ as an algebra by ii). Therefore we have that

$$A_{2^n m, L}^{(v\lambda)} \simeq A_{2^n, L}^{(v\lambda)} \otimes kC_m, \quad \text{as Hopf algebras.}$$

iv) Let $2^n = N$. Applying Lemma 3.6 to $A_{NL}^{(v\lambda)}$, we may assume

$$A_{NL}^{(v\lambda)} = \langle C_{st} \rangle \otimes kF,$$

for some $0 \leq s \leq N-1$, $1 \leq t \leq L-1$, (abelian)subgroup $F \subset G(A_{NL}^{(v\lambda)})$.

Since $A_{NL}^{(v\lambda)}$ is non-commutative, so is $\langle C_{st} \rangle$. This means that t is odd. By Theorem 3.5, $\langle C_{st} \rangle \simeq A_{N_0 L_0}^{(v\lambda)}$.

Comparing the dimensions, we have that $|F| = m_N m_L$.

Counting the number of 4-dimensional simple subcolagebras, we have the following:

$$\begin{aligned} N(L-1) &= N_0(L_0-1) \cdot |F| \\ &= N_0(L_0-1)m_N m_L \\ &= N(L-m_L). \end{aligned}$$

Thus we have that $m_L = 1$.

On the other hand, it follows that $m_N = 1$ since $2s + t$ is odd and N is a power of 2.

Thus we have that $F = \langle 1 \rangle$. □

Next we show that we can obtain all braidings on $A_{NL}^{(v\lambda)}$. See [GW], [G]. We identify $C < A_{NL}^{(v\lambda)}$ as in Remark 3.4. Note that any braiding on $A_{NL}^{(v\lambda)}$ is

determined on $C \otimes C$. If a bilinear map τ on C extends to a braiding on $A_{NL}^{(v\lambda)}$, we denote the braiding by $\tilde{\tau}$.

Recall YB-forms $\sigma_{\alpha\beta}, \tau_{\alpha\beta}^{(\lambda)}$ on C .

CLAIM 3.8. *Let σ be a braiding on $A_{NL}^{(v\lambda)}$.*

i) *If $L \geq 3$, $\sigma|_{C \otimes C}$ coincides with $\sigma_{\alpha\beta}$ for some $\alpha, \beta \in k^\times$ such that $(\alpha\beta)^N = v$, $(\alpha\beta^{-1})^L = \lambda$.*

ii) *If $L = 2$, $\sigma|_{C \otimes C}$ coincides with either $\sigma_{\alpha\beta}$ for some $\alpha, \beta \in k^\times$ such that $(\alpha\beta)^N = v$, $(\alpha\beta^{-1})^2 = \lambda$ or $\tau_{\gamma\delta}^{(\lambda)}$ for some $\gamma, \delta \in k^\times$ such that $\delta^2 = \gamma^2$, $\gamma^{2N} = 1$.*

PROOF. i) Assume that $L \geq 3$.

The subcoalgebra $C \cdot C$ of $A_{NL}^{(v\lambda)}$ has a basis

$$\{x_{11}^2, x_{12}^2, x_{11}x_{22}, x_{22}x_{11}, x_{12}x_{21}, x_{21}x_{12}\}.$$

We have similarly as in Proposition 2.9,

$$\sigma|_{C \otimes C} = \sigma_{\alpha\beta} \quad \text{for some } \alpha, \beta \in k^\times.$$

Moreover σ satisfies the following:

$$\begin{aligned} 0 &= \sigma(1 - (x_{11}^{2N} + vx_{12}^{2N}), x_{11}) \\ &= 1 - v\{\sigma_{\alpha\beta}(x_{12}, x_{12})\sigma_{\alpha\beta}(x_{12}, x_{21})\}^N \\ &= 1 - v(\alpha\beta)^N. \end{aligned}$$

Thus it follows that $(\alpha\beta)^N = v$.

Observe that when L is even,

$$\begin{aligned} 0 &= \sigma(\overbrace{x_{21}x_{12} \cdots x_{12}}^L - \lambda \overbrace{x_{12}x_{21} \cdots x_{21}}^L, x_{22}) \\ &= \alpha^L - \lambda\beta^L, \end{aligned}$$

and that when L is odd,

$$\begin{aligned} 0 &= \sigma(\overbrace{x_{21}x_{12} \cdots x_{21}}^L - \lambda \overbrace{x_{12}x_{21} \cdots x_{12}}^L, x_{21}) \\ &= \alpha^L - \lambda\beta^L. \end{aligned}$$

Thus in either case, it follows that $\alpha^L = \lambda\beta^L$, or $(\alpha\beta^{-1})^L = \lambda$.

ii) Assume that $L = 2$.

The subcoalgebra $C \cdot C$ of $A_{N^2}^{(\nu\lambda)}$ has a basis

$$\{x_{11}^2, x_{12}^2, x_{11}x_{22}, x_{12}x_{21}\}.$$

As in the proof of Proposition 2.9, we have the following:

$$\begin{aligned}\sigma(x_{ij}, x_{lm})x_{jj}x_{mm} &= x_{ll}x_{ii}\sigma(x_{ij}, x_{lm}), \\ \sigma(x_{i,j+1}, x_{l,m+1})x_{j+1,j}x_{m+1,m} &= x_{l,l+1}x_{i,i+1}\sigma(x_{i+1,j}, x_{l+1,m}).\end{aligned}$$

Using these relations, we have the following with $\alpha, \beta, \gamma, \delta \in k$,

σ	X_{11}	X_{12}	X_{21}	X_{22}
X_{11}	γ	0	0	δ
X_{12}	0	α	β	0
X_{21}	0	β	α	0
X_{22}	$\lambda\delta$	0	0	γ .

Moreover σ satisfies the following equations:

$$\begin{aligned}0 &= \sigma(x_{11}x_{12}, x_{12}) = \gamma\alpha, \\ 0 &= \sigma(x_{11}x_{21}, x_{12}) = \gamma\beta, \\ 0 &= \sigma(x_{11}x_{12}, x_{21}) = \delta\beta, \\ 0 &= \sigma(x_{11}x_{21}, x_{21}) = \delta\alpha.\end{aligned}$$

So it follows that either $\gamma = 0 = \delta$ or $\alpha = 0 = \beta$.

Thus $\sigma|_{C \otimes C}$ is either $\sigma_{\alpha\beta}$ or $\tau_{\gamma\delta}^{(\lambda)}$, for $\alpha, \beta, \gamma, \delta \in k^\times$.

If $\sigma|_{C \otimes C} = \sigma_{\alpha\beta}$, then the relations on α, β follow similarly as in the proof of i).

Let $\sigma|_{C \otimes C} = \tau_{\gamma\delta}^{(\lambda)}$. Observe that

$$\begin{aligned}0 &= \sigma(x_{11}^2 - x_{22}^2, x_{22}) \\ &= \tau_{\gamma\delta}^{(\lambda)}(x_{11}, x_{22})^2 - \tau_{\gamma\delta}^{(\lambda)}(x_{22}, x_{22})^2 \\ &= \delta^2 - \gamma^2, \\ 0 &= \sigma(1 - (x_{11}^{2N} - \nu x_{12}^{2N}), x_{11}) \\ &= 1 - \tau_{\gamma\delta}^{(\lambda)}(x_{11}, x_{11})^{2N} \\ &= 1 - \gamma^{2N}.\end{aligned}$$

Thus it follows that $\delta^2 = \gamma^2$, $\gamma^{2N} = 1$. □

CLAIM 3.9.

- i) The YB-form $\sigma_{\alpha\beta}$ extends to a braiding on $A_{NL}^{(v\lambda)}$ if $(\alpha\beta)^N = v$, $(\alpha\beta^{-1})^L = \lambda$.
- ii) The YB-form $\tau_{\gamma\delta}^{(\lambda)}$ extends to a braiding on $A_{N2}^{(v\lambda)}$ if $\delta^2 = \gamma^2$, $\gamma^{2N} = 1$.

PROOF. Recall that B has braidings $\{\tilde{\sigma}_{\alpha\beta} | \alpha, \beta \in k^\times\}$ and that $B^{(\lambda)}$ has braidings $\{\tilde{\tau}_{\gamma\delta}^{(\lambda)} | \gamma, \delta \in k^\times\}$.

i) It is easy to see by Proposition 1.1 that $\tilde{\sigma}_{\alpha\beta} : B \otimes B \rightarrow k$ induces a braiding on $A_{NL}^{(v\lambda)}$ iff

$$\begin{cases} (\alpha\beta)^N = v, \\ (\alpha\beta^{-1})^L = \lambda. \end{cases}$$

ii) Recall that $A_{N2}^{(v\lambda)} = B^{(\lambda)} / (X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2, 1 - (X_{11}^{2N} + vX_{12}^{2N}))$.

It follows that $\tilde{\tau}_{\gamma\delta}^{(\lambda)}$ induces a braiding on $B^{(\lambda)} / (X_{11}^2 - X_{22}^2, X_{12}^2 - X_{21}^2)$ iff $\delta^2 = \gamma^2$, and that $\tilde{\tau}_{\gamma\delta}^{(\lambda)}$ induces a braiding on $A_{N2}^{(v\lambda)}$ iff $\delta^2 = \gamma^2$, $\gamma^{2N} = 1 = \delta^{2N}$. \square

PROPOSITION 3.10.

i) The set of braidings on $A_{NL}^{(v\lambda)}$ is given as follows:

$$\begin{aligned} & \{\tilde{\sigma}_{\alpha\beta} | (\alpha\beta)^N = v, (\alpha\beta^{-1})^L = \lambda\}, \quad \text{if } L \geq 3, \\ & \{\tilde{\sigma}_{\alpha\beta}, \tilde{\tau}_{\gamma\delta}^{(\lambda)} | (\alpha\beta)^N = v, (\alpha\beta^{-1})^2 = \lambda, \delta^2 = \gamma^2, \gamma^{2N} = 1\}, \quad \text{if } L = 2. \end{aligned}$$

ii) $A_{NL}^{(v\lambda)}$ is, in fact, a braided Hopf algebra.

If $chk \nmid NL$, the number of braidings on $A_{NL}^{(v\lambda)}$ is

$$\begin{cases} 2NL, & \text{if } L \geq 3, \\ 8N, & \text{if } L = 2. \end{cases}$$

iii) The number of symmetric braidings on $A_{NL}^{(v\lambda)}$ is given as follows;

When $L \geq 3$,

N	L	(v, λ)	$\tilde{\sigma}$
odd	odd	$(\pm 1, \pm 1)$	2
		$(\pm 1, \mp 1)$	0
odd	even	$(v, +1)$	2
		$(v, -1)$	0
even	odd	$(+1, \lambda)$	2
		$(-1, \lambda)$	0
even	even	$(+1, +1)$	4
		otherwise	0.

When $L = 2$,

N	(v, λ)	$\tilde{\sigma}$	$\tilde{\tau}^{(\lambda)}$
<i>odd</i>	$(v, +1)$	2	4
	$(v, -1)$	0	0
<i>even</i>	$(+1, +1)$	4	4
	$(+1, -1)$	0	0
	$(-1, +1)$	0	4
	$(-1, -1)$	0	0.

PROOF. i) This follows from Claim 3.8 and 3.9.

ii) There is a surjective map

$$\{(p, q) \in k \times k \mid p^{2N} = v, q^{2L} = \lambda\} \rightarrow \{(\alpha, \beta) \in k \times k \mid (\alpha\beta)^N = v, (\alpha\beta^{-1})^L = \lambda\},$$

$$(p, q) \mapsto (pq, pq^{-1}).$$

Set $(p, q) \sim (p', q') \Leftrightarrow (p, q) = \pm(p', q')$. It is an equivalence relation, which induces the bijection

$$\{(p, q) \mid p^{2N} = v, q^{2L} = \lambda\} / \sim \approx \{(\alpha, \beta) \mid (\alpha\beta)^N = v, (\alpha\beta^{-1})^L = \lambda\}.$$

Let $chk \not\sim NL$. Then it follows that $|\{\tilde{\sigma}\}| = 2N \cdot 2L \cdot \frac{1}{2} = 2NL$. For $\tilde{\tau}^{(\lambda)}$, since $\gamma^{2N} = 1$ and $\delta^2 = \gamma^2$, it follows that $|\{\tilde{\tau}^{(\lambda)}\}| = 2N \cdot 2 = 4N$.

iii) Recall that $chk \neq 2$. On $A_{NL}^{(v\lambda)}$, $\tilde{\sigma}_{\alpha\beta}$ is symmetric iff $\alpha^2 = 1 = \beta^2$ and $(\alpha\beta)^N = v, (\alpha\beta^{-1})^L = \lambda$.

On $A_{N2}^{(v\lambda)}$, $\tilde{\tau}_{\gamma\delta}^{(\lambda)}$ is symmetric iff $\gamma^2 = 1, \delta^2 = \lambda$ and $\gamma^{2N} = 1, \delta^2 = \gamma^2$. \square

REMARK 3.11. The algebra map $\theta : A_{NL}^{(v\lambda)} \rightarrow A_{NL}^{(v\lambda)cop}$, $x_{ij} \mapsto x_{ji}$, is a bijective Hopf algebra map. Define $\langle a, b \rangle = \tilde{\sigma}_{\alpha\beta}(\theta(a), b)$ for $a, b \in A_{NL}^{(v\lambda)}$.

The linear map $\langle , \rangle : A_{NL}^{(v\lambda)} \otimes A_{NL}^{(v\lambda)} \rightarrow k$ is a non-trivial Hopf paring.

Using Proposition 3.10, we have the following indispensable proposition.

PROPOSITION 3.12. $A_{N_1 L_1}^{(v_1 \lambda_1)} \simeq A_{N_2 L_2}^{(v_2 \lambda_2)}$ if and only if both $(N_1, L_1) = (N_2, L_2)$ and

$$\begin{cases} (v_2, \lambda_2) = \pm(v_1, \lambda_1), & (\text{case } N_1, L_1 \text{ odd}); \\ \lambda_2 = \lambda_1, & (\text{case } N_1 \text{ odd}, L_1 \text{ even}); \\ v_2 = v_1, & (\text{case } N_1 \text{ even}, L_1 \text{ odd}); \\ (v_2, \lambda_2) = (v_1, \lambda_1), & (\text{case } N_1, L_1 \text{ even}). \end{cases}$$

PROOF. For a fixed $\sqrt{-1}$, we can define a bialgebra map $\xi : B \rightarrow B$,

$$\begin{aligned}\xi : X_{ii} &\mapsto X_{ii}, \\ X_{12} &\mapsto \sqrt{-1}X_{12}, \\ X_{21} &\mapsto -\sqrt{-1}X_{21}.\end{aligned}$$

Let

$$\check{A}_{NL}^{(v\lambda)} = \begin{cases} A_{NL}^{(-v,-\lambda)}, & \text{if } N, L \text{ are odd,} \\ A_{NL}^{(-v,\lambda)}, & \text{if } N \text{ is odd, } L \text{ is even,} \\ A_{NL}^{(v,-\lambda)}, & \text{if } N \text{ is even, } L \text{ is odd,} \\ A_{NL}^{(v\lambda)}, & \text{if } N, L \text{ are even.} \end{cases}$$

Then the following diagram commutes:

$$\begin{array}{ccc} B & \xrightarrow{\xi} & B \\ \pi \downarrow & & \downarrow \pi \\ A_{NL}^{(v\lambda)} & \xrightarrow{\simeq} & \check{A}_{NL}^{(v\lambda)}. \end{array}$$

Thus by Proposition 3.10.iii), if N or L is odd, then the statement follows.

Assume that both N and L are even. Then

$$(v_1, \lambda_1) = \begin{cases} (++) \Rightarrow \text{by Prop. 3.10.iii), } (v_2, \lambda_2) = (++) . \\ (-+) \Rightarrow \text{by Prop. 3.3.ii), } G(A_{N_1 L_1}^{(v_1 \lambda_1)}) \text{ is central so } \lambda_2 = +1 . \\ \quad \text{By Prop. 3.10.iii), } v_2 = -1 \text{ so } (v_2, \lambda_2) = (-+) . \\ (+-) \Rightarrow \text{by Prop. 3.3.ii), } kG(A_{N_1 L_1}^{(v_1 \lambda_1)}) \cap Z(A_{N_1 L_1}^{(v_1 \lambda_1)}) = K \text{ so } \lambda_2 = -1 . \\ \quad \text{By Prop. 3.3.i), } v_2 = +1 \text{ so } (v_2, \lambda_2) = (+-) . \\ (--) \Rightarrow \text{it follows that } (v_2, \lambda_2) = (--) . \end{cases}$$

This completes the proof. □

REMARK 3.13 ([Mas2], [F]). The “non-trivial” 8-dimensional semisimple Hopf algebra is given by

$$A_{1,2}^{(+-)} \simeq A_{1,2}^{(--)}$$

Let $chk \neq 3$. The two “non-trivial” 12-dimensional semisimple Hopf algebras are given by

$$A_{1,3}^{(++)} \simeq A_{1,3}^{(--)} \quad \text{and} \quad A_{1,3}^{(+-)} \simeq A_{1,3}^{(-+)}.$$

Recall that H is a Hopf closure of B and that $A_{NL}^{(\nu\lambda)}$ is a Hopf algebra which is a quotient of B through π . So there is a Hopf algebra map $\tilde{\pi} : H \rightarrow A_{NL}^{(\nu\lambda)}$ such that $\tilde{\pi} = \pi|_B$.

H is a right $A_{NL}^{(\nu\lambda)}$ -comodule algebra via $\tilde{\pi}$. See [DT]. Then

PROPOSITION 3.14. *H is a cleft $A_{NL}^{(\nu\lambda)}$ -comodule algebra. Namely there is an invertible comodule map $\phi : A_{NL}^{(\nu\lambda)} \rightarrow H$.*

PROOF. Recall the basis $\{x_{11}^s \cdot \overbrace{x_{22}x_{11} \cdots}^t, x_{12}^s \cdot \overbrace{x_{21}x_{12} \cdots}^t \mid 1 \leq s \leq 2N, 0 \leq t \leq L-1\}$. This can be written as follows:

$$\left(\begin{array}{cccc} & x_{11}^{2(s+1)} & & x_{12}^{2(s+1)} \\ x_{11}^{2s} \cdot x_{11} & x_{11}^{2s} \cdot x_{22} & x_{12}^{2s} \cdot x_{12} & x_{12}^{2s} \cdot x_{21} \\ x_{11}^{2s} \cdot x_{11}x_{22} & x_{11}^{2s} \cdot x_{22}x_{11} & x_{12}^{2s} \cdot x_{12}x_{21} & x_{12}^{2s} \cdot x_{21}x_{12} \\ \vdots & \vdots & \vdots & \vdots \\ x_{11}^{2s} \cdot \overbrace{x_{11}x_{22} \cdots}^{L-1} & x_{11}^{2s} \cdot \overbrace{x_{22}x_{11} \cdots}^{L-1} & x_{12}^{2s} \cdot \overbrace{x_{12}x_{21} \cdots}^{L-1} & x_{12}^{2s} \cdot \overbrace{x_{21}x_{12} \cdots}^{L-1} \\ x_{11}^{2s} \cdot \overbrace{x_{11}x_{22} \cdots x_{LL}}^L & & x_{12}^{2s} \cdot \overbrace{x_{12}x_{21} \cdots x_{L,L+1}}^L & \end{array} \right)$$

for $0 \leq s \leq N-1$.

We use it. Define, for example, a linear map $\phi : A_{NL}^{(\nu\lambda)} \rightarrow B \rightarrow H$ by the small letters to its capital letters, i.e., x_{ij} to X_{ij} , etc. Then ϕ is a right $A_{NL}^{(\nu\lambda)}$ -comodule map.

We define another linear map $\psi : A_{NL}^{(\nu\lambda)} \rightarrow H$ as follows:

On the bottom row,

$$\psi : x_{11}^{2s} \cdot \overbrace{x_{11}x_{22} \cdots x_{LL}}^L \mapsto \overbrace{(X_{LL} \cdots X_{22}X_{11})}^L \cdot X_{11}^{2s} \left(\frac{1}{d_+} \right)^{2s+L},$$

$$x_{12}^{2s} \cdot \overbrace{x_{12}x_{21} \cdots x_{L,L+1}}^L \mapsto \lambda \overbrace{(X_{L,L+1} \cdots X_{21}X_{12})}^L \cdot X_{12}^{2s} \left(\frac{1}{d_+} \right)^{2s+L},$$

and on the other rows,

$$\psi = S \circ \phi.$$

Then we have $\psi = \phi^{-1}$, so ϕ is invertible.

Therefore H is a cleft $A_{NL}^{(\nu\lambda)}$ -comodule algebra. \square

Added in Proof

The group $G = G(A_{NL}^{(\nu\lambda)})$ is abelian, and the type is given as follows. The case that L is even:

$$G = \langle h_+, h_- \rangle \times \langle h_+^{-L/2} g \rangle$$

$$= \begin{cases} (C_N \times C_2) \times C_2, & \text{if } (N, \nu) = (\text{even}, +1); \\ (C_{2N}) \times C_2, & \text{otherwise.} \end{cases}$$

The case that L is odd:

$$G = \begin{cases} \langle h_\lambda^{(1-L)/2} g \rangle = C_{4N} & \text{if } \nu = -\lambda^N; \\ \langle h_\lambda^{(1-L)/2} g \rangle \times \langle h_+^{-1} h_- \rangle = C_{2N} \times C_2, & \text{if } \nu = \lambda^N. \end{cases}$$

Proposition 3.12 follows from this and Proposition 3.3.

References

- [D] Y. Doi, Braided bialgebras and quadratic bialgebras, *Comm. Algebra* **21(5)**, 1731–1749.
- [DT] Y. Doi and M. Takeuchi, Cleft comodule algebras for a bialgebra, *Comm. Algebra* **14** (1986), 801–817.
- [F] N. Fukuda, Semisimple Hopf algebras of dimension 12 (to appear).
- [GW] S. Gelaki and S. Westreich, On the quasitriangularity of $U_q(\mathfrak{sl}_n)'$, preprint.
- [G] S. Gelaki, Quantum groups of dimension pq^2 , preprint.
- [H] T. Hayashi, Quantum groups and quantum determinants, *J. Algebra* **152** (1992), 146–165.
- [Man] Yu. Manin, Quantum groups and non-commutative geometry, U. of Montreal Lectures, 1988.
- [Mas1] A. Masuoka, Coideal subalgebras in finite Hopf algebras, *J. Algebra* **163** (1994), 819–831.
- [Mas2] ———, Semisimple Hopf algebras of dimension 6, 8, *Israel J. Math.* **92** (1995), 361–373.
- [M] S. Montgomery, Hopf algebras and their actions on rings, American Mathematical Society, Providence, 1993.
- [S] M. Sweedler, *Hopf Algebras*, Benjamin, New York, 1969.
- [T1] M. Takeuchi, A two-parameter quantization of $GL(n)$, *Proc. Japan Acad.* **66. Ser. A** (1990), 112–114.
- [T2] ———, Matric bialgebras and quantum groups, *Israel J. Math.* **72** (1990), 232–251.

Institute of Mathematics
University of Tsukuba
Ibaraki 305, Japan