

ON THE MICROLOCAL STRUCTURE OF REGULAR
SIMPLE PREHOMOGENEOUS VECTOR SPACE
 $(\mathrm{GL}(1)^2 \times \mathrm{SL}(7), \Lambda_3 + \Lambda_1^*)$

By

Shin-ichi KASAI

Abstract. The purpose of this paper is to calculate the b -function of the regular simple prehomogeneous vector space $(\mathrm{GL}(1)^2 \times \mathrm{SL}(7), \Lambda_3 + \Lambda_1^*)$ by the aid of microlocal method using the holonomy diagram of relative invariants.

1. Main Results


In the present paper, we consider a special reducible prehomogeneous vector space, namely the regular simple prehomogeneous vector space $(\mathrm{GL}(1)^2 \times \mathrm{SL}(7), \Lambda_3 + \Lambda_1^*)$ ($\mathrm{A}(14)$ of §3 in [5]). Following the ideas of [7], we shall determine the holonomy diagram and compute the b -function. From the result of Kimura [5], there are twelve regular simple prehomogeneous vector spaces which have two algebraically independent relative invariants. The b -functions of four of them are reduced to the case of irreducible prehomogeneous vector spaces. In [1, 2], we have studied three of the remaining nontrivial eight cases. We use the same notations in [1, 4, 7].

By the microlocal calculus on analysis of prehomogeneous vector spaces, we obtain the holonomy diagram (Figure 1) and the following theorem.

THEOREM. *The b -function of $(\mathrm{GL}(1)^2 \times \mathrm{SL}(7), \Lambda_3 + \Lambda_1^*)$ is given as follows.*

$$b_\chi(s_1, s_2) = [s_1 + 1]_{n_1} [s_1 + 2]_{n_2} \left[s_1 + \frac{5}{2} \right]_{n_1} [s_1 + s_2 + 3]_{n_1+n_2}$$

Received April 28, 1999
Revised November 12, 1999

Figure 1. The holonomy diagram of $(GL(1)^2 \times SL(7), \Lambda_3 + \Lambda_1^*)$.

$$\begin{aligned}
& \left[s_1 + \frac{7}{2} \right]_{n_1} [s_1 + s_2 + 4]_{n_1+n_2} [s_1 + s_2 + 5]_{n_1+n_2} \\
& [s_2 + 1]_{n_2} \left[s_2 + \frac{7}{2} \right]_{n_2}
\end{aligned}$$

for $\chi = \chi_1^{n_1} \chi_2^{n_2}$ ($n_i > 0$) where $[\alpha]_k = \alpha(\alpha + 1) \cdots (\alpha + k - 1)$.

2. Preliminary results

In the following, we denote by G the group $GL(1)^2 \times SL(7)$ and by ρ the representation $\Lambda_3 + \Lambda_1^*$ of G . We define an element e_i of \mathbf{C}^7 by $e_i = {}^t(0, \dots, 0, \overset{i}{1}, 0, \dots, 0)$ for $0 \leq i \leq 7$. The representation space is identified with $V = \{\tilde{x} = (x, y); x = \sum_{1 \leq i < j < k \leq 7} x_{ijk} e_i \wedge e_j \wedge e_k \in \bigwedge^3 \mathbf{C}^7, y = \sum_{l=1}^7 y_l e_l \in \mathbf{C}^7\}$. Then the action ρ is given by $\rho(\tilde{g})\tilde{x} = (\alpha\rho_3(g)x, \beta^t g^{-1}y)$ for $\tilde{g} = (\alpha, \beta; g) \in G = GL(1)^2 \times SL(7)$ and $\tilde{x} = (x, y) \in V$ where $\rho_3(g)x = \sum x_{ijk}(ge_i) \wedge (ge_j) \wedge (ge_k)$.

We define $\partial/\partial e_l$ by $\partial/\partial e_l(e_i \wedge e_j \wedge e_k) = \delta_{il}e_j \wedge e_k$ for $j, k \neq l$. Let $\varphi(x) = (\varphi_{ij}(x))$ be the 7×7 symmetric matrix obtained by $\varphi_{ij}(x)e_1 \wedge \dots \wedge e_7 = x \wedge \partial x/\partial e_i \wedge \partial x/\partial e_j$ ($i, j = 1, \dots, 7$). Then $\varphi_{ij}(x)$ is a homogeneous polynomial of degree 3 and $\varphi(\rho_3(g)x) = g\varphi(x){}^t g$ for $g \in SL(7)$ and $x \in \bigwedge^3 \mathbf{C}^7$. We define $f_{jk}^i(x)$ by $f_{jk}^i e_1 \wedge \dots \wedge e_7 = x \wedge \partial x/\partial e_i \wedge e_j \wedge e_k$ ($i, j, k = 1, \dots, 7$). Let $\varphi^*(x) = (\varphi_{ij}^*(x))$ be the 7×7 symmetric matrix where $\varphi_{ij}^*(x) = \sum_{s,t=1}^7 f_{it}^s(x) f_{sj}^t(x)$. Then $\varphi_{ij}^*(x)$ is homogeneous polynomial of degree 4 and $\varphi^*(\rho_3(g)x) = {}^t g^{-1} \varphi^*(x) g^{-1}$ for $g \in SL(7)$ and $x \in \bigwedge^3 \mathbf{C}^7$ (see [3]).

- PROPOSITION 1([5]).** (1) *The triplet (G, ρ, V) is a regular P.V.*
 (2) *The algebraically independent relative invariants are given by $f_1(\tilde{x}) = \text{tr}\varphi(x)\varphi^*(x)$ and $f_2(\tilde{x}) = {}^t y\varphi(x)y$ for $\tilde{x} = (x, y) \in V$ (See [3]).*
 (3) *Let χ_i be the caharacter of f_i . Then their infinitesimal characters are given by $\delta\chi_1(\tilde{A}) = 7\alpha$ and $\delta\chi_2(\tilde{A}) = 3\alpha + 2\beta$ for $\tilde{A} = (\alpha, \beta; A) \in \mathfrak{gl}(1)^2 \oplus \mathfrak{sl}(7)$.*

Let Λ be the conormal bundle of an orbit S in V and Λ^* that of an orbit S^* in V^* . When $\Lambda = \Lambda^*$, we say that S and S^* are the dual orbits of each other. We identify the dual space V^* with V as usual. Since G is reductive, we have $(G, \rho, V) \cong (G, \rho^*, V^*)$ and hence (G, ρ, V) and (G, ρ^*, V^*) have the same number of G -orbits.

Put $x_0 = 234 + 567 + 1(25 + 36 + 47)$, $x_1 = 235 + 346 + 1(27 - 45)$, $x_4 = 134 + 256 + 127$, $x_7 = 234 + 1(25 + 36 + 47)$, $x_9 = 123 + 456$, $x_{10} = 126 - 135 + 234$, $x_{14} = 1(25 + 36 + 47)$, $x_{15} = 1(24 + 35)$, $x_{22} = 123$ and $x_{35} = 0$ where ijk stands for $e_i \wedge e_j \wedge e_k$.

PROPOSITION 2([6]). *The triplet (G, ρ, V) has the following thirty-eight orbits.*

Representative point	Codim	Dual orbit
(1) $\tilde{x}_1 = (x_0, e_1)$	0	\tilde{x}_{38}
(2) $\tilde{x}_2 = (x_0, e_2)$	1	\tilde{x}_{35}

Representative point	Codim	Dual orbit
(3) $\tilde{x}_3 = (x_1, e_1 + e_4)$	1	\tilde{x}_{36}
(4) $\tilde{x}_4 = (x_1, e_1)$	2	\tilde{x}_{31}
(5) $\tilde{x}_5 = (x_4, e_1 + e_2)$	4	\tilde{x}_{33}
(6) $\tilde{x}_6 = (x_1, e_5)$	5	\tilde{x}_{21}
(7) $\tilde{x}_7 = (x_4, e_1)$	5	\tilde{x}_{28}
(8) $\tilde{x}_8 = (x_1, e_6)$	6	\tilde{x}_{29}
(9) $\tilde{x}_9 = (x_4, e_3 + e_5)$	6	\tilde{x}_{19}
(10) $\tilde{x}_{10} = (x_0, 0)$	7	\tilde{x}_{37}
(11) $\tilde{x}_{11} = (x_7, e_1)$	7	\tilde{x}_{30}
(12) $\tilde{x}_{12} = (x_1, 0)$	8	\tilde{x}_{34}
(13) $\tilde{x}_{13} = (x_4, e_3)$	8	\tilde{x}_{20}
(14) $\tilde{x}_{14} = (x_7, e_2)$	8	\tilde{x}_{16}
(15) $\tilde{x}_{15} = (x_9, e_1 + e_4)$	9	\tilde{x}_{32}
(16) $\tilde{x}_{16} = (x_4, e_7)$	10	\tilde{x}_{14}
(17) $\tilde{x}_{17} = (x_{10}, e_1)$	10	\tilde{x}_{22}
(18) $\tilde{x}_{18} = (x_4, 0)$	11	\tilde{x}_{26}
(19) $\tilde{x}_{19} = (x_7, e_5)$	11	\tilde{x}_9
(20) $\tilde{x}_{20} = (x_9, e_1)$	12	\tilde{x}_{13}
(21) $\tilde{x}_{21} = (x_{10}, e_4)$	13	\tilde{x}_6
(22) $\tilde{x}_{22} = (x_7, 0)$	14	\tilde{x}_{17}
(23) $\tilde{x}_{23} = (x_{14}, e_1)$	14	\tilde{x}_{27}
(24) $\tilde{x}_{24} = (x_9, e_7)$	15	\tilde{x}_{25}
(25) $\tilde{x}_{25} = (x_{14}, e_2)$	15	\tilde{x}_{24}
(26) $\tilde{x}_{26} = (x_{15}, e_1)$	15	\tilde{x}_{18}
(27) $\tilde{x}_{27} = (x_9, 0)$	16	\tilde{x}_{23}
(28) $\tilde{x}_{28} = (x_{10}, e_7)$	16	\tilde{x}_7
(29) $\tilde{x}_{29} = (x_{15}, e_2)$	16	\tilde{x}_8
(30) $\tilde{x}_{30} = (x_{10}, 0)$	17	\tilde{x}_{11}
(31) $\tilde{x}_{31} = (x_{15}, e_6)$	20	\tilde{x}_4
(32) $\tilde{x}_{32} = (x_{14}, 0)$	21	\tilde{x}_{15}
(33) $\tilde{x}_{33} = (x_{15}, 0)$	22	\tilde{x}_5
(34) $\tilde{x}_{34} = (x_{22}, e_1)$	22	\tilde{x}_{12}
(35) $\tilde{x}_{35} = (x_{22}, e_4)$	25	\tilde{x}_2
(36) $\tilde{x}_{36} = (x_{22}, 0)$	29	\tilde{x}_3
(37) $\tilde{x}_{37} = (x_{35}, e_1)$	35	\tilde{x}_{10}
(38) $\tilde{x}_{38} = (x_{35}, 0)$	42	\tilde{x}_1

3. Holonomy diagram

For a point \tilde{x} of V , let $G_{\tilde{x}} = \{g \in G; \rho(g)\tilde{x} = \tilde{x}\}$ be the isotropy subgroup of G at \tilde{x} , and let $V_{\tilde{x}}^*$ be the conormal vector space. Then $G_{\tilde{x}}$ acts on $V_{\tilde{x}}^*$ by $\rho_{\tilde{x}} = \rho^*|_{G_{\tilde{x}}}$. If the triplet $(G_{\tilde{x}}, \rho_{\tilde{x}}, V_{\tilde{x}}^*)$ is a P.V., then we denote by y_0 its generic point, and if there is one one-codimensional orbit, then y_1 denotes a point of that orbit. When there exist several one-codimensional orbits, we denote representative points of their orbits by y_1, y'_1 , etc. Let $\mathfrak{G}_{\tilde{x}}$ (resp. \mathfrak{G}) be the Lie algebra of $G_{\tilde{x}}$ (resp. G), and $d\rho_{\tilde{x}}$ the infinitesimal representation of $\rho_{\tilde{x}}$. We denote by A_0 an element of $\mathfrak{G}_{\tilde{x}} = \{A \in \mathfrak{G}; d\rho(A)\tilde{x} = 0\}$ such that $d\rho^*(A_0)y_0 = y_0$. We denote by Λ_i the conormal bundle $T(\rho(G)\tilde{x}_i)^\perp$ of a orbit $\rho(G)\tilde{x}_i$.

(1) The case for \tilde{x}_1 . Since \tilde{x}_1 is a generic point of (G, ρ, V) , the isotropy subalgebra $\mathfrak{G}_{\tilde{x}_1}$ is isomorphic to $\mathfrak{sl}(3)$ (see $A(14)$ of §3 in [5]). Since $\Lambda_1 = V \times \{0\}$, we have $\text{ord}_{\Lambda_1} f^s = 0$.

(2) The case for \tilde{x}_2 . $V_{\tilde{x}_2}^* = \mathbf{C}\langle v_1 \rangle$ where $v_1 = (267, e_5)$. $y_0 = v_1 \in \rho^*(G)\tilde{x}_{35}$, $y_1 = 0 \in \rho^*(G)\tilde{x}_{38}$. $\text{ord}_{\Lambda_2} f^s = -s_2 - 1/2$.

(3) The case for \tilde{x}_3 . $V_{\tilde{x}_3}^* = \mathbf{C}\langle v_1 \rangle$ where $v_1 = (567, 0)$. $y_0 = v_1 \in \rho^*(G)\tilde{x}_{36}$, $y_1 = 0 \in \rho^*(G)\tilde{x}_{38}$. $\text{ord}_{\Lambda_3} f^s = -s_1 - 1/2$.

(4) The case for \tilde{x}_4 . $V_{\tilde{x}_4}^* = \mathbf{C}\langle v_1, v_2 \rangle$ where $v_1 = (567, 0)$ and $v_2 = (136, -e_4)$. $y_0 = v_1 + v_2 \in \rho^*(G)\tilde{x}_{31}$. $y_1 = v_1 \in \rho^*(G)\tilde{x}_{36}$, $y'_1 = v_2 \in \rho^*(G)\tilde{x}_{35}$. $\text{ord}_{\Lambda_4} f^s = -s_1 - s_2 - 1$.

(5) The case for \tilde{x}_5 . $V_{\tilde{x}_5}^* = \mathbf{C}\langle v_1, \dots, v_4 \rangle$ where $v_1 = (357, 0)$, $v_2 = (367, 0)$, $v_3 = (457, 0)$ and $v_4 = (467, 0)$. $(G_{\tilde{x}_5}, \rho_{\tilde{x}_5}, V_{\tilde{x}_5}^*) \cong (SL(2) \times GL(2), \Lambda_1 \otimes \Lambda_1, V(2) \otimes V(2))$. $y_0 = v_1 + v_4 \in \rho^*(G)\tilde{x}_{33}$. $y_1 = v_1 \in \rho^*(G)\tilde{x}_{36}$. $\text{ord}_{\Lambda_5} f^s = -2s_1 - 2$.

(6) The case for \tilde{x}_6 . $V_{\tilde{x}_6}^* = \mathbf{C}\langle v_1, \dots, v_5 \rangle$ where $v_1 = (567, 0)$, $v_2 = (-257 - 467, e_1)$, $v_3 = (157 + 367, e_2)$, $v_4 = (267 - 456, e_3)$ and $v_5 = (-167 + 356, e_4)$. $y_0 = v_3 + v_4 \in \rho^*(G)\tilde{x}_{21}$, $y_1 = v_3 + v_5 \in \rho^*(G)\tilde{x}_{31}$. $\text{ord}_{\Lambda_6} f^s = -2s_1 - 2s_2 - 7/2$.

(7) The case for \tilde{x}_7 . $V_{\tilde{x}_7}^* = \mathbf{C}\langle v_1, \dots, v_5 \rangle$ where $v_1 = (357, 0)$, $v_2 = (367, 0)$, $v_3 = (457, 0)$, $v_4 = (467, 0)$ and $v_5 = (156, e_2)$. $(G_{\tilde{x}_7}, \rho_{\tilde{x}_7}, V_{\tilde{x}_7}^*) \cong (GL(1) \times SO(4) \times GL(1), \Lambda_1 \otimes \Lambda_1 \otimes 1 + 1 \otimes 1 \otimes \Lambda_1, V(4) + V(1))$. $y_0 = v_1 + v_4 + v_5 \in \rho^*(G)\tilde{x}_{28}$. $y_1 = v_1 + v_4 \in \rho^*(G)\tilde{x}_{33}$, $y'_1 = v_1 + v_5 \in \rho^*(G)\tilde{x}_{31}$. $\text{ord}_{\Lambda_7} f^s = -2s_1 - s_2 - 5/2$.

(8) The case for \tilde{x}_8 . $V_{\tilde{x}_8}^* = \mathbf{C}\langle v_1, \dots, v_6 \rangle$ where $v_1 = (567, 0)$, $v_2 = (-256, e_3)$, $v_3 = (156, e_4)$, $v_4 = (-267, e_1)$, $v_5 = (167, e_2)$ and $v_6 = (126, e_7)$. $y_0 = v_1 + v_6 \in \rho^*(G)\tilde{x}_{29}$. $y_1 = v_6 \in \rho^*(G)\tilde{x}_{35}$, $y'_1 = v_3 + v_4 \in \rho^*(G)\tilde{x}_{29}$. $\text{ord}_{\Lambda_8} f^s = -s_1 - 2s_2 - 3$.

(9) The case for \tilde{x}_9 . $V_{\tilde{x}_9}^* = \mathbf{C}\langle v_1, \dots, v_6 \rangle$ where $v_1 = (357, 0)$, $v_2 = (367, 0)$, $v_3 = (457, 0)$, $v_4 = (467, 0)$, $v_5 = (-237 + 345, e_1)$ and $v_6 = (157 + 356, e_2)$. $y_0 = v_4 + v_5 + v_6 \in \rho^*(G)\tilde{x}_{19}$. $y_1 = v_5 + v_6 \in \rho^*(G)\tilde{x}_{21}$, $y'_1 = v_4 + v_5 \in \rho^*(G)\tilde{x}_{28}$ and $y''_1 = v_4 + v_6 \in \rho^*(G)\tilde{x}_{28}$. $ord_{\Lambda_9} f^s = -3s_1 - 2s_2 - 5$.

(10) The case for \tilde{x}_{10} . $V_{\tilde{x}_{10}}^* = \mathbf{C}\langle (0, e_i); 1 \leq i \leq 7 \rangle$. $(G_{\tilde{x}_{10}}, \rho_{\tilde{x}_{10}}, V_{\tilde{x}_{10}}^*) \cong (GL(1) \times G_2, \Lambda_1 \otimes \Lambda_2, V(7))$. $y_0 = (0, e_1) \in \rho^*(G)\tilde{x}_{37}$, $y_1 = (0, e_2) \in \rho^*(G)\tilde{x}_{37}$. $ord_{\Lambda_{10}} f^s = -2s_2 - 7/2$.

(11) The case for \tilde{x}_{11} . $V_{\tilde{x}_{11}}^* = \mathbf{C}\langle v_1, \dots, v_7 \rangle$ where $v_1 = (567, 0)$, $v_2 = (267, 0)$, $v_3 = (456, 0)$, $v_4 = (-357, 0)$, $v_5 = (257 - 367, 0)$, $v_6 = (457 - 356, 0)$ and $v_7 = (256 + 467, 0)$. $y_0 = v_2 + v_3 - v_4 \in \rho^*(G)\tilde{x}_{30}$, $y_1 = v_2 + v_3 \in \rho^*(G)\tilde{x}_{33}$. $ord_{\Lambda_{11}} f^s = -3s_1 - 4$.

(12) The case for \tilde{x}_{12} . $V_{\tilde{x}_{12}}^* = \mathbf{C}\langle v_1, \dots, v_8 \rangle$ where $v_1 = (567, 0)$ and $v_{1+i} = (0, e_i)$, $1 \leq i \leq 7$. $y_0 = v_1 + v_6 \in \rho^*(G)\tilde{x}_{34}$. $y_1 = v_6 \in \rho^*(G)\tilde{x}_{31}$, $y'_1 = v_1 + v_7 \in \rho^*(G)\tilde{x}_{37}$. $ord_{\Lambda_{12}} f^s = -s_1 - 3s_2 - 6$.

(13) The case for \tilde{x}_{13} . $V_{\tilde{x}_{13}}^* = \mathbf{C}\langle v_1, \dots, v_8 \rangle$ where $v_1 = (357, 0)$, $v_2 = (367, 0)$, $v_3 = (457, 0)$, $v_4 = (467, 0)$, $v_5 = (-237, e_1)$, $v_6 = (356, e_2)$, $v_7 = (236, e_5)$ and $v_8 = (-235, e_6)$. $y_0 = v_3 + v_7 \in \rho^*(G)\tilde{x}_{20}$. $y_1 = v_1 + v_4 + v_7 \in \rho^*(G)\tilde{x}_{21}$. $A_0 = (\alpha, \alpha; diag(-3\alpha - 2, 2, \alpha, \alpha + 2, -\alpha/2 - 1, -\alpha/2 - 1, 2\alpha)) \in \mathfrak{G}_{\tilde{x}_{13}}$. Then $\delta\chi_1(A_0) = 7\alpha$ and $\delta\chi_2(A_0) = 5\alpha$.

(14) The case for \tilde{x}_{14} . $V_{\tilde{x}_{14}}^* = \mathbf{C}\langle v_1, \dots, v_8 \rangle$ where $v_1 = (567, 0)$, $v_2 = (267, 0)$, $v_3 = (256 + 467, 0)$, $v_4 = (257 - 367, 0)$, $v_5 = (456, 0)$, $v_6 = (457 - 356, 0)$, $v_7 = (357, 0)$ and $v_8 = (-167 + 236 + 247, 2e_1)$. $y_0 = v_6 + v_8 \in \rho^*(G)\tilde{x}_{16}$. $y_1 = v_7 + v_8 \in \rho^*(G)\tilde{x}_{19}$, $y'_1 = v_2 + v_6 \in \rho^*(G)\tilde{x}_{30}$. $ord_{\Lambda_{14}} f^s = -4s_1 - 2s_2 - 7$.

(15) The case for \tilde{x}_{15} . $V_{\tilde{x}_{15}}^* = \mathbf{C}\langle v_1, \dots, v_9 \rangle$ where $v_1 = (147, 0)$, $v_2 = (-347, 0)$, $v_3 = (247, 0)$, $v_4 = (-167, 0)$, $v_5 = (157, 0)$, $v_6 = (257, 0)$, $v_7 = (267, 0)$, $v_8 = (357, 0)$ and $v_9 = (367, 0)$. $y_0 = v_1 + v_6 + v_9 \in \rho^*(G)\tilde{x}_{32}$. $y_1 = v_6 + v_9 \in \rho^*(G)\tilde{x}_{33}$. $ord_{\Lambda_{15}} f^s = -3s_1 - s_2 - 9/2$.

(16) The case for \tilde{x}_{16} . $V_{\tilde{x}_{16}}^* = \mathbf{C}\langle v_1, \dots, v_{10} \rangle$ where $v_1 = (357, 0)$, $v_2 = (367, 0)$, $v_3 = (457, 0)$, $v_4 = (467, 0)$, $v_5 = (347, e_1)$, $v_6 = (567, e_2)$, $v_7 = (-147 + 456, e_3)$, $v_8 = (137 - 356, e_4)$, $v_9 = (-267 - 346, e_5)$ and $v_{10} = (257 + 345, e_6)$. $y_0 = v_4 + v_8 + v_{10} \in \rho^*(G)\tilde{x}_{14}$. $y_1 = v_8 + v_{10} \in \rho^*(G)\tilde{x}_{21}$. $ord_{\Lambda_{16}} f^s = -3s_1 - 3s_2 - 7$.

(17) The case for \tilde{x}_{17} . $V_{\tilde{x}_{17}}^* = \mathbf{C}\langle v_1, \dots, v_{10} \rangle$ where $v_1 = (-467, 0)$, $v_2 = (457, 0)$, $v_3 = (567, 0)$, $v_4 = (147, 0)$, $v_5 = (-167 - 347, 0)$, $v_6 = (157 + 247, 0)$, $v_7 = (-267 - 357, 0)$, $v_8 = (367, 0)$, $v_9 = (257, 0)$ and $v_{10} = (456, 0)$. $y_0 = v_4 - v_7 +$

$v_{10} \in \rho^*(G)\tilde{x}_{22}$. $y_1 = -v_7 + v_{10} \in \rho^*(G)\tilde{x}_{30}$, $y'_1 = v_4 + v_8 + v_9 \in \rho^*(G)\tilde{x}_{32}$. $\text{ord}_{\Lambda_{17}} f^s = -4s_1 - s_2 - 13/2$.

(18) The case for \tilde{x}_{18} . $V_{\tilde{x}_{18}}^* = \mathbf{C}\langle v_1, \dots, v_{11} \rangle$ where $v_1 = (357, 0)$, $v_2 = (367, 0)$, $v_3 = (457, 0)$, $v_4 = (467, 0)$ and $v_{4+i} = (0, e_i)$, $1 \leq i \leq 7$. $y_0 = v_1 + v_4 + v_{11} \in \rho^*(G)\tilde{x}_{26}$. $y_1 = v_1 + v_4 + v_7 + v_9 \in \rho^*(G)\tilde{x}_{29}$, $y'_1 = v_1 + v_{11} \in \rho^*(G)\tilde{x}_{34}$. $\text{ord}_{\Lambda_{18}} f^s = -2s_1 - 4s_2 - 19/2$.

(19) The case for \tilde{x}_{19} . $V_{\tilde{x}_{19}}^* = \mathbf{C}\langle v_1, \dots, v_{11} \rangle$ where $v_1 = (567, 0)$, $v_2 = (456, 0)$, $v_3 = (356 - 457, 0)$, $v_4 = (357, 0)$, $v_5 = (256 + 467, 0)$, $v_6 = (257 - 367, 0)$, $v_7 = (267, 0)$, $v_8 = (-457, e_1)$, $v_9 = (345, e_2)$, $v_{10} = 1/2(156 - 245 + 346, 2e_3)$ and $v_{11} = 1/2(157 + 235 + 347, 2e_4)$. $y_0 = v_4 + v_7 + v_{10} \in \rho^*(G)\tilde{x}_9$. $y_1 = v_4 + v_{10} \in \rho^*(G)\tilde{x}_{14}$, $y'_1 = v_7 + v_{10} \in \rho^*(G)\tilde{x}_{19}$. $\text{ord}_{\Lambda_{19}} f^s = -4s_1 - 3s_2 - 17/2$.

(20) The case for \tilde{x}_{20} . $V_{\tilde{x}_{20}}^* = \mathbf{C}\langle v_1, \dots, v_{12} \rangle$ where $v_1 = (147, 0)$, $v_2 = (157, 0)$, $v_3 = (167, 0)$, $v_4 = (247, 0)$, $v_5 = (257, 0)$, $v_6 = (267, 0)$, $v_7 = (347, 0)$, $v_8 = (357, 0)$, $v_9 = (367, 0)$, $v_{10} = (156, e_4)$, $v_{11} = (-146, e_5)$ and $v_{12} = (145, e_6)$. $y_0 = v_4 + v_8 + v_{10} \in \rho^*(G)\tilde{x}_{13}$. There is no one-codimensional orbit. $A_0 \in \mathfrak{G}_{\tilde{x}_{20}}$ with $\beta = -5\alpha - 4$, $A = \text{diag}(-5\alpha - 4, 2(\alpha + 1), 2(\alpha + 1), -5\alpha - 3, -5\alpha - 3, 9\alpha + 6, 2\alpha)$. $\delta\chi_1(A_0) = 7\alpha$, $\delta\chi_2(A_0) = -7\alpha - 8$.

(21) The case for \tilde{x}_{21} . $V_{\tilde{x}_{21}}^* = \mathbf{C}\langle v_1, \dots, v_{13} \rangle$ where $v_1 = (467, 0)$, $v_2 = (-457, 0)$, $v_3 = (567, 0)$, $v_4 = (456, 0)$, $v_5 = (147, 0)$, $v_6 = (-167 - 347, 0)$, $v_7 = (157 + 247, 0)$, $v_8 = (367, 0)$, $v_9 = (-267 - 357, 0)$, $v_{10} = (257, 0)$, $v_{11} = (146, e_2)$, $v_{12} = (-145, e_3)$ and $v_{13} = 1/2(-156 - 246 + 345, 2e_1)$. $y_0 = v_5 + v_9 + v_{13} \in \rho^*(G)\tilde{x}_6$. $y_1 = v_5 + v_{10} + v_{13} \in \rho^*(G)\tilde{x}_9$, $y'_1 = v_9 + v_{13} \in \rho^*(G)\tilde{x}_{16}$, and $y''_1 = v_9 + v_{10} + v_{11} \in \rho^*(G)\tilde{x}_{13}$. $\text{ord}_{\Lambda_{21}} f^s = -5s_1 - 3s_2 - 21/2$.

(22) The case for \tilde{x}_{22} . $V_{\tilde{x}_{22}}^* = \mathbf{C}\langle v_1, \dots, v_{14} \rangle$ where $v_1 = (567, 0)$, $v_2 = (267, 0)$, $v_3 = (456, 0)$, $v_4 = (-357, 0)$, $v_5 = (257 - 367, 0)$, $v_6 = (457 - 356, 0)$, $v_7 = (256 + 467, 0)$ and $v_{7+i} = (0, e_i)$, $1 \leq i \leq 7$. $y_0 = v_2 + v_3 - v_4 + v_{12} \in \rho^*(G)\tilde{x}_{17}$. $y_1 = v_2 + v_3 + v_{13} \in \rho^*(G)\tilde{x}_{26}$, $y'_1 = v_2 + v_3 - v_4 + v_{12} + \sqrt{-1}v_{13} \in \rho^*(G)\tilde{x}_{17}$. $\text{ord}_{\Lambda_{22}} f^s = -3s_1 - 4s_2 - 10$.

(23) The case for \tilde{x}_{23} . $V_{\tilde{x}_{23}}^* = \mathbf{C}\langle v_1, \dots, v_{14} \rangle$ where $v_1 = (234, 0)$, $v_2 = (567, 0)$, $v_3 = (345, 0)$, $v_4 = (267, 0)$, $v_5 = (246, 0)$, $v_6 = (357, 0)$, $v_7 = (237, 0)$, $v_8 = (456, 0)$, $v_9 = (236 - 247, 0)$, $v_{10} = (356 - 457, 0)$, $v_{11} = (235 + 347, 0)$, $v_{12} = (256 + 467, 0)$, $v_{13} = (245 - 346, 0)$ and $v_{14} = (257 - 367, 0)$. $(G_{\tilde{x}_{23}}, \rho_{\tilde{x}_{23}}, V_{\tilde{x}_{23}}^*) \cong (GL(1) \times Sp(3), \Lambda_1 \otimes \Lambda_3, V(14))$. $y_0 = v_1 + v_2 \in \rho^*(G)\tilde{x}_{27}$, $y_1 = v_7 + v_{13} \in \rho^*(G)\tilde{x}_{30}$. $\text{ord}_{\Lambda_{23}} f^s = -4s_1 - 7$.

(24) The case for \tilde{x}_{24} . $V_{\tilde{x}_{24}}^* = \mathbf{C}\langle v_1, \dots, v_{15} \rangle$ where $v_1 = (147, 0)$, $v_2 = (157, 0)$, $v_3 = (167, 0)$, $v_4 = (247, 0)$, $v_5 = (257, 0)$, $v_6 = (267, 0)$, $v_7 = (347, 0)$, $v_8 = (357, 0)$, $v_9 = (367, 0)$, $v_{10} = (237, e_1)$, $v_{11} = (-137, e_2)$, $v_{12} = (127, e_3)$, $v_{13} = (567, e_4)$, $v_{14} = (-467, e_5)$ and $v_{15} = (457, e_6)$. $(G_{\tilde{x}_{24}}, \rho_{\tilde{x}_{24}}, V_{\tilde{x}_{24}}^* \cong (GL(1) \times SL(3) \times SL(3), \Lambda_1 \otimes \Lambda_1^* \otimes \Lambda_1^* + \Lambda_1 \otimes \Lambda_1 \otimes 1 + \Lambda_1 \otimes 1 \otimes \Lambda_1, V(9) + V(3) + V(3))$. Thus the triplet $(G_{\tilde{x}_{24}}, \rho_{\tilde{x}_{24}}, V_{\tilde{x}_{24}}^*)$ is a non P.V.

(25) The case for \tilde{x}_{25} . $V_{\tilde{x}_{25}}^* = \mathbf{C}\langle v_1, \dots, v_{15} \rangle$ where $v_1 = (236 - 247, 0)$, $v_2 = (234, 0)$, $v_3 = (237, 0)$, $v_4 = (246, 0)$, $v_5 = (267, 0)$, $v_6 = (235 + 347, 0)$, $v_7 = (245 - 346, 0)$, $v_8 = (256 + 467, 0)$, $v_9 = (257 - 367, 0)$, $v_{10} = (356 - 457, 0)$, $v_{11} = (-345, 0)$, $v_{12} = (357, 0)$, $v_{13} = (456, 0)$, $v_{14} = (-567, 0)$ and $v_{15} = (247, e_1)$. The triplet $(G_{\tilde{x}_{25}}, \rho_{\tilde{x}_{25}}, V_{\tilde{x}_{25}}^*)$ is a non P.V.

(26) The case for \tilde{x}_{26} . $V_{\tilde{x}_{26}}^* = \mathbf{C}\langle v_1, \dots, v_{15} \rangle$ where $v_1 = (267, 0)$, $v_2 = (367, 0)$, $v_3 = (467, 0)$, $v_4 = (567, 0)$, $v_5 = (167, 0)$, $v_6 = (246 - 356, 0)$, $v_7 = (236, 0)$, $v_8 = (256, 0)$, $v_9 = (346, 0)$, $v_{10} = (456, 0)$, $v_{11} = (247 - 357, 0)$, $v_{12} = (237, 0)$, $v_{13} = (257, 0)$, $v_{14} = (347, 0)$ and $v_{15} = (457, 0)$. $y_0 = v_5 + v_9 + v_{13} \in \rho^*(G)\tilde{x}_{18}$, $y_1 = v_5 + v_6 + v_{12} \in \rho^*(G)\tilde{x}_{22}$, $y'_1 = v_9 + v_{13} \in \rho^*(G)\tilde{x}_{27}$. $ord_{\Lambda_{26}} f^s = -5s_1 - s_2 - 19/2$.

(27) The case for \tilde{x}_{27} . $V_{\tilde{x}_{27}}^* = \mathbf{C}\langle v_1, \dots, v_{16} \rangle$ where $\{v_1, \dots, v_9\} = \{(ij7, 0); 1 \leq i \leq 3, 4 \leq j \leq 6\}$ and $v_{9+i} = (0, e_i)$, $1 \leq i \leq 7$. $y_0 = (147 + 257 + 367, e_7) \in \rho^*(G)\tilde{x}_{23}$. $y_1 = (147 + 257, e_7) \in \rho^*(G)\tilde{x}_{26}$, $y'_1 = (147 + 257 + 367, e_1 + e_4) \in \rho^*(G)\tilde{x}_{25}$. $ord_{\Lambda_{27}} f^s = -3s_1 - 5s_2 - 14$.

(28) The case for \tilde{x}_{28} . $V_{\tilde{x}_{28}}^* = \mathbf{C}\langle v_1, \dots, v_{16} \rangle$ where $v_1 = (567, 0)$, $v_2 = (-467, 0)$, $v_3 = (457, 0)$, $v_4 = (456, 0)$, $v_5 = (147, 0)$, $v_6 = (257, 0)$, $v_7 = (367, 0)$, $v_8 = (267 + 357, 0)$, $v_9 = (167 + 347, 0)$, $v_{10} = (157 + 247, 0)$, $v_{11} = \frac{1}{2}(267 - 357, 2e_1)$, $v_{12} = 1/2(-167 + 247, 2e_2)$, $v_{13} = \frac{1}{2}(157 - 247, 2e_3)$, $v_{14} = (237, e_4)$, $v_{15} = (-137, e_5)$ and $v_{16} = (127, e_6)$. $y_0 = v_4 + v_5 + v_{14} \in \rho^*(G)\tilde{x}_7$, $y_1 = v_4 + v_{14} \in \rho^*(G)\tilde{x}_9$, $y'_1 = v_1 + v_5 + v_{14} \in \rho^*(G)\tilde{x}_{25}$. $ord_{\Lambda_{28}} f^s = -5s_1 - 4s_2 - 13$.

(29) The case for \tilde{x}_{29} . $V_{\tilde{x}_{29}}^* = \mathbf{C}\langle v_1, \dots, v_{16} \rangle$ where $v_1 = (267, 0)$, $v_2 = (236, 0)$, $v_3 = (237, 0)$, $v_4 = (256, 0)$, $v_5 = (257, 0)$, $v_6 = (567, 0)$, $v_7 = (-367, 0)$, $v_8 = (467, 0)$, $v_9 = (167, 0)$, $v_{10} = (-247 + 357, 0)$, $v_{11} = (246 - 356, 0)$, $v_{12} = (456, 0)$, $v_{13} = (457, 0)$, $v_{14} = (346, 0)$, $v_{15} = (347, 0)$ and $v_{16} = (235, e_1)$. $y_0 = v_9 + v_{12} + v_{15} + v_{16} \in \rho^*(G)\tilde{x}_8$, $y_1 = v_3 + v_9 + v_{12} + v_{15} \in \rho^*(G)\tilde{x}_{18}$, $y'_1 = v_9 + v_{10} + v_{14} + v_{16} \in \rho^*(G)\tilde{x}_{18}$. $ord_{\Lambda_{29}} f^s = -6s_1 - 3s_2 - 27/2$.

(30) The case for \tilde{x}_{30} . $V_{\tilde{x}_{30}}^* = \mathbf{C}\langle v_1, \dots, v_{17} \rangle$ where $v_1 = (567, 0)$, $v_2 = (-467, 0)$, $v_3 = (457, 0)$, $v_4 = (147, 0)$, $v_5 = (257, 0)$, $v_6 = (367, 0)$, $v_7 = (267 + 357, 0)$, $v_8 = (167 + 347, 0)$, $v_9 = (157 + 247, 0)$, $v_{10} = (456, 0)$ and $v_{10+i} =$

$(0, e_i), \quad 1 \leq i \leq 7.$ $y_0 = v_4 + v_8 + v_{10} + v_{17} \in \rho^*(G)\tilde{x}_{11}, \quad y_1 = v_4 + v_8 + v_{10} + v_{14} \in \rho^*(G)\tilde{x}_{14}, \quad y'_1 = v_8 + v_{10} + v_{17} \in \rho^*(G)\tilde{x}_{17}, \quad \text{and} \quad y''_1 = v_4 + v_8 + v_{17} \in \rho^*(G)\tilde{x}_{23}.$ $\text{ord}_{\Lambda_{30}} f^s = -4s_1 - 5s_2 - 29/2.$

(31) The case for \tilde{x}_{31} . $V_{\tilde{x}_{31}}^* = \mathbf{C}\langle v_1, \dots, v_{20} \rangle$ where $v_1 = (-467, 0), v_2 = (-567, 0), v_3 = (267, 0), v_4 = (367, 0), v_5 = (167, 0), v_6 = (246 - 356, 0), v_7 = (236, 0), v_8 = (256, 0), v_9 = (346, 0), v_{10} = (456, 0), v_{11} = (247 - 357, 0), v_{12} = (237, 0), v_{13} = (257, 0), v_{14} = (347, 0), v_{15} = (457, 0), v_{16} = (356, e_1), v_{17} = (-146, e_2), v_{18} = (-156, e_3), v_{19} = (126, e_4)$ and $v_{20} = (136, e_5).$ $y_0 = v_8 + v_9 + v_{12} + v_{15} + v_{17} \in \rho^*(G)\tilde{x}_4, y_1 = v_8 + v_9 + v_{11} + v_{19} + v_{20} \in \rho^*(G)\tilde{x}_{14}, y'_1 = v_7 + v_8 + v_{14} + v_{15} + v_{20} \in \rho^*(G)\tilde{x}_7,$ and $y''_1 = v_8 + v_{12} + v_{15} + v_{17} \in \rho^*(G)\tilde{x}_4.$ $\text{ord}_{\Lambda_{31}} f^s = -6s_1 - 4s_2 - 15.$

(32) The case for \tilde{x}_{32} . $V_{\tilde{x}_{32}}^* = \mathbf{C}\langle v_1, \dots, v_{21} \rangle$ where $v_1 = (234, 0), v_2 = (567, 0), v_3 = (345, 0), v_4 = (267, 0), v_5 = (246, 0), v_6 = (357, 0), v_7 = (237, 0), v_8 = (456, 0), v_9 = (236 - 247, 0), v_{10} = (356 - 457, 0), v_{11} = (235 + 347, 0), v_{12} = (256 + 467, 0), v_{13} = (245 - 346, 0), v_{14} = (257 - 367, 0)$ and $v_{14+i} = (0, e_i), 1 \leq i \leq 7.$ $y_0 = v_1 + v_2 + v_{16} + v_{19} \in \rho^*(G)\tilde{x}_{15}, y_1 = v_7 + v_{13} + v_{18} \in \rho^*(G)\tilde{x}_{17}, y'_1 = v_1 + v_2 + v_{16} + v_{20} \in \rho^*(G)\tilde{x}_{15}.$ $\text{ord}_{\Lambda_{32}} f^s = -4s_1 - 4s_2 - 23/2.$

(33) The case for \tilde{x}_{33} . $V_{\tilde{x}_{33}}^* = \mathbf{C}\langle v_1, \dots, v_{22} \rangle$ where $v_1 = (267, 0), v_2 = (367, 0), v_3 = (467, 0), v_4 = (567, 0), v_5 = (167, 0), v_6 = (246 - 356, 0), v_7 = (236, 0), v_8 = (256, 0), v_9 = (346, 0), v_{10} = (456, 0), v_{11} = (247 - 357, 0), v_{12} = (237, 0), v_{13} = (257, 0), v_{14} = (347, 0), v_{15} = (457, 0)$ and $v_{15+i} = (0, e_i), 1 \leq i \leq 7.$ $y_0 = v_5 + v_9 + v_{13} + v_{21} + v_{22} \in \rho^*(G)\tilde{x}_5.$ $y_1 = v_5 + v_9 + v_{13} + v_{21} \in \rho^*(G)\tilde{x}_7, y'_1 = v_5 + v_6 + v_{12} + v_{21} \in \rho^*(G)\tilde{x}_{11}$ and $y''_1 = v_9 + v_{13} + v_{21} + v_{22} \in \rho^*(G)\tilde{x}_{15}.$ $\text{ord}_{\Lambda_{33}} f^s = -5s_1 - 5s_2 - 16.$

(34) The case for \tilde{x}_{34} . $V_{\tilde{x}_{34}}^* = \mathbf{C}\langle v_1, \dots, v_{22} \rangle$ where $v_1 = (567, 0), v_2 = (-467, 0), v_3 = (457, 0), v_4 = (-456, 0), v_5 = (145, 0), v_6 = (146, 0), v_7 = (147, 0), v_8 = (167, 0), v_9 = (-157, 0), v_{10} = (156, 0), v_{11} = (245, 0), v_{12} = (246, 0), v_{13} = (247, 0), v_{14} = (267, 0), v_{15} = (-257, 0), v_{16} = (256, 0), v_{17} = (345, 0), v_{18} = (346, 0), v_{19} = (347, 0), v_{20} = (367, 0), v_{21} = (-357, 0)$ and $v_{22} = (356, 0).$ $y_0 = v_5 + v_8 + v_{12} - v_{21} \in \rho^*(G)\tilde{x}_{12}.$ $y_1 = v_8 + v_{11} + v_{12} - v_{21} \in \rho^*(G)\tilde{x}_{18}.$ $\text{ord}_{\Lambda_{34}} f^s = -6s_1 - 2s_2 - 13.$

(35) The case for \tilde{x}_{35} . $V_{\tilde{x}_{35}}^* = \mathbf{C}\langle v_1, \dots, v_{25} \rangle$ where $v_1 = (467, 0), v_2 = (-457, 0), v_3 = (456, 0), v_4 = (567, 0), v_5 = (145, 0), v_6 = (146, 0), v_7 = (147, 0), v_8 = (245, 0), v_9 = (246, 0), v_{10} = (247, 0), v_{11} = (345, 0), v_{12} = (346, 0), v_{13} = (347, 0), v_{14} = (167, 0), v_{15} = (-157, 0), v_{16} = (156, 0), v_{17} = (267, 0), v_{18} =$

$(-257, 0)$, $v_{19} = (256, 0)$, $v_{20} = (367, 0)$, $v_{21} = (-357, 0)$, $v_{22} = (356, 0)$, $v_{23} = (234, e_1)$, $v_{24} = (-134, e_2)$ and $v_{25} = (124, e_3)$. $y_0 = v_5 + v_{14} + v_{18} + v_{22} + v_{23} \in \rho^*(G)\tilde{x}_2$. $y_1 = v_5 + v_{14} + v_{18} + v_{23} \in \rho^*(G)\tilde{x}_4$, $y'_1 = v_{14} + v_{18} + v_{22} + v_{23} \in \rho^*(G)\tilde{x}_8$. $\text{ord}_{\Lambda_{35}} f^s = -7s_1 - 4s_2 - 18$.

(36) The case for \tilde{x}_{36} . $V_{\tilde{x}_{36}}^* = \mathbf{C}\langle v_1, \dots, v_{29} \rangle$ where $v_1 = (567, 0)$, $v_2 = (-467, 0)$, $v_3 = (457, 0)$, $v_4 = (-456, 0)$, $v_5 = (145, 0)$, $v_6 = (146, 0)$, $v_7 = (147, 0)$, $v_8 = (167, 0)$, $v_9 = (-157, 0)$, $v_{10} = (156, 0)$, $v_{11} = (245, 0)$, $v_{12} = (246, 0)$, $v_{13} = (247, 0)$, $v_{14} = (267, 0)$, $v_{15} = (-257, 0)$, $v_{16} = (256, 0)$, $v_{17} = (345, 0)$, $v_{18} = (346, 0)$, $v_{19} = (347, 0)$, $v_{20} = (367, 0)$, $v_{21} = (-357, 0)$, $v_{22} = (356, 0)$ and $v_{22+i} = (0, e_i)$, $1 \leq i \leq 7$. $y_0 = v_5 + v_8 + v_{12} - v_{21} + v_{27} + v_{28} \in \rho^*(G)\tilde{x}_3$. $y_1 = v_5 + v_8 + v_{12} - v_{21} + v_{27} \in \rho^*(G)\tilde{x}_4$, $y'_1 = v_8 + v_{11} + v_{12} - v_{21} + v_{28} + v_{29} \in \rho^*(G)\tilde{x}_5$. $\text{ord}_{\Lambda_{36}} f^s = -6s_1 - 5s_2 - 18$.

(37) The case for \tilde{x}_{37} . $V_{\tilde{x}_{37}}^* = \mathbf{C}\langle (ijk, 0); 1 \leq i < j < k \leq 7 \rangle$. $y_0 = (125 + 136 + 147 + 234 + 567, 0) \in \rho^*(G)\tilde{x}_{10}$. $y_1 = (125 + 136 + 234 + 567, 0) \in \rho^*(G)\tilde{x}_{12}$, $y'_1 = (145 + 167 + 234 + 256 + 357, 0) \in \rho^*(G)\tilde{x}_{12}$. $\text{ord}_{\Lambda_{37}} f^s = -7s_1 - 3s_2 - 35/2$.

(38) The case for \tilde{x}_{38} . $(G_{\tilde{x}_{38}}, \rho_{\tilde{x}_{38}}, V_{\tilde{x}_{38}}^*) \cong (G, \rho, V)$. $y_0 = \tilde{x}_1$, $y_1 = \tilde{x}_2$, $y'_1 = \tilde{x}_3$. $\text{ord}_{\Lambda_{38}} f^s = -7s_1 - 5s_2 - 21$.

Hence, we obtain the holonomy diagram (Figure 1) where we denote by



the conormal bundle of the orbit $\rho(G)\tilde{x}_i$ which is j -codimensional.

Acknowledgement

The author would like to express his hearty thanks to Professor Tatsuo Kimura for his invaluable advice and encouragement.

References

- [1] Kasai, S., The b -function and the holonomy diagram of a regular simple prehomogeneous vector space $(GL(1)^2 \times Spin(10)$, half-spin rep. + vector rep.), to appear in J. Algebra.
- [2] Kasai, S., The b -functions of regular simple prehomogeneous vector spaces $(GL(1)^3 \times SL(2m)$, $\Lambda_2 + (\Lambda_1 + \Lambda_1)^{(*)})$
- [3] Kimura, T., Remark on some combinatorial construction of relative invariants, Tsukuba J. Math. **5** (1981), 101–115.
- [4] Kimura, T., The b -functions and holonomy diagrams of irreducible regular prehomogeneous vector spaces, Nagoya Math. J. **85** (1982), 1–80.

- [5] Kimura, T., A classification of prehomogeneous vector spaces of simple algebraic groups with scalar multiplications, *J. Algebra* **83** (1983), 72–100.
- [6] Kimura, T. and Kasai, S., The orbital decomposition of some prehomogeneous vector spaces, *Adv. Stud. Pure Math.* **6** (1985), 437–480.
- [7] Sato, M., Kashiwara, M., Kimura, T. and Oshima, T., Micro-local analysis of prehomogeneous vector spaces, *Inv. Math.* **62** (1980), 117–179.
- [8] Sato, M. and Kimura, T., A classification of irreducible prehomogeneous vector spaces and their relative invariants, *Nagoya Math. J.* **65** (1977), 1–155.

Department of Mathematics,
Faculty of Education,
Yamaguchi University,
Yamaguchi, 753-8513, Japan
e-mail address: kasai@edu.yamaguchi-u.ac.jp