

FIXED POINTS SUBGROUPS BY TWO INVOLUTIVE AUTOMORPHISMS σ, γ OF COMPACT EXCEPTIONAL LIE GROUPS F_4, E_6 AND E_7

By

Toshikazu MIYASHITA

Introduction

For simply connected compact exceptional Lie groups $G = F_4, E_6$ and E_7 , we consider two involutions σ, γ and determine the group structure of subgroups $G^{\sigma, \gamma}$ of G which are the intersection $G^\sigma \cap G^\gamma$ of the fixed points subgroups of G^σ and G^γ . The motivation is as follows. In [1], we determine the group structure of $(F_4)^{\sigma, \sigma'}$, $(E_6)^{\sigma, \sigma'}$ and $(E_7)^{\sigma, \sigma'}$, and in [2], we also determine the group structure of $(G_2)^{\gamma, \gamma'}$, $(F_4)^{\gamma, \gamma'}$ and $(E_6)^{\gamma, \gamma'}$. So, in this paper, we try to determine the type of groups $(F_4)^{\sigma, \gamma}$, $(E_6)^{\sigma, \gamma}$ and $(E_7)^{\sigma, \gamma}$. Our results are the following second columns. The first columns are already known in [3], [4] or [5] and these play an important role to obtain our results. In Table 1, the results of the group structure of $G^{\sigma, \gamma}$ are obtained by the result of G^γ and in Table 2, ones are obtained by the result of G^σ . In this paper, we show the proof of the results of the first and the second line of Table 1 and the third line of Table 2.

Acknowledgment

The author is grateful to Professor Ichiro Yokota for his valuable comments. As for the group $(E_8)^{\sigma, \gamma}$, we can not realize explicitly, however we conjecture

$$(E_8)^{\sigma, \gamma} \cong (Spin(4) \times Spin(12)) / (\mathbf{Z}_2 \times \mathbf{Z}_2).$$

REMARK. In E_7 , since γ is conjugate to $-\sigma$, we have $(E_7)^\gamma \cong (E_7)^\sigma$. (In detail, see [4].) Note that the results of Table 1 and Table 2 are the same as a set, however they are different as realizations.

Table 1

G	G^γ	$G^{\sigma,\gamma}$
F_4	$(Sp(1) \times Sp(3))/\mathbf{Z}_2$	$(Sp(1) \times Sp(1) \times Sp(2))/\mathbf{Z}_2$
E_6	$(Sp(1) \times SU(6))/\mathbf{Z}_2$	$(Sp(1) \times S(U(2) \times U(4)))/\mathbf{Z}_2$
E_7	$(SU(2) \times Spin(12))/\mathbf{Z}_2$	$(SU(2) \times Spin(4) \times Spin(8))/(\mathbf{Z}_2 \times \mathbf{Z}_2)$

Table 2

G	G^σ	$G^{\sigma,\gamma}$
F_4	$Spin(9)$	$(Spin(4) \times Spin(5))/\mathbf{Z}_2$
E_6	$(U(1) \times Spin(10))/\mathbf{Z}_4$	$(U(1) \times Spin(4) \times Spin(6))/\mathbf{Z}_2$
E_7	$(SU(2) \times Spin(12))/\mathbf{Z}_2$	$(SU(2) \times Spin(4) \times Spin(8))/(\mathbf{Z}_2 \times \mathbf{Z}_2)$

Notation

- (1) For a group G and an element s of G , we denote $\{g \in G \mid sg = gs\}$ by G^s .
- (2) For a transformation group G of a space M , the isotropy subgroup of G at $m_1, \dots, m_k \in M$ is denoted by $G_{m_1, \dots, m_k} = \{g \in G \mid gm_1 = m_1, \dots, gm_k = m_k\}$.
- (3) For a \mathbf{R} -vector space V , its complexification $\{u + iv \mid u, v \in V\}$ is denoted by V^C . The complex conjugation in V^C is denoted by $\tau : \tau(u + iv) = u - iv$. In particular, the complexification of \mathbf{R} is briefly denoted by $C : \mathbf{R}^C = C$.
- (4) For a Lie group G , the Lie algebra of G is denoted by the corresponding German small letter \mathfrak{g} . For example, $\mathfrak{so}(n)$ is the Lie algebra of the group $SO(n)$.
- (5) Although we will give all definitions used in the following Sections, if in case of insufficiency, refer to [3], [4] or [5].

1. Group F_4

We use the same notation as in [1], [2] or [5] (however, some will be rewritten). For example, the Cayley algebra $\mathfrak{C} = \mathbf{H} \oplus \mathbf{H}e_4$, the exceptional Jordan algebra $\mathfrak{J} = \{X \in M(3, \mathfrak{C}) \mid X^* = X\}$, the Jordan multiplication $X \circ Y$, the inner product (X, Y) and the elements $E_1, E_2, E_3 \in \mathfrak{J}$, the group $F_4 = \{\alpha \in \text{Iso}_{\mathbf{R}}(\mathfrak{J}) \mid \alpha(X \circ Y) = \alpha X \circ \alpha Y\}$.

We define \mathbf{R} -linear transformations σ and γ of \mathfrak{J} by

$$\sigma X = \sigma \begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix} = \begin{pmatrix} \xi_1 & -x_3 & -\bar{x}_2 \\ -\bar{x}_3 & \xi_2 & x_1 \\ -x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}, \quad \gamma X = \begin{pmatrix} \xi_1 & \gamma x_3 & \overline{\gamma x_2} \\ \overline{\gamma x_3} & \xi_2 & \gamma x_1 \\ \gamma x_2 & \overline{\gamma x_1} & \xi_3 \end{pmatrix},$$

respectively, where $\gamma x_k = \gamma(m_k + a_k e_4) = m_k - a_k e_4$, $x_k = m_k + a_k e_4 \in \mathbf{H} \oplus \mathbf{H}e_4 = \mathfrak{C}$. Then, $\sigma, \gamma \in F_4$ and $\sigma^2 = \gamma^2 = 1$. σ and γ are commutative. From $\sigma\gamma = \gamma\sigma$, we have

$$(F_4)^\sigma \cap (F_4)^\gamma = ((F_4)^\sigma)^\gamma = ((F_4)^\gamma)^\sigma.$$

Hence, this group will be denoted briefly by $(F_4)^{\sigma, \gamma}$.

PROPOSITION 1.1. $(F_4)^\gamma \cong (Sp(1) \times Sp(3))/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{(1, E), (-1, -E)\}$.

PROOF. The isomorphism is induced by the homomorphism $\varphi : Sp(1) \times Sp(3) \rightarrow (F_4)^\gamma$, $\varphi(p, A)(M + \mathbf{a}) = AMA^* + p\mathbf{a}A^*$, $M + \mathbf{a} \in \mathfrak{J}(3, \mathbf{H}) \oplus \mathbf{H}^3 = \mathfrak{J}$. (In detail, see [3], [5].)

LEMMA 1.2. $\varphi : Sp(1) \times Sp(3) \rightarrow (F_4)^\gamma$ of Proposition 1.1 satisfies $\sigma\varphi(p, A)\sigma = \varphi(p, I_1 A I_1)$, where $I_1 = \text{diag}(-1, 1, 1)$.

PROOF. From $\sigma = \varphi(-1, I_1)$, we have the required one.

Now, we shall determine the group structure of $(F_4)^{\sigma, \gamma} = ((F_4)^\gamma)^\sigma = ((F_4)^\sigma)^\gamma = (F_4)^\sigma \cap (F_4)^\gamma$.

THEOREM 1.3. $(F_4)^{\sigma, \gamma} \cong (Sp(1) \times Sp(1) \times Sp(2))/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{(1, 1, E), (-1, -1, -E)\}$.

PROOF. We define a map $\varphi_4 : Sp(1) \times Sp(1) \times Sp(2) \rightarrow (F_4)^{\sigma, \gamma}$ by

$$\varphi_4(p, q, B)(M + \mathbf{a}) = \begin{pmatrix} q & 0 & 0 \\ 0 & & B \\ 0 & & \end{pmatrix} M \begin{pmatrix} q & 0 & 0 \\ 0 & & B \\ 0 & & \end{pmatrix}^* + p\mathbf{a} \begin{pmatrix} q & 0 & 0 \\ 0 & & B \\ 0 & & \end{pmatrix}^*,$$

$M + \mathbf{a} \in \mathfrak{J}(3, \mathbf{H}) \oplus \mathbf{H}^3 = \mathfrak{J}$, as the restriction of Proposition 1.1. By Lemma 1.2, φ_4 is well-defined and a homomorphism. We shall show that φ_4 is onto. Let $\alpha \in (F_4)^{\sigma, \gamma}$. Since $(F_4)^{\sigma, \gamma} \subset (F_4)^\gamma$, there exist $p \in Sp(1)$ and $A \in Sp(3)$ such that $\alpha = \varphi(p, A)$ (Proposition 1.1). From $\sigma\alpha\sigma = \alpha$, we have $\varphi(p, I_1 A I_1) = \varphi(p, A)$ (Lemma 1.2). Hence,

$$\begin{cases} p = p \\ I_1 A I_1 = A \end{cases} \quad \text{or} \quad \begin{cases} p = -p \\ I_1 A I_1 = -A \end{cases}.$$

The latter case is impossible because $p = 0$ is false. In the former case, from

$I_1 A I_1 = A$, we have $A = \left(\begin{array}{c|cc} q & 0 & 0 \\ \hline 0 & & B \\ 0 & & \end{array} \right)$, $q \in Sp(1)$, $B \in Sp(2)$. Hence, $\alpha = \varphi(q, \left(\begin{array}{c|cc} q & 0 & 0 \\ \hline 0 & & B \\ 0 & & \end{array} \right)) = \varphi_4(p, q, B)$, that is, φ_4 is onto. And $\text{Ker } \varphi_4 = \{(1, 1, E), (-1, -1, -E)\} = \mathbf{Z}_2$. Thus, we have the required isomorphism $(Sp(1) \times Sp(1) \times Sp(2))/\mathbf{Z}_2 \cong (F_4)^{\sigma, \gamma}$.

2. Group E_6

We use the same notation as in [1], [2] or [5] (however, some will be rewritten). For example, the complex exceptional Jordan algebra $\mathfrak{J}^C = \{X \in M(3, \mathbb{C}^C) \mid X^* = X\}$, the Freudenthal multiplication $X \times Y$ and the Hermitian inner product $\langle X, Y \rangle$, the group $E_6 = \{\alpha \in \text{Iso}_C(\mathfrak{J}^C) \mid \alpha X \times \alpha Y = \tau \alpha \tau(X \times Y), \langle \alpha X, \alpha Y \rangle = \langle X, Y \rangle\}$, and the natural inclusion $F_4 \subset E_6$.

PROPOSITION 2.1. $(E_6)^\gamma \cong (Sp(1) \times SU(6))/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{(1, E), (-1, -E)\}$.

PROOF. The isomorphism is induced by the homomorphism $\varphi : Sp(1) \times SU(6) \rightarrow (E_6)^\gamma$, $\varphi(p, A)(M + \mathbf{a}) = k_J^{-1}(A k_J(M)^t A) + p a k^{-1}(A^*)$, $M + \mathbf{a} \in \mathfrak{J}(3, \mathbf{H})^C \oplus (\mathbf{H}^3)^C = \mathfrak{J}^C$. (In detail, see [3], [5].)

LEMMA 2.2. $\varphi : Sp(1) \times SU(6) \rightarrow (E_6)^\gamma$ of Proposition 2.1 satisfies $\sigma \varphi(p, A) \sigma = \varphi(p, I_2 A I_2)$, where $I_2 = \text{diag}(-1, -1, 1, 1, 1, 1)$.

PROOF. From $\sigma = \varphi(-1, I_2)$, we have the required one.

Now, we shall determine the group structure of $(E_6)^{\sigma, \gamma} = ((E_6)^\gamma)^\sigma = ((E_6)^\sigma)^\gamma = (E_6)^\sigma \cap (E_6)^\gamma$.

THEOREM 2.3. $(E_6)^{\sigma, \gamma} \cong (Sp(1) \times S(U(2) \times U(4)))/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{(1, E), (-1, -E)\}$.

PROOF. We define a map $\varphi_6 : Sp(1) \times S(U(2) \times U(4)) \rightarrow (E_6)^{\sigma, \gamma}$ by

$$\varphi_6(p, A)(M + \mathbf{a}) = k_J^{-1}(A k_J(M)^t A) + p a k^{-1}(A^*),$$

$M + \mathbf{a} \in \mathfrak{J}(3, \mathbf{H})^C \oplus (\mathbf{H}^3)^C = \mathfrak{J}^C$, as the restriction of φ of Proposition 2.1. By Lemma 2.2, φ_6 is well-defined and a homomorphism. We shall show that φ_6 is

onto. Let $\alpha \in (E_6)^{\sigma, \gamma}$. Since $(E_6)^{\sigma, \gamma} \subset (E_6)^\gamma$, there exist $p \in Sp(1)$ and $A \in SU(6)$ such that $\alpha = \varphi(p, A)$ (Proposition 2.1). From $\sigma\alpha\sigma = \alpha$, we have $\varphi(p, I_2AI_2) = \varphi(p, A)$ (Lemma 2.2). Hence,

$$\begin{cases} p = p \\ I_2AI_2 = A \end{cases} \quad \text{or} \quad \begin{cases} p = -p \\ I_2AI_2 = -A \end{cases}.$$

The latter case is impossible because $p = 0$ is false. In the former case, we have $A \in S(U(2) \times U(4))$. Therefore, φ_6 is onto. $\text{Ker } \varphi_6 = \{(1, E), (-1, -E)\} = \mathbf{Z}_2$. Thus, we have the required isomorphism $(Sp(1) \times S(U(2) \times U(4))) / \mathbf{Z}_2 \cong (E_6)^{\sigma, \gamma}$.

3. Group E_7

We use the same notation as in [1], [4] or [5] (however, some will be rewritten). For example, the Freudenthal C -vector space $\mathfrak{P}^C = \mathfrak{J}^C \oplus \mathfrak{J}^C \oplus C \oplus C$, the Hermitian inner product $\langle P, Q \rangle$, the C -linear map $P \times Q : \mathfrak{P}^C \rightarrow \mathfrak{P}^C$ ($P, Q \in \mathfrak{P}^C$), the group $E_7 = \{\alpha \in \text{Iso}_C(\mathfrak{P}^C) \mid \alpha(X \times Y)\alpha^{-1} = \alpha P \times \alpha Q, \langle \alpha P, \alpha Q \rangle = \langle P, Q \rangle\}$, the natural inclusion $E_6 \subset E_7$ and elements $\sigma, \sigma' \in F_4 \subset E_6 \subset E_7$, $\lambda \in E_7$.

We shall consider the following subgroup of F_4 .

$$((F_4)^{\sigma, \gamma})_{F_1(h)} = \{\alpha \in (F_4)^{\sigma, \gamma} \mid \alpha F_1(h) = F_1(h) \text{ for all } h \in \mathbf{H}\}.$$

PROPOSITION 3.1. $((F_4)^{\sigma, \gamma})_{F_1(h)} \cong Sp(1) \times Sp(1)$ ($=Spin(4)$).

PROOF. We define a map $\varphi : Sp(1) \times Sp(1) \rightarrow ((F_4)^{\sigma, \gamma})_{F_1(h)}$ by

$$\varphi(p, q)(M + \mathbf{a}) = \begin{pmatrix} q & 0 & 0 \\ 0 & & E \\ 0 & & \end{pmatrix} M \begin{pmatrix} q & 0 & 0 \\ 0 & & E \\ 0 & & \end{pmatrix}^* + p\mathbf{a} \begin{pmatrix} q & 0 & 0 \\ 0 & & E \\ 0 & & \end{pmatrix}^*,$$

as the restriction of φ_4 of Theorem 1.3. By $F_1(h) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & h \\ 0 & \bar{h} & 0 \end{pmatrix} + \mathbf{O}$, φ is well-defined and homomorphism. We shall show that φ is onto. Let $\alpha \in ((F_4)^{\sigma, \gamma})_{F_1(h)}$. Since $((F_4)^{\sigma, \gamma})_{F_1(h)} \subset (F_4)^{\sigma, \gamma}$, there exist $p, q \in Sp(1)$ and $B \in Sp(2)$ such that $\alpha = \varphi_4(p, q, B)$ (Theorem 1.3). From $\alpha F_1(h) = F_1(h)$, we have $B \begin{pmatrix} 0 & h \\ \bar{h} & 0 \end{pmatrix} B^* = \begin{pmatrix} 0 & h \\ \bar{h} & 0 \end{pmatrix}$, so that

$$\alpha = \varphi_4(p, q, E) \quad \text{or} \quad \alpha = \varphi_4(p, q, -E).$$

In the former case, we have $\alpha = \varphi_4(p, q, E) = \varphi(p, q)$. In the latter case, we have

$$\begin{aligned}\alpha &= \varphi_4(p, q, -E) = \varphi_4(-p, -q, E)\varphi_4(-1, -1, -E) \\ &= \varphi_4(-p, -q, E)1 = \varphi(-p, -q).\end{aligned}$$

Hence, φ is onto. $\text{Ker } \varphi = \{(1, 1)\}$. Thus, we have the required isomorphism $Sp(1) \times Sp(1) \cong ((F_4)^{\sigma, \gamma})_{F_1(h)}$.

Hereafter, in \mathfrak{B}^C , we use the following notations.

$$\begin{aligned}(F_1(h), 0, 0, 0) &= \dot{F}_1(h), & (0, E_1, 0, 1) &= \tilde{E}_1, \\ (0, E_1, 0, -1) &= \tilde{E}_{-1}, & (E_2 + E_3, 0, 0, 0) &= \dot{E}_{23}.\end{aligned}$$

We shall consider a subgroup $((E_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}}$ of E_7 .

LEMMA 3.2. *The Lie algebra $((e_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}}$ of the group $((E_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}}$ is given by*

$$\begin{aligned}& ((e_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}} \\ &= \left\{ \Phi \left(\left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & D'_4 \end{array} \right), 0, 0, 0 \right) \mid \left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & D'_4 \end{array} \right) \in \mathfrak{so}(8), D'_4 \in \mathfrak{so}(4) \right\}.\end{aligned}$$

In particular, we have

$$\dim(((e_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}}) = 6.$$

Hereafter, $\left(\begin{array}{c|c} 0 & 0 \\ \hline 0 & D'_4 \end{array} \right)$ will be denoted by D'_4 , and also $\Phi(D'_4, 0, 0, 0)$ will be denoted by Φ_4 .

PROPOSITION 3.3. $((E_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}} = ((F_4)^{\sigma, \gamma})_{F_1(h)}$.

PROOF. Let $\alpha \in ((F_4)^{\sigma, \gamma})_{F_1(h)}$. Since $((F_4)^{\sigma, \gamma})_{F_1(h)} \subset (F_4)^\sigma = (F_4)_{E_1}$ (as for $(F_4)^\sigma = (F_4)_{E_1}$, see [3], [5]), we see $\alpha E_1 = E_1$. As a result, because κ and μ are defined using by E_1 (see [1], [4] or [5]), we see that $\kappa\alpha = \alpha\kappa$ and $\mu\alpha = \alpha\mu$. From $\alpha E = E$ (see [3], [5]), we have $\alpha(E_2 + E_3) = E_2 + E_3$. Hence, $\alpha\dot{E}_{23} = \dot{E}_{23}$. Moreover, from $\alpha(0, 0, 0, 1) = (0, 0, 0, 1)$ (see [4], [5]), we have $\alpha\tilde{E}_1 = \tilde{E}_1$ and $\alpha\tilde{E}_{-1} = \tilde{E}_{-1}$. Obviously $\alpha\dot{F}_1(h) = \dot{F}_1(h)$. Thus, $\alpha \in (((E_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}})$. Conversely, let $\alpha \in (((E_7)^{\kappa, \mu})_{\dot{F}_1(h), \tilde{E}_1, \tilde{E}_{-1}, \dot{E}_{23}})$. From $\alpha\tilde{E}_1 = \tilde{E}_1$ and $\alpha\tilde{E}_{-1} = \tilde{E}_{-1}$, we have

$\alpha(0, E_1, 0, 0) = (0, E_1, 0, 0)$ and $\alpha(0, 0, 0, 1) = (0, 0, 0, 1)$. Hence, $\alpha \in ((E_6)^\gamma)_{F_1(h), E_1, E_2+E_3}$ (see [4], [5]). Thus, $((F_4)_{E_1})^\gamma_{F_1(h)} = ((F_4)^{\sigma, \gamma})_{F_1(h)}$. Therefore, the proof of this proposition is completed.

Next, we shall consider the following subgroup of F_4 .

$$((F_4)^{\sigma, \gamma})_{F_1(he_4)} = \{\alpha \in (F_4)^{\sigma, \gamma} \mid \alpha F_1(he_4) = F_1(he_4) \text{ for all } h \in H\}.$$

PROPOSITION 3.4. $((F_4)^{\sigma, \gamma})_{F_1(he_4)} \cong Sp(2) (=Spin(5))$.

PROOF. We define a map $\varphi : Sp(2) \rightarrow ((F_4)^{\sigma, \gamma})_{F_1(he_4)}$ by

$$\varphi(B)(M + \mathbf{a}) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & & B \\ 0 & & \end{pmatrix} M \begin{pmatrix} 1 & 0 & 0 \\ 0 & & B \\ 0 & & \end{pmatrix}^* + \mathbf{a} \begin{pmatrix} 1 & 0 & 0 \\ 0 & & B \\ 0 & & \end{pmatrix}^*,$$

as the restriction of φ_4 of Theorem 1.3. Obviously φ is well-defined and homomorphism. We shall show that φ is onto. Let $\alpha \in ((F_4)^{\sigma, \gamma})_{F_1(he_4)}$. Since $((F_4)^{\sigma, \gamma})_{F_1(he_4)} \subset (F_4)^{\sigma, \gamma}$, there exist $p, q \in Sp(1)$ and $B \in Sp(2)$ such that $\alpha = \varphi_4(p, q, B)$ (Theorem 1.3). From $\alpha F_1(he_4) = F_1(he_4) (=O + (h, 0, 0))$, we have $ph\bar{q} = h$ ($h \in H$), so that

$$\alpha = \varphi_4(1, 1, B) \quad \text{or} \quad \alpha = \varphi_4(-1, -1, B).$$

In the former case, we have $\alpha = \varphi_4(1, 1, B) = \varphi(B)$. In the latter case, we have

$$\begin{aligned} \alpha &= \varphi_4(-1, -1, B) = \varphi_4(1, 1, -B)\varphi_4(-1, -1, -E) \\ &= \varphi_4(1, 1, -B)1 = \varphi(-B). \end{aligned}$$

Hence, φ is onto. $\text{Ker } \varphi = \{E\}$. Thus, we have the required isomorphism $Sp(2) \cong ((F_4)^{\sigma, \gamma})_{F_1(he_4)}$.

Then, we have the following proposition.

PROPOSITION 3.5. $((E_7)^{\kappa, \mu})^\gamma_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}, \tilde{E}_{23}} = ((F_4)^{\sigma, \gamma})_{F_1(he_4)}$.

PROOF. This proof is in the way similar to Proposition 3.3.

We shall consider the subgroup $((E_7)^{\kappa, \mu})^\gamma_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ of E_7 .

LEMMA 3.6. *The Lie algebra $((e_7)^{\kappa, \mu})^\gamma_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ of the group $((E_7)^{\kappa, \mu})^\gamma_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ is given by*

$$\begin{aligned}
& (((\mathbf{e}_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(\mathbf{he}_4), \dot{E}_1, \dot{E}_{-1}} \\
&= \left\{ \Phi \left(\left(\begin{array}{c|c} D_4 & 0 \\ \hline 0 & 0 \end{array} \right) + \tilde{A}_1(p) + i \begin{pmatrix} 0 & 0 & 0 \\ 0 & \varepsilon & q \\ 0 & \bar{q} & -\varepsilon \end{pmatrix}, 0, 0, 0 \right) \middle| \left(\begin{array}{c|c} D_4 & 0 \\ \hline 0 & 0 \end{array} \right) \in \mathfrak{so}(8), \right. \\
&\quad \left. D_4 \in \mathfrak{so}(4), \varepsilon \in \mathbf{R}, p, q \in \mathbf{H} \right\}.
\end{aligned}$$

In particular, we have

$$\dim(((\mathbf{e}_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(\mathbf{he}_4), \dot{E}_1, \dot{E}_{-1}} = 15.$$

Hereafter, $\left(\begin{array}{c|c} D_4 & 0 \\ \hline 0 & 0 \end{array} \right)$ will be denoted by D_4 .

LEMMA 3.7. (1) For $a \in \mathbf{H}$, we define a map $\tilde{\alpha}_1(a)$ of \mathfrak{J}^C by

$$\begin{cases} \xi'_1 = \xi_1 \\ \xi'_2 = \frac{\xi_2 - \xi_3}{2} + \frac{\xi_2 + \xi_3}{2} \cos|a| + i \frac{(a, x_1)}{|a|} \sin|a| \\ \xi'_3 = -\frac{\xi_2 - \xi_3}{2} + \frac{\xi_2 + \xi_3}{2} \cos|a| + i \frac{(a, x_1)}{|a|} \sin|a| \\ x'_1 = x_1 + i \frac{(\xi_2 + \xi_3)a}{|a|} \sin|a| - \frac{2(a, x_1)a}{|a|^2} \left(\sin \frac{|a|}{2} \right)^2 \\ x'_2 = x_2 \cos \frac{|a|}{2} + i \frac{\bar{x}_3 a}{|a|} \sin \frac{|a|}{2} \\ x'_3 = x_3 \cos \frac{|a|}{2} + i \frac{\bar{a} x_2}{|a|} \sin \frac{|a|}{2} \end{cases}$$

Then, $\tilde{\alpha}_1(a) \in (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(\mathbf{he}_4), \dot{E}_1, \dot{E}_{-1}}$.

(2) For $t \in \mathbf{R}$, we define a map $\tilde{\alpha}_{23}(t)$ of \mathfrak{J}^C by

$$\tilde{\alpha}_{23}(t) \begin{pmatrix} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{pmatrix} = \begin{pmatrix} \xi_1 & e^{it/2} x_3 & e^{-it/2} \bar{x}_2 \\ e^{it/2} \bar{x}_3 & e^{it} \xi_2 & x_1 \\ e^{-it/2} x_2 & \bar{x}_1 & e^{-it} \xi_3 \end{pmatrix}.$$

Then, $\tilde{\alpha}_{23}(t) \in (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(\mathbf{he}_4), \dot{E}_1, \dot{E}_{-1}}$.

PROOF. (1) For $a \in \mathbf{H}$, we have $i\tilde{F}_1(a) \in (((\mathbf{e}_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(\mathbf{he}_4), \dot{E}_1, \dot{E}_{-1}}$ (Lemma 3.6). Hence, $\tilde{\alpha}_1(a) = \exp i\tilde{F}_1(a) \in (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(\mathbf{he}_4), \dot{E}_1, \dot{E}_{-1}}$.

(2) For $t \in \mathbf{R}$, we have $it(E_2 - E_3)^\sim \in (((e_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ (Lemma 3.6). Hence, $\tilde{\alpha}_{23}(t) = \exp it(E_2 - E_3)^\sim \in (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$.

We define a 6 dimensional \mathbf{R} -vector space V^6 by

$$V^6 = \{P \in \mathfrak{B}^C \mid \kappa P = P, \mu\tau\lambda P = P, \gamma P = P, \langle P, \tilde{E}_1 \rangle = 0, \langle P, \tilde{E}_{-1} \rangle = 0\}$$

$$= \left\{ P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, 0, 0, 0 \right) \mid \xi \in C, h \in \mathbf{H} \right\}$$

with the norm (see [5] for the definition of $\{, \}$'s)

$$(P, P)_\mu = \frac{1}{2} \{ \mu P, P \} = \frac{1}{2} (\mu P, \lambda P) = (\tau\xi)\xi + \bar{h}h.$$

Then, $S^5 = \{P \in V^6 \mid (P, P)_\mu = 1\}$ is a 5 dimensional sphere.

LEMMA 3.8. $(((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}} / Spin(5) \simeq S^5$. In particular, $(((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ is connected.

PROOF. Since E_7 is commutative with $\tau\lambda$, the group $(((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ acts on S^5 . We shall show that this action is transitive. To show this, it is sufficient to show that any element $P \in S^5$ can be transformed to $(i(E_2 + E_3), 0, 0, 0) \in S^5$ under the action of $(((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$. Now, for a given

$$P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, 0, 0, 0 \right) \in S^5,$$

choose $t \in \mathbf{R}$ such that $e^{it\xi} \in \mathbf{R}$. For this $t \in \mathbf{R}$, operate $\tilde{\alpha}_{23}(t)$ (Lemma 3.7(2)) $\in (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ on P . Then, we have

$$\tilde{\alpha}_{23}(t)P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & r & h \\ 0 & \bar{h} & -r \end{pmatrix}, 0, 0, 0 \right) = P_1, \quad r \in \mathbf{R}.$$

In the case of $h \neq 0$, operate $\tilde{\alpha}_1(\pi h/2|h|)$ (Lemma 3.7(1)) $\in (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4), \tilde{E}_1, \tilde{E}_{-1}}$ on P_1 . Then, we have

$$\tilde{\alpha}_1\left(\frac{\pi h}{2|h|}\right)P_1 = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi' & 0 \\ 0 & 0 & -\tau\xi' \end{pmatrix}, 0, 0, 0 \right) = P_2 \in S^5, \quad \xi' \in C.$$

Here, from $(\tau\xi')\xi' = 1$, $\xi' \in C$, we can put $\xi' = e^{i\theta}$, $0 \leq \theta < 2\pi$. Operate $\tilde{\alpha}_{23}(-\theta)$ on P_2 . Then,

$$\tilde{\alpha}_{23}(-\theta)P_2 = (E_2 - E_3, 0, 0, 0) = P_3.$$

Moreover, operate $\tilde{\alpha}_{23}(\pi/2)$ on P_3 ,

$$\tilde{\alpha}_{23}\left(\frac{\pi}{2}\right)P_3 = (i(E_2 + E_3), 0, 0, 0) = i\dot{E}_{23}.$$

This shows the transitivity. The isotropy subgroup $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}}$ at \dot{E}_{23} is $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}, \dot{E}_{23}} = Sp(2)$ (Propositions 3.4, 3.5) = $Spin(5)$. Therefore, we have the homeomorphism $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}}/Spin(5) \simeq S^5$.

PROPOSITION 3.9. $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}} \cong Spin(6)$.

PROOF. Since $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}}$ is connected (Lemma 3.8), we can define a homomorphism $\pi : ((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}} \rightarrow SO(6) = SO(V^6)$ by

$$\pi(\alpha) = \alpha|V^6.$$

It is not difficult to see that $\text{Ker } \pi = \{1, \sigma\} = \mathbf{Z}_2$. Since $\dim(((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}}) = 15$ (Lemma 3.6) = $\dim(\mathfrak{so}(6))$, π is onto. Hence, $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}}/\mathbf{Z}_2 \cong SO(6)$. Therefore, $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1, \dot{E}_{-1}}$ is isomorphism to $Spin(6)$ as a double covering group of $SO(6)$.

We shall consider a subgroup $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1}$ of E_7 .

LEMMA 3.10. *The Lie algebra $((\mathfrak{e}_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1}$ of the group $((E_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1}$ is given by*

$$\begin{aligned} & ((\mathfrak{e}_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1} \\ &= \left\{ \Phi \left(D_4 + \tilde{A}_1(p) + i \begin{pmatrix} 0 & 0 & 0 \\ 0 & \varepsilon & q \\ 0 & \bar{q} & -\varepsilon \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & \alpha & ix \\ 0 & \bar{ix} & \tau\alpha \end{pmatrix}, -\tau \begin{pmatrix} 0 & 0 & 0 \\ 0 & \alpha & ix \\ 0 & \bar{ix} & \tau\alpha \end{pmatrix}, 0 \right) \right. \\ & \quad \left. | D_4 \in \mathfrak{so}(4) \subset \mathfrak{so}(8), \varepsilon \in \mathbf{R}, \alpha \in C, p, q, x \in \mathbf{H} \right\}. \end{aligned}$$

In particular, we have

$$\dim(((\mathfrak{e}_7)^{\kappa, \mu})^{\gamma}_{\dot{F}_1(\dot{h}e_4), \dot{E}_1}) = 21.$$

LEMMA 3.11. *For $a \in \mathbf{R}$, we define maps $\alpha_k(a)$, $k = 2, 3$ of \mathfrak{P}^C by*

$$\alpha_k(a) \begin{pmatrix} X \\ Y \\ \xi \\ \eta \end{pmatrix} = \begin{pmatrix} (1 + (\cos a - 1)p_k)X - 2(\sin a)E_k \times Y + \eta(\sin a)E_k \\ 2(\sin a)E_k \times X + (1 + (\cos a - 1)p_k)Y - \xi(\sin a)E_k \\ ((\sin a)E_k, Y) + (\cos a)\xi \\ -(\sin a)E_k, X) + (\cos a)\eta \end{pmatrix},$$

where $p_k : \mathfrak{J}^C \rightarrow \mathfrak{J}^C$ is defined by

$$p_k(X) = (X, E_k)E_k + 4E_k \times (E_k \times X), \quad X \in \mathfrak{J}^C.$$

Then, $\alpha_k \in E_7$ and $\alpha_2(a), \alpha_3(b)$ ($a, b \in \mathbf{R}$) commute with each other.

PROOF. For $\Phi_k(a) = \Phi(0, aE_k, -aE_k, 0) \in \mathfrak{e}_7$, we have $\alpha_k(a) = \exp \Phi_k(a) \in E_7$. Since $[\Phi_2(a), \Phi_3(b)] = 0$, $\alpha_2(a)$ and $\alpha_3(b)$ are commutative.

We define a 7 dimensional \mathbf{R} -vector space V^7 by

$$V^7 = \left\{ P \in \mathfrak{B}^C \mid \kappa P = P, \mu\tau\lambda P = P, \gamma P = P, \langle P, \tilde{E}_1 \rangle = 0 \right\}$$

$$= \left\{ P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, \begin{pmatrix} i\eta & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, 0, -i\eta \right) \mid \xi \in C, h \in \mathbf{H}, \eta \in \mathbf{R} \right\}$$

with the norm

$$(P, P)_\mu = \frac{1}{2}(\mu P, \lambda P) = (\tau\xi)\xi + \bar{h}h + \eta^2.$$

Then, $S^6 = \{P \in V^7 \mid (P, P)_\mu = 1\}$ is a 6 dimensional sphere.

LEMMA 3.12. $((E_7)^{\kappa, \mu})_{\dot{F}_1(he_4), \tilde{E}_1}^\gamma / Spin(6) \simeq S^6$. In particular, $((E_7)^{\kappa, \mu})_{\dot{F}_1(he_4), \tilde{E}_1}^\gamma$ is connected.

PROOF. The group $((E_7)^{\kappa, \mu})_{\dot{F}_1(he_4), \tilde{E}_1}^\gamma$ acts on S^6 . We shall show that this action is transitive. To show this, it is sufficient to show that any element $P \in S^6$ can be transformed to $(0, -iE_1, 0, i) \in S^6$ under the action of $((E_7)^{\kappa, \mu})_{\dot{F}_1(he_4), \tilde{E}_1}^\gamma$. Now, for a given

$$P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, \begin{pmatrix} i\eta & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, 0, -i\eta \right) \in S^6,$$

choose $a \in \mathbf{R}$, $0 \leq a < \pi/2$ such that $\tan 2a = \frac{i2\eta}{\tau\xi - \xi}$ (if $\tau\xi - \xi = 0$, then let

$a = \pi/4$). Operate $\alpha_{23}(a) := \alpha_2(a)\alpha_3(a) = \exp(\Phi(0, a(E_2 + E_3), -a(E_2 + E_3), 0))$ (Lemma 3.11) $\in (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1}$ (Lemma 3.10) on P . Then, the η -term of $\alpha_{23}(a)P$ is $(1/2)(\xi - \tau\zeta) \sin 2a + i\eta \cos 2a = 0$. Hence,

$$\alpha_{23}(a)P = \left(\left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \zeta & m \\ 0 & \bar{m} & -\tau\zeta \end{pmatrix}, 0, 0, 0 \right), 0, 0, 0 \right) = P_1 \in S^5 \subset S^6.$$

Since $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1, \tilde{E}_{-1}} (\subset (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1}))$ acts transitively on S^5 (Lemma 3.8), there exist $\beta \in (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1, \tilde{E}_{-1}}$ such that $\beta P_1 = (i(E_2 + E_3), 0, 0, 0) = P_2 \in S^5 \subset S^6$. Moreover, operate $\alpha_{23}(-\pi/4)$ on P_2 ,

$$\alpha_{23}\left(-\frac{\pi}{4}\right)P_2 = (0, -iE_1, 0, i) = -i\tilde{E}_{-1}.$$

This shows the transitivity. The isotropy subgroup $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1})$ at \tilde{E}_{-1} is $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1, \tilde{E}_{-1}} = \text{Spin}(6)$ (Proposition 3.9). Thus, we have the homeomorphism $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} / \text{Spin}(6) \simeq S^6$.

PROPOSITION 3.13. $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} \cong \text{Spin}(7)$.

PROOF. Since $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1})$ is connected (Lemma 3.12), we can define a homomorphism $\pi : (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} \rightarrow SO(7) = SO(V^7)$ by

$$\pi(\alpha) = \alpha|_{V^7}.$$

It is not difficult to see that $\text{Ker } \pi = \{1, \sigma\} = \mathbf{Z}_2$. Since $\dim(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} = 21$ (Lemma 3.10) $= \dim(\mathfrak{so}(7))$, π is onto. Hence, $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} / \mathbf{Z}_2 \cong SO(7)$. Therefore, $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1})$ is isomorphism to $\text{Spin}(7)$ as a double covering group of $SO(7)$.

We shall consider the subgroup $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)})$ of E_7 .

LEMMA 3.14. *The Lie algebra $(((\mathfrak{e}_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)}$ of the group $((((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)})$ is given by*

$$\begin{aligned} & ((\mathfrak{e}_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)} \\ &= \left\{ \Phi \left(D_4 + \tilde{A}_1(p) + i \begin{pmatrix} \varepsilon_1 & 0 & 0 \\ 0 & \varepsilon_2 & q \\ 0 & \bar{q} & \varepsilon_3 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & \alpha_2 & x \\ 0 & \bar{x} & \alpha_3 \end{pmatrix}, -\tau \begin{pmatrix} 0 & 0 & 0 \\ 0 & \alpha_2 & x \\ 0 & \bar{x} & \alpha_3 \end{pmatrix}, -\frac{3}{2}i\varepsilon_1 \right) \right. \\ & \left. | D_4 \in \mathfrak{so}(4) \subset \mathfrak{so}(8), \alpha_k \in \mathbf{C}, p, q \in \mathbf{H}, x \in \mathbf{H}^{\mathbf{C}}, \varepsilon_k \in \mathbf{R}, \varepsilon_1 + \varepsilon_2 + \varepsilon_3 = 0 \right\}. \end{aligned}$$

In particular, we have

$$\dim(\left(\left(\left(\mathfrak{e}_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}) = 28.$$

Hereafter, any element of the Lie algebra $\left(\left(\left(\mathfrak{e}_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$ will be denoted by Φ_8 .

LEMMA 3.15. For $t \in \mathbf{R}$, we define a map $\alpha(t)$ of \mathfrak{B}^C by

$$\alpha(t)(X, Y, \xi, \eta) = \left(\left(\begin{pmatrix} e^{2it}\xi_1 & e^{it}x_3 & e^{it}\bar{x}_2 \\ e^{it}\bar{x}_3 & \xi_2 & x_1 \\ e^{it}x_2 & \bar{x}_1 & \xi_3 \end{pmatrix}, \begin{pmatrix} e^{-2it}\eta_1 & e^{-it}y_3 & e^{-it}\bar{y}_2 \\ e^{-it}\bar{y}_3 & \eta_2 & y_1 \\ e^{-it}y_2 & \bar{y}_1 & \eta_3 \end{pmatrix}, e^{-2it}\xi, e^{2it}\eta \right).$$

Then, $\alpha(t) \in \left(\left(\left(E_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$.

PROOF. For $\Phi = \Phi(2itE_1 \vee E_1, 0, 0, -2it) \in \left(\left(\left(\mathfrak{e}_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$ (Lemma 3.14), we have $\alpha(t) = \exp \Phi \in \left(\left(\left(E_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$ by $E_1 \vee E_1 = (1/3)(2E_1 - E_2 - E_3)^\sim$.

We define an 8 dimensional \mathbf{R} -vector space V^8 by

$$V^8 = \{P \in \mathfrak{B}^C \mid \kappa P = P, \mu\tau\lambda P = P, \gamma P = P\} \\ = \left\{ P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, \begin{pmatrix} \eta & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, 0, \tau\eta \right) \mid \xi, \eta \in \mathbf{C}, h \in \mathbf{H} \right\}$$

with the norm

$$(P, P)_\mu = \frac{1}{2}(\mu P, \lambda P) = (\tau\xi)\xi + \bar{h}h + (\tau\eta)\eta.$$

Then, $S^7 = \{P \in V^8 \mid (P, P)_\mu = 1\}$ is a 7 dimensional sphere.

LEMMA 3.16. $\left(\left(\left(E_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}/Spin(7) \simeq S^7$. In particular, $\left(\left(\left(E_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$ is connected.

PROOF. The group $\left(\left(\left(E_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$ acts on S^7 . We shall show that this action is transitive. To show this, it is sufficient to show that any element $P \in S^7$ can be transformed to $(0, E_1, 0, 1) \in S^7$ under the action of $\left(\left(\left(E_7\right)^{\kappa,\mu}\right)^\gamma\right)_{\hat{F}_1(\mathfrak{he}_4)}$. Now, for a given

$$P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, \begin{pmatrix} \eta & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, 0, \tau\eta \right) \in S^7,$$

choose $t \in \mathbf{R}$ such that $e^{-2it}\eta \in i\mathbf{R}$. Operate $\alpha(t)$ (Lemma 3.15) $\in (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)}$ on P . Then,

$$\alpha(t)P = \left(\begin{pmatrix} 0 & 0 & 0 \\ 0 & \xi & h \\ 0 & \bar{h} & -\tau\xi \end{pmatrix}, \begin{pmatrix} i\eta' & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, 0, -i\eta' \right) = P_1 \in S^6 \subset S^7, \quad \eta' \in \mathbf{R}$$

Since $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} (\subset ((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)}$ acts transitively on S^6 (Lemma 3.12), there exists $\beta \in (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1}$ such that $\beta P_1 = (0, -iE_1, 0, i) = P_2 \in S^6 \subset S^7$. Moreover, operate $\alpha(-\pi/4)$ (Lemma 3.15) on P_2 ,

$$\alpha\left(-\frac{\pi}{4}\right)P_2 = (0, E_1, 0, 1) = \tilde{E}_1.$$

This shows the transitivity. The isotropy subgroup $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)}$ at \tilde{E}_1 is $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4), \tilde{E}_1} = \text{Spin}(7)$ (Proposition 3.12). Thus, we have the homeomorphism $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)} / \text{Spin}(7) \simeq S^7$.

PROPOSITION 3.17. $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)} \cong \text{Spin}(8)$.

PROOF. Since $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)}$ is connected (Lemma 3.16), we can define a homomorphism $\pi : (((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)} \rightarrow \text{SO}(8) = \text{SO}(V^8)$ by

$$\pi(\alpha) = \alpha|V^8.$$

It is not difficult to see that $\text{Ker } \pi = \{1, \sigma\} = \mathbf{Z}_2$. Since $\dim(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)} = 28$ (Lemma 3.14) $= \dim(\mathfrak{so}(8))$, π is onto. Hence, $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)} / \mathbf{Z}_2 \cong \text{SO}(8)$. Therefore, $(((E_7)^{\kappa, \mu})^\gamma)_{\tilde{F}_1(\text{he}_4)}$ is isomorphism to $\text{Spin}(8)$ as a double covering group of $\text{SO}(8)$.

We shall determine the group structure of $(E_7)^{\kappa, \mu}$.

LEMMA 3.18. *The Lie algebra $((e_7)^{\kappa, \mu})^\gamma$ of the group $(E_7)^{\kappa, \mu}$ is given by*

$$\begin{aligned} ((e_7)^{\kappa, \mu})^\gamma = & \left\{ \Phi \left(D_4 + D'_4 + \tilde{A}_1(p) + i \begin{pmatrix} \varepsilon_1 & 0 & 0 \\ 0 & \varepsilon_2 & q \\ 0 & \bar{q} & \varepsilon_3 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & \alpha_2 & x \\ 0 & \bar{x} & \alpha_3 \end{pmatrix}, \right. \\ & \left. -\tau \begin{pmatrix} 0 & 0 & 0 \\ 0 & \alpha_2 & x \\ 0 & \bar{x} & \alpha_3 \end{pmatrix}, -\frac{3}{2}i\varepsilon_1 \right) \mid D_4, D'_4 \in \mathfrak{so}(4) \subset \mathfrak{so}(8), \alpha_k \in \mathbf{C}, p, q \in \mathbf{H}, \\ & \left. x \in \mathbf{H}^{\mathbf{C}}, \varepsilon_k \in \mathbf{R}, \varepsilon_1 + \varepsilon_2 + \varepsilon_3 = 0 \right\}. \end{aligned}$$

In particular, we have

$$\dim(((e_7)^{\kappa, \mu})^\gamma) = 34.$$

PROPOSITION 3.19. $((E_7)^{\kappa, \mu})^\gamma \cong (Spin(4) \times Spin(8))/Z_2$, $Z_2 = \{(1, 1), (-1, -1)\}$.

PROOF. For $Spin(4) = Sp(1) \times Sp(1) = (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(h), \dot{E}_1, \dot{E}_{-1}, \dot{E}_{23}}$ (Propositions 3.1, 3.3) and $Spin(8) = (((E_7)^{\kappa, \mu})^\gamma)_{\dot{F}_1(he_4)}$ (Proposition 3.17), we define a map $\phi_1 : Spin(4) \times Spin(8) \rightarrow ((E_7)^{\kappa, \mu})^\gamma$ by

$$\phi_1(\alpha, \beta) = \alpha\beta.$$

Then, ϕ_1 is well-defined. For $\Phi_4 \in \mathfrak{spin}(4)$ (Lemma 3.2) and $\Phi_8 \in \mathfrak{spin}(8)$ (Lemma 3.14), since $[\Phi_4, \Phi_8] = 0$, we have $\alpha\beta = \beta\alpha$. Hence, ϕ_1 is a homomorphism. It is not difficult to see that $\text{Ker } \phi_1 = \{(1, 1), (-1, -1)\} = Z_2$. Since $((E_7)^{\kappa, \mu})^\gamma (\cong Spin(12))^\gamma$. (see [4], [5]) is connected and $\dim(((E_7)^{\kappa, \mu})^\gamma) = 34$ (Lemma 3.18) $= 6 + 28 = \dim(\mathfrak{spin}(4) \oplus \mathfrak{spin}(8))$, ϕ_1 is onto. Thus, we have the required isomorphism $(Spin(4) \times Spin(8))/Z_2 \cong ((E_7)^{\kappa, \mu})^\gamma$.

Now, we shall determine the group structure of $(E_7)^{\sigma, \gamma}$.

LEMMA 3.20. The Lie algebra $(e_7)^{\sigma, \gamma}$ of the group $(E_7)^{\sigma, \gamma}$ is given by

$$(e_7)^{\sigma, \gamma} = \left\{ \Phi \left(D_4 + D'_4 + \tilde{A}_1(p) + i \begin{pmatrix} \varepsilon_1 & 0 & 0 \\ 0 & \varepsilon_2 & q \\ 0 & \bar{q} & \varepsilon_3 \end{pmatrix} \right), \begin{pmatrix} \alpha_1 & 0 & 0 \\ 0 & \alpha_2 & x \\ 0 & \bar{x} & \alpha_3 \end{pmatrix}, \right. \\ \left. -\tau \begin{pmatrix} \alpha_1 & 0 & 0 \\ 0 & \alpha_2 & x \\ 0 & \bar{x} & \alpha_3 \end{pmatrix}, \nu \right\} \mid D_4, D'_4 \in \mathfrak{so}(4) \subset \mathfrak{so}(8), \alpha_k \in \mathbf{C}, p, q \in \mathbf{H}, x \in \mathbf{H}^C, \\ \varepsilon_k \in \mathbf{R}, \varepsilon_1 + \varepsilon_2 + \varepsilon_3 = 0, \nu \in i\mathbf{R} \left. \right\}.$$

In particular, we have

$$\dim((e_7)^{\sigma, \gamma}) = 37.$$

PROPOSITION 3.21. For $A \in SU(2) = \{A \in M(2, \mathbf{C}) \mid (\tau^t A)A = E, \det A = 1\}$, we define \mathbf{C} -linear transformations $\phi(A)$ of \mathfrak{B}^C by

$$\begin{aligned}
\phi(A) & \left(\left(\begin{array}{ccc} \xi_1 & x_3 & \bar{x}_2 \\ \bar{x}_3 & \xi_2 & x_1 \\ x_2 & \bar{x}_1 & \xi_3 \end{array} \right), \left(\begin{array}{ccc} \eta_1 & y_3 & \bar{y}_2 \\ \bar{y}_3 & \eta_2 & y_1 \\ y_2 & \bar{y}_1 & \eta_3 \end{array} \right), \xi, \eta \right) \\
& = \left(\left(\begin{array}{ccc} \xi'_1 & x'_3 & \bar{x}'_2 \\ \bar{x}'_3 & \xi'_2 & x'_1 \\ x'_2 & \bar{x}'_1 & \xi'_3 \end{array} \right), \left(\begin{array}{ccc} \eta'_1 & y'_3 & \bar{y}'_2 \\ \bar{y}'_3 & \eta'_2 & y'_1 \\ y'_2 & \bar{y}'_1 & \eta'_3 \end{array} \right), \xi', \eta' \right), \\
\begin{pmatrix} \xi'_1 \\ \eta' \end{pmatrix} & = A \begin{pmatrix} \xi_1 \\ \eta \end{pmatrix}, \quad \begin{pmatrix} \xi' \\ \eta'_1 \end{pmatrix} = A \begin{pmatrix} \xi \\ \eta_1 \end{pmatrix}, \quad \begin{pmatrix} \eta'_2 \\ \xi'_3 \end{pmatrix} = A \begin{pmatrix} \eta_2 \\ \xi_3 \end{pmatrix}, \\
\begin{pmatrix} \eta'_3 \\ \xi'_2 \end{pmatrix} & = A \begin{pmatrix} \eta_3 \\ \xi_2 \end{pmatrix}, \quad \begin{pmatrix} x'_1 \\ y'_1 \end{pmatrix} = (\tau A) \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \\
\begin{pmatrix} x'_2 \\ y'_2 \end{pmatrix} & = \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, \quad \begin{pmatrix} x'_3 \\ y'_3 \end{pmatrix} = \begin{pmatrix} x_3 \\ y_3 \end{pmatrix}.
\end{aligned}$$

Then, $\phi(A) \in (E_7)^{\sigma, \gamma}$.

PROOF. Let $\Phi = \Phi(2\nu E_1 \vee E_1, aE_1, -\tau aE_1, \nu)$, $a \in C$, $\nu \in i\mathbf{R}$. Then, $\Phi \in (\mathfrak{e}_7)^{\sigma, \gamma}$ (Lemma 3.20). Therefore, for $A = \exp \begin{pmatrix} \nu & a \\ -\tau a & -\nu \end{pmatrix} \in SU(2)$, we have $\phi(A) = \exp \Phi \in (E_7)^{\sigma, \gamma}$.

PROPOSITION 3.22. $(E_7)^\sigma \cong (SU(2) \times Spin(12))/\mathbf{Z}_2$, $\mathbf{Z}_2 = \{(E, 1), (-E, -\sigma)\}$.

PROOF. The isomorphism is induced by the homomorphism $\varphi_1 : SU(2) \times Spin(12) \rightarrow (E_7)^\sigma$ by $\varphi_1(A, \delta) = \phi(A)\delta$. (In detail, see [4], [5].)

THEOREM 3.23. $(E_7)^{\sigma, \gamma} \cong (SU(2) \times Spin(4) \times Spin(8))/(\mathbf{Z}_2 \times \mathbf{Z}_2)$, $\mathbf{Z}_2 \times \mathbf{Z}_2 = \{(E, 1, 1), (E, \sigma, \sigma)\} \times \{(E, 1, 1), (-E, \gamma, -\sigma\gamma)\}$.

PROOF. For $SU(2)$ (Proposition 3.21), $Spin(4) = (((E_7)^{\kappa, \mu})^\gamma)_{\hat{F}_1(h), \hat{E}_1, \hat{E}_{-1}, \hat{E}_{23}}$ (Propositions 3.1, 3.3) and $Spin(8) = (((E_7)^{\kappa, \mu})^\gamma)_{\hat{F}_1(he_4)}$ (Proposition 3.17), we define a map $\varphi : SU(2) \times Spin(4) \times Spin(8) \rightarrow (E_7)^{\sigma, \gamma}$ by

$$\varphi(A, \alpha, \beta) = \phi(A)\alpha\beta.$$

Then, φ is well-defined. From Propositions 3.19, 3.22, φ is a homomorphism. We shall show that φ is onto. Let $\rho \in (E_7)^{\sigma, \gamma}$. Since $(E_7)^{\sigma, \gamma} \subset (E_7)^\sigma$, there exist $A \in SU(2)$ and $\delta \in Spin(12)$ such that $\rho = \varphi_1(A, \delta)$ (Proposition 3.22). Now, from $\gamma\rho\gamma = \rho$, we have $\phi(A)(\gamma\delta\gamma) = \phi(A)\delta$. Hence,

$$\begin{cases} A = A \\ \gamma\delta\gamma = \delta \end{cases} \quad \text{or} \quad \begin{cases} A = -A \\ \gamma\delta\gamma = -\sigma\delta \end{cases}.$$

The latter case is impossible because $A = 0$ is false. In the former case, from Proposition 3.19, there exist $\alpha \in Spin(4)$ and $\beta \in Spin(8)$ such that $\delta = \phi_1(\alpha, \beta)$. Hence, we have

$$\begin{aligned} \rho &= \phi_1(A, \delta) = \phi(A)\delta = \phi(A)\phi_1(\alpha, \beta) \\ &= \phi(A)\alpha\beta = \varphi(A, \alpha, \beta). \end{aligned}$$

It is not difficult to see that

$$\begin{aligned} \text{Ker } \varphi &= \{(E, 1, 1), (E, \sigma, \sigma), (-E, \gamma, -\sigma\gamma), (-E, \sigma\gamma, -\gamma)\} \\ &= \{(E, 1, 1), (E, \sigma, \sigma)\} \times \{(E, 1, 1), (-E, \gamma, -\sigma\gamma)\} \\ &= \mathbf{Z}_2 \times \mathbf{Z}_2. \end{aligned}$$

Thus, we have the required isomorphism $(SU(2) \times Spin(4) \times Spin(8))/(\mathbf{Z}_2 \times \mathbf{Z}_2) \cong (E_7)^{\sigma, \gamma}$.

References

- [1] T. Miyashita and I. Yokota, Fixed points subgroups $G^{\sigma, \sigma'}$ by two involutive automorphisms σ, σ' of compact exceptional Lie group $G = F_4, E_6$ and E_7 , (2001), preprint.
- [2] T. Miyashita and I. Yokota, Fixed points subgroups $G^{\gamma, \gamma'}$ by two involutive automorphisms γ, γ' of compact exceptional Lie group $G = G_2, F_4$ and E_6 (in Japanese), (2001), preprint.
- [3] I. Yokota, Realizations of involutive automorphisms σ and G^σ of exceptional linear Lie groups G , Part I, $G = G_2, F_4$ and E_6 , Tsukuba J. Math., **4** (1990), 185–223.
- [4] I. Yokota, Realizations of involutive automorphisms σ and G^σ of exceptional linear Lie groups G , Part II, $G = E_7$, Tsukuba J. Math., **14** (1990), 379–404.
- [5] I. Yokota, Exceptional simple Lie groups (in Japanese), Gendaisuugakusya, Kyoto, 1992.

Toshikazu Miyashita
Nagano Higashi High School
Nagano, 381-0022, Japan
E-mail: spin15ss16@ybb.ne.jp