

## PARABOLIC HARNACK INEQUALITY FOR THE MIXTURE OF BROWNIAN MOTION AND STABLE PROCESS

RENMING SONG\* AND ZORAN VONDRAČEK\*\*

(Received July 11, 2005, revised March 22, 2006)

**Abstract.** Let  $X$  be a mixture of independent Brownian motion and symmetric stable process. In this paper we establish sharp bounds for transition density of  $X$ , and prove a parabolic Harnack inequality for nonnegative parabolic functions of  $X$ .

**1. Introduction.** Let  $W = (W_t : t \geq 0)$  be a Brownian motion in Euclidean  $d$ -space  $\mathbf{R}^d$ , and let  $Y = (Y_t : t \geq 0)$  be a rotationally invariant  $\alpha$ -stable process in  $\mathbf{R}^d$ , where  $0 < \alpha < 2$ . Suppose that  $W$  and  $Y$  are independent and define the process  $X = (X_t : t \geq 0)$  by  $X_t = W_t + Y_t$ . The law of  $X$  started from  $x \in \mathbf{R}^d$  will be denoted by  $\mathbf{P}^x$ . We will call the process  $X$  the mixture of the Brownian motion  $W$  and the stable process  $Y$ . Although  $X$  is a Lévy process with explicitly known generator and Lévy measure, until recently not much was known about the Green function and transition density of this process. The main difficulty in studying the process is the fact that it runs on two different scales. By realizing the process  $X$  as a subordinate Brownian motion and using Tauberian theorems, the asymptotic behaviors of the Green function of  $X$  near zero and infinity were established in [7]. These asymptotics were used in proving an elliptic Harnack inequality for the nonnegative harmonic functions of  $X$ . The study of elliptic Harnack inequality for purely discontinuous processes was initiated only recently by Bass and Levin in [1] whose approach was also used in [7].

Parabolic Harnack inequality for nonnegative parabolic functions of purely discontinuous symmetric Markov processes was established by Chen and Kumagai in [4] based on the ideas developed in [2]. The processes they studied have a scaling property that was essentially used in their argument. In a work in progress Chen and Kumagai were able to extend the parabolic Harnack inequality to a more general class of purely discontinuous symmetric Markov processes including sums of independent stable processes with different scales. Their work so far does not include the process  $X$  described in the paragraph above.

The goal of this paper is to establish a parabolic Harnack inequality for the nonnegative parabolic functions of the process  $X$ . In order to do this, we first establish sharp upper and lower bounds for the transition density of  $X$ . Although our proof of these bounds is elementary

---

2000 *Mathematics Subject Classification.* Primary 60J45; Secondary 60J25, 60J35, 60J75.

*Key words and phrases.* Parabolic Harnack inequality, parabolic functions, Brownian motion, stable processes, transition density.

\*The research supported in part by a joint US-Croatia grant INT 0302167.

\*\*The research supported in part by MZOS grant 0037107 of the Republic of Croatia and in part by a joint US-Croatia grant INT 0302167.

and does not extend to general Markov processes which have both a continuous component and a discontinuous component, these bounds can serve as guidelines for the general case.

The content of this paper is organized as follows. The upper and lower bounds on the density of  $X$  are established in Section 2. In Section 3 we establish a lower bound for the transition density of the process  $X$  killed upon exiting a ball, and in Section 4 we prove the parabolic Harnack inequality.

**2. Bounds for transition densities of the mixture.** Let  $p^{(2)}(t, x)$  be the transition density of  $W$ , and  $p^{(\alpha)}(t, x)$  the transition density of  $Y$ . Then

$$p^{(2)}(t, x) = (4\pi t)^{-d/2} \exp\left(-\frac{|x|^2}{4t}\right),$$

while it follows from [3] that there are positive constants  $C_1, C_2$  such that for all  $t > 0$  and  $x \in \mathbf{R}^d$ ,

$$(1) \quad C_1 \min(t^{-d/\alpha}, t|x|^{-d-\alpha}) \leq p^{(\alpha)}(t, x) \leq C_2 \min(t^{-d/\alpha}, t|x|^{-d-\alpha}).$$

The transition density  $p(t, x)$  of  $X$  is given by

$$p(t, x) := \int_{\mathbf{R}^d} p^{(2)}(t, x-y) p^{(\alpha)}(t, y) dy.$$

The purpose of this section is to obtain sharp bounds on  $p(t, x)$ . In order to do this, we will need to compare  $p^{(2)}(t, x)$  and  $p^{(\alpha)}(t, x)$ .

LEMMA 2.1. *Let  $\gamma > 0$ .*

(i) *There exists a positive constant  $c > 0$  such that for all  $x \in \mathbf{R}^d$  and all  $t > 0$  satisfying  $|x| \leq 1 \leq t$ , it holds that*

$$t^{-d/\alpha} \leq ct^{-d/2} \exp\left(-\frac{|x|^2}{\gamma t}\right).$$

(ii) *For all  $x \in \mathbf{R}^d$  and all  $t \in (0, 1)$ , it holds that*

$$t^{-d/\alpha} \geq t^{-d/2} \exp\left(-\frac{|x|^2}{\gamma t}\right).$$

(iii) *There exists a positive constant  $c > 0$  such that for all  $t > 0$  and all  $|x| \geq 1$ , it holds that*

$$(2) \quad t^{-d/2} \exp\left(-\frac{|x|^2}{\gamma t}\right) \leq ct|x|^{-d-\alpha}.$$

PROOF. We omit the easy proofs of (i) and (ii), and only give a proof of (iii).

For fixed  $x \neq 0$ , define  $f : (0, \infty) \rightarrow (0, \infty)$  by

$$f(t) := t^{-d/2-1} \exp(-|x|^2/(\gamma t)).$$

Then  $f(0+) = f(+\infty) = 0$ . Further,

$$f'(t) = f(t)t^{-2}(-(d/2 + 1)t + |x|^2/\gamma).$$

This derivative is zero for

$$t_0 = \frac{|x|^2}{(d/2 + 1)\gamma},$$

positive for  $t < t_0$ , and negative for  $t > t_0$ . Thus  $f$  attains its maximum value at  $t_0$ , and

$$\begin{aligned} \max f = f(t_0) &= \left( \frac{|x|^2}{(d/2 + 1)\gamma} \right)^{-d/2-1} \exp\left( -\frac{|x|^2}{\gamma} \frac{(d/2 + 1)\gamma}{|x|^2} \right) \\ &= ((d/2 + 1)\gamma)^{d/2+1} \exp(-d/2 - 1) |x|^{-d-2} = c|x|^{-d-2}. \end{aligned}$$

It follows that for all  $t > 0$

$$t^{-d/2} \exp(-|x|^2/(\gamma t)) \leq t f(t) \leq t c |x|^{-d-2} = c t |x|^{-d-\alpha} |x|^{\alpha-2} \leq c t |x|^{-d-\alpha}$$

since  $|x| \geq 1$ .  $\square$

REMARK 2.2. Note that the proof of (iii) shows that for  $|x| < 1$  there does not exist a positive constant  $c$  independent of  $x$  such that (2) holds. Clearly, the reverse inequality cannot be true either.

Now we will establish upper bounds for  $p(t, x)$ .

LEMMA 2.3. *There exists a positive constant  $c$  such that for  $t \geq |x|^\alpha$ ,*

$$p(t, x) \leq c p^{(\alpha)}(t, x).$$

PROOF. For all  $y \in \mathbf{R}^d$ ,  $p^{(\alpha)}(t, y) \leq C_2 t^{-d/\alpha}$ . For  $t > |x|^\alpha$ , we have that  $p^{(\alpha)}(t, x) \geq C_1 t^{-d/\alpha}$ . Hence,

$$\begin{aligned} p(t, x) &= \int_{\mathbf{R}^d} p^{(2)}(t, x - y) p^{(\alpha)}(t, y) dy \\ &\leq C_2 t^{-d/\alpha} \int_{\mathbf{R}^d} p^{(2)}(t, x - y) dy \leq (C_2/C_1) p^{(\alpha)}(t, x). \end{aligned} \quad \square$$

LEMMA 2.4. *There exists a positive constant  $c$  such that for  $t \geq |x|^2$ ,*

$$p(t, x) \leq c p^{(2)}(t, x).$$

PROOF.

$$\begin{aligned} p(t, x) &= \int_{\mathbf{R}^d} p^{(2)}(t, x - y) p^{(\alpha)}(t, y) dy \\ &= p^{(2)}(t, x) \int_{\mathbf{R}^d} \frac{p^{(2)}(t, x - y)}{p^{(2)}(t, x)} p^{(\alpha)}(t, y) dy \\ &= p^{(2)}(t, x) \int_{\mathbf{R}^d} \exp\left( \frac{1}{4t} (|x|^2 - |x - y|^2) \right) p^{(\alpha)}(t, y) dy \\ &\leq p^{(2)}(t, x) \int_{\mathbf{R}^d} \exp(|x|^2/(4t)) p^{(\alpha)}(t, y) dy \\ &= p^{(2)}(t, x) \exp(|x|^2/(4t)) \leq e^{1/4} p^{(2)}(t, x). \end{aligned} \quad \square$$

LEMMA 2.5. *Let  $\tilde{p}^{(2)}(t, x) := (4\pi t)^{-d/2} \exp\{-|x|^2/(16t)\}$ . There exists a positive constant  $c$  such that for all  $|x| \leq 1$  and all  $t < |x|^2$ ,*

$$p(t, x) \leq C \max(\tilde{p}^{(2)}(t, x), p^{(\alpha)}(t, x)).$$

PROOF.

$$\begin{aligned} p(t, x) &= \int_{\mathbf{R}^d} p^{(2)}(t, x-y) p^{(\alpha)}(t, y) dy \\ &= \int_{|y| < \frac{t^{1/\alpha}}{2}} + \int_{\frac{t^{1/\alpha}}{2} \leq |y| < \frac{t^{1/2}}{2}} + \int_{|y| \geq \frac{t^{1/2}}{2}, |y-x| > |x|/2} + \int_{|y| \geq \frac{t^{1/2}}{2}, |y-x| \leq |x|/2} \\ &=: I_1 + I_2 + I_3 + I_4. \end{aligned}$$

(i) For  $|y| < t^{1/\alpha}/2$  and  $t^{1/2} < |x| \leq 1$ , it holds that  $2|y| < t^{1/\alpha} < t^{1/2} < |x|$ . Hence  $|x-y| > |x|/2$ , and so  $\exp\{-|x-y|^2/(4t)\} \leq \exp\{-|x|^2/(16t)\}$ . Clearly,  $p^{(\alpha)}(t, y) \leq C_2 t^{-d/\alpha}$ . Therefore,

$$\begin{aligned} I_1 &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t^{-d/\alpha} \int_{|y| < \frac{t^{1/\alpha}}{2}} dy \\ &= c_1 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)). \end{aligned}$$

(ii) For  $|y| < t^{1/2}/2$  we have that  $2|y| < t^{1/2} < |x|$ , and so again  $|x-y| > |x|/2$ . Clearly,  $p^{(\alpha)}(t, y) \leq C_2 t|y|^{-d-\alpha}$ . Therefore,

$$\begin{aligned} I_2 &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) \int_{\frac{t^{1/\alpha}}{2} \leq |y| < \frac{t^{1/2}}{2}} t|y|^{-d-\alpha} dy \\ &= c_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t \int_{\frac{t^{1/\alpha}}{2}}^{\frac{t^{1/2}}{2}} r^{d-1} r^{-d-\alpha} dr \\ &= c_3 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t ((t^{1/\alpha})^{-\alpha} - (t^{1/2})^{-\alpha}) \\ &= c_3 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) (1 - t^{1-\alpha/2}) \\ &\leq c_3 (4\pi t)^{-d/2} \exp\{-|x|^2/(16t)\}. \end{aligned}$$

(iii) Similarly as in (ii),

$$\begin{aligned} I_3 &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) \int_{|y| \geq \frac{t^{1/2}}{2}, |y-x| > |x|/2} t|y|^{-d-\alpha} dy \\ &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t \int_{|y| \geq \frac{t^{1/2}}{2}} |y|^{-d-\alpha} dy \\ &= c_4 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t (t^{1/2})^{-\alpha} \\ &\leq c_5 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)), \end{aligned}$$

since  $t^{1-\alpha/2} \leq 1$  for  $t \leq 1$ .

(iv) For  $|y - x| \leq |x|/2$ , we have  $|y| \geq |x|/2$ , and hence  $|y|^{-d-\alpha} \leq 2^{d+\alpha}|x|^{-d-\alpha}$ . Further, since  $t < |x|^2 < |x|^\alpha$  for  $|x| \leq 1$ , it holds that  $t|x|^{-d-\alpha} \leq (1/C_1)p^{(\alpha)}(t, x)$ . Thus

$$\begin{aligned} I_4 &\leq C_2 \int_{|y| \geq \frac{t^{1/2}}, |y-x| \leq |x|/2} p^{(2)}(t, x-y)t|y|^{-d-\alpha} dy \\ &\leq c_6 \int_{|y| \geq \frac{t^{1/2}}, |y-x| \leq |x|/2} p^{(2)}(t, x-y)t|x|^{-d-\alpha} dy \\ &\leq c_6 t|x|^{-d-\alpha} \int_{\mathbb{R}^d} p^{(2)}(t, x-y) dy \leq c_7 p^{(\alpha)}(t, x). \end{aligned}$$

From the estimates above it follows that

$$p(t, x) \leq c_8 \tilde{p}^{(2)}(t, x) + c_7 p^{(\alpha)}(t, x) \leq c_9 \max(\tilde{p}^{(2)}(t, x), p^{(\alpha)}(t, x)). \quad \square$$

LEMMA 2.6. *There exists a positive constant  $c$  such that for all  $t < |x|^\alpha$  and  $|x| \geq 1$ , it holds that*

$$p(t, x) \leq c p^{(\alpha)}(t, x).$$

PROOF.

$$\begin{aligned} p(t, x) &= \int_{\mathbb{R}^d} p^{(2)}(t, x-y)p^{(\alpha)}(t, y) dy \\ &= \int_{|x-y| \leq |x|/2} + \int_{|x-y| > |x|/2, |y| \geq t^{1/\alpha}} + \int_{|x-y| > |x|/2, |y| < t^{1/\alpha}} \\ &=: I_1 + I_2 + I_3. \end{aligned}$$

(i) For  $|x - y| \leq |x|/2$  we have  $|y| \geq |x|/2$ . Hence  $p^{(\alpha)}(t, y) \leq C_2 t|y|^{-d-\alpha} \leq C_2 2^{d+\alpha} t|x|^{-d-\alpha} = c_{10} t|x|^{-d-\alpha}$ . Also,  $p^{(\alpha)}(t, x) \geq C_1 t|x|^{-d-\alpha}$ . Therefore,

$$\begin{aligned} I_1 &\leq c_{10} t|x|^{-d-\alpha} \int_{|x-y| \leq |x|/2} p^{(2)}(t, x-y) dy \\ &\leq c_{10} t|x|^{-d-\alpha} \leq c_{11} p^{(\alpha)}(t, x). \end{aligned}$$

(ii) For  $|x - y| \geq |x|/2$  we have  $\exp\{-|x - y|^2/(4t)\} \leq \exp(-|x|^2/(16t))$ . Also,  $p^{(\alpha)}(t, y) \leq C_2 t|y|^{-d-\alpha}$ . Therefore,

$$\begin{aligned} I_2 &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) \int_{|x-y| \geq |x|/2, |y| \geq t^{1/\alpha}} t|y|^{-d-\alpha} dy \\ &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t \int_{|y| \geq t^{1/\alpha}} |y|^{-d-\alpha} dy \\ &\leq c_{12} (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) t (t^{1/\alpha})^{-\alpha} \\ &= c_{13} (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) \leq c_{14} p^{(\alpha)}(t, x), \end{aligned}$$

where in the last inequality we used Lemma 2.1 (iii).

(iii) For  $|x - y| \geq |x|/2$  we have  $\exp(-|x - y|^2/(4t)) \leq \exp\{-|x|^2/(16t)\}$ . Also,  $p^{(\alpha)}(t, y) \leq C_2 t^{-d/\alpha}$ . Therefore,

$$\begin{aligned} I_3 &\leq C_2 (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) \int_{|y| < t^{1/\alpha}} t^{-d/\alpha} dy \\ &\leq c_{15} (4\pi t)^{-d/2} \exp(-|x|^2/(16t)) \leq c_{16} p^{(\alpha)}(t, x), \end{aligned}$$

where in the last inequality we used Lemma 2.1 (iii).

From the estimates above it follows that  $p(t, x) \leq c_{17} p^{(\alpha)}(t, x)$ .  $\square$

REMARK 2.7. Suppose that  $t < |x|^\alpha$  and  $|x| \geq R$ , where  $0 < R < 1$ . Then  $p(t, x) \leq c R^{\alpha-2} p^{(\alpha)}(t, x)$ , where the constant  $c$  does not depend on  $R$ . This can be proved by changing the estimates for  $I_2$  and  $I_3$ , by using a modification of Lemma 2.1 (iii).

Next we establish lower bounds for  $p(t, x)$ .

LEMMA 2.8. Let  $\hat{p}^{(2)}(t, x) := (4\pi t)^{-d/2} \exp(-|x|^2/t)$ . There exists a positive constant  $c$  such that for all  $t \leq |x|^\alpha$ ,

$$p(t, x) \geq c \hat{p}^{(2)}(t, x).$$

PROOF. For  $|y| \leq |x|$  we have that  $|y - x| \leq 2|x|$ , and hence  $\exp(-|x - y|^2/(4t)) \geq \exp(-|x|^2/t)$ . Therefore,

$$\begin{aligned} p(t, x) &\geq \int_{B(0, |x|)} p^{(2)}(t, x - y) p^{(\alpha)}(t, y) dy \\ &\geq (4\pi t)^{-d/2} \exp(-|x|^2/t) \int_{B(0, |x|)} p^{(\alpha)}(t, y) dy \\ &= (4\pi t)^{-d/2} \exp(-|x|^2/t) \int_{B(0, |x|)} t^{-d/\alpha} p^{(\alpha)}(1, t^{-1/\alpha} y) dy \\ &= (4\pi t)^{-d/2} \exp(-|x|^2/t) \int_{B(0, t^{-1/\alpha} |x|)} p^{(\alpha)}(1, u) du \\ &\geq (4\pi t)^{-d/2} \exp(-|x|^2/t) \int_{B(0, 1)} p^{(\alpha)}(1, u) du \\ &= c_1 (4\pi t)^{-d/2} \exp(-|x|^2/t). \quad \square \end{aligned}$$

LEMMA 2.9. There exists a positive constant  $c$  such that for every  $x \in \mathbf{R}^d$  and every  $y \in B(x, |x|/2)$ , it holds that

$$\frac{p^{(\alpha)}(t, y)}{p^{(\alpha)}(t, x)} \geq c.$$

PROOF. This result can be easily proved by looking at the following four cases. Case 1:  $t > |x|^\alpha, t > |y|^\alpha$ ; Case 2:  $t \leq |x|^\alpha, t \leq |y|^\alpha$ ; Case 3:  $t > |x|^\alpha, t \leq |y|^\alpha$  and Case 4:  $t \leq |x|^\alpha, t > |y|^\alpha$ . We omit the details.  $\square$

LEMMA 2.10. *There exists a positive constant  $c$  such that for all  $t \leq |x|^2$ ,*

$$p(t, x) \geq cp^{(\alpha)}(t, x).$$

PROOF.

$$\begin{aligned} p(t, x) &= \int_{\mathbf{R}^d} p^{(\alpha)}(t, y) p^{(2)}(t, x - y) dy \\ &= p^{(\alpha)}(t, x) \int_{\mathbf{R}^d} \frac{p^{(\alpha)}(t, y)}{p^{(\alpha)}(t, x)} p^{(2)}(t, x - y) dy \\ &\geq c_1 p^{(\alpha)}(t, x) \int_{B(x, |x|/2)} p^{(2)}(t, x - y) dy \\ &= c_1 p^{(\alpha)}(t, x) \int_{B(0, |x|/2)} p^{(2)}(t, y) dy \\ &= c_1 p^{(\alpha)}(t, x) \int_{B(0, t^{-1/2}|x|/2)} p^{(2)}(1, u) du \\ &\geq c_1 p^{(\alpha)}(t, x) \int_{B(0, 1/2)} p^{(2)}(1, u) du = c_2 p^{(\alpha)}(t, x), \end{aligned}$$

where the third line follows from Lemma 2.9.  $\square$

LEMMA 2.11. *There exists a positive constant  $c$  such that for all  $t \geq 1$  and all  $|x|^\alpha < t$  we have*

$$p(t, x) \geq cp^{(\alpha)}(t, x).$$

PROOF. For  $|x|^\alpha < t$ ,  $p^{(\alpha)}(t, x) \leq C_2 t^{-d/\alpha}$ . If  $|y - x| \leq t^{1/\alpha}$ , then  $|y| \leq |x - y| + |x| \leq 2t^{1/\alpha}$ . If  $|y| \leq t^{1/\alpha}$ , then  $p^{(\alpha)}(t, y) \geq C_1 t^{-d/\alpha}$ . If  $t^{1/\alpha} < |y| \leq 2t^{1/\alpha}$ , then  $p^{(\alpha)}(t, y) \geq C_1 t|y|^{-d-\alpha} \geq C_1 2^{-d-\alpha} t^{-d/\alpha}$ . Therefore,

$$\begin{aligned} p(t, x) &\geq \int_{|x-y| \leq t^{1/\alpha}} p^{(2)}(t, x - y) p^{(\alpha)}(t, y) dy \\ &\geq c_3 t^{-d/\alpha} \int_{|x-y| \leq t^{1/\alpha}} p^{(2)}(t, x - y) dy \\ &= c_3 t^{-d/\alpha} \int_{|y| \leq t^{1/\alpha}} p^{(2)}(t, y) dy = c_3 t^{-d/\alpha} \int_{|y| \leq t^{-1/2} t^{1/\alpha}} p^{(2)}(1, u) du \\ &\geq c_3 t^{-d/\alpha} \int_{|y| \leq 1} p^{(2)}(1, u) du = c_4 t^{-d/\alpha} \geq c_5 p^{(\alpha)}(t, x). \quad \square \end{aligned}$$

LEMMA 2.12. *There exists a positive constant  $c$  such that for all  $|x|^\alpha \leq t \leq 1$  it holds that*

$$p(t, x) \geq cp^{(2)}(t, x).$$

PROOF. If  $|y| < t^{1/\alpha}$ , then  $p^{(\alpha)}(t, y) \geq C_1 t^{-d/\alpha}$ . Also,  $|x - y| \leq |x| + |y| \leq 2t^{1/\alpha}$  implying  $\exp(-|x - y|^2/(4t)) \geq \exp(-4t^{2/\alpha}/(4t)) = \exp(-t^{2/\alpha-1}) \geq e^{-1}$ , since  $t^{2/\alpha-1} \leq 1$

for  $t \leq 1$ . Therefore,

$$\begin{aligned}
p(t, x) &\geq \int_{|y| < t^{1/\alpha}} p^{(2)}(t, x - y) p^{(\alpha)}(t, y) dy \\
&\geq C_1 t^{-d/\alpha} \int_{|y| < t^{1/\alpha}} p^{(2)}(t, x - y) dy \\
&\geq C_1 t^{-d/\alpha} (4\pi t)^{-d/2} e^{-1} \int_{|y| < t^{1/\alpha}} dy \\
&= c_6 (4\pi t)^{-d/2} \geq c_6 (4\pi t)^{-d/2} \exp(-|x|^2/(4t)) \\
&= c_6 p^{(2)}(t, x). \quad \square
\end{aligned}$$

By collecting the results from previous lemmas, we obtain the lower and upper bounds for the transition density  $p(t, x)$ . In order to briefly state the result, we define

$$q_1(t, x) = \begin{cases} \hat{p}^{(2)}(t, x), & |x|^2 < t < |x|^\alpha \leq 1, \\ \max(\hat{p}^{(2)}(t, x), p^{(\alpha)}(t, x)), & t < |x|^2 \leq 1, \\ p^{(2)}(t, x), & |x|^\alpha \leq t \leq 1, \\ p^{(\alpha)}(t, x), & t \geq 1 \text{ or } |x| \geq 1, \end{cases}$$

and

$$q_2(t, x) = \begin{cases} p^{(2)}(t, x), & |x|^2 < t < |x|^\alpha \leq 1, \\ \max(\tilde{p}^{(2)}(t, x), p^{(\alpha)}(t, x)), & t < |x|^2 \leq 1, \\ p^{(2)}(t, x), & |x|^\alpha \leq t \leq 1, \\ p^{(\alpha)}(t, x), & t \geq 1 \text{ or } |x| \geq 1. \end{cases}$$

**THEOREM 2.13.** *There exists a positive constant  $C_3$  such that*

$$C_3^{-1} q_1(t, x) \leq p(t, x) \leq C_3 q_2(t, x).$$

**3. Lower bounds for transition densities of the killed process.** In this section we will establish a lower bound for the transition density of the process  $X$  killed upon exiting a ball of radius  $R$ . Let  $p(t, x, y) := p(t, y - x)$ . We first need the following lemma.

**LEMMA 3.1.** *There exists a constant  $C_4 > 0$  such that for every  $R > 0$ , every  $x \in \mathbf{R}^d$  and every  $t > 0$ ,*

$$(3) \quad \mathbf{P}^x(\tau_{B(x, R)} \leq t) \leq \frac{C_4 t}{R^2 \wedge R^\alpha},$$

where  $\tau_{B(x, R)} = \inf\{s > 0; X_t \notin B(x, R)\}$ .

This result for  $R \in (0, 1]$  appears as Lemma 2.1 in [8]. By a slight modification of the proof, the result follows for  $R \geq 1$  as well.

Let  $R > 0$ ,  $B = B(0, R)$ , and let  $\tau_B$  denote the first exit time of  $X$  from  $B$ . Let  $X^B$  denote the process  $X$  killed upon exiting  $B$ . The transition density of  $X^B$  is given by

$$p^B(t, x, y) = p(t, x, y) - r^B(t, x, y), \quad x, y \in B,$$

where

$$r^B(t, x, y) = \mathbf{E}^x[p(t - \tau_B, X_{\tau_B}, y) \mathbf{1}_{(t > \tau_B)}].$$



LEMMA 3.2. *There exist constants  $C_5 > 0$  and  $C_6 \in (0, 1/10)$  such that:*

(i) *For every  $R > 0$ , for all  $x, y \in B(0, 2R/3)$  and all  $0 < t < 1$  satisfying  $|x - y|^2 < t \leq C_6(R^2 \wedge R^\alpha)$  it holds that*

$$p^B(t, x, y) \geq C_5 t^{-d/2}.$$

(ii) *For every  $R \geq 1$ , for all  $x, y \in B(0, 2R/3)$  and all  $t \geq 1$  satisfying  $|x - y|^\alpha < t \leq C_6 R^\alpha$  it holds that*

$$p^B(t, x, y) \geq C_5 t^{-d/\alpha}.$$

(iii) *For every  $R \geq 1$ , for all  $x, y \in B(0, 2R/3)$  and all  $t \geq 1$  satisfying  $|x - y|^\alpha \geq t$  and  $t \leq C_6 R^\alpha$  it holds that*

$$p^B(t, x, y) \geq C_5 t R^{-d-\alpha}.$$

PROOF. We first find an upper bound for  $r^B(t, x, y)$ . Suppose that  $0 < R \leq 1$ . Note that by combining Lemma 2.6 with Remark 2.7, if  $|y - z| > R/3$  and  $t < |y - z|^2$ , then  $p(t, z, y) \leq c(R/3)^{\alpha-2} p^{(\alpha)}(t, z, y) \leq cC_2(R/3)^{\alpha-2} t |z - y|^{-d-\alpha}$ . Let  $x, y \in B(0, 2R/3)$  and choose  $t \leq R^2/10$ . Then  $|X_{\tau_B} - y| > R/3$  and  $t \leq R^2/10 < |X_{\tau_B} - y|^2$ , so on  $\{t > \tau_B\}$

$$\begin{aligned} p(t - \tau_B, X_{\tau_B}, y) &\leq cC_2(R/3)^{\alpha-2} (t - \tau_B) |X_{\tau_B} - y|^{-d-\alpha} \\ &\leq cC_2(R/3)^{\alpha-2} t (R/3)^{-d-\alpha} \leq c_1 R^{-d-2} t. \end{aligned}$$

Note further that for  $x \in B(0, 2R/3)$  it holds that  $\mathbf{P}^x(\tau_B < t) \leq \mathbf{P}^x(\tau_{B(x, R/3)} < t) \leq C_4 t / (R/3)^2$  by Lemma 3.1. Therefore, for all  $x, y \in B(0, 2R/3)$  and all  $t \leq R^2/10$ ,

$$\begin{aligned} r^B(t, x, y) &= \mathbf{E}^x[p(t - \tau_B, X_{\tau_B}, y) 1_{\{t > \tau_B\}}] \\ &\leq c_1 R^{-d-2} t \mathbf{P}^x(t > \tau_B) \\ &\leq 9c_1 R^{-d-2} t C_4 t R^{-2} = c_2 t^2 R^{-d-4}. \end{aligned}$$

Suppose now that  $R \geq 1$ . If  $1 > |y - z| \geq R/3 \geq 1/3$ , and  $t < |y - z|^\alpha$ , it holds by Remark 2.7 that  $p(t, y, z) \leq c(1/3)^{\alpha-2} p^{(\alpha)}(t, y, z)$ . If  $1 > |y - z| \geq R/3 \geq 1/3$ , and  $t \geq |y - z|^\alpha$ , then by Lemma 2.3,  $p(t, y, z) \leq c p^{(\alpha)}(t, y, z)$ . If  $|y - z| \geq 1$ , then the estimate  $p(t, y, z) \leq c p^{(\alpha)}(t, y, z)$  follows from Theorem 2.13. Therefore, whenever  $|y - z| \geq R/3$  it holds that for all  $t > 0$ ,  $p(t, y, z) \leq c p^{(\alpha)}(t, y, z) \leq cC_2 t |y - z|^{-\alpha-d} \leq cC_2 t (R/3)^{-\alpha-d}$ . Let  $x, y \in B(0, 2R/3)$ . Then on  $\{t > \tau_B\}$

$$p(t - \tau_B, X_{\tau_B}, y) \leq cC_2 t (R/3)^{-d-\alpha} \leq c_3 R^{-d-\alpha} t.$$

Again by Lemma 3.1,  $\mathbf{P}^x(\tau_B < t) \leq \mathbf{P}^x(\tau_{B(x, R/3)} < t) \leq C_4 t / (R/3)^\alpha$ . It follows that

$$\begin{aligned} r^B(t, x, y) &= \mathbf{E}^x[p(t - \tau_B, X_{\tau_B}, y) 1_{\{t > \tau_B\}}] \\ &\leq c_3 R^{-d-\alpha} t \mathbf{P}^x(t > \tau_B) \\ &\leq 9c_3 R^{-d-\alpha} t C_4 t R^{-\alpha} = c_4 t^2 R^{-d-2\alpha}. \end{aligned}$$

(i) Suppose first that  $0 < t < 1$  and  $R \in (0, 1]$ . For all  $x, y \in \mathbf{R}^d$  such that  $|x - y| \leq 1$ , we have by Theorem 2.13 that  $p(t, x, y) \geq c_5 \hat{p}^{(2)}(t, y - x)$ . Therefore, for  $x, y \in B(0, 2R/3)$

and  $|x - y|^2 \leq t \leq R^2/10$  it follows that

$$p(t, x, y) \geq c_5(4\pi t)^{-d/2} e^{-|x-y|^2/t} \geq c_6 e^{-1} t^{-d/2} = c_7 t^{-d/2}.$$

It follows that for  $x, y \in B(0, 2R/3)$  and  $|x - y|^2 \leq t \leq R^2/10$

$$\begin{aligned} p^B(t, x, y) &= p(t, x, y) - r^B(t, x, y) \\ &\geq c_7 t^{-d/2} - c_2 t^2 R^{-d-4} \\ &= c_7 t^{-d/2} \left( 1 - \frac{c_2}{c_7} t^{(d+4)/2} R^{-d-4} \right) \geq \frac{c_7}{2} t^{-d/2} \end{aligned}$$

provided that  $1 - (c_2/c_7)t^{(d+4)/2}R^{-d-4} > 1/2$ . This last condition is satisfied if

$$t < \left( \frac{c_7}{2c_2} \right)^{2/(d+4)} R^2 = c_8 R^2.$$

Suppose now that  $0 < t < 1$  and  $R \geq 1$ . The same argument as above shows that for  $x, y \in B(0, 2R/3)$  and  $|x - y|^2 \leq t \leq R^2/10$  it holds that  $p(t, x, y) \geq c_7 t^{-d/2}$ , and consequently

$$\begin{aligned} p^B(t, x, y) &= p(t, x, y) - r^B(t, x, y) \\ &\geq c_7 t^{-d/2} - c_2 t^2 R^{-d-2\alpha} \\ &= c_7 t^{-d/2} \left( 1 - \frac{c_2}{c_7} t^{(d+4)/2} R^{-d-2\alpha} \right) \geq \frac{c_7}{2} t^{-d/2} \end{aligned}$$

provided that  $1 - (c_2/c_7)t^{(d+4)/2}R^{-d-2\alpha} > 1/2$ . This last condition is satisfied if

$$t < \left( \frac{c_7}{2c_2} \right)^{2/(d+4)} R^{2(d+2\alpha)/(d+4)} = c_9 R^{2(d+2\alpha)/(d+4)}.$$

Note that  $\alpha < 2(d+2\alpha)/(d+4)$ . Therefore, if  $t < c_9 R^\alpha$ , then  $t < c_9 R^{2(d+2\alpha)/(d+4)}$ , and consequently  $p^B(t, x, y) \geq (c_7/2)t^{-d/2}$ .

Choose  $c_{11} = \min(1/10, c_8, c_9)$ . Then we have proved that for every  $R > 0$ , for all  $x, y \in B(0, 2R/3)$  and all  $0 < t < 1$  satisfying  $|x - y|^2 < t \leq c_{11}(R^2 \wedge R^\alpha)$  it holds that

$$p^B(t, x, y) \geq \frac{c_7}{2} t^{-d/2}.$$

(ii) Let  $R \geq 1$  and  $x, y \in B(0, 2R/3)$ . Suppose that  $|x - y|^\alpha < t$ . By Theorem 2.13,  $p(t, x, y) \geq C_3^{-1} p^{(\alpha)}(t, x, y) \geq C_3^{-1} C_2 \min(t^{-d/\alpha}, t|x - y|^{-d-\alpha}) \geq c_{12} t^{-d/\alpha}$ . By combining with the upper bound for  $r^B(t, x, y)$ , it follows that

$$\begin{aligned} p^B(t, x, y) &= p(t, x, y) - r^B(t, x, y) \\ &\geq c_{12} t^{-d/\alpha} - c_4 t^2 R^{-d-2\alpha} \\ &= c_{12} t^{-d/\alpha} \left( 1 - \frac{c_4}{c_{12}} t^{d/\alpha+2} R^{-d-2\alpha} \right) \geq \frac{c_{12}}{2} t^{-d/\alpha} \end{aligned}$$

provided that  $1 - (c_4/c_{12})t^{d/\alpha+2}R^{-d-2\alpha} > 1/2$ . This last condition is satisfied if

$$t < \left(\frac{c_{12}}{2c_4}\right)^{\alpha/(d+2\alpha)} R^\alpha = c_{13}R^\alpha.$$

(iii) Let  $R \geq 1$  and  $x, y \in B(0, 2R/3)$ . Suppose that  $|x - y|^\alpha \geq t$ . Again by Theorem 2.13,  $p(t, x, y) \geq C_3^{-1}p^{(\alpha)}(t, x, y) \geq C_3^{-1}C_2 \min(t^{-d/\alpha}, t|x - y|^{-d-\alpha}) = C_3^{-1}C_2t|x - y|^{-d-\alpha} \geq c_{14}tR^{-d-\alpha}$ . By combining with the upper bound for  $r^B(t, x, y)$ , it follows that

$$\begin{aligned} p^B(t, x, y) &= p(t, x, y) - r^B(t, x, y) \\ &\geq c_{14}tR^{-d-\alpha} - c_4t^2R^{-2\alpha-d} \\ &\geq c_{14}tR^{-d-\alpha} \left(1 - \frac{c_4}{c_{14}}tR^{-\alpha}\right) \\ &\geq \frac{c_{14}}{2}tR^{-d-\alpha} \end{aligned}$$

provided that  $1 - (c_4/c_{14})tR^{-\alpha} \geq 1/2$ . This is satisfied if

$$t < \frac{c_{14}}{2c_4}R^\alpha = c_{15}R^\alpha.$$

We finish the proof of the lemma by choosing  $C_5 = \min(c_5/2, c_{12}/2, c_{14}/2)$  and  $C_6 = \min(c_{11}, c_{13}, c_{15})$ .  $\square$

Let  $N = \lfloor 2/C_6 \rfloor$  where  $C_6$  is the constant from Lemma 3.2, and  $\lfloor \cdot \rfloor$  denotes the smallest integer function. The proof of the next result follows the proof of Theorem 2.7 in [5].

**PROPOSITION 3.3.** *Let  $\delta \in (0, 1)$ . There exists a constant  $C_7 = C_7(\delta) > 0$  such that for all  $0 < R \leq (2N/\delta)^{1/\alpha}$ , all  $x, y \in B(0, R/2)$  and all  $0 < t \leq R^2 \wedge R^\alpha$  it holds that*

$$p^B(t, x, y) \geq C_7t^{-d/2}e^{-|x-y|^2/(C_7t)}.$$

**PROOF.** Let  $R \leq (2N/\delta)^{1/\alpha}$ ,  $x, y \in B(0, R/2)$  and  $t \leq R^2 \wedge R^\alpha$ . Suppose first that  $t < |x - y|^2$  and define  $k = \lfloor 4N|x - y|^2/(\delta t) \rfloor$ . Then  $k \geq 3N/\delta \geq 4/(C_6\delta)$ , and therefore  $t/k \leq (C_6\delta t)/2 \leq (1/2)C_6\delta(R^2 \wedge R^\alpha) \leq C_6(R^2 \wedge R^\alpha)$ . Moreover,

$$\frac{C_6\delta}{4}(R^2 \wedge R^\alpha) \leq \frac{C_6\delta}{4} \left[ \left( \frac{2N}{\delta} \right)^{1/\alpha} \right]^\alpha \leq \frac{C_6\delta}{2} \frac{N}{\delta} \leq 1,$$

implying that  $t/k \leq 1$ . For  $l = 1, 2, \dots, k-1$  let

$$z_l = x + \frac{l}{k}(y - x).$$

From  $k \geq 2N|x - y|^2/(\delta t)$  it follows that  $|x - y|^2 \leq \delta kt/(2N)$ . Therefore

$$|z_l - z_{l-1}| = \frac{|x - y|}{k} \leq \sqrt{\frac{\delta}{2}} \frac{\sqrt{kt/N}}{k} = \sqrt{\frac{\delta}{2N}} \sqrt{\frac{t}{k}}.$$

Define

$$S = \prod_{l=1}^{k-1} B\left(z_l, \sqrt{\frac{\delta}{2N}} \sqrt{\frac{t}{k}}\right).$$

Note that

$$\sqrt{\frac{\delta}{2N}}\sqrt{\frac{t}{k}} \leq \sqrt{\frac{\delta}{2N}}\sqrt{C_6 R^2} \leq C_6 R \leq \frac{1}{10}R,$$

implying that for every  $l = 1, 2, \dots, k-1$ ,

$$B\left(z_l, \sqrt{\frac{\delta}{2N}}\sqrt{\frac{t}{k}}\right) \subset B(0, 2R/3).$$

For  $\zeta_l \in B(z_l, \sqrt{\delta/(2N)}\sqrt{t/k})$  and  $\zeta_{l-1} \in B(z_{l-1}, \sqrt{\delta/(2N)}\sqrt{t/k})$ ,  $l = 2, 3, \dots, k-1$ , we have that  $|\zeta_l - \zeta_{l-1}| \leq |\zeta_l - z_l| + |z_l - z_{l-1}| + |z_{l-1} - \zeta_{l-1}| \leq 3\sqrt{\delta/(2N)}\sqrt{t/k} \leq \sqrt{t/k}$ . Therefore,

$$|\zeta_l - \zeta_{l-1}|^2 \leq \frac{t}{k} \leq C_6(R^2 \wedge R^\alpha),$$

implying by Lemma 3.2(i) that  $p^B(t/k, \zeta_{l-1}, \zeta_l) \geq C_5(t/k)^{-d/2}$ . Hence

$$\begin{aligned} p^B(t, x, y) &= \int_B \int_B \dots \int_B p^B\left(\frac{t}{k}, x, \zeta_1\right) p^B\left(\frac{t}{k}, \zeta_1, \zeta_2\right) \dots p^B\left(\frac{t}{k}, \zeta_{k-1}, y\right) d\zeta_1 d\zeta_2 \dots d\zeta_{k-1} \\ &\geq \int \dots \int_S p^B\left(\frac{t}{k}, x, \zeta_1\right) p^B\left(\frac{t}{k}, \zeta_1, \zeta_2\right) \dots p^B\left(\frac{t}{k}, \zeta_{k-1}, y\right) d\zeta_1 d\zeta_2 \dots d\zeta_{k-1} \\ &\geq |S| \left(C_5 \left(\frac{t}{k}\right)^{-d/2}\right)^k = \left(|B(0, 1)| \left(\frac{\delta t}{2Nk}\right)^{d/2}\right)^{k-1} \left(C_5 \left(\frac{t}{k}\right)^{-d/2}\right)^k \\ &= \left(\frac{C_5 \delta^{d/2} |B(0, 1)|}{2^{d/2} N^{d/2}}\right)^k \frac{2^{d/2} N^{d/2}}{\delta^{d/2} |B(0, 1)|} k^{d/2} t^{-d/2} \\ &\geq \exp\left(-k \log\left(\frac{2^{d/2} N^{d/2}}{C_5 \delta^{d/2} |B(0, 1)|}\right)\right) \frac{2^{d/2} N^{d/2}}{\delta^{d/2} |B(0, 1)|} N^{d/2} t^{-d/2} \\ &\geq c_{16} \exp\left(-\log\left(\frac{2^{d/2} N^{d/2}}{C_5 \delta^{d/2} |B(0, 1)|}\right) \frac{4N|x-y|^2}{\delta t}\right) t^{-d/2} \\ &\geq c_{17} t^{-d/2} \exp\left(-\frac{|x-y|^2}{c_{17} t}\right). \end{aligned}$$

Assume now that  $t \geq |x-y|^2$  and define  $k = \lfloor 4N/\delta \rfloor$ . Then again  $k \geq 3N/\delta \geq 4/(C_6\delta)$  implying  $t/k \leq C_6 t \leq C_6(R^2 \wedge R^\alpha)$  and  $t/k \leq 1$ . The same argument as above gives the following estimate

$$\begin{aligned} p^B(t, x, y) &\geq \left(\frac{C_5 \delta^{d/2} |B(0, 1)|}{2^{d/2} N^{d/2}}\right)^k \frac{2^{d/2} N^{d/2}}{\delta^{d/2} |B(0, 1)|} k^{d/2} t^{-d/2} \\ &\geq \left(\frac{C_5 \delta^{d/2} |B(0, 1)|}{2^{d/2} N^{d/2}}\right)^{2N} \frac{2^{d/2} N^{d/2}}{\delta^{d/2} |B(0, 1)|} (2N)^{d/2} t^{-d/2} \\ &= c_{18} t^{-d/2} \geq c_{18} t^{-d/2} \exp\left(-\frac{|x-y|^2}{c_{18} t}\right). \quad \square \end{aligned}$$

The claim follows by taking  $C_7 = \min(c_{17}, c_{18})$ .

PROPOSITION 3.4. *Let  $\delta \in (0, 1)$ . There exists a constant  $C_8 = C_8(\delta) > 0$  such that for all  $R \geq (2N/\delta)^{1/\alpha}$ , all  $x, y \in B(0, R/2)$  and all  $t$  satisfying  $\delta R^\alpha < t < R^\alpha$  it holds that*

$$p^B(t, x, y) \geq C_8 t^{-d/\alpha}.$$

PROOF. Let  $R \geq (2N/\delta)^{1/\alpha}$  and  $\delta R^\alpha < t < R^\alpha$ . First note that  $\delta R^\alpha \geq 2N$ , implying  $t \geq 2N$ . Define  $k = N$ . Then clearly  $t/k = t/N \geq 1$ . Also, since  $N > 1/C_6$ , we have that  $t/k \leq C_6 t \leq C_6 R^\alpha$ .

Suppose that  $u, v \in B(0, R/2)$ . If  $|u - v|^\alpha \leq t/k$ , then by Lemma 3.2(ii) it follows that

$$p^B(t/k, u, v) \geq C_5 \left(\frac{t}{k}\right)^{-d/\alpha} \geq C_5 t^{-d/\alpha}.$$

If  $|u - v|^\alpha \geq t/k$ , then by Lemma 3.2(iii) it follows that  $p^B(t/k, u, v) \geq C_5(t/k)R^{-d-\alpha}$ . But since  $\delta R^\alpha < t$ , we have that  $R < (t/\delta)^{1/\alpha}$ , implying

$$p^B(t/k, u, v) \geq C_5 \frac{t}{k} \left(\frac{t}{\delta}\right)^{-(d+\alpha)/\alpha} = C_5 \frac{\delta^{1+d/\alpha}}{N} t^{-d/\alpha} = c_{19} t^{-d/\alpha}.$$

Hence, for any  $u, v \in B(0, R/2)$  it holds that  $p^B(t/k, u, v) \geq c_{19} t^{-d/\alpha}$ . Define

$$S = \prod_{l=1}^{k-1} B\left(0, \frac{t^{1/\alpha}}{k}\right).$$

Note that  $B(0, t^{1/\alpha}/k) \subset B(0, R/2)$  since  $t < R^\alpha$ . Hence

$$\begin{aligned} p^B(t, x, y) &= \int_B \int_B \dots \int_B p^B\left(\frac{t}{k}, x, \zeta_1\right) p^B\left(\frac{t}{k}, \zeta_1, \zeta_2\right) \dots p^B\left(\frac{t}{k}, \zeta_{k-1}, y\right) d\zeta_1 d\zeta_2 \dots d\zeta_{k-1} \\ &\geq \int \dots \int_S p^B\left(\frac{t}{k}, x, \zeta_1\right) p^B\left(\frac{t}{k}, \zeta_1, \zeta_2\right) \dots p^B\left(\frac{t}{k}, \zeta_{k-1}, y\right) d\zeta_1 d\zeta_2 \dots d\zeta_{k-1} \\ &\geq |S| \left(c_{19} \left(\frac{t}{k}\right)^{-d/\alpha}\right)^k = \left(|B(0, 1)| \left(\frac{t^{1/\alpha}}{k}\right)^d\right)^{k-1} \left(c_{19} \left(\frac{t}{k}\right)^{-d/\alpha}\right)^k \\ &= |B(0, 1)|^{k-1} c_{19}^k k^{d(k(1/\alpha-1)+d)} t^{-d/\alpha} = c_{20} t^{-d/\alpha}, \end{aligned}$$

where  $c_{20}$  depends on  $\delta$ . Choose  $C_8 = c_{20}$ .  $\square$

COROLLARY 3.5. *Let  $\delta \in (0, 1)$ . There exists a constant  $C_9 = C_9(\delta) > 0$  such that for all  $R > 0$ , all  $x, y \in B(0, R/2)$  and all  $t \in (\delta(R^2 \wedge R^\alpha), R^2 \wedge R^\alpha)$  it holds that*

$$p^B(t, x, y) \geq \frac{C_9}{R^d}.$$

PROOF. Suppose first that  $R \leq 1$ . By Proposition 3.3,

$$p^B(t, x, y) \geq C_7 t^{-d/2} e^{-|x-y|^2/(C_7 t)}.$$

We use that  $|x - y|^2/t \leq 1/\delta$  and  $t^{-d/2} \geq R^{-d}$  to obtain the estimate with  $c_{21} = C_7 \exp(-1/(C_7 \delta))$ . If  $1 \leq R \leq (2N/\delta)^{1/\alpha}$ , then again by Proposition 3.3,

$$p^B(t, x, y) \geq C_7 t^{-d/2} e^{-|x-y|^2/(C_7 t)}.$$

Now we use the estimate

$$\frac{|x - y|^2}{t} \leq \frac{R^2}{\delta R^\alpha} \leq \frac{1}{\delta} \left( \frac{2N}{\delta} \right)^{(2-\alpha)/\alpha} = c_{22},$$

and  $t \leq R^\alpha \leq R^2$ , to obtain that

$$p^B(t, x, y) \geq C_7 e^{-c_{22}/C_7} R^{-d} = c_{23} R^{-d}.$$

Finally, let  $R \geq (2N/\delta)^{1/\alpha}$ . Then by Proposition 3.4 it holds that

$$p^B(t, x, y) \geq C_8 t^{-d/\alpha} \geq C_8 R^{-d}.$$

The proof is finished by choosing  $C_9 = \min(c_{21}, c_{22}, C_8)$ .  $\square$

**4. Parabolic Harnack inequality.** In this section we are going to prove the parabolic Harnack inequality following closely the approach from [4]. Let us first introduce the space-time process  $Z_s = (T_0 + s, X_s)$ . The law of the space-time process starting from  $(t, x) \in [0, \infty) \times \mathbf{R}^d$  will be denoted by  $\mathbf{P}^{(t,x)}$ .

DEFINITION 4.1. Let  $(t, x) \in [0, \infty) \times \mathbf{R}^d$  and let  $r_1, r_2 > 0$ . We say that a nonnegative function  $q : [0, \infty) \times \mathbf{R}^d \rightarrow [0, \infty)$  is *parabolic* in  $[t, t + r_1] \times B(x, r_2)$  if for any  $[s_1, s_2] \subset [t, t + r_1)$  and  $B(y, r) \subset \overline{B(y, r)} \subset B(x, r_2)$  we have

$$q(s, z) = \mathbf{E}^{(s,z)}(q(Z_{\tau_{[s_1, s_2] \times B(y, r)}})), \quad (s, z) \in [s_1, s_2] \times B(y, r),$$

where  $\tau_{[s_1, s_2] \times B(y, r)} = \inf\{s; Z_s \notin [s_1, s_2] \times B(y, r)\}$ .

For  $t \geq 0, x \in \mathbf{R}^d$  and  $R > 0$ , define

$$Q(t, x, R) = [t, t + (R^2 \wedge R^\alpha)] \times B(x, R).$$

For  $A \subset [0, \infty) \times \mathbf{R}^d$ , let  $\sigma_A = \inf\{t > 0; Z_t \in A\}$  and  $A_s = \{y \in \mathbf{R}^d; (s, y) \in A\}$ .

The idea for the proof of the next result comes from [6].

LEMMA 4.2. Let  $\delta \in (0, 1]$ . There exists a constant  $C_{10} = C_{10}(\delta) > 0$  such that for all  $R > 0$ , any  $z \in \mathbf{R}^d$ , any  $v \in B(z, R/3)$  and any  $A \subset Q(0, z, R/2) \cap ([\delta(R^2 \wedge R^\alpha), \infty) \times \mathbf{R}^d)$ ,

$$\mathbf{P}^{(0,v)}(\sigma_A < \tau_R) \geq C_{10} \frac{|A|}{R^d (R^2 \wedge R^\alpha)},$$

where  $\tau_R = \tau_{Q(0,z,R)}$ .

PROOF. We are going to estimate the expected time that the space-time process  $Z$  spends in  $A$  before exiting  $Q(0, z, R)$ . Let  $X^{B(z, R)}$  denote the process  $X$  killed upon exiting the ball  $B(z, R)$  and let  $p^{B(z, R)}$  be its transition density. Then

$$\begin{aligned}
 \mathbf{E}^{(0, v)} \int_0^{\tau_R} 1_A(s, X_s) ds &= \mathbf{E}^{(0, v)} \int_0^\infty 1_A(s, X_s^{B(z, R)}) ds \\
 &= \int_0^{R^2 \wedge R^\alpha} \mathbf{P}^{(0, v)}((s, X_s^{B(z, R)}) \in A) ds \\
 &= \int_0^{R^2 \wedge R^\alpha} \mathbf{P}^v(X_s^{B(z, R)} \in A_s) ds \\
 &= \int_{\delta(R^2 \wedge R^\alpha)}^{R^2 \wedge R^\alpha} \int_{A_s} p^{B(z, R)}(s, v, y) dy ds \\
 &\geq \int_{\delta(R^2 \wedge R^\alpha)}^{R^2 \wedge R^\alpha} \int_{A_s} \frac{C_9}{R^d} dy ds = C_9 \frac{|A|}{R^d},
 \end{aligned}$$

where the inequality follows from Corollary 3.5 by using that  $s \in (\delta(R^2 \wedge R^\alpha), R^2 \wedge R^\alpha)$  and  $v, y \in B(z, R/2)$ . On the other hand,

$$\begin{aligned}
 \mathbf{E}^{(0, v)} \int_0^{\tau_R} 1_A(s, X_s) ds &= \int_0^\infty \mathbf{P}^{(0, v)}\left(\int_0^{\tau_R} 1_A(s, X_s) ds > u\right) du \\
 &= \int_0^{R^2 \wedge R^\alpha} \mathbf{P}^{(0, v)}\left(\int_0^{\tau_R} 1_A(s, X_s) ds > u\right) du \\
 &\leq \int_0^{R^2 \wedge R^\alpha} \mathbf{P}^{(0, v)}\left(\int_0^{\tau_R} 1_A(s, X_s) ds > 0\right) du \\
 &\leq (R^2 \wedge R^\alpha) \mathbf{P}^{(0, v)}(\sigma_A < \tau_R).
 \end{aligned}$$

The last two displays prove the lemma.  $\square$

Define  $U(t, x, r) = \{t\} \times B(x, r)$ .

LEMMA 4.3. *Let  $\delta \in (0, 1)$ . There exists  $C_{11} = C_{11}(\delta) > 0$  such that for all  $R > 0$ , any  $z \in \mathbf{R}^d$ ,  $(t, x) \in Q(0, z, R/3)$ ,  $v \in B(z, R/3)$ ,  $r \leq R/4$  and  $t \geq \delta(R^2 \wedge R^\alpha)$ ,*

$$\mathbf{P}^{(0, v)}(\sigma_{U(t, x, r/3)} < \tau_{Q(0, z, R)}) \geq C_{11} \frac{(r/3)^{d+2}}{R^{d+2}}.$$

PROOF. Note that

$$\begin{aligned}
 \mathbf{P}^{(0, v)}(\sigma_{U(t, x, r/3)} < \tau_{Q(0, z, R)}) &= \mathbf{P}^v(X_t^{B(z, R)} \in B(x, r/3)) \\
 &= \int_{B(x, r/3)} p^{B(z, R)}(t, v, y) dy \geq C_9 \frac{|B(x, r/3)|}{R^d} \\
 &= C_{11} \frac{(r/3)^d}{R^d} \geq C_{11} \frac{(r/3)^{d+2}}{R^{d+2}},
 \end{aligned}$$

which proves the corollary. Note that the first inequality above follows from Corollary 3.5, because  $v, y \in B(z, R/2)$  and  $t \geq \delta(R^2 \wedge R^\alpha)$ .  $\square$

LEMMA 4.4. *There exists a constant  $C_{12}$  such that for any  $x \in \mathbf{R}^d$ ,  $r > 0$ ,  $y \in B(x, r/3)$  and any bounded non-negative function  $h$  on  $[0, \infty) \times \mathbf{R}^d$  that is supported in  $[0, \infty) \times B(x, 2r)^c$ ,*

$$(4) \quad \mathbf{E}^{(0,x)}[h(\tau_r, X_{\tau_r})] \leq C_{12} \mathbf{E}^{(0,y)}[h(\tau_r, X_{\tau_r})],$$

where  $\tau_r = \tau_{Q(0,x,r)}$ .

PROOF. A Lévy system  $(N, H)$  of the process  $X$  is given by  $N(x, dy) = cdy/|x - y|^{d+\alpha}$  and  $H_t = t$  for some positive constant  $c$ . Thus the proof of the lemma is the same as that of Lemma 4.9 in [4]. The fact that our process  $X$  has a continuous component does not play any role since the function  $h$  is supported in  $[0, \infty) \times B(x, 2r)^c$ .  $\square$

With these lemmas observed, the next theorem can be proved in a manner similar to that in Proposition 4.3 in [4].

THEOREM 4.5. *For every  $\delta \in (0, 1/18)$  there exists a constant  $C_{13} = C_{13}(\delta) > 0$  such that for all  $R > 0$ , for every  $z \in \mathbf{R}^d$  and every non-negative function  $q$  on  $[0, \infty) \times \mathbf{R}^d$  that is parabolic and bounded on  $[0, 4(R^2 \wedge R^\alpha)] \times B(z, 2R)$ ,*

$$\sup_{(t,y) \in Q(\delta(R^2 \wedge R^\alpha), z, R/3)} q(t, y) \leq C_{13} \inf_{y \in B(z, R/3)} q(0, y).$$

PROOF. Without loss of generality we may assume that

$$\inf_{y \in B(z, R/3)} q(0, y) = 1/2.$$

Let  $v \in B(z, R/3)$  be such that  $q(0, v) \leq 1$ . For any  $x \in \mathbf{R}^d$  and  $t \geq 0$ , consider  $Q(t, x, r)$  for  $r \leq R/4$  and let  $\tau_r = \tau_{Q(t,x,r)}$ . Suppose that  $C \subset Q^\delta(t, x, r/3) := Q(t, x, r/3) \cap ([t + \delta(r^2 \wedge r^\alpha), \infty) \times \mathbf{R}^d) = [t + \delta(r^2 \wedge r^\alpha), t + (r/3)^2 \wedge (r/3)^\alpha] \times B(x, r/3)$ . Then by Lemma 4.2,

$$\mathbf{P}^{(t,x)}(\sigma_C < \tau_r) \geq C_{10} \frac{|C|}{r^d (r^2 \wedge r^\alpha)}.$$

Note that there exists a constant  $c_0$  such that  $c_0 r^d (r^2 \wedge r^\alpha) \leq |Q^\delta(t, x, r/3)| \leq c_0^{-1} r^d (r^2 \wedge r^\alpha)$ . Hence, there exists a constant  $c_1 = c_1(\delta) > 0$  such that for all  $C \subset Q^\delta(t, x, r/3)$  satisfying  $|C|/|Q^\delta(t, x, r/3)| \geq 2/3$  we have

$$(5) \quad \mathbf{P}^{(t,x)}(\sigma_C < \tau_r) \geq c_1.$$

Define

$$(6) \quad \eta = \frac{c_1}{3} \quad \text{and} \quad \xi = \frac{1}{3} \wedge (C_{12}\eta),$$

where  $C_{12}$  is the constant from Lemma 4.4.



Suppose there is some point  $(t, x) \in Q(\delta(R^2 \wedge R^\alpha), z, R/3)$  such that  $q(t, x) > K$ , where  $K$  is a constant to be determined later. Define

$$c_2 = \max\left(\left(\frac{3}{c_0 C_{10} \xi}\right)^{1/(d+\alpha)}, \left(\frac{3}{c_0 C_{10} \xi}\right)^{1/(d+2)}, \frac{3 \cdot 2^{1/(d+2)}}{(C_{11} \xi)^{1/(d+2)}}\right),$$

where  $C_{10}$  and  $C_{11}$  are constants from Lemma 4.2 and Lemma 4.3 respectively. Choose

$$(7) \quad r = c_2 R K^{-1/(d+2)}.$$

Then an easy computation shows that

$$(8) \quad \frac{|Q^\delta(0, x, r/3)|}{R^d(R^2 \wedge R^\alpha)} \geq \frac{3}{C_{10} \xi K}, \quad \frac{r^{d+2}}{R^{d+2}} \geq \frac{2 \cdot 3^{d+2}}{C_{11} \xi K}.$$

Let  $U = \{t\} \times B(x, r/3)$ . Suppose that  $q \geq \xi K$  on  $U$ . Let  $Q = Q(0, z, R)$ . Then by Lemma 4.3

$$1 \geq q(0, v) = \mathbf{E}^{(0,v)}[q(Z_{\sigma_U \wedge \tau_R})] \geq \xi K \mathbf{P}^{(0,v)}(\sigma_U < \tau_R) \geq \xi K \frac{C_{11}(r/3)^{d+2}}{R^{d+2}},$$

which contradicts the choice of  $r$  in the second inequality in (8). Thus, there exists at least one point in  $U$  at which  $q$  takes a value less than  $\xi K$ .

We next claim that

$$(9) \quad \mathbf{E}^{(t,x)}[q(\tau_r, X_{\tau_r}) : X_{\tau_r} \notin B(x, 2r)] \leq \eta K,$$

where  $\tau_r = \tau_{Q(t,x,r)}$ . If not, by Lemma 4.4, for all  $y \in B(x, r/3)$ ,

$$\begin{aligned} q(t, y) &\geq \mathbf{E}^{(t,y)}[q(\tau_r, X_{\tau_r}) : X_{\tau_r} \notin B(x, 2r)] \\ &\geq C_{12}^{-1} \mathbf{E}^{(t,x)}[q(\tau_r, X_{\tau_r}) : X_{\tau_r} \notin B(x, 2r)] \\ &\geq C_{12}^{-1} \eta K \geq \xi K. \end{aligned}$$

But this contradicts the already proven fact that there exists at least one point in  $U$  at which  $q$  takes a value less than  $\xi K$ . Therefore, (9) holds true.

Let  $A$  be any compact subset of

$$\tilde{A} := \{(s, y) \in Q^\delta(t, x, r/3) : q(s, y) \geq \xi K\}.$$

Note that  $\tilde{A} \subset Q(0, z, R/2)$ . By Lemma 4.2

$$1 \geq q(0, v) \geq \mathbf{E}^{(0,v)}[q(Z_{\sigma_A}) : \sigma_A < \tau_Q] \geq \xi K \mathbf{P}^{(0,v)}(\sigma_A < \tau_Q) \geq \xi K \frac{C_{10}|A|}{R^d(R^2 \wedge R^\alpha)}.$$

By the first inequality in (8)

$$(10) \quad \frac{|A|}{|Q^\delta(t, x, r/3)|} \leq \frac{R^d(R^2 \wedge R^\alpha)}{C_{10}|Q^\delta(t, x, r/3)|\xi K} \leq \frac{1}{3}.$$

Since (10) holds for every compact subset  $A$  of  $\tilde{A}$ , it holds for  $\tilde{A}$  in place of  $A$ .

Let  $C := Q^\delta(t, x, r/3) \setminus \tilde{A}$ . Then by (10),  $|C|/|Q^\delta(t, x, r/3)| \geq 2/3$ . Let

$$M = \sup_{(s,y) \in Q(t,x,2r)} q(s, y).$$

Then

$$\begin{aligned} q(t, x) &= \mathbf{E}^{(t,x)}[q(\sigma_C, X_{\sigma_C}) : \sigma_C < \tau_r] \\ &\quad + \mathbf{E}^{(t,x)}[q(\sigma_C, X_{\sigma_C}) : \tau_r \leq \sigma_C, X_{\tau_r} \notin B(x, 2r)] \\ &\quad + \mathbf{E}^{(t,x)}[q(\sigma_C, X_{\sigma_C}) : \tau_r \leq \sigma_C, X_{\tau_r} \in B(x, 2r)]. \end{aligned}$$

The first term on the right is bounded by  $\xi K \mathbf{P}^{(t,x)}(\sigma_C < \tau_r)$ , the second term is according to (9) bounded by  $\eta K$ , and the third term is bounded by  $M \mathbf{P}^{(t,x)}(\sigma_C \geq \tau_r)$ . Therefore,

$$K \leq q(t, x) \leq \xi K \mathbf{P}^{(t,x)}(\sigma_C < \tau_r) + \eta K + M \mathbf{P}^{(t,x)}(\sigma_C \geq \tau_r).$$

Note that by (5),  $\mathbf{P}^{(t,x)}(\sigma_C < \tau_r) \geq c_1$ . Hence by use of (6),

$$\frac{M}{K} \geq \frac{1 - \eta - \xi \mathbf{P}^{(t,x)}(\sigma_C < \tau_r)}{\mathbf{P}^{(t,x)}(\sigma_C \geq \tau_r)} \geq \frac{1 - \eta - \xi c_1}{1 - c_1} \geq \frac{1 - 2c_1/3}{1 - c_1} = 1 + 2\beta,$$

where  $\beta = c_1/6(1 - c_1)$ . Hence, there exists a point  $(t_1, x_1) \in Q(t, x, 2r) \subset \hat{Q}(0, z, R) := [0, 3(R^2 \wedge R^\alpha)] \times B(z, R)$  such that  $q(t_1, x_1) \geq (1 + \beta)K =: K_1$ . Note that  $0 \leq t_1 - t \leq (2r)^2 \wedge (2r)^\alpha$  and  $|x_1 - x| \leq 2r$ .

Iterate the above procedure to obtain a sequence of points  $\{(t_k, x_k)\}$  in the following way. Using above argument (with  $(t_1, x_1)$  and  $K_1$  instead of  $(t, x)$  and  $K$ ), there exists  $(t_2, x_2) \in Q(t_1, x_1, 2r_1)$  such that  $q(t_2, x_2) \geq (1 + \beta)K_1 =: K_2$ . Continue this procedure to obtain a sequence of points  $\{(t_k, x_k)\}$  such that  $(t_{k+1}, x_{k+1}) \in Q(t_k, x_k, 2r_k)$  and  $q(t_{k+1}, x_{k+1}) \geq (1 + \beta)^{k+1} K_1 =: K_{k+1}$ . We have that  $0 \leq t_{k+1} - t_k \leq (2r_k)^2 \wedge (2r_k)^\alpha$ ,  $|x_{k+1} - x_k| \leq 2r_k$ . Moreover, by (7),

$$r_k \leq c_2 R K_k^{-1/(d+2)} \leq c_2 (1 + \beta)^{-k/(d+2)} K^{-1/(d+2)} R.$$

Note that

$$\begin{aligned} \sum_k r_k &\leq c_2 K^{-1/(d+2)} R \sum_k ((1 + \beta)^{-1/(d+2)})^k = \frac{c_2 K^{-1/(d+2)} R}{1 - (1 + \beta)^{-1/(d+2)}}, \\ \sum_k (2r_k)^2 &\leq (2c_2 K^{-1/(d+2)} R)^2 \sum_k ((1 + \beta)^{-2/(d+2)})^k = \frac{(2c_2 K^{-1/(d+2)} R)^2}{1 - (1 + \beta)^{-2/(d+2)}}, \\ \sum_k (2r_k)^\alpha &\leq (2c_2 K^{-1/(d+2)} R)^\alpha \sum_k ((1 + \beta)^{-2/(d+2)})^k = \frac{(2c_2 K^{-1/(d+2)} R)^\alpha}{1 - (1 + \beta)^{-2/(d+2)}}. \end{aligned}$$

Therefore, we may choose  $K$  large enough so that  $(t_k, x_k) \in \hat{Q}(0, z, R)$  for all  $k$ . This is a contradiction because  $q(t_k, x_k) \geq (1 + \beta)^k K$  goes to infinity as  $k \rightarrow \infty$ .  $\square$

*Acknowledgment.* We are grateful to an anonymous referee for suggesting a more general version of Theorem 4.5. We also thank Z.-Q. Chen for helpful discussion leading to Lemma 2.8.

#### REFERENCES

- [ 1 ] R. F. BASS AND D. A. LEVIN, Harnack inequalities for jump processes, *Potential Anal.* 17 (2002), 375–388.
- [ 2 ] R. F. BASS AND D. A. LEVIN, Transition probabilities for symmetric jump processes, *Trans. Amer. Math. Soc.* 354 (2002), 2933–2953.

- [ 3 ] R. M. BLUMENTHAL AND R. K. GETTOOR, Some theorems on stable processes, *Trans. Amer. Math. Soc.* 95 (1960), 263–273.
- [ 4 ] Z.-Q. CHEN AND T. KUMAGAI, Heat kernel estimates for stable-like processes on  $d$ -sets, *Stochastic Processes. Appl.* 108 (2003), 27–62.
- [ 5 ] E. B. FABES AND D. W. STROOCK, A new proof of Moser’s parabolic Harnack inequality using old ideas of Nash, *Arch. Rational Mech. Anal.* 96 (1986), 327–338.
- [ 6 ] R. MIKULEVICIUS AND H. PRAGARAUSKAS, Green measures of Itô processes, *Litovsk. Mat. Sb.* 27 (1987), 711–723.
- [ 7 ] M. RAO, R. SONG AND Z. VONDRAČEK, Green function estimates and Harnack inequality for subordinate Brownian motions, *Potential Anal.* 25 (2006), 1–27.
- [ 8 ] R. SONG AND Z. VONDRAČEK, Harnack inequality for some discontinuous Markov processes with a Diffusion Part, *Glas. Mat. Ser. III*, 40 (2005), 177–187.

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF ILLINOIS  
URBANA, IL 61801  
USA

*E-mail address:* rsong@math.uiuc.edu

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF ZAGREB  
ZAGREB  
CROATIA

*E-mail address:* vondra@math.hr