

PERIODIC SOLUTIONS OF THE NAVIER-STOKES EQUATIONS IN UNBOUNDED DOMAINS

Dedicated to Professor Kôji Kubota on his sixtieth birthday

HIDEO KOZONO AND MITSUHIRO NAKAO

(Received August 29, 1994, revised February 21, 1995)

Abstract. We shall construct a periodic strong solution of the Navier-Stokes equations for the prescribed external force in unbounded domains.

Introduction. The purpose of this paper is to show that if the incompressible fluid in *unbounded* domains is governed by the periodic external force, the Navier-Stokes equations have a *periodic strong* solution with the same period as the external force. Let Ω be a domain in \mathbf{R}^n ($n \geq 3$), not necessarily bounded, with smooth boundary $\partial\Omega$. Consider the following Navier-Stokes equations in Ω :

$$(N-S) \quad \begin{cases} \frac{\partial u}{\partial t} - \Delta u + u \cdot \nabla u + \nabla p = f, & x \in \Omega, \quad t \in \mathbf{R}, \\ \operatorname{div} u = 0, & x \in \Omega, \quad t \in \mathbf{R}, \\ u|_{\partial\Omega} = 0, \end{cases}$$

where $u = u(x, t) = (u^1(x, t), \dots, u^n(x, t))$ and $p = p(x, t)$ denote the unknown velocity vector and pressure of the fluid at point $(x, t) \in \Omega \times \mathbf{R}$, respectively; while $f = f(x, t) = (f^1(x, t), \dots, f^n(x, t))$ is the given periodic external force.

Under some restrictive conditions, Serrin [20] gave a criterion for the existence of periodic solutions of (N-S) when Ω is a three-dimensional bounded domain whose boundary moves periodically in time. Kaniel-Shinbrot [11] considered a simpler case such as bounded domains whose boundary is fixed in time and realized the criterion of Serrin. Having introduced the notion of reproductive property, they showed the existence of periodic strong solutions with periodic small forces f . In two-dimensional bounded domains, Takeshita [23] obtained the same result as Kaniel-Shinbrot [11] without assuming the smallness of f . The original problem posed by Serrin had been treated by Morimoto [19] and Miyakawa-Teramoto [18] who showed the existence of periodic weak solutions. Later on, Teramoto [25] constructed periodic strong solutions in a situation such that the boundary moves slowly in time.

All of these results are obtained in two- or three-dimensional *bounded* domains. On the other hand, few results are known in *unbounded* domains. Recently, Maremonti

[15], [16] showed the existence of periodic strong solutions in the three-dimensional whole space \mathbf{R}^3 and the half space \mathbf{R}_+^3 , respectively. However, the result corresponding to exterior domains has not been obtained up to the present. The main difficulty in unbounded domains stems from the lack of exponential decay in time for solutions to the initial value problem of (N-S). Indeed, Serrin [20] and Kaniel-Shinbrot [11] made full use of the fact that $\|u(t)\|_2$ and $\|\nabla u(t)\|_2$ decay exponentially in t provided the initial data at $t=0$ are prescribed. Such a decay property is due to the Poincaré inequality in bounded domains, and invertibility of the Stokes operator in L^2 makes it easy to obtain better asymptotic behaviour of solutions at $t \rightarrow \infty$.

To overcome this difficulty, Maremonti [15], [16] first showed the algebraic decay rates in time of strong solutions for initial value problem of (N-S) in \mathbf{R}^3 and in \mathbf{R}_+^3 . As a by-product, he constructed periodic strong solutions for periodic small external forces. His method is based on the skillful energy estimates in L^2 for higher derivatives of solutions. Although our results are not altogether new, our approach is different and gives more results than those by Maremonti [15], [16]. We do not employ the energy estimates in L^2 but the L^p -theory of the Stokes operator. Making use of L^p - L^r estimates for the semigroup generated by the Stokes operator, we shall show the existence and uniqueness of *periodic strong* solutions more directly than Maremonti [15], [16]. Compared with the energy estimates in L^2 , our L^p method can cover also the higher dimensional cases. Unfortunately, we cannot obtain the same result in three-dimensional exterior domains because the corresponding L^p - L^r estimate is still an open problem.

We shall first reduce our problem to an integral equation, the solution of which is necessarily periodic with the same period as the external force. The solution will be constructed in the class of functions defined on the whole interval \mathbf{R} with values in $L^n(\Omega)$. Then by a regularity criterion similar to Serrin's [21], we shall show that our solution is actually a strong solution. For that purpose, we shall estimate a time-interval of the existence of local strong solutions for the initial-boundary value problem to (N-S) in terms of the given data. Our estimate extends the result obtained by Giga [7, Theorem 4]. The stability of periodic solutions will be discussed in a forthcoming paper.

1. Results. Before stating our results, we need to impose the following assumption on the domain Ω :

ASSUMPTION 1. (Case I) Ω is the whole space \mathbf{R}^n or the half-space \mathbf{R}_+^n , where $n \geq 3$.

(Case II) Ω is an exterior domain in \mathbf{R}^n with $C^{2+\mu}$ ($\mu > 0$)-boundary $\partial\Omega$, where $n \geq 4$.

The reason why we exclude three-dimensional exterior domains in (Case II) is due to the restriction on gradient bounds for the Stokes semigroup in L^p (see Lemma 2.1 (2) below).

We shall next introduce some notation and function spaces. Let $C_{0,\sigma}^\infty$ denote the set of all real vector C^∞ -functions $\phi = (\phi^1, \dots, \phi^n)$ with compact support in Ω such

that $\operatorname{div} \phi = 0$. L'_σ is the closure of $C^\infty_{0,\sigma}$ with respect to the L^r -norm $\|\cdot\|_r$; (\cdot, \cdot) denotes the duality pairing between L^r and $L^{r'}$, where $1/r + 1/r' = 1$. L^r stands for the usual (vector-valued) L^r -space over Ω , $1 < r < \infty$. When X is a Banach space, its norm is denoted by $\|\cdot\|_X$. Then $C^m([t_1, t_2]; X)$ is the usual Banach space, where $m = 0, 1, 2, \dots$ and t_1 and t_2 are real numbers such that $t_1 < t_2$. $BC^m([t_1, t_2]; X)$ is the set of all functions $u \in C^m([t_1, t_2]; X)$ such that $\sup_{t_1 < t < t_2} \|d^m u(t)/dt^m\|_X < \infty$.

Let us recall the Helmholtz decomposition:

$$L^r = L'_\sigma \oplus G^r \text{ (direct sum)}, \quad 1 < r < \infty,$$

where $G^r = \{\nabla p \in L^r; p \in L^r_{\text{loc}}(\bar{\Omega})\}$. For the proof, see Fujiwara-Morimoto [4], Miyakawa [17] and Simader-Sohr [22]. P_r denotes the projection operator from L^r onto L'_σ along G^r . The Stokes operator A_r on L'_σ is then defined by $A_r = -P_r \Delta$ with domain $D(A_r) = \{u \in H^{2,r}(\Omega); u|_{\partial\Omega} = 0\} \cap L'_\sigma$. It is known that the dual space $(L'_\sigma)^*$ of L'_σ and the adjoint operator A_r^* of A_r are respectively

$$(L'_\sigma)^* = L^{r'}, \quad A_r^* = A_{r'},$$

where $1/r + 1/r' = 1$. Moreover, we have:

PROPOSITION 1 (Giga [5], Giga-Sohr [9]). *Let $1 < r < \infty$. Then $-A_r$ generates a uniformly bounded holomorphic semigroup $\{e^{-tA_r}\}_{t \geq 0}$ of class C^0 in L'_σ .*

Applying the projection operator P_r to both sides of the first equation of (N-S), we have

$$(E) \quad \frac{du}{dt} + A_r u + P_r(u \cdot \nabla u) = P_r f \quad \text{on } L'_\sigma, \quad t \in \mathbf{R}.$$

The above (E) can be further transformed to the following integral equation:

$$(I.E.) \quad u(t) = \int_{-\infty}^t e^{-(t-s)A_r} P_r f(s) ds - \int_{-\infty}^t e^{-(t-s)A_r} P_r(u \cdot \nabla u)(s) ds.$$

Concerning the external force f , we impose the following assumption:

ASSUMPTION 2. Let the exponents r and q be according to the (Case I) and (Case II) of Assumption 1 as

- (Case I) $2 < r < n, n/2 < q < n$;
- (Case II) $2n/(n-1) \leq r < n, n/2 < q < n$.

For such r and q , we assume that f belongs to the class

$$(I.1) \quad f \in BC(\mathbf{R}; L^p \cap L^l)$$

for $1 < p, l < \infty$ with $1/r + 2/n < 1/p, 1/q < 1/l < 1/q + 1/n$ provided $n \geq 4$ in both (Case I) and (Case II).

If $n = 3$ in (Case I), assume moreover that

$$(1.2) \quad P_p f(s) = A_p^\delta g(s) \quad (s \in \mathbf{R}) \quad \text{with some } g \in BC(\mathbf{R}; D(A_p^\delta)) \text{ and } f \in BC(\mathbf{R}; L^1)$$

for $1 < p < \min\{r, q\}$ and $\delta > 0$ satisfying $3/2p + \delta > 1 + \max\{1 + 3/2r, 1/2 + 3/2q\}$ and for $1/q < 1/l < 1/q + 1/3$.

Our result now reads:

THEOREM 1. *Let Ω and f satisfy Assumption 1 and Assumption 2, respectively. Suppose that $f(t) = f(t + \omega)$ for all $t \in \mathbf{R}$ with some $\omega > 0$. Then there is a constant $\eta = \eta(n, r, q, p, l, \delta) > 0$ such that if*

$$\sup_{s \in \mathbf{R}} \|P_p f(s)\|_p + \sup_{s \in \mathbf{R}} \|P_l f(s)\|_l \leq \eta \quad \text{for } n \geq 4 \text{ in (Case I) and (Case II),}$$

$$\sup_{s \in \mathbf{R}} \|g(s)\|_p + \sup_{s \in \mathbf{R}} \|P_l f(s)\|_l \leq \eta \quad \text{for } n = 3 \text{ in (Case I),}$$

we have a periodic solution u of (I.E.) with the same period ω as f in the class $BC(\mathbf{R}; L'_\sigma)$ with $\nabla u \in BC(\mathbf{R}; L^q)$.

Such a solution u is unique within this class provided $\sup_{s \in \mathbf{R}} \|u(s)\|_r + \sup_{s \in \mathbf{R}} \|\nabla u(s)\|_q$ is sufficiently small.

Concerning the existence of solutions to (E), we have:

THEOREM 2. *In addition to the hypotheses of Theorem 1, let us assume furthermore that f is a Hölder continuous function on \mathbf{R} with values in L^n . Then the periodic solution u given by Theorem 1 has the following additional properties:*

- (i) $u \in BC(\mathbf{R}; L^n_\sigma) \cap C^1(\mathbf{R}; L^n_\sigma)$;
- (ii) $u(t) \in D(A_n)$ for all $t \in \mathbf{R}$ and $A_n u \in C(\mathbf{R}; L^n_\sigma)$;
- (iii) u satisfies (E) in L^n_σ for all $t \in \mathbf{R}$.

REMARKS. (1) Taking $n = 3$, $2 < r < 3$ and $q = 2$ in (Case I), our theorems include Maremonti [15, Theorem 1] and [16, Theorem 2].

(2) The first condition of (1.2) seems to be artificial, but it may be replaced by $f(s) = \operatorname{div} F(s)$ with some $F = \{F_{i,j}\}_{i,j=1,2,3} \in BC(\mathbf{R}; H^{1,p}(\Omega))$ for $1 < p < \infty$ satisfying $1/r + 1/3 < 1/p$.

(3) When Ω is a bounded domain in \mathbf{R}^n ($n \geq 2$), the above results also hold and we can relax the assumption on the external force. Indeed, it suffices to assume that $f \in BC(\mathbf{R}; L^r)$ with $\sup_{s \in \mathbf{R}} \|P f(s)\|_r$ small for $r > n/2$. Under such a hypothesis, there is a periodic solution u of (I.E.) in the class $u \in BC(\mathbf{R}; D(A_r^{1/2}))$.

2. Preliminaries. Throughout this paper, we shall denote by C various constants. In particular, $C = C(*, \dots, *)$ will denote the constants which depend only on the quantities appearing in parentheses. Since $P_r u = P_q u$ for all $u \in L^r \cap L^q$ and since $A_r u = A_q u$ for all $u \in D(A_r) \cap D(A_q)$, for simplicity, we shall abbreviate $P_r u$, $P_q u$ as Pu for $u \in L^r \cap L^q$ and $A_r u$, $A_q u$ as Au for $u \in D(A_r) \cap D(A_q)$, respectively. Let us first recall the following

L^p - L^r estimates for the semigroup $\{e^{-tA}\}_{t \geq 0}$:

LEMMA 2.1 (Kato [12], Ukai [26], Giga-Sohr [9], Iwashita [10], Borchers-Miyakawa [1], [2]).

(1) Let Ω be as in (Case I) of Assumption 1. Then there holds

$$\begin{aligned} \|e^{-tA}a\|_r &\leq Ct^{-n(1/p-1/r)/2} \|a\|_p, & 1 < p \leq r < \infty \\ \|\nabla e^{-tA}a\|_r &\leq Ct^{-n(1/p-1/r)/2-1/2} \|a\|_p, & 1 < p \leq r < \infty \end{aligned}$$

for all $a \in L^p_\sigma$ and all $t > 0$, where $C = C(n, p, r)$.

(2) Let Ω be as in (Case II) of Assumption 1. Then there holds

$$\begin{aligned} \|e^{-tA}a\|_r &\leq Ct^{-n(1/p-1/r)/2} \|a\|_p, & 1 < p \leq r < \infty \\ \|\nabla e^{-tA}a\|_r &\leq Ct^{-n(1/p-1/r)/2-1/2} \|a\|_p, & 1 < p \leq r \leq n \end{aligned}$$

for all $a \in L^p_\sigma$ and all $t > 0$, where $C = C(n, p, r)$.

Using this lemma, we shall estimate the nonlinear term of (I.E.).

LEMMA 2.2. Let r and q be as in Theorem 1 according to the (Case I) and (Case II) of Assumption 1. Define a function space Y and a bilinear operator $G(\cdot, \cdot)$ on Y by

$$Y \equiv \{u \in BC(\mathbf{R}; L^r_\sigma); \nabla u \in BC(\mathbf{R}; L^q)\},$$

$$G(u, v)(t) \equiv - \int_{-\infty}^t e^{-(t-s)A} P(u \cdot \nabla v)(s) ds \quad u, v \in Y,$$

respectively. Then we have $G(u, v) \in Y$ with

$$(2.1) \quad \sup_{s \in \mathbf{R}} \|G(u, v)(s)\|_r \leq C_1 \left(\sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|v(s)\|_r + \sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q \right)$$

$$(2.2) \quad \sup_{s \in \mathbf{R}} \|\nabla G(u, v)(s)\|_q \leq C_1 \left(\sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q + \sup_{s \in \mathbf{R}} \|\nabla u(s)\|_q \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q \right)$$

for all $u, v \in Y$, where $C_1 = C_1(n, r, q)$.

PROOF. Set

$$\begin{aligned} (2.3) \quad G(u, v)(t) &= - \int_{-\infty}^{t-1} e^{-(t-s)A} P(u \cdot \nabla v)(s) ds - \int_{t-1}^t e^{-(t-s)A} P(u \cdot \nabla v)(s) ds \\ &\equiv -I_1(t) - I_2(t). \end{aligned}$$

By integration by parts and Lemma 2.1 we have

$$\begin{aligned}
|(I_1(t), \phi)| &= \left| - \int_{-\infty}^{t-1} (u(s) \cdot \nabla e^{-(t-s)A} \phi, v(s)) ds \right| \\
&\leq \int_{-\infty}^{t-1} \|\nabla e^{-(t-s)A} \phi\|_{(r/2)'} \|u(s) \cdot v(s)\|_{r/2} ds \\
&\leq C \int_{-\infty}^{t-1} (t-s)^{-n(1/r' - 1/(r/2)')/2 - 1/2} \|\phi\|_{r'} \|u(s)\|_r \|v(s)\|_r ds \\
&\leq C \sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|v(s)\|_r \int_{-\infty}^{t-1} (t-s)^{-n/2r - 1/2} ds \|\phi\|_{r'} \\
&\leq C \sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|v(s)\|_r \|\phi\|_{r'}.
\end{aligned}$$

for all $\phi \in C_{0,\sigma}^\infty$ and all $t \in \mathbf{R}$, where $C = C(n, r)$. Note that, in (Case II), on account of such restriction as Lemma 2.2 (2), we need $(r/2)' \leq n$, i.e., $2n' \leq r$. By duality there holds

$$(2.4) \quad \|I_1(t)\|_r \leq C \sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|v(s)\|_r \quad \text{for all } t \in \mathbf{R},$$

where $C = C(n, r)$. Similarly, Lemma 2.1 and the Hölder inequality yield

$$\begin{aligned}
\|I_2(t)\|_r &\leq \int_{t-1}^t \|e^{-(t-s)A} P(u \cdot \nabla v)(s)\|_r ds \\
(2.5) \quad &\leq \int_{t-1}^t (t-s)^{-n(1/r + 1/q - 1/r)/2} \|u(s)\|_r \|\nabla v(s)\|_q ds \\
&\leq \sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q
\end{aligned}$$

for all $t \in \mathbf{R}$ with $C = C(n, r, q)$. Now (2.1) follows from (2.3)–(2.5).

To show (2.2), we make use of the following inequality of the Sobolev type due to Giga-Sohr [9, Corollary 2.2 (ii)]:

$$(2.6) \quad \|\phi\|_{q^*} \leq C \|\nabla \phi\|_q \quad \text{with } q^* = nq/(n-q) \text{ for all } \phi \in L^r \text{ with } \nabla \phi \in L^q,$$

where $C = C(n, q)$. It should be noted that (2.6) holds even though ϕ does not vanish on the boundary $\partial\Omega$. Now it follows from Lemma 2.1 and (2.6) that

$$\begin{aligned}
\|\nabla G(u, v)(t)\|_q &\leq \int_{-\infty}^{t-1} \|\nabla e^{-(t-s)A} P(u \cdot \nabla v)(s)\|_q ds + \int_{t-1}^t \|\nabla e^{-(t-s)A} P(u \cdot \nabla v)(s)\|_q ds \\
&\leq C \int_{-\infty}^{t-1} (t-s)^{-n(1/r + 1/q - 1/q)/2 - 1/2} \|u(s)\|_r \|\nabla v(s)\|_q ds
\end{aligned}$$

$$\begin{aligned}
 & + C \int_{t-1}^t (t-s)^{-n(1/q^*+1/q-1/q)/2-1/2} \|u(s)\|_{q^*} \|\nabla v(s)\|_q ds \\
 & \leq C \sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q \int_{-\infty}^{t-1} (t-s)^{-n/2r-1/2} ds \\
 & \quad + C \sup_{s \in \mathbf{R}} \|u(s)\|_{q^*} \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q \int_{t-1}^t (t-s)^{-n/2q} ds \\
 & \leq C \left(\sup_{s \in \mathbf{R}} \|u(s)\|_r \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q + \sup_{s \in \mathbf{R}} \|\nabla u(s)\|_q \sup_{s \in \mathbf{R}} \|\nabla v(s)\|_q \right)
 \end{aligned}$$

for all $t \in \mathbf{R}$ with $C = C(n, r, q)$, which yields (2.2). This completes the proof of Lemma 2.2.

We shall next show bounds for the external force.

LEMMA 2.3. *Let f be as in (1.1) and (1.2) of Assumption 2. Let*

$$F(t) \equiv \int_{-\infty}^t e^{-(t-s)A} P f(s) ds, \quad t \in \mathbf{R}.$$

Then we have $F \in Y$ and the following estimates hold:

$$(2.7) \quad \left. \begin{array}{l} \sup_{s \in \mathbf{R}} \|F(s)\|_r \\ \sup_{s \in \mathbf{R}} \|\nabla F(s)\|_q \end{array} \right\} \leq C \left(\sup_{s \in \mathbf{R}} \|P f(s)\|_p + \sup_{s \in \mathbf{R}} \|P f(s)\|_l \right)$$

with $C = C(n, r, q, p, l)$ provided f satisfies (1.1);

$$(2.8) \quad \left. \begin{array}{l} \sup_{s \in \mathbf{R}} \|F(s)\|_r \\ \sup_{s \in \mathbf{R}} \|\nabla F(s)\|_q \end{array} \right\} \leq C \left(\sup_{s \in \mathbf{R}} \|g(s)\|_p + \sup_{s \in \mathbf{R}} \|P f(s)\|_l \right)$$

with $C = C(n, r, q, p, l, \delta)$ provided f satisfies (1.2).

PROOF. If f satisfies (1.1), we have by Lemma 2.1

$$\begin{aligned}
 \|F(t)\|_r & \leq \int_{-\infty}^{t-1} \|e^{-(t-s)A} P f(s)\|_r ds + \int_{t-1}^t \|e^{-(t-s)A} P f(s)\|_r ds \\
 & \leq C \sup_{s \in \mathbf{R}} \|P f(s)\|_p \int_{-\infty}^{t-1} (t-s)^{-n(1/p-1/r)/2} ds \\
 & \quad + C \sup_{s \in \mathbf{R}} \|P f(s)\|_l \int_{t-1}^t (t-s)^{-n(1/l-1/r)/2} ds
 \end{aligned}$$

for all $t \in \mathbf{R}$, where $C = C(n, r, q, p, l)$. Since $n/2 < q$ and $r < n$, we have by hypothesis on

l that $1/l < 2/n + 1/r$ and hence the second integral above converges. So does the first integral by hypothesis on p and we obtain (2.7) for $\sup_{s \in \mathbf{R}} \|F(s)\|_r$. Similarly, we have by Lemma 2.1

$$\begin{aligned} \|\nabla F(t)\|_q &\leq C \sup_{s \in \mathbf{R}} \|Pf(s)\|_p \int_{-\infty}^{t-1} (t-s)^{-n(1/p-1/q)/2-1/2} ds \\ &\quad + C \sup_{s \in \mathbf{R}} \|Pf(s)\|_l \int_{t-1}^t (t-s)^{-n(1/l-1/q)/2-1/2} ds \end{aligned}$$

for all $t \in \mathbf{R}$ with $C = C(n, r, q, p, l)$. Since $1/p > 1/r + 2/n > 1/n + 1/q$, the first integral above converges and we obtain (2.7) for $\sup_{s \in \mathbf{R}} \|\nabla F(s)\|_q$.

To show (2.8), we shall make use of the estimate

$$(2.9) \quad \|A^\delta e^{-tA} a\|_p \leq C t^{-\delta} \|a\|_p \quad \text{for all } a \in L^p_\sigma, \quad t > 0$$

with $C = C(p, \delta)$. This is an immediate consequence of Proposition 1. Therefore, if f satisfies (1.2), it follows from Lemma 2.1 and (2.9) that

$$\begin{aligned} \|F(t)\|_r &\leq \int_{-\infty}^{t-1} \|e^{-(t-s)A} A^\delta g(s)\|_r ds + \int_{t-1}^t \|e^{-(t-s)A} Pf(s)\|_r ds \\ &\leq C \int_{-\infty}^{t-1} (t-s)^{-3(1/p-1/r)/2} \|A^\delta e^{-(t-s)A/2} g(s)\|_p ds \\ &\quad + C \int_{t-1}^t (t-s)^{-3(1/l-1/r)/2} \|Pf(s)\|_l ds \\ &\leq C \int_{-\infty}^{t-1} (t-s)^{-3(1/p-1/r)/2-\delta} \|g(s)\|_p ds \\ &\quad + C \sup_{s \in \mathbf{R}} \|Pf(s)\|_l \int_{t-1}^t (t-s)^{-3(1/l-1/r)/2} ds \\ &\leq C \left(\sup_{s \in \mathbf{R}} \|g(s)\|_p + \sup_{s \in \mathbf{R}} \|Pf(s)\|_l \right) \end{aligned}$$

for all $t \in \mathbf{R}$ with $C = C(n, r, q, p, l, \delta)$. This yields (2.8) for $\sup_{s \in \mathbf{R}} \|F(s)\|_r$. Similarly, we can deal with ∇F to show (2.8) for $\sup_{s \in \mathbf{R}} \|\nabla F(s)\|_q$ and the proof of Lemma 2.3 is complete.

3. Existence of periodic solution; Proof of Theorem 1. Using Lemmas 2.2 and 2.3, we shall prove the existence and uniqueness of solution to the integral equation (I.E.) by successive approximation. Let us recall the function space Y and the bilinear operator $G(\cdot, \cdot)$ on Y introduced in Lemma 2.2. Equipped with the norm $\|\cdot\|_Y$ defined by

$$\|u\|_Y \equiv \sup_{s \in \mathbf{R}} \|u(s)\|_r + \sup_{s \in \mathbf{R}} \|\nabla u(s)\|_q,$$

Y is a Banach space. We construct a periodic solution of (I.E.) according to the scheme

$$(3.1) \quad u_0(t) = \int_{-\infty}^t e^{-(t-s)A} Pf(s) ds,$$

$$(3.2) \quad u_{m+1}(t) = u_0(t) + G(u_m, u_m)(t), \quad m = 0, 1, \dots$$

By Lemma 2.3, we have $u_0 \in Y$ with

$$(3.3) \quad \|u_0\|_Y \leq C \left(\sup_{s \in \mathbf{R}} \|Pf(s)\|_p + \sup_{s \in \mathbf{R}} \|Pf(s)\|_l \right)$$

provided f satisfies (1.1);

$$(3.4) \quad \|u_0\|_Y \leq C \left(\sup_{s \in \mathbf{R}} \|g(s)\|_p + \sup_{s \in \mathbf{R}} \|Pf(s)\|_l \right)$$

provided f satisfies (1.2). Since f is a periodic function with period ω , we can easily verify that u_0 is also periodic with the same period ω . By induction and Lemma 2.2, so is u_m for all $m = 0, 1, \dots$. Moreover, it follows from (2.1) and (2.2) that

$$\|u_{m+1}\|_Y \leq \|u_0\|_Y + \|G(u_m, u_m)\|_Y \leq \|u_0\|_Y + C_* \|u_m\|_Y^2, \quad m = 0, 1, \dots,$$

where $C_* = 2C_1$. Hence if

$$(3.5) \quad \|u_0\|_Y < \frac{1}{4C_*},$$

then there holds

$$(3.6) \quad \|u_m\|_Y \leq \frac{1 - \sqrt{1 - 4C_* \|u_0\|_Y}}{2C_*} \equiv K < \frac{1}{2C_*} \quad \text{for all } m = 0, 1, \dots$$

By (3.3) and (3.4), we can take the constant η in Theorem 1 so that the condition (3.5) is satisfied.

Now assume (3.5). Setting $w_m \equiv u_m - u_{m-1}$ ($u_{-1} \equiv 0$), we have

$$w_{m+1}(t) = G(u_m, u_m)(t) - G(u_{m-1}, u_{m-1})(t) = G(w_m, u_m)(t) - G(u_{m-1}, w_m)(t)$$

and Lemma 2.2 and (3.6) yield

$$(3.7) \quad \begin{aligned} \|w_{m+1}\|_Y &\leq \|G(w_m, u_m)\|_Y + \|G(u_{m-1}, w_m)\|_Y \\ &\leq C_*(\|w_m\|_Y \|u_m\|_Y + \|u_{m-1}\|_Y \|w_m\|_Y) \leq 2C_* K \|w_m\|_Y \\ &\leq \dots \leq (2C_* K)^{m+1} \|u_0\|_Y \end{aligned}$$

for all $m=0, 1, \dots$. Since $u_m(t) = \sum_{j=0}^m w_j(t)$, we see by (3.6) and (3.7) that there exists a function $u \in Y$ such that

$$(3.8) \quad u_m \rightarrow u \quad \text{in } Y \text{ as } m \rightarrow \infty .$$

Clearly, such a limit $u(t)$ is also periodic in t with the same period ω as f . As in (3.7), we have by (3.6) that

$$(3.9) \quad \begin{aligned} \|G(u_m, u_m) - G(u, u)\|_Y &\leq \|G(u_m - u, u_m)\|_Y + \|G(u, u_m - u)\|_Y \\ &\leq C_* \|u_m - u\|_Y \|u_m\|_Y + C_* \|u\|_Y \|u_m - u\|_Y \\ &< \|u_m - u\|_Y \end{aligned}$$

for all m , from which follows that

$$(3.10) \quad G(u_m, u_m) \rightarrow G(u, u) \quad \text{in } Y \text{ as } m \rightarrow \infty .$$

Now letting $m \rightarrow \infty$ in (3.2), we see by (3.8) and (3.10) that u is a desired periodic solution of the integral equation (I.E.).

It remains to show the uniqueness. Suppose that $v \in Y$ is another solution of (I.E.) with $\|v\|_Y \leq K$, where K is the same constant as in (3.6). Then we have as in (3.9) that

$$\|u - v\|_Y \leq C_* (\|u - v\|_Y \|u\|_Y + \|v\|_Y \|u - v\|_Y) \leq 2C_* K \|u - v\|_Y .$$

Since $2C_* K < 1$, there holds $u = v$ and the assertion on uniqueness follows. This proves Theorem 1.

4. Regularity of solutions to (I.E.); Proof of Theorem 2. In this section, we shall show that the periodic solution u constructed in the preceding section is actually a solution of the differential equation (E). To this end, we need the local existence of strong solutions to the initial-boundary value problem for (N-S). In particular, it is important to give the time-interval of existence in terms of the prescribed data. Here we follow the argument of Kato [12] and Giga [7].

Let us first define the strong solution of the initial value problem for (N-S).

DEFINITION. Let $a \in L_\sigma^n$ and let $Pf \in C((t_0, t_1); L_\sigma^n)$, where $t_0 < t_1$. Then a measurable function v on $\Omega \times (t_0, t_1)$ is called a *strong* solution of (N-S) on (t_0, t_1) with the initial data a at t_0 if

- (i) $v \in BC([t_0, t_1]; L_\sigma^n) \cap C^1((t_0, t_1); L_\sigma^n)$;
- (ii) $v(t) \in D(A_n)$ for $t_0 < t < t_1$ and $A_n v \in C((t_0, t_1); L_\sigma^n)$;
- (iii)

$$\begin{cases} \frac{dv}{dt} + Av + P(v \cdot \nabla v) = Pf & \text{in } L_\sigma^n \text{ for } t_0 < t < t_1, \\ v(t_0) = a. \end{cases}$$

Our result on the local existence of strong solutions now reads:

LEMMA 4.1. *Let $n/2 < q < n$ and let $1 < l < \infty$ satisfy $1/q < 1/l < 1/q + 1/n$. Assume that $a \in L^n_\sigma \cap L^{q^*}_\sigma$ with $q^* = nq/(n-q)$, $f \in BC(\mathbf{R}; L^l)$ and that $Pf(\cdot)$ is a Hölder continuous function on \mathbf{R} with values in L^n_σ . Then there exists $T > 0$ such that for every $t_0 \in \mathbf{R}$, we have a unique strong solution v of (N-S) on $(t_0, t_0 + T)$ with the initial data a at t_0 . Moreover, v has the additional property $v \in BC([t_0, t_0 + T]; L^{q^*}_\sigma)$ with*

$$(4.1) \quad \sup_{t_0 \leq t < t_0 + T} \|v(t)\|_{q^*} \leq C_2,$$

where $C_2 = C_2(\|a\|_{q^*}, \|Pf\|_{BC(\mathbf{R}; L^l)})$ is independent of t_0 . Here T is estimated as

$$(4.2) \quad T = C_3(\|a\|_{q^*} + \|Pf\|_{BC(\mathbf{R}; L^l)})^{-2q^*/(q^* - n)}$$

with $C_3 = C_3(n, q, l)$ independent of a, f and t_0 .

REMARK. When $f \equiv 0$, Giga [7, Theorem 4] obtained (4.2) by making use of the fact that $A_r^{-1/2} P_r(\partial/\partial x_j)$ ($j = 1, \dots, n$) is a bounded operator from L^r into L^r_σ for $1 < r < \infty$. Our proof below seems to be rather elementary; we use Lemma 2.1 and integration by parts.

PROOF OF LEMMA 4.1. The proof is similar to that of Kato [12] and Giga [7]. However, we give it for completeness. It suffices only to construct the solution v of the integral equation:

$$(4.3) \quad v(t) = e^{-(t-t_0)A} a + \int_{t_0}^t e^{-(t-s)A} Pf(s) ds - \int_{t_0}^t e^{-(t-s)A} P(v \cdot \nabla v)(s) ds, \quad t_0 < t < t_0 + T$$

in the class

$$(4.4) \quad v \in BC([t_0, t_0 + T]; L^n_\sigma \cap L^{q^*}_\sigma) \quad \text{with} \quad (t-t_0)^{1/2} \nabla v \in BC([t_0, t_0 + T]; L^n).$$

Indeed, with the aid of Kozono-Ogawa [13, Lemma A.4], the assumption on Pf and a general theory of holomorphic semigroup guarantee that the solution v of (4.3) in the class (4.4) satisfies the properties (i), (ii) and (iii) of Lemma 4.1 (see, e.g., Tanabe [24, Theorem 3.3.4]).

Since this lemma deals with only local existence of solutions, we may assume that $0 < T \leq 1$. Let us solve (4.3) by successive approximation:

$$(4.5) \quad v_0(t) = e^{-(t-t_0)A} a + \int_{t_0}^t e^{-(t-s)A} Pf(s) ds,$$

$$(4.6) \quad v_{m+1}(t) = v_0(t) - \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds.$$

Taking $\alpha = n/q - 1$, we have by assumption $0 < \alpha < 1$ and $q^* = n/\alpha$. Let us first show

$$(4.7) \quad \sup_{t_0 < t < t_0 + T} (t-t_0)^{(1-\alpha)/2} \|v_m(t)\|_{n/\alpha} \leq K_m, \quad m = 0, 1, \dots$$

with some constant K_m . In fact, by Lemma 2.1 and the Sobolev inequality (2.6), we have

$$\begin{aligned}
\|v_0(t)\|_{n/\alpha} &\leq \|e^{-(t-t_0)A}a\|_{n/\alpha} + \int_{t_0}^t \|e^{-(t-s)A}Pf(s)\|_{n/\alpha} ds \\
&\leq C\|a\|_{n/\alpha} + C \int_{t_0}^t \|\nabla e^{-(t-s)A}Pf(s)\|_q ds \\
&\leq C\|a\|_{n/\alpha} + C \int_{t_0}^t (t-s)^{-n(1/l-1/q)/2-1/2} \|Pf(s)\|_l ds \\
&\leq C\|a\|_{n/\alpha} + C\|Pf\|_{BC(\mathbf{R}; L^1_{\sigma})}(t-t_0)^{-(1-\alpha)/2+3/2-n/2l}
\end{aligned}$$

for all $t_0 < t < t_0 + T$ with $C = C(n, q, l)$ independent of t_0 . Since $1/l < 1/q + 1/n$, we have $(1-\alpha)/2 < (3-n/l)/2$ and hence the above estimate yields

$$\sup_{t_0 < t < t_0 + T} (t-t_0)^{(1-\alpha)/2} \|v_0(t)\|_{n/\alpha} \leq C_4 T^{(1-\alpha)/2} (\|a\|_{n/\alpha} + \|Pf\|_{BC(\mathbf{R}; L^1_{\sigma})}).$$

Then K_0 may be chosen as

$$(4.8) \quad K_0 = C_4 T^{(1-\alpha)/2} (\|a\|_{n/\alpha} + \|Pf\|_{BC(\mathbf{R}; L^1_{\sigma})}),$$

where $C_4 = C_4(n, q, l)$ is independent of t_0 . Suppose that (4.7) is true. By Lemma 2.1 and integration by parts there holds

$$\begin{aligned}
\left| \left(- \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds, \phi \right) \right| &= \left| \int_{t_0}^t (v_m(s) \cdot \nabla e^{-(t-s)A} \phi, v_m(s)) ds \right| \\
&\leq \int_{t_0}^t \|v_m \otimes v_m(s)\|_{n/2\alpha} \|\nabla e^{-(t-s)A} \phi\|_{n/(n-2\alpha)} ds \\
&\leq C \int_{t_0}^t (t-s)^{-\alpha/2-1/2} \|v_m(s)\|_{n/\alpha}^2 ds \cdot \|\phi\|_{n/(n-\alpha)} \\
&\leq CB((1-\alpha)/2, \alpha) K_m^2 (t-t_0)^{-(1-\alpha)/2} \|\phi\|_{n/(n-\alpha)}
\end{aligned}$$

for all $\phi \in C_{0,\sigma}^\infty$ and all $t \in (t_0, t_0 + T)$, where $B(\cdot, \cdot)$ denotes the beta function and $C = C(n, q)$. By duality we have

$$\left\| \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds \right\|_{n/\alpha} \leq C_5 K_m^2 (t-t_0)^{-(1-\alpha)/2}, \quad t_0 < t < t_0 + T$$

and hence we may define K_{m+1} as

$$(4.9) \quad K_{m+1} \equiv K_0 + C_5 K_m^2,$$

where $C_5 = C_5(n, q)$ is independent of t_0 . An elementary consideration shows that if

$$(4.10) \quad K_0 < \frac{1}{4C_5},$$

then there holds

$$(4.11) \quad K_m \leq \frac{1 - \sqrt{1 - 4C_5 K_0}}{2C_5} \equiv k < \frac{1}{2C_5} \quad \text{for all } m = 0, 1, \dots$$

Assume (4.10) for a moment. Then in the same way as in (3.8), the uniform estimate (4.11) with respect to m yields a function v with $(t - t_0)^{(1-\alpha)/2} v(\cdot) \in BC([t_0, t_0 + T]; L_\sigma^{n/\alpha})$ such that

$$(4.12) \quad \lim_{m \rightarrow \infty} \sup_{t_0 < t < t_0 + T} (t - t_0)^{(1-\alpha)/2} \|v_m(t) - v(t)\|_{n/\alpha} = 0.$$

We shall next show that if K_0 is sufficiently small, then the limit v also satisfies $(t - t_0)^{1/2} \nabla v(\cdot) \in BC([t_0, t_0 + T]; L^n)$ with

$$(4.13) \quad \lim_{m \rightarrow \infty} \sup_{t_0 < t < t_0 + T} (t - t_0)^{1/2} \|\nabla v_m(t) - \nabla v(t)\|_n = 0.$$

To this end, let us prove that

$$(4.14) \quad \sup_{t_0 < t < t_0 + T} (t - t_0)^{1/2} \|\nabla v_m(t)\|_n \leq L_m \quad \text{for } m = 0, 1, \dots$$

By Lemma 2.1, there holds

$$\|\nabla v_0\|_n \leq C(t - t_0)^{-1/2} \|a\|_n + C(t - t_0)^{1/2} \|Pf\|_{BC((t_0, t_0 + T); L_\sigma^n)} \quad \text{for all } t \in (t_0, t_0 + T)$$

and hence we may take L_0 as

$$L_0 \equiv C(\|a\|_n + \|Pf\|_{BC((t_0, t_0 + T); L_\sigma^n)}),$$

where $C = C(n)$ is independent of t_0 . Moreover, it follows from (4.7), (4.11) and (4.14) that

$$\begin{aligned} \left\| \nabla \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds \right\|_n &\leq \int_{t_0}^t (t-s)^{-n(1/n + \alpha/n - 1/n)/2 - 1/2} \|v_m(s)\|_{n/\alpha} \|\nabla v_m(s)\|_n ds \\ &\leq CK_m L_m \int_{t_0}^t (t-s)^{-\alpha/2 - 1/2} (s-t_0)^{\alpha/2 - 1} ds \\ &\leq C_6 k L_m (t-t_0)^{-1/2} \end{aligned}$$

for all $t_0 < t < t_0 + T$, where $C_6 = C_6(n, q)$ is independent of t_0 . Hence we may take L_{m+1} as

$$L_{m+1} \equiv L_0 + C_6 k L_m,$$

which shows that $\{L_m\}_{m=0}^\infty$ is a *linear recurrence*. If

$$(4.15) \quad k < 1/C_6,$$

then we have a uniform bound of $\{L_m\}_{m=0}^\infty$ as

$$L_m \leq \frac{L_0}{1 - C_6 k} \equiv L \quad \text{for all } m=0, 1, \dots$$

Assume (4.15) for a moment. Then it is easy to see that the limit v satisfies (4.13).

To prove $v \in BC([t_0, t_0 + T]; L_\sigma^n \cap L_\sigma^{q^*})$, we need to show

$$(4.16) \quad \sup_{t_0 < t < t_0 + T} \|v_m(t)\|_{n/\mu} \leq M_{\mu,m} \quad (m=0, 1, \dots) \quad \text{for } \mu=\alpha \text{ and } \mu=1.$$

Calculation similar to (4.7) and (4.14) yields $M_{\mu,0}$ as

$$M_{\alpha,0} = C(\|a\|_{n/\alpha} + \|Pf\|_{BC(\mathbf{R}; L_\sigma^1)}), \quad M_{1,0} = C(\|a\|_n + \|Pf\|_{BC([t_0, t_0 + T]; L_\sigma^n)}),$$

where $C = C(n, l, \mu)$ is independent of t_0 . Notice that $0 < T \leq 1$. Suppose that (4.16) is true. Then by Lemma 2.1, (4.11) and integration by parts, we have

$$\begin{aligned} & \left| \left(- \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds, \phi \right) \right| \\ & \leq \left| \int_{t_0}^t (v_m(s) \cdot \nabla e^{-(t-s)A} \phi, v_m(s)) ds \right| \\ & \leq \int_{t_0}^t \|v_m(s)\|_{n/\alpha} \|v_m(s)\|_{n/\mu} \|\nabla e^{-(t-s)A} \phi\|_{n/(n-\alpha-\mu)} ds \\ & \leq CK_m M_{\mu,m} \int_{t_0}^t (t-s)^{-\alpha/2-1/2} (s-t_0)^{-(1-\alpha)/2} ds \cdot \|\phi\|_{n/(n-\mu)} \\ & \leq Ck M_{\mu,m} B((1-\alpha)/2, (1+\alpha)/2) \|\phi\|_{n/(n-\mu)} \end{aligned}$$

for all $\phi \in C_{0,\sigma}^\infty$ and all $t_0 < t < t_0 + T$, where $C = C(n, q, \mu)$. By duality we may take $M_{\mu,m+1}$ as

$$M_{\mu,m+1} = M_{\mu,0} + C_7 k M_{\mu,m} \quad \text{for } \mu=\alpha, 1,$$

where $C_7 = C_7(n, q, \mu)$ is independent of t_0 . If

$$(4.17) \quad k < 1/C_7,$$

then there holds

$$M_{\mu,m} \leq \frac{M_{\mu,0}}{1 - C_7 k} \quad (\mu=\alpha, 1) \quad \text{for all } m=0, 1, \dots,$$

which yields $v \in BC([t_0, t_0 + T]; L_\sigma^n \cap L_\sigma^{n/\alpha})$ with

$$(4.18) \quad \lim_{m \rightarrow \infty} \sup_{t_0 < t < t_0 + T} \|v_m(t) - v(t)\|_{n/\mu} = 0 \quad \text{for } \mu = \alpha, 1.$$

In particular, the constant C_2 in (4.1) can be given as $C_2 = M_{\alpha,0}/(1 - C_7k)$. Now we see that under the conditions (4.10), (4.15) and (4.17), the limit v belongs to the class in (4.4). Moreover, there holds

$$(4.19) \quad \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds \rightarrow \int_{t_0}^t e^{-(t-s)A} P(v \cdot \nabla v)(s) ds \quad \text{in } L_\sigma^n$$

uniformly in $t \in [t_0, t_0 + T)$ as $m \rightarrow \infty$. Indeed, by Lemma 2.1, (4.11) and (4.14) we have

$$\begin{aligned} & \left\| \int_{t_0}^t e^{-(t-s)A} P(v_m \cdot \nabla v_m)(s) ds - \int_{t_0}^t e^{-(t-s)A} P(v \cdot \nabla v)(s) ds \right\|_n \\ & \leq \int_{t_0}^t \|e^{-(t-s)A} P((v_m - v) \cdot \nabla v_m)(s)\|_n ds + \int_{t_0}^t \|e^{-(t-s)A} P(v \cdot \nabla (v_m - v))(s)\|_n ds \\ & \leq \int_{t_0}^t (t-s)^{-n(\alpha/n + 1/n - 1/n)/2} \|v_m(s) - v(s)\|_{n/\alpha} \|\nabla v_m(s)\|_n ds \\ & \quad + \int_{t_0}^t (t-s)^{-n(\alpha/n + 1/n - 1/n)/2} \|v(s)\|_{n/\alpha} \|\nabla v_m(s) - \nabla v(s)\|_n ds \\ & \leq CB(1 - \alpha/2, \alpha/2) \left(L \sup_{t_0 < s < t_0 + T} (s - t_0)^{(1-\alpha)/2} \|v_m(s) - v(s)\|_{n/\alpha} \right. \\ & \quad \left. + k \sup_{t_0 < s < t_0 + T} (s - t_0)^{1/2} \|\nabla v_m(s) - \nabla v(s)\|_n \right) \end{aligned}$$

for all $t_0 < t < t_0 + T$, from which and (4.12–13) we obtain (4.19). Now, letting $m \rightarrow \infty$ in (4.6), we see by (4.18) and (4.19) that v is a solution of (4.3). The proof for uniqueness is standard, so we may omit it (see [3], [8]).

It remains to estimate the time-interval T of existence in terms of the prescribed data. Since k is determined by (4.11), there exists a constant $\kappa = \kappa(n, q, l)$ independent of t_0 such that if $K_0 \leq \kappa$, then all conditions (4.10), (4.15) and (4.17) are satisfied. Now from (4.8) we see that T may be chosen as

$$T \equiv \left(\frac{\kappa}{C_4(\|a\|_{n/\alpha} + \|Pf\|_{BC(\mathbf{R}; L_\sigma^l)})} \right)^{2/(1-\alpha)},$$

which shows (4.2) and proves Lemma 4.1.

PROOF OF THEOREM 2. Let u be the periodic solution of the integral equation (I.E.) given by Theorem 1. Since $u \in Y$, we have by (2.6) that $u \in BC(\mathbf{R}; L_\sigma^n \cap L_\sigma^{q^*})$, where $q^* = nq/(n - q)$. Let

$$T = C_3 (\|u\|_{BC(\mathbf{R}; L_{q^*}^s)} + \|Pf\|_{BC(\mathbf{R}; L_{q^*}^1)})^{-2q^*/(q^*-n)},$$

where C_3 is the same constant as in (4.2). Then by Lemma 4.1, for every $t_0 \in \mathbf{R}$ there exists a unique strong solution v of (N-S) on $(t_0, t_0 + T)$ with the initial data $u(t_0)$. By (3.6) and (4.1), we have

$$(4.20) \quad \sup_{t_0 < s < t_0 + T} \|v(s)\|_{q^*} + \sup_{t_0 < s < t_0 + T} \|\nabla u(s)\|_q \leq C_2 + K \equiv C_8,$$

where C_8 is independent of t_0 . By (4.3) with a replaced by $u(t_0)$ and by (I.E.), it is easy to see

$$(4.21) \quad \begin{aligned} u(t) - v(t) &= - \int_{t_0}^t e^{-(t-s)A} P(u \cdot \nabla u)(s) ds + \int_{t_0}^t e^{-(t-s)A} P(v \cdot \nabla v)(s) ds \\ &= - \int_{t_0}^t e^{-(t-s)A} P((u-v) \cdot \nabla u)(s) ds - \int_{t_0}^t e^{-(t-s)A} P(v \cdot \nabla(u-v))(s) ds \\ &\equiv J_1(t) + J_2(t), \quad t_0 < t < t_0 + T. \end{aligned}$$

By Lemma 2.1 there holds

$$(4.22) \quad \begin{aligned} \|J_1(t)\|_n &\leq C \int_{t_0}^t (t-s)^{-n(1/n+1/q-1/n)/2} \|u(s) - v(s)\|_n \|\nabla u(s)\|_q ds \\ &\leq C \sup_{s \in \mathbf{R}} \|\nabla u(s)\|_q \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n (t-t_0)^{1-n/2q}, \end{aligned}$$

for all $t_0 < t < t_0 + T$, where $C = C(n, q)$ is independent of t_0 . By integration by parts we have

$$\begin{aligned} |(J_2(t), \phi)| &= \left| \int_{t_0}^t (v(s) \cdot \nabla e^{-(t-s)A} \phi, u(s) - v(s)) ds \right| \\ &\leq C \int_{t_0}^t \|v(s)\|_{q^*} \|\nabla e^{-(t-s)A} \phi\|_{q'} \|u(s) - v(s)\|_n ds \\ &\leq C \sup_{t_0 < s < t_0 + T} \|v(s)\|_{q^*} \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n \int_{t_0}^t (t-s)^{-n(1/n'-1/q')/2-1/2} ds \|\phi\|_n \\ &\leq C \sup_{t_0 < s < t_0 + T} \|v(s)\|_{q^*} \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n (t-t_0)^{1-n/2q} \|\phi\|_n, \end{aligned}$$

for all $\phi \in C_{0,\sigma}^\infty$ and all $t_0 < t < t_0 + T$, where $C = C(n, q)$. By duality,

$$(4.23) \quad \|J_2(t)\|_n \leq C \sup_{t_0 < s < t_0 + T} \|v(s)\|_{q^*} \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n (t-t_0)^{1-n/2q}$$

for all $t_0 < t < t_0 + T$. Now it follows from (4.20–23) that

$$\|u(t) - v(t)\|_n \leq C_9 \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n (t - t_0)^{1 - n/2q}, \quad t_0 < t < t_0 + T$$

with C_9 independent of t_0 . Defining $\tau \equiv \min\{(1/2C_9)^{2q/(2q-n)}, T\}$, we obtain from the above estimate that

$$\begin{aligned} \|u(t) - v(t)\|_n &\leq C_9 \tau^{1 - n/2q} \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n \\ &\leq \frac{1}{2} \sup_{t_0 < s < t_0 + t} \|u(s) - v(s)\|_n \end{aligned}$$

for all $t_0 \leq t \leq t_0 + \tau$, which yields

$$u \equiv v \quad \text{on} \quad [t_0, t_0 + \tau].$$

Since τ can be taken independently of t_0 , we have

$$u \equiv v \quad \text{on} \quad [t_0, t_0 + T].$$

Now, since t_0 is arbitrary, it follows from Lemma 4.1 that u has the desired properties (i), (ii) and (iii) in Theorem 2. \square

REFERENCES

- [1] W. BORCHERS AND T. MIYAKAWA, L^2 decay for Navier-Stokes flow in half spaces, *Math. Ann.* 282 (1988), 139–155.
- [2] W. BORCHERS AND T. MIYAKAWA, Algebraic L^2 decay for Navier-Stokes flows in exterior domains, *Acta Math.* 165 (1990), 189–227.
- [3] H. FUJITA AND T. KATO, On the Navier-Stokes initial value problem I, *Arch. Rational Mech. Anal.* 46 (1964), 269–315.
- [4] D. FUJIWARA AND H. MORIMOTO, An L^r -theorem of the Helmholtz decomposition of vector fields, *J. Fac. Sci. Univ. Tokyo, Sec. IA* 24 (1977), 685–700.
- [5] Y. GIGA, Analyticity of the semigroup generated by the Stokes operator in L_r spaces, *Math. Z.* 187 (1981), 297–329.
- [6] Y. GIGA, Domains of fractional powers of the Stokes operator in L_r spaces, *Arch. Rational Mech. Anal.* 89 (1985), 251–265.
- [7] Y. GIGA, Solutions for semilinear parabolic equations in L^p and regularity of weak solutions of the Navier-Stokes system, *J. Differential Equations* 62 (1986), 186–212.
- [8] Y. GIGA AND T. MIYAKAWA, Solution in L_r of the Navier-Stokes initial value problem, *Arch. Rational Mech. Anal.* 89 (1985), 267–281.
- [9] Y. GIGA AND H. SOHR, On the Stokes operator in exterior domains, *J. Fac. Sci. Univ. Tokyo, Sec. IA.* 36 (1989), 103–130.
- [10] H. IWASHITA, $L_q - L_r$ estimates for solutions of the nonstationary Stokes equations in an exterior domain and the Navier-Stokes initial value problems in L_q spaces, *Math. Ann.* 285 (1989), 265–288.
- [11] S. KANIEL AND M. SHINBROT, A reproductive property of the Navier-Stokes equations, *Arch. Rational Mech. Anal.* 24 (1967), 265–288.
- [12] T. KATO, Strong L^p -solution of the Navier-Stokes equation in \mathbf{R}^m , with applications to weak solutions,

- Math. Z. 187 (1984), 471–480.
- [13] H. KOZONO AND T. OGAWA, Some L^p estimate for the exterior Stokes flow and an application to the nonstationary Navier-Stokes equations, *Indiana Univ. Math. J.* 41 (1992), 789–808.
- [14] O. A. LADYZHENSKAYA, *The Mathematical Theory of Viscous Incompressible Flow*, Gordon and Breach, New York, 1969.
- [15] P. MAREMONTI, Existence and stability of time periodic solution of the Navier-Stokes equations in the whole space, *Nonlinearity* 4 (1991), 503–529.
- [16] P. MAREMONTI, Some theorems of existence for solutions of the Navier-Stokes equations with slip boundary conditions in half-space, *Rich. Mat.* 40 (1991), 81–135.
- [17] T. MIYAKAWA, On nonstationary solutions of the Navier-Stokes equations in an exterior domain, *Hiroshima Math. J.* 12 (1982), 115–140.
- [18] T. MIYAKAWA AND Y. TERAMOTO, Existence and periodicity of weak solutions of the Navier-Stokes equations in a time dependent domain, *Hiroshima Math. J.* 12 (1982), 513–528.
- [19] H. MORIMOTO, On existence of periodic weak solutions of the Navier-Stokes equations in regions with periodically moving boundaries, *J. Fac. Sci. Univ. Tokyo Sec. IA* 18 (1970/1971), 499–524.
- [20] J. SERRIN, A note on the existence of periodic solutions of the Navier-Stokes equations, *Arch. Rational Mech. Anal.* 3 (1959), 120–122.
- [21] J. SERRIN, The initial value problem for the Navier-Stokes equations, *Nonlinear Problem*, (R. Langer, ed.), The University of Wisconsin Press, Madison, 1960, pp. 69–98.
- [22] G. G. SIMADER AND H. SOHR, A new approach to the Helmholtz decomposition and the Neumann problem in L^q -spaces for bounded and exterior domains, “*Mathematical Problems Relating to the Navier-Stokes Equations*”, series on *Advanced in Mathematics for Applied Sciences* (G. P. Galdi, ed.), (1992), World Scientific, Singapore-New Jersey-London-Hong Kong, 1–35.
- [23] A. TAKESHITA, On the reproductive property of 2-dimensional Navier-Stokes equations, *J. Fac. Sci. Univ. Tokyo Sec. IA* 16 (1970), 297–311.
- [24] H. TANABE, *Equations of Evolution*, Pitman, London, 1979.
- [25] Y. TERAMOTO, On the stability of periodic solutions of the Navier-Stokes equations in a noncylindrical domain, *Hiroshima Math. J.* 13 (1983), 607–625.
- [26] S. UKAI, A solution formula for the Stokes equation in \mathbb{R}_+^n , *Comm. Pure Appl. Math.* 40 (1987), 611–621.
- [27] W. VON WAHL, The equations of Navier-Stokes and abstract parabolic equations, Vieweg, Braunschweig-Wiessbaden, 1985.

GRADUATE SCHOOL OF POLYMATHEMATICS
 NAGOYA UNIVERSITY
 NAGOYA 464-01
 JAPAN

GRADUATE SCHOOL OF MATHEMATICS
 KYUSHU UNIVERSITY
 ROPPONMATSU
 FUKUOKA 810
 JAPAN