

STABILITY OF A STATIONARY SOLUTION FOR THE LUGIATO-LEFEVER EQUATION

TOMOYUKI MIYAJI, ISAMU OHNISHI AND YOSHIO TSUTSUMI

(Received January 26, 2011, revised May 6, 2011)

Abstract. We study the stability of a stationary solution for the Lugiato-Lefever equation with the periodic boundary condition in one space dimension, which is a damped and driven nonlinear Schrödinger equation introduced to model the optical cavity. In this paper, we prove the Strichartz estimates for the linear damped Schrödinger equation with potential and external forcing and investigate the stability of certain stationary solutions under the initial perturbation within the framework of L^2 .

1. Introduction and main theorem. We consider the stability of a stationary solution for the nonlinear Schrödinger equation with damping and spatially homogeneous forcing terms:

$$(1) \quad \frac{\partial}{\partial t} A = -(1 + i\theta)A + ib^2 \frac{\partial^2}{\partial x^2} A + i|A|^2 A + F, \\ t > 0, \quad x \in T = \mathbf{R}/2\pi\mathbf{Z}.$$

Here, A denotes the slowly varying envelope of the electric field, $\theta > 0$ denotes the detuning parameter, and $b > 0$ denotes the diffraction parameter. Let $F > 0$ be the spatially homogeneous input field. In [12], Lugiato and Lefever present the equation (1) to model the so-called cavity soliton in the ring or the Fabry-Pérot cavity oscillator (see also [1], [2], [3], [9]). The existence and the stability of spatially nonhomogeneous stationary solutions for (1) have been studied by the authors [13]. In this paper, we prove the Strichartz estimate for the linear Schrödinger equation with potential and external forcing and investigate the stability of certain stationary solutions for (1) given in [13] under the L^2 perturbation. We decompose the solution $A(t)$ of (1) into effective dynamical components, following Buslaev and Perel'man [6] (see also Soffer and Weinstein [15]) and show the a priori estimates of those effective dynamical components, which ensure the asymptotic stability of stationary solution for (1). There are many papers on the asymptotic stability of a family of equilibria in the setting of nonlinear parabolic equation (see, e.g., Exercise 6 in Henry [11, Section 5.1]). In the case of nonlinear parabolic equation, the smoothing property and the fractional power of infinitesimal generator for the holomorphic semigroup are useful. But, in the case of nonlinear Schrödinger equation, the Strichartz estimate plays a crucial role, which enables us to treat the rougher perturbation than the H^1 perturbation in the previous papers (see, e.g., Ghidaglia [10], X.-M. Wang [16], and Miyaji, Ohnishi and Tsutsumi [13]).

THEOREM 1.1. *Assume D is a stationary solution of (1) such that the spectrum of the linearized operator around D for the stationary equation associated with (1) lies in $\{z \in \mathbf{C}; \operatorname{Re} z \leq -\alpha\} \cup \{0\}$ for some $\alpha > 0$ and the eigenspace corresponding to the zero eigenvalue is a one dimensional subspace in L^2 spanned by $\partial_x D$. Let $A_0 \in L^2$ and let $\varepsilon > 0$. For $c \in \mathbf{R}$, we put*

$$D_c(x) = D(x + c).$$

Then, there exist $\delta > 0$ and $0 \leq c_0 < 2\pi$ such that if the initial data A_0 satisfies

$$\|A_0 - D\|_{L^2} < \delta,$$

we have

$$\begin{aligned} \sup_{t \geq 0} [\inf_{0 \leq c < 2\pi} \|A(t) - D_c\|_{L^2}] &< \varepsilon, \\ \|A(t) - D_{c_0}\|_{L^2} &\rightarrow 0 \quad (t \rightarrow \infty), \end{aligned}$$

where A is a solution of (1) with $A(0) = A_0$.

REMARK 1.1. (i) It is known that there exists a spatially nonhomogeneous stationary solution of (1) satisfying all the assumptions in Theorem 1.1 (see Theorems 2.1 and 2.2 in Section 2).

(ii) If we consider the equation (1) and the initial perturbation within the framework of H^1 , it would be slightly easier to prove Theorem 1.1. Indeed, $H^1 \subset L^\infty$ in the one dimensional case and the a priori estimates in H^1 needed for the proof of Theorem 1.1 follows from the energy estimates only. We note that Theorem 1.1 can cover the L^2 perturbation, which belongs to a bigger class than H^1 .

The present paper is organized as follows. In Section 2, we summarize results on the existence of a stationary solution and the spectral analysis of the linearized operator, which are mainly proved by the authors [13]. In Section 3, we show the Strichartz estimate of the linear Schrödinger equation with complex potential and shift terms, which plays a crucial role in the proof of Theorem 1.1. Finally, in Section 4, we give a sketch of the proof of Theorem 1.1.

2. Existence of a stationary solution and linear stability. In this section, we summarize the results on the existence of a stationary solution and the spectrum of the linearized operator, which have been obtained by the authors [13]. Let the spatially homogeneous stationary solutions A_S be defined as follows.

$$(2) \quad A_S = \frac{F}{1 + i(\theta - \alpha)}, \quad \alpha = |A_S|^2,$$

where α is uniquely determined by the condition

$$(3) \quad F^2 = \alpha\{1 + (\alpha - \theta)^2\}, \quad \theta < \sqrt{3}.$$

We now consider the stationary Lugiato-Lefever Equation and change the unknown function A to B as follows. Set $A = A_S(1 + B)$. Then, B satisfies

$$(4) \quad 0 = -(1 + i\theta)B + ib^2\partial_x^2 B + i\alpha(2B + \bar{B} + B^2 + 2|B|^2 + |B|^2 B), \quad x \in T.$$

REMARK 2.1. Instead of F , we regard α as a bifurcation parameter. In that case, A_S and F are determined for given α through (2), (3).

We first state the existence theorem of spatially nonhomogeneous stationary solutions which bifurcate from the spatially homogeneous stationary solution A_S .

THEOREM 2.1. *There exist $b > 0$, $\sqrt{3} > \theta > 0$, $\eta > 0$, $n \in \mathbf{N}$, $B_0 \in \mathbf{C}$ such that the equation (4) has a family of solutions $\{(\alpha(s), B(s)) \in \mathbf{R} \times H^2; -\eta < s < \eta\}$ satisfying the following conditions*

$$\begin{aligned} B(s) &= sB_0 \cos(2\pi nx) + r(s), \quad s \in (-\eta, \eta), \\ \|r(s)\|_{H^2} &= o(s) \quad (s \rightarrow 0), \\ r(s) &\perp \text{span}\{\cos(2\pi nx)\}, \\ r(\cdot, x) &= r(\cdot, -x), \quad x \in T, \\ \alpha(s) &= 1 - \frac{30\theta - 41}{9(2 - \theta)^2} s^2 + o(s^2) \quad (s \rightarrow 0). \end{aligned}$$

For the proof of Theorem 2.1, see Theorem 3.1 in [13, pp. 2071, 2072].

We next consider the spectrum of the linearized operator and the linear stability of stationary solutions given by Theorem 2.1. We set

$$A_S(1 + B(s)) = w(s) + iz(s), \quad s \in (-\eta, \eta),$$

where w and z are real-valued functions.

Let L be the linearized operator around (w, z) for (1):

$$(5) \quad L = \begin{pmatrix} -1 - 2wz & -\Delta_{b,\theta} - 2V_+ + V_- \\ \Delta_{b,\theta} + 2V_+ + V_- & -1 + 2wz \end{pmatrix},$$

where

$$\begin{aligned} \Delta_{b,\theta} &= b^2\partial_x^2 - \theta, \\ V_{\pm} &= w^2 \pm z^2. \end{aligned}$$

THEOREM 2.2. *Assume $0 < \theta < 41/30$. Then, there exist $\eta' > 0$ and $\gamma \in C((-\eta', \eta'); \mathbf{R})$ such that $\gamma(s) > 0$ ($0 < |s| < \eta'$), $\gamma(0) = 0$ and*

$$\sigma(L) \subset \{z \in \mathbf{C}; \text{Re } z \leq -\gamma\} \cup \{0\} \quad (0 < |s| < \eta').$$

In addition, when $\eta' > |s| > 0$, the eigenspace belonging to the zero eigenvalue of L consists of the derivative of the stationary solution (w, z) , which is an odd function.

For the proof of Theorem 2.2, see [13, pp. 2071, 2072].

REMARK 2.2. We can completely analyze the spectrum of the linearized operator near the bifurcation point. Because at the bifurcation point, the linearized operator around the homogeneous stationary solution is reduced to the Sturm-Liouville operator with constant coefficients.

REMARK 2.3. Theorem 2.2 implies the linear stability of stationary solutions given by Theorem 2.1 for $\theta < 41/30$ within the framework of even functions (in fact, the nonlinear stability is proved in [13]). In the ODE case, it is well known that when all eigenvalues of the linearized operator have negative real part except for the zero eigenvalue and every orbit starting from a neighborhood of stationary solution has a non-empty ω -limit set, the stationary solution is nonlinearly stable (see, e.g., [4, Proposition 1.1]). In the nonlinear PDE case, the situations are more complicated, but suitable a priori estimates often ensure the similar conclusion as the ODE case.

REMARK 2.4. It can be proved that when $\theta > 41/30$, the stationary solution given by Theorem 2.1 is nonlinearly unstable (see [13]).

3. Strichartz estimate. In this section, we prove the global Strichartz estimate in time for the Schrödinger equation on one-dimensional torus with linear time-independent potential and external forcing.

$$(6) \quad i\partial_t u = -\partial_x^2 u + Vu + f, \quad t > 0, x \in \mathbf{T},$$

$$(7) \quad u(0, x) = u_0(x).$$

We assume the following two hypotheses:

(A1) V is a complex-valued function in $L^\infty(\mathbf{T})$,

(A2) There exists $\gamma > 0$ such that

$$\sigma(i(\partial_x^2 - V)) \subset \{z \in \mathbf{C}; \operatorname{Re} z \leq -\gamma\}.$$

REMARK 3.1. (i) We note that (A2) is equivalent to the condition that there exists $\gamma > 0$ such that

$$\operatorname{Re}(-i(\partial_x^2 - V)v, v) \geq \gamma \|v\|_{L^2}^2, \quad v \in H^1.$$

This implies that the potential V has a damping effect, which yields the global Strichartz estimate in time for (6).

(ii) The linearized equation of (1) has the form : $i\partial_t u = -\partial_x^2 u + V_1 u + V_2 \bar{u}$. This includes the conjugate linear term $V_2 \bar{u}$. But the Strichartz estimate of this equation can be proved in the same way, if we replace (A2) by the condition that there exists $\gamma > 0$ such that

$$\operatorname{Re}(-i(\partial_x^2 - V_1)v + iV_2 \bar{v}, v) \geq \gamma \|v\|_{L^2}^2, \quad v \in H^1.$$

We set

$$U_0(t) = e^{it\partial_x^2}, \quad U(t) = e^{it(\partial_x^2 - V)}, \quad \mathbf{R}_+ = (0, \infty).$$

We first give two lemmas, which are useful for the proof of the Strichartz estimate of (6), (7). We begin with the Christ-Kiselev lemma (see [8, Theorem 1.2]).

LEMMA 3.1. *Let X, Y be Banach spaces and let $T > 0$. Assume $K(t, s)$ is continuous from $[0, T] \times [0, T]$ to $B(X, Y)$ and that $1 \leq p < q \leq \infty$. We put*

$$Sf(t) = \int_0^T K(t, s)f(s) ds,$$

$$\tilde{S}f(t) = \int_0^t K(t, s) f(s) ds .$$

Assume that

$$\|Sf\|_{L^q((0,T);Y)} \leq C \|f\|_{L^p((0,T);X)} .$$

Then, we have

$$\|\tilde{S}f\|_{L^q((0,T);Y)} \leq \tilde{C} \|f\|_{L^p((0,T);X)} .$$

For the proof of Lemma 3.1, see [14, Lemma 3.1].

The following lemma is concerned with the Strichartz estimate of (6), (7) without potential (see [5, Propositions 2.1, 2.33]), which is traced back to Zygmund [17].

LEMMA 3.2. *We have*

$$\begin{aligned} \|U_0(\cdot)u_0\|_{L^4((0,2\pi)\times T)} &\leq C \|u_0\|_{L^2(T)} , \\ \left\| \int_{t_0}^t U_0(t-s)f(s) ds \right\|_{L^4((t_0,t_0+2\pi)\times T)} &\leq C \|f\|_{L^{4/3}((t_0,t_0+2\pi)\times T)} , \end{aligned}$$

where t_0 is an arbitrary real number.

For the convenience of the reader, we present the proof of Lemma 3.2.

PROOF. We show the first inequality. By the Fourier expansion, we have

$$\begin{aligned} u_0 &= \frac{1}{\sqrt{2\pi}} \sum_{m=-\infty}^{\infty} \hat{u}_0(m) e^{imx} , \\ \hat{u}_0(m) &= \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} u_0(x) e^{-imx} dx . \end{aligned}$$

Then, since we have

$$U_0(t)u_0 = \frac{1}{\sqrt{2\pi}} \sum_{m=-\infty}^{\infty} \hat{u}_0(m) e^{-i(m^2t-mx)} ,$$

a simple computation yields

$$\begin{aligned} &\int_T \int_0^{2\pi} |U_0(t)u_0|^4 dt dx \\ (8) \quad &= \frac{1}{(2\pi)^2} \times \int_T \int_0^{2\pi} \sum_{\substack{m_1, m_2, \\ m_3, m_4}} \hat{u}_0(m_1) \hat{u}_0(m_2) \overline{\hat{u}_0(m_3) \hat{u}_0(m_4)} \\ &\quad \times e^{-i(m_1^2+m_2^2-m_3^2-m_4^2)t} e^{i(m_1+m_2-m_3-m_4)x} dt dx . \end{aligned}$$

On the right-hand side of (8), the integrals vanish unless

$$\begin{aligned} m_1^2 + m_2^2 - m_3^2 - m_4^2 &= 0 , \quad m_1 + m_2 - m_3 - m_4 = 0 \\ \iff \begin{cases} m_1 = m_3 , \\ m_2 = m_4 , \end{cases} \quad \text{or} \quad \begin{cases} m_1 = m_4 , \\ m_2 = m_3 . \end{cases} \end{aligned}$$

Thus, the right-hand side of (8) is equal to:

$$2\left(\sum_{m_1} |\hat{u}_0(m_1)|^2\right)\left(\sum_{m_2} |\hat{u}_0(m_2)|^2\right) = 2\|u_0\|_{L^2(\mathcal{T})}^4.$$

We next show the second inequality. We put

$$G = \int_{t_0}^{t_0+2\pi} U_0(t-s)f(s) ds.$$

The first inequality proved above yields

$$(9) \quad \|G\|_{L^4((t_0, t_0+2\pi) \times \mathcal{T})} \leq C \left\| \int_{t_0}^{t_0+2\pi} U_0(-s)f(s) ds \right\|_{L^2(\mathcal{T})}.$$

On the other hand,

$$\begin{aligned} & \left\| \int_{t_0}^{t_0+2\pi} U_0(-s)f(s) ds \right\|_{L^2(\mathcal{T})}^2 \\ &= \left\langle f(s), \int_{t_0}^{t_0+2\pi} U_0(s-s')f(s') ds' \right\rangle \\ &\leq C \|f\|_{L^{4/3}((t_0, t_0+2\pi) \times \mathcal{T})} \left\| \int_{t_0}^{t_0+2\pi} U_0(s-s')f(s') ds' \right\|_{L^4((t_0, t_0+2\pi) \times \mathcal{T})}, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ denotes the inner product of $L^2((t_0, t_0 + 2\pi) \times \mathcal{T})$. Combining this inequality and (9), we have

$$\|G\|_{L^4((t_0, t_0+2\pi) \times \mathcal{T})} \leq C \|f\|_{L^{4/3}((t_0, t_0+2\pi) \times \mathcal{T})}.$$

This and Lemma 3.1 imply the second inequality. □

We first consider the Strichartz estimate for (6), (7) without external forcing term.

THEOREM 3.3. *Assume (A1) and (A2). Let $0 < \gamma' < \gamma$. Then, we have*

$$\|e^{\gamma' t} U(t)u_0\|_{L^\infty(\mathcal{R}_+; L^2(\mathcal{T}))} \leq C \|u_0\|_{L^2(\mathcal{T})},$$

$$\|e^{\gamma' t} U(t)u_0\|_{L^4(\mathcal{R}_+ \times \mathcal{T})} \leq C \|u_0\|_{L^2(\mathcal{T})}.$$

PROOF OF THEOREM 3.3. The first inequality follows immediately from (A2) and the standard L^2 inequality. For the proof of the second inequality, we consider the Cauchy problem of (6) with initial data prescribed at $t = t_0$, where $t_0 \geq 0$. By Duhamel’s principle,

$$u(t) = U_0(t - t_0)u(t_0) - iF(t), \quad t \geq t_0,$$

$$F(t) = \int_{t_0}^t U_0(t - s)Vu(s)ds.$$

If we can prove

$$(10) \quad \|F\|_{L^4((t_0, t_0+2\pi) \times \mathcal{T})} \leq C \|u\|_{L^\infty((t_0, t_0+2\pi); L^2(\mathcal{T}))},$$

then, for any $n \in \mathbb{N} \cup \{0\}$, we have by Lemma 3.2

$$(11) \quad \|u\|_{L^4((2\pi n, 2\pi(n+1)) \times \mathcal{T})} \leq C \|u(2\pi n)\|_{L^2(\mathcal{T})} + C \|u\|_{L^\infty((2\pi n, 2\pi(n+1)); L^2(\mathcal{T}))}.$$

On the other hand, for $0 < \gamma' < \gamma$, (A2) yields

$$\|U(t)u_0\|_{L^2(\mathbf{T})} \leq C \exp\left(-\frac{\gamma + \gamma'}{2}t\right) \|u_0\|_{L^2(\mathbf{T})}, \quad t \geq 0.$$

We use this inequality to bound the two terms on the right-hand side of (11) by

$$C \exp(-\pi(\gamma + \gamma')n) \|u_0\|_{L^2(\mathbf{T})}.$$

Accordingly, we conclude that for $0 < \gamma' < \gamma$,

$$\begin{aligned} \|e^{\gamma't}u\|_{L^4(\mathbf{R}_+ \times \mathbf{T})} &\leq \sum_{n=0}^{\infty} e^{2\pi\gamma'n} \|u\|_{L^4((2\pi n, 2\pi(n+1)) \times \mathbf{T})} \\ &\leq C \sum_{n=0}^{\infty} e^{-\pi(\gamma - \gamma')n} \|u_0\|_{L^2(\mathbf{T})} \\ &\leq C \|u_0\|_{L^2(\mathbf{T})}, \end{aligned}$$

which implies Theorem 3.3.

It remains only to prove the estimate (10). We easily see by Lemma 3.2 that

$$\begin{aligned} &\left\| \int_{t_0}^{t_0+2\pi} U_0(t-s)f(s) ds \right\|_{L^4((t_0, t_0+2\pi) \times \mathbf{T})} \\ &= \left\| U_0(t) \int_{t_0}^{t_0+2\pi} U_0(-s)f(s) ds \right\|_{L^4((t_0, t_0+2\pi) \times \mathbf{T})} \\ &\leq C \left\| \int_{t_0}^{t_0+2\pi} U_0(-s)f(s) ds \right\|_{L^2(\mathbf{T})} \\ &\leq C \|f\|_{L^1((t_0, t_0+2\pi); L^2(\mathbf{T}))}. \end{aligned}$$

Lemma 3.1 ensures that the integral operator

$$\int_{t_0}^t U_0(t-s)f(s) ds$$

has the same estimate as above. Therefore, we obtain

$$\begin{aligned} \|F\|_{L^4((t_0, t_0+2\pi) \times \mathbf{T})} &\leq C \|Vu\|_{L^1((t_0, t_0+2\pi); L^2(\mathbf{T}))} \\ &\leq C \|u\|_{L^\infty((t_0, t_0+2\pi); L^2(\mathbf{T}))}, \end{aligned}$$

which yields inequality (10). □

We next consider the Strichartz estimate for (6), (7) with $u_0 = 0$.

THEOREM 3.4. *Assume (A1) and (A2). Let u be the solution of (6), (7) with $u_0 = 0$. Then, for any γ' with $0 < \gamma' < \gamma$, we have*

$$\begin{aligned} \|e^{\gamma't}u\|_{L^\infty(\mathbf{R}_+; L^2(\mathbf{T}))} &\leq C \|e^{\gamma't}f\|_{L^{4/3}(\mathbf{R}_+ \times \mathbf{T})}, \\ \|e^{\gamma't}u\|_{L^4(\mathbf{R}_+ \times \mathbf{T})} &\leq C \|e^{\gamma't}f\|_{L^{4/3}(\mathbf{R}_+ \times \mathbf{T})}. \end{aligned}$$

REMARK 3.2. It seems likely that $e^{\gamma t}$ can be replaced by $e^{\gamma' t}$ on the right-hand sides of the two inequalities in Theorem 3.4.

PROOF OF THEOREM 3.4. We begin with the proof of the second inequality. We follow the same strategy as in the case of the homogeneous Schrödinger equation. We take the L^2 inner product of (6) and $e^{(\gamma'+\gamma)t}u$ and integrate the resulting equation in t to have by (A2)

$$(12) \quad \|e^{(\gamma'+\gamma)t/2}u(t)\|_{L^2(\mathbf{T})}^2 \leq C \|e^{\gamma' \cdot} f\|_{L^{4/3}((0,t)\times\mathbf{T})} \times \|e^{\gamma' \cdot} u\|_{L^4((0,t)\times\mathbf{T})}, \quad t > 0.$$

We note that Duhamel’s principle yields

$$(13) \quad \begin{aligned} u(t) &= U(t - t_0)u(t_0) + G(t), \\ G(t) &= -i \int_{t_0}^t U(t - s)f(s) ds \end{aligned}$$

for any $t_0 \geq 0$. If we can prove

$$(14) \quad \|G\|_{L^4((t_0,t_0+2\pi)\times\mathbf{T})} \leq C \|f\|_{L^{4/3}((t_0,t_0+2\pi)\times\mathbf{T})},$$

then, for any $n \in \mathbf{N} \cup \{0\}$, we have by Theorem 3.3

$$(15) \quad \|u\|_{L^4((2\pi n,2\pi(n+1))\times\mathbf{T})} \leq C \|u(2\pi n)\|_{L^2(\mathbf{T})} + C \|f\|_{L^{4/3}((2\pi n,2\pi(n+1))\times\mathbf{T})}.$$

Inequalities (12) and (15) yield the following estimate

$$\begin{aligned} \|u\|_{L^4((2\pi n,2\pi(n+1))\times\mathbf{T})} &\leq C e^{-\pi(\gamma'+\gamma)n} \|e^{\gamma' t} f\|_{L^{4/3}((0,2\pi n)\times\mathbf{T})}^{1/2} \|e^{\gamma' t} u\|_{L^4((0,2\pi n)\times\mathbf{T})}^{1/2} \\ &\quad + C \|f\|_{L^{4/3}((2\pi n,2\pi(n+1))\times\mathbf{T})}. \end{aligned}$$

Accordingly, we conclude that for $0 < \gamma' < \gamma$,

$$\begin{aligned} \|e^{\gamma' t} u\|_{L^4(\mathbf{R}_+\times\mathbf{T})} &\leq \sum_{n=0}^{\infty} e^{2\pi\gamma'(n+1)} \|u\|_{L^4((2\pi n,2\pi(n+1))\times\mathbf{T})} \\ &\leq C \sum_{n=0}^{\infty} e^{-\pi(\gamma-\gamma')n} \left[\|e^{\gamma' t} f\|_{L^{4/3}((0,2\pi n)\times\mathbf{T})}^{1/2} \|e^{\gamma' t} u\|_{L^4((0,2\pi n)\times\mathbf{T})}^{1/2} \right. \\ &\quad \left. + \|e^{(\gamma'+\gamma)t/2} f\|_{L^{4/3}((2\pi n,2\pi(n+1))\times\mathbf{T})} \right] \\ &\leq C \|e^{\gamma' t} f\|_{L^{4/3}(\mathbf{R}_+\times\mathbf{T})} + \frac{1}{2} \|e^{\gamma' t} u\|_{L^4(\mathbf{R}_+\times\mathbf{T})}. \end{aligned}$$

This shows Theorem 3.4.

It remains only to prove (14). We first note that G satisfies the following conditions

$$(16) \quad \begin{aligned} i\partial_t G &= -\partial_x^2 G + VG + f, \quad t > t_0, \quad x \in \mathbf{T}, \\ G(t_0, x) &= 0, \quad x \in \mathbf{T}. \end{aligned}$$

Therefore, Duhamel’s principle yields

$$\begin{aligned} G(t) &= -iG_1(t) - iG_2(t), \quad t \geq t_0, \\ G_1(t) &= \int_{t_0}^t U_0(t - s)VG(s) ds, \quad G_2(t) = \int_{t_0}^t U_0(t - s)f(s) ds. \end{aligned}$$

Furthermore, by (16) and (A2), we have

$$(17) \quad \|G(t)\|_{L^2(\mathbf{T})}^2 \leq C \|f\|_{L^{4/3}((t_0,t) \times \mathbf{T})} \|G\|_{L^4((t_0,t) \times \mathbf{T})}, \quad t > t_0.$$

On the other hand, by Lemma 3.2, we have

$$\|G\|_{L^4((t_0,t_0+2\pi) \times \mathbf{T})} \leq C [\|G\|_{L^\infty((t_0,t_0+2\pi); L^2(\mathbf{T}))} + \|f\|_{L^{4/3}((t_0,t_0+2\pi); L^2(\mathbf{T}))}].$$

By combining this inequality and (17), we obtain (14).

The first inequality follows from (17) and the second inequality was proved above. \square

REMARK 3.3. We note that Theorems 3.3 and 3.4 also hold for the linear Schrödinger equation with shift term:

$$(18) \quad \begin{aligned} i \partial_t u &= -\partial_x^2 u - i \dot{c}(t) \partial_x u + V u + f, \quad t > 0, \quad x \in \mathbf{T}, \\ u(0, x) &= u_0(x), \end{aligned}$$

where $c(t)$ is a continuously differentiable real-valued function. In fact, if we put

$$U_1(t, s) = e^{i[(t-s)\partial_x^2 - (c(t) - c(s))\partial_x]} \quad (t \geq s \geq 0),$$

then we have Lemma 3.2 without any change for $U_1(t, 0)$. Because the shift term $-i \dot{c}(t) \partial_x u$ only leads to the spatial translation of solutions. The norms appearing in Lemma 3.2 are invariant under the spatial translation and so the spatial translation caused by the shift term has no influence on the Strichartz estimate for the Schrödinger equation:

$$i \partial_t u = -\partial_x^2 u - i \dot{c}(t) \partial_x u + f, \quad t > 0, \quad x \in \mathbf{T}.$$

Furthermore, the shift term $-i \dot{c}(t) \partial_x u$ has no influence on the exponential decay in $L^2(\mathbf{T})$ of solution, either. This enables us to prove Theorems 3.3 and 3.4.

4. Proof of Theorem 1.1. In this section, we describe the proof of Theorem 1.1. Let L be the linearized operator around $D(x)$ defined as in (5). We denote the subspace $\text{span}\{\partial_x D(x)\}$ and its complementary subspace in $L^2(\mathbf{T})$ by L_0^2 and L_-^2 , respectively. We choose L_-^2 so that L_-^2 is an invariant subspace of L . Let $\widetilde{\partial_x D}$ denote the normalization in L^2 of $\partial_x D$. Let Q and P be the projections from $L^2(\mathbf{T})$ to L_-^2 and from $L^2(\mathbf{T})$ to L_0^2 , respectively. The projections Q and P are explicitly expressed as follows.

$$Qf = f - (f, E) \widetilde{\partial_x D}, \quad Pf = (f, E) \widetilde{\partial_x D},$$

where E is the normalized eigenfunction belonging to the zero eigenvalue of the adjoint operator of L . We choose $2\pi > c_1 \geq 0$ such that

$$\|A_0(\cdot + c_1) - D(\cdot)\|_{L^2} = \min_{2\pi > c \geq 0} \|A_0(\cdot + c) - D(\cdot)\|_{L^2}.$$

Without loss of generality, we may change the initial data $A_0(x)$ to $A_0(x + c_1)$, since the equation (1) is invariant under the spatial translation. We denote $A_0(x + c_1)$ by $A_0(x)$ again and we have

$$(19) \quad \text{Re}(A_0, \partial_x D) = 0.$$

It is expedient to work with the real and the imaginary parts of complex-valued functions and to regard the space $L^2(\mathbf{T})$ as a real Hilbert space with inner product $\text{Re}(\cdot, \cdot)$. In that case, (19) implies that A_0 is orthogonal to the subspace spanned by $\partial_x D$. Let w and z denote the real and the imaginary parts of the stationary solution $D(x)$, respectively. We now decompose the solution A into effective dynamical components, following Buslaev and Perel'man [6] and Soffer and Weinstein [15] (for the case of the nonlinear parabolic equation, see Exercise 6 in Henry [11, Section 5.1]). For the solution A of (1), we make an ansatz as follows.

$$(20) \quad A(t, x) = D(x + c(t)) + u(t, x + c(t)) + i v(t, x + c(t)), \quad u, v \in L^2_-,$$

where $u(t, x)$ and $v(t, x)$ are real-valued functions, and $c(t)$ is a continuously differentiable function with $c(0) = 0$ to be determined later. If we insert the ansatz (20) into (1) and remove the spatial translation $c(t)$ by the change of variables, then we rewrite the equation (1) as in the following form.

$$(21) \quad \partial_t T - LT + \dot{c}(t)Q\partial_x T = Q\mathcal{F}(x, T), \quad t > 0, \quad x \in \mathbf{T},$$

$$(22) \quad \dot{c}(t) = \frac{(\mathcal{F}(x, T), E)}{a - (T, \partial_x E)}, \quad t > 0, \quad a = (\partial_x D, E),$$

$$(23) \quad T(0, x) = T_0(x) \quad (x \in \mathbf{T}), \quad c(0) = 0,$$

where

$$T(t, x) = \begin{pmatrix} u(t, x) \\ v(t, x) \end{pmatrix} \in L^2_-, \quad T_0(x) = \begin{pmatrix} \text{Re}(A_0(x) - D(x)) \\ \text{Im}(A_0(x) - D(x)) \end{pmatrix},$$

$$\mathcal{F} \in C^2(\mathbf{T} \times \mathbf{R}^2; \mathbf{R}^4),$$

$$|\mathcal{F}(x, T_1) - \mathcal{F}(x, T_2)| \leq C(|T_1| + |T_2| + |T_1|^2 + |T_2|^2)|T_1 - T_2|$$

$$(x \in \mathbf{T}, \quad T_1, T_2 \in \mathbf{R}^2),$$

$$|\partial_T \mathcal{F}(x, T_1) - \partial_T \mathcal{F}(x, T_2)| \leq C(1 + |T_1| + |T_2|)|T_1 - T_2|$$

$$(x \in \mathbf{T}, \quad T_1, T_2 \in \mathbf{R}^2).$$

Here, $|T| = \sqrt{u^2 + v^2}$ for $T = (u, v) \in \mathbf{R}^2$. Let $U_c(t, s)$ ($t \geq s \geq 0$) denote the evolution operator associated with the infinitesimal generator $L - \dot{c}(t)\partial_x$ for each $c(t)$. We note that the Strichartz estimates such as Theorems 3.3 and 3.4 are applicable to the first and the second components of the solution T of (21) (see Remark 3.1 (ii) and Remark 3.3). In fact, we have

$$\dot{c}(t)Q\partial_x T = \dot{c}(t)\partial_x T - \dot{c}(t)P\partial_x T,$$

and the term $\dot{c}(t)P\partial_x T$ can be regarded as a small regular perturbation as long as $\dot{c}(t)$ is small. Furthermore, we note that the unique global existence of the solution $A(t)$ on the time interval $(-\eta, \infty)$ for the Cauchy problem of (1) with initial data in $L^2(\mathbf{T})$ follows from the result by Bourgain [5], where η is a small positive constant depending only on the initial data. For a given solution $A(t)$, by (20), we set

$$T(t, x) = A(t, x - c(t)) - D(x).$$

We insert T into (22) to have by (23)

$$(24) \quad c(t) = \int_0^t \frac{\mathcal{F}(x, T), E}{a - (T, \partial_x E)} ds, \quad t > -\eta.$$

For a short time, we have the solution $c(t)$ of (24) by the implicit function theorem as long as T is small. Because a direct computation yields

$$\begin{aligned} & \frac{\partial}{\partial c} \left(c - \int_0^t \frac{\mathcal{F}(x, T), E}{a - (T, \partial_x E)} ds \right) \\ &= 1 - \int_0^t \frac{(\partial_T \mathcal{F}(x, T) \partial_x A(s, x - c), E)}{a - (T, \partial_x E)} ds \\ & \quad + \int_0^t \frac{(\partial_x A(s, x - c), \partial_x E)(\mathcal{F}(x, T), E)}{(a - (T, \partial_x E))^2} ds \\ &= 1 + \int_0^t \frac{(\mathcal{F}(x, T), \partial_x E) + (\partial_x \mathcal{F}(x, T), E) - (\partial_x D, E)}{a - (T, \partial_x E)} ds \\ & \quad - \int_0^t \frac{(T, \partial_x^2 E)(\mathcal{F}(x, T), E) - (\partial_x D, \partial_x E)}{(a - (T, \partial_x E))^2} ds. \end{aligned}$$

The right-hand side of this formula does not vanish at $(t, c) = (0, 0)$. We note that the right-hand side of this formula makes sense for $|t| < \tau$, provided that for some $\tau > 0$, T is in $L^\infty((-\tau, \tau); L^2(\mathbf{T})) \cap L^4((-\tau, \tau) \times \mathbf{T})$ and T is small in $L^\infty((-\tau, \tau); L^2(\mathbf{T}))$. These facts and the implicit function theorem imply the existence of $c(t)$ for a short time. From the above construction of the function $c(t)$, it automatically follows that $T(t)$ must satisfy (21). Indeed, the L^2_0 -component of T satisfies

$$\frac{d}{dt}(PT) = 0, \quad t > 0, \quad (PT)(0) = 0,$$

which implies that $T(t) \in L^2_-$ for each $t > 0$. Therefore, if we have proved the a priori estimates of $(T(t), c(t))$, then we obtain Theorem 1.1.

By the L^2 estimate, Theorem 3.3 and Remark 3.3, for some $0 < \gamma' < \gamma$, we have

$$\begin{aligned} & \|e^{\gamma' t}(U_c(t, 0)T_0)\|_{L^\infty(\mathbf{R}_+; L^2)} \leq C \|T_0\|_{L^2}, \\ & \|e^{\gamma' t}(U_c(t, 0)T_0)\|_{L^4(\mathbf{R}_+ \times \mathbf{T})} \leq C \|T_0\|_{L^2}. \end{aligned}$$

We put $\delta = C \|T_0\|_{L^2}$. Later, we choose T_0 so small in $L^2(\mathbf{T})$ that δ is sufficiently small. We define the space X_δ by

$$\begin{aligned} & \{(T, c) \in (L^\infty(\mathbf{R}_+; L^2) \cap L^4(\mathbf{R}_+ \times \mathbf{T}))^2 \times C_b([0, \infty)); \\ & \quad c(0) = 0, \quad \|(T, c)\|_{X_\delta} \leq 2\delta\}, \end{aligned}$$

where $C_b([0, \infty))$ denotes the space of all bounded continuous functions defined on $[0, \infty)$ and

$$\|(T, c)\|_{X_\delta} = \max\{\|e^{\gamma' t} T\|_{L^\infty(\mathbf{R}_+; L^2)}, \|e^{\gamma' t} T\|_{L^4(\mathbf{R}_+ \times \mathbf{T})}, \|e^{\gamma' t} \dot{c}\|_{L^2((0, \infty); \mathbf{R})}\}.$$

Let $(T, c) \in X_\delta$ be the solution of (21)–(23). We now prove the a priori estimates for (T, c) , which ensure the global existence of (T, c) and there exists $c_0 \in \mathbf{R}$ such that

$$(25) \quad \|T(t)\|_{L^2(T)} \rightarrow 0,$$

$$(26) \quad c(t) \rightarrow c_0 \quad (t \rightarrow \infty).$$

We first show that if $\delta > 0$ is sufficiently small, (T, c) is small in X_δ . We apply Theorems 3.3 and 3.4, together with Remark 3.3, to (21) and have by the Hölder inequality and the assumption $\gamma' < \gamma$

$$\begin{aligned} \|e^{\gamma' t} T\|_{L^\infty(\mathbf{R}_+; L^2)} &\leq \delta + C(\|e^{\gamma' t} T\|_{L^4(\mathbf{R}_+ \times T)}^2 + \|e^{\gamma' t} T\|_{L^4(\mathbf{R}_+ \times T)}^3) \\ &\leq \delta + C(\delta + \delta^2)\delta, \end{aligned}$$

$$\begin{aligned} \|e^{\gamma' t} T\|_{L^4(\mathbf{R}_+ \times T)} &\leq \delta + C(\|e^{\gamma' t} T\|_{L^4(\mathbf{R}_+ \times T)}^2 + \|e^{\gamma' t} T\|_{L^4(\mathbf{R}_+ \times T)}^3) \\ &\leq \delta + C(\delta + \delta^2)\delta \end{aligned}$$

for $T \in X_\delta$. We easily see that

$$\|e^{\gamma' t} \dot{c}\|_{C_b([0, \infty))} \leq C \frac{(\delta + \delta^2)\delta}{a - C\delta}.$$

Here, if we choose $\delta > 0$ such that

$$C(\delta + \delta^2) \leq 1, \quad C \frac{(\delta + \delta^2)}{a - C\delta} < 1, \quad a - C\delta > 0,$$

then we can conclude that

$$\|(T, c)\|_{X_\delta} \leq 2\delta.$$

This implies (25) and (26), that is, the asymptotic stability of D .

Acknowledgment. The third author, Tsutsumi is grateful to Professor Kenji Nakanishi for fruitful discussions.

REFERENCES

- [1] N. AKHMEDIEV AND A. ANKIEWICZ, Dissipative soliton, Lecture Notes in Phys. 661, Springer-Verlag, Berlin, Heidelberg, 2005.
- [2] N. AKHMEDIEV AND A. ANKIEWICZ, Dissipative solitons: From optics to biology and medicine, Lecture Notes in Phys. 751, Springer-Verlag, Berlin, Heidelberg, 2008.
- [3] T. ACKEMANN AND W. J. FIRTH, Dissipative solitons in pattern-forming nonlinear optical systems, Lecture Notes in Phys. 661 (2005), 55–100.
- [4] B. AULBACH, Continuous and discrete dynamics near manifolds of equilibria, Lecture Notes in Math. 1058, Springer-Verlag, Berlin, 1984.
- [5] J. BOURGAIN, Fourier restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, Part I: Schrödinger equations, Geom. Funct. Anal. 3 (1993), 107–156.
- [6] V. S. BUSLAEV AND G. S. PEREL'MAN, Scattering for the nonlinear Schrödinger equation: states close to a soliton, St. Petersburg Math. J. 4 (1993), 1111–1142.
- [7] T. CAZENAVE, Semilinear Schrödinger equations, Courant Lect. Notes Math. 10, New York University, Courant Institute of Mathematical Sciences, New York, American Mathematical Society, Providence, RI, 2003.

- [8] M. CHRIST AND A. KISELEV, Maximal functions associated to filtrations, *J. Funct. Anal.* 179 (2001), 409–425.
- [9] W. J. FIRTH, G. K. HARKNESS, A. LORD AND J. M. MCSLOY, Dynamical properties of two-dimensional Kerr cavity solitons, *J. Opt. Soc. Amer. B Opt. Phys.* 19 (2002), 747–752.
- [10] J.-M. GHIDAGLIA, Finite-dimensional behavior for weakly damped driven nonlinear Schrödinger equations, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 5 (1988), 365–405.
- [11] D. HENRY, Geometric theory of semilinear parabolic equations, *Lecture Notes in Math.* 840, Springer-Verlag, Berlin-New York, 1981.
- [12] L. A. LUGIATO AND R. LEFEVER, Spatial dissipative structures in passive optical systems, *Phys. Rev. Lett.* 58 (1987), 2209–2211.
- [13] T. MIYAJI, I. OHNISHI AND Y. TSUTSUMI, Bifurcation analysis to the Lugiato-Lefever equation in one space dimension, *Phys. D* 239 (2010), 2066–2083.
- [14] H. F. SMITH AND C. D. SOGGE, Global Strichartz estimates for nontrapping perturbations of the Laplacian, *Comm. Partial Differential Equations* 25 (2000), 2171–2183.
- [15] A. SOFFER AND M. I. WEINSTEIN, Multichannel nonlinear scattering for nonintegrable equations, *Comm. Math. Phys.* 133 (1990), 119–146.
- [16] X.-M. WANG, An energy equation for the weakly damped driven nonlinear Schrödinger equations and its application to their attractors, *Phys. D* 88 (1995), 167–175.
- [17] A. ZYGMUND, On Fourier coefficients and transforms of functions of two variables, *Studia Math.* 50 (1974), 189–201.

RESEARCH INSTITUTE FOR
MATHEMATICAL SCIENCES
KYOTO UNIVERSITY
KYOTO 606–8502
JAPAN

E-mail address: tmiyajji@kurims.kyoto-u.ac.jp

DEPARTMENT OF MATHEMATICAL AND
LIFE SCIENCES
GRADUATE SCHOOL OF SCIENCE
HIROSHIMA UNIVERSITY
HIGASHI-HIROSHIMA 739–8526
JAPAN

E-mail address: isamu_o@math.sci.hiroshima-u.ac.jp

DEPARTMENT OF MATHEMATICS
GRADUATE SCHOOL OF SCIENCE
KYOTO UNIVERSITY
KYOTO 606–8502
JAPAN

E-mail address: tsutsumi@math.kyoto-u.ac.jp