

WAVELETS IN WEIGHTED NORM SPACES

KAZAROS S. KAZARIAN, SAMVEL S. KAZARYAN AND ANGEL SAN ANTOLÍN

(Received November 24, 2015, revised July 25, 2016)

Abstract. We give a complete characterization of the classes of weight functions for which the higher rank Haar wavelet systems are unconditional bases in weighted norm Lebesgue spaces. Particularly it follows that higher rank Haar wavelets are unconditional bases in the weighted norm spaces with weights which have strong zeros at some points. This shows that the class of weight functions for which higher rank Haar wavelets are unconditional bases is much richer than it was supposed.

1. Introduction. The wavelet analysis, since its creation, has been used in many areas of applied mathematics. The main idea is simple: find a function (*wavelet or wavelet function*) defined in \mathbb{R} or in \mathbb{R}^d so that the system of its dilations and translations constitute a complete orthonormal system (ONS) in $L^2(\mathbb{R})$ or in $L^2(\mathbb{R}^d)$. Previously that idea was used by A. Haar for constructing a complete ONS in $L^2([0, 1])$ such that the expansion of any continuous function on $[0, 1]$ converges uniformly. In the univariate case the simplest dilation is the dyadic dilation. Thus for the simplest case a function $g \in L^2(\mathbb{R})$ is a wavelet if $\{g_{k,j} : k, j \in \mathbb{Z}\}$, where $g_{k,j}(x) := 2^{k/2}g(2^kx - j)$ is a complete ONS in the space $L^2(\mathbb{R})$.

Weighted norm inequalities have been studied by Hardy and Littlewood [17], Babenko [3], Hirschman [22], Gaposhkin [13], Edwards [9], Chen [4], Helson and Szegő [19] and others. Probably M. Rosenblum [40] was the first one who obtained a complete characterization of weight functions for which a certain weighted norm inequality holds. Afterwards B. Muckenhoupt [38] proved that the Hardy-Littlewood maximal function is a bounded operator $L^p(\mathbb{R}, \psi) \rightarrow L^p(\mathbb{R}, \psi)$, $1 \leq p < \infty$ if and only if ψ satisfies condition (A_p) : for every interval $I \subset \mathbb{R}$

$$|I|^{-1} \int_I \psi dt \left[|I|^{-1} \int_I \psi^{-1/(p-1)} dt \right]^{p-1} \leq B_p,$$

B_p is independent of I . The last result boosts investigations in the harmonic analysis and related areas, several important weighted norm inequalities were obtained for weights which satisfy (A_p) condition. It is clear that if ψ satisfies condition (A_p) , $p > 1$ then the function $\psi^{-1/(p-1)}$ is locally integrable, hence, if $f \in L^p(\mathbb{R}, \psi)$ is locally integrable. The characterization of weight functions w for which families of some operators related with generalized Fourier series are uniformly bounded in weighted norm $L^p(w)$ spaces gives classes of weight functions which are different from (A_p) (see [29]). Particularly, $w^{-1/(p-1)}$ are not integrable

2010 *Mathematics Subject Classification.* Primary 41A65; Secondary 41A25, 41A46, 46B20.

Key words and phrases. Wavelet, high rank Haar wavelet, complete orthonormal system, weighted norm space, basis, unconditional basis, weights with singularities.

on some intervals. Interestingly discussed cases came together when we study the class of weight functions $w \geq 0$ for which a given wavelet system $\{g_{k,j} : k, j \in \mathbb{Z}\}$ is a basis in a certain sense or unconditional basis in the weighted norm space $L^p(\mathbb{R}, w)$, $1 \leq p < \infty$. Let us distinguish two cases:

- (α): $w^{-1/(p-1)}$ is locally integrable.
- (β): $w^{-1/(p-1)}$ is not locally integrable.

In terms of the above cases we can classify the publications related with the study of weight functions for which a given wavelet system is an unconditional basis in the weighted norm space L^p into two main groups. In [10], [11],[12], [1], [2] and others the study is done with some restrictions which reduce it to the case (α). While in [35], [45] and [32] the aim is to give a complete description without any additional restrictions.

Unfortunately in these papers the case (β) is not considered. It should be mentioned that in [11] the description of the class of all weight functions w for which the dyadic Haar wavelet system is an unconditional basis in weighted norm space $L^p(\mathbb{R}, w)$, $1 \leq p < \infty$ was formulated. The proof of the mentioned result follows easily from the results obtained in [28]. Remarkably that class of weight functions contain weights which correspond to both cases (α) and (β). At this point it would be appropriate to bring a citation from [1]: “we have to say that, as it was pointed out by our referee in a previous version of this article, we are unable to prove, without some extra condition on the weight, the implication (D1) \Rightarrow (D3) in [35] which gives A_p as necessary condition for the fact that the Daubechies wavelet system is an unconditional basis for $L^p(d\mu)$. Actually for the Daubechies case our result is contained in Corollary 2 and we are only able to show that A_p is necessary under the extra assumption of the local integrability of $w^{-1/(p-1)}$.” The purpose of the present paper is to shed some light on the doubts expressed by authors in [1].

To give a preliminary idea about the main subject of our study let us suppose that for a given weight function w there exist some functions h such that

- (a) $\int_{\mathbb{R}} g_{k,j}(x)h(x)dx = 0$ for all $k, j \in \mathbb{Z}$;
- (b) $\frac{h}{w} \in L^{p'}(\mathbb{R}, w)$, $1/p + 1/p' = 1$.

Then any non trivial function $\frac{h}{w}$ as linear continuous functional will be a non trivial element in the dual space $L^{p'}(\mathbb{R}, w)$. Moreover, this functional vanishes on all $g_{k,j} : k, j \in \mathbb{Z}$. Hence our wavelet system is not complete in the space $L^p(\mathbb{R}, w)$. A necessary condition for completeness of the system $\{g_{k,j} : k, j \in \mathbb{Z}\}$ in the space $L^p(\mathbb{R}, w)$ is the following condition: $\frac{h}{w} \notin L^{p'}(\mathbb{R}, w)$. Thus if we are going to describe all weight functions w for which the system $\{g_{k,j} : k, j \in \mathbb{Z}\}$ is a basis in a certain sense in $L^p(\mathbb{R}, w)$, we have to consider all those weight functions w for which the condition (b) is not true. For our study we will use the technique by which similar questions were studied for incomplete systems in the weighted norm spaces (see [26], [27], [28]). We also will show that the conditions (a) and (b) are not hypothetical cases.

If we are going to study the wavelet system in $L^p(\mathbb{R}, w)$ it will be natural to suppose that

$$g \in L^1(\mathbb{R}) \cap L^{\max(p,p')}(\mathbb{R}).$$

On the other hand the purpose of the present paper is not to obtain the most general results. Hence, instead of the last restriction we will suppose that $g \in L^1(\mathbb{R}) \cap L^\infty(\mathbb{R})$. It is well known that if a wavelet $g \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ then its Fourier transform \widehat{g} should vanish at the origin and thus the constant functions satisfy to condition (a). Which means that if someone has studied the formulated question without describing the class of functions for which the condition (a) holds and without considering weighted norm spaces $L^p(\mathbb{R}, w)$ with weights w which does not satisfy the condition (b), then his proof is not complete.

In order to show that the question under consideration is not a technical problem we revise the study done in [32] consisting of characterization of all weight functions w for which the Haar wavelet system for m -dilations, $m = 2, 3, \dots$ is an unconditional basis in $L^p(\mathbb{R}, w)$. We have modified a little bit the definition of the higher rank Haar wavelets given in [32] to make it real valued. It should be considered that we are complementing the results obtained in [32] with results which correspond to the case (β). From the corollary of the main theorem of the last section it follows that higher rank Haar wavelets are unconditional bases in the weighted norm spaces $L^p(\mathbb{R}, w)$, where $w(x) = |x|^r$, $r > p - 1$, which shows that higher rank Haar wavelets are unconditional bases for weight functions from the case (β). For the convenience of the reader we give complete proofs of all results giving preference to classical methods.

We would like to add one more remark related with the given above citation from [1]. One can say that Daubechies wavelets are smooth so probably the question which we study for the higher rank Haar wavelets is different for the Daubechies wavelets. Here we would like to share our intuition about this question because at this moment we have no intention to continue the research in this area. It should be mentioned that in [10], [11] there were obtained all necessary tools for the study of the problem for the spline wavelets. Particularly it was obtained the subspace of the functions h for which (a) holds which depends on the smoothness of the splines. The last fact gives some hint for the Daubechies wavelets: it seems that the solution depends also from the smoothness. The present paper is not a survey article and we do not pretend to give a detailed review of all publications related with weighted norm inequalities. Some publications have been dedicated to the decompositions in terms of scaling and wavelet functions (e.g. [23], [24] and others). In those cases the weight function w satisfies the condition (α) and the phenomenon which we are studying is not present. Review of some other results on weighted norm inequalities can be find in [39].

In Section 2 we prove an inequality for the orthogonal wavelet systems which particularly shows that in the case of general orthogonal wavelet systems the set of nontrivial functions h for which the condition (a) holds is not empty. In Section 3 we give all preliminary results which will be used for our study. Next two sections are dedicated to the m th rank Haar system on $[0, 1]$. The results of these sections have certain interest.

It should be mentioned that they are used for proofs given in the last section, where the main results for the higher rank Haar wavelets are obtained. As in the case of the dyadic Haar wavelet the solution of the problem is obtained combining two cases on the interval $[0, 1]$: the description of weight functions w for which the m th rank Haar system by $\{h_l(x)\}_{l=0}^\infty$ and

$\{h_l(x)\}_{l=1}^\infty$ are unconditional bases in $L^p([0, 1], w)$, $1 < p < \infty$.

2. An inequality for wavelet type systems. If $w \geq 0$ be a weight function on \mathbb{R} , i.e. a non negative locally integrable function then we write $\phi \in L^p(\mathbb{R}, w)$, $1 \leq p < \infty$ if $\phi : \mathbb{R} \rightarrow \mathbb{C}$ is measurable on \mathbb{R} and the norm is defined by

$$\|\phi\|_{L^p(\mathbb{R}, w)} := \left(\int_{\mathbb{R}} |\phi(t)|^p w(t) dt \right)^{\frac{1}{p}} < +\infty.$$

For a $g \in L^2(\mathbb{R})$ and $m = 2, 3, \dots$ we will denote

$$(1) \quad g_{k,j,m}(x) := m^{k/2} g(m^k x - j), \quad k, j \in \mathbb{Z}.$$

In this paper we will use a slightly modified version of the classical definition of the Fourier transform. For a function $f \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ we put

$$\widehat{f}(y) = \int_{\mathbb{R}} f(x) e^{-2\pi ixy} dx.$$

The characteristic function of a set E is denoted by χ_E and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$.

The following lemma is a well known result (cf. [8], p. 132; [20], p. 71).

LEMMA 2.1. *The system $\{h(\cdot - j) : j \in \mathbb{Z}\}$, where $h \in L^2(\mathbb{R})$, is an orthonormal system if and only if*

$$\sum_{j \in \mathbb{Z}} |\widehat{h}(t + j)|^2 = 1 \quad \text{for a.e. } t \in \mathbb{R}.$$

As an obvious corollary of the above lemma we have that if $g \in L^2(\mathbb{R})$ is a wavelet then $|\widehat{g}(t)| \leq 1$ a.e. on \mathbb{R} .

THEOREM 2.2. *Let $g \in L^2(\mathbb{R})$ and $m = 2, 3, \dots$. Suppose that the system $\{g_{k,j,m} : k \in \mathbb{N}_0, j \in \mathbb{Z}\}$ is orthonormal. Then*

$$(2) \quad \sum_{k=0}^\infty |\widehat{g}(m^{-k}x)|^2 \leq 1.$$

PROOF. It is easy to check that

$$(3) \quad \widehat{g_{k,j,m}}(y) = m^{-k/2} \widehat{g}(m^{-k}y) e^{-2\pi ijm^{-k}y}.$$

It is well known that for any interval $I \subset \mathbb{R}$, $|I| = 1$ the trigonometric system $\{e^{-2\pi i j y}\}_{j \in \mathbb{Z}}$ is a complete orthonormal system in $L^2(I)$. Hence, for any $k \in \mathbb{Z}$ and $\Delta \subset \mathbb{R}$, $|\Delta| = m^k$ the system $\{m^{-k/2} e^{-2\pi i j m^{-k} y}\}_{j \in \mathbb{Z}}$ will be a complete orthonormal system in $L^2(\Delta)$. Thus for any $f \in L^2(\mathbb{R})$ such that $\text{supp } \widehat{f} \subseteq I$, $|I| = 1$ we will have that

$$S_0(f, x) = \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}} f(t) \overline{g_{0,j,m}(t)} dt g_{0,j,m}(x) = \sum_{j \in \mathbb{Z}} \int_I \widehat{f}(t) \overline{\widehat{g}(t)} e^{2\pi i j t} dt g_{0,j,m}(x).$$

Which yields

$$\widehat{S_0(f, \cdot)}(y) = \sum_{j \in \mathbb{Z}} \int_I \widehat{f}(t) \overline{\widehat{g}(t)} e^{2\pi i j t} dt \widehat{g}(y) e^{-2\pi i j y} = \widehat{f}(y) |\widehat{g}(y)|^2$$

if $y \in I$. It should be observed that the last equality holds because $\widehat{f}(\cdot) \overline{\widehat{g}(\cdot)} \in L^2(I)$ which is true because of Lemma 2.1. If for any $k \in \mathbb{N}$ we put

$$S_k(f, x) = \sum_{j \in \mathbb{Z}} \int_{\mathbb{R}} f(t) \overline{g_{k,j,m}(t)} dt g_{k,j,m}(x)$$

in the same way we obtain that

$$(4) \quad \widehat{S_k(f, \cdot)}(y) = \widehat{f}(y) |\widehat{g}(m^{-k}y)|^2 \quad \text{if } y \in I.$$

By the orthogonality of the system $\{g_{k,j,m} : k \in \mathbb{N}_0, j \in \mathbb{Z}\}$ we have that

$$S_k(f, \cdot) \perp S_{k'}(f, \cdot) \quad \text{in } L^2(\mathbb{R}) \quad \text{if } k \neq k'.$$

Hence,

$$\begin{aligned} \int_{\mathbb{R}} \left| \sum_{k=0}^l S_k(f, x) \right|^2 dx &= \sum_{k=0}^l \int_{\mathbb{R}} |S_k(f, x)|^2 dx = \sum_{k=0}^l \int_{\mathbb{R}} |\widehat{S_k(f, \cdot)}(y)|^2 dy \\ &= \int_{\mathbb{R}} \sum_{k=0}^l |\widehat{S_k(f, \cdot)}(y)|^2 dy = \int_{\mathbb{R}} |\widehat{f}(y)|^2 \left| \sum_{k=0}^l |\widehat{g}(m^{-k}y)|^2 \right|^2 dy. \end{aligned}$$

By (4) we have that

$$\begin{aligned} \int_I |\widehat{f}(y)|^2 \left| \sum_{k=0}^l |\widehat{g}(m^{-k}y)|^2 \right|^2 dy &= \int_I \sum_{k=0}^l |\widehat{S_k(f, \cdot)}(y)|^2 dy \\ &\leq \int_{\mathbb{R}} \sum_{k=0}^l |\widehat{S_k(f, \cdot)}(y)|^2 dy. \end{aligned}$$

By the above relations and Bessel's inequality we obtain that

$$\begin{aligned} \int_I |\widehat{f}(y)|^2 \left| \sum_{k=0}^l |\widehat{g}(m^{-k}y)|^2 \right|^2 dy &\leq \int_{\mathbb{R}} \left| \sum_{k=0}^l S_k(f, x) \right|^2 dx \\ &\leq \|f\|_{L^2(\mathbb{R})}^2 = \int_I |\widehat{f}(y)|^2 dy. \end{aligned}$$

The last inequality can be interpreted as follows. Let

$$\mu_l(y) = \sum_{k=0}^l |\widehat{g}(m^{-k}y)|^2 \quad \text{if } y \in I.$$

Then for any $l \in \mathbb{N}$ the multiplicative operator $T_l(\phi)(y) = \mu_l(y)\phi(y)$ is a bounded operator $L^2(I) \rightarrow L^2(I)$ with the norm less than or equal to 1. Which is true if and only if $\mu_l(y) \leq 1$. □

By a simple modification of the last part, related with the application of the Bessel inequality and the proof of Theorem 2.2 we obtain the following

THEOREM 2.3. *Let $h^{(\nu)} \in L^2(\mathbb{R}), 1 \leq \nu \leq \mu$ and $m = 2, 3, \dots$. Suppose that the system $\{h_{k,j,m}^{(\nu)} : k \in \mathbb{N}_0, j \in \mathbb{Z}, 1 \leq \nu \leq \mu\}$ is orthonormal. Then*

$$\sum_{k=0}^{\infty} \sum_{\nu=1}^{\mu} |\widehat{h^{(\nu)}}(m^{-k}x)|^2 \leq 1.$$

We formulate the following corollary for $g \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$. In the general case a similar result can be proved using the concept of points of approximate continuity (cf. [5], [6]).

COROLLARY 2.4. *Let $g \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ and $m = 2, 3, \dots$. Suppose that the system $\{g_{k,j,m} : k \in \mathbb{N}_0, j \in \mathbb{Z}\}$ is orthonormal. Then the continuous function \widehat{g} vanishes at the origin, $\widehat{g}(0) = 0$.*

PROOF. Let $\widehat{g}(0) \neq 0$. Without loss in generality we can suppose that $\widehat{g}(0) > 0$. Which yields that $\widehat{g}(y)$ is greater than $\widehat{g}(0)/2$ in a neighborhood of the origin. The last condition contradicts to (2). □

By Corollary 2.4 we have that when for $g \in L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ and $m = 2, 3, \dots$ the system $\{g_{k,j,m} : k \in \mathbb{N}_0, j \in \mathbb{Z}\}$ is orthonormal then the set of nontrivial functions h defined on \mathbb{R} such that

$$(a)_m \quad \int_{\mathbb{R}} g_{k,j,m}(x)h(x)dx = 0 \quad \text{for all } k, j \in \mathbb{Z}$$

is not empty. Hence, having in mind that the constant function belongs to $L^\infty(\mathbb{R})$ we obtain

COROLLARY 2.5. *Let $\{g^{(\nu)}\}_{\nu=1}^{\mu} \subset L^1(\mathbb{R}) \cap L^2(\mathbb{R})$ and $m = 2, 3, \dots$. If the system $\{g_{k,j,m}^{(\nu)} : k \in \mathbb{Z}, j \in \mathbb{Z}, 1 \leq \nu \leq \mu\}$ is orthonormal then it cannot be complete in $L^1(\mathbb{R})$.*

3. Preliminary results.

3.1. On \mathcal{M} -sets. Further in this section we will consider that $m \geq 2$ is a fixed natural number. Let $\mathcal{M} = \mathcal{M}(m) := \{[\frac{j-1}{m^k}, \frac{j}{m^k}] : k \in \mathbb{Z}, j \in \mathbb{Z}\}$. Further, the parameter m will be omitted to make the notation understandable. We will assume that any m -adic rational point $\xi = \frac{j}{m^k}, k \in \mathbb{Z}, j \in \mathbb{Z}$ is “split” into two distinct points ξ_l and ξ_r characterized by the following conditions: for any $-\infty < a < \xi < b < +\infty$ we have

$$\xi_l \in (a, \xi), \quad \xi_l \notin [\xi, b), \quad \text{and} \quad \xi_r \in [\xi, b), \quad \xi_r \notin (a, \xi).$$

Hence, there can be easily established one to one correspondence between any $y \in \mathbb{R}$ and the sequences $\{\Delta_j(y)\}_{j=-\infty}^{\infty} \subset \mathcal{M}$ such that

$$\Delta_{j+1}(y) \subset \Delta_j(y) \quad \text{for all } j \in \mathbb{Z} \quad \text{and} \quad |\Delta_j(y)| = m^{-j}.$$

When talking about the neighborhoods of the points ξ_l and ξ_r , we will understand some intervals (a, ξ) and (ξ, b) , respectively. The measure of the set of all m -adic rational points is

equal to zero, hence, this assumption will not affect the results that we are going to consider. In the last section we need concept of \mathcal{M} -neighborhoods of $+\infty$ and $-\infty$. For $j \in \mathbb{N}_0$ we put

$$(5) \quad \Delta_j(+\infty) = \mathbb{R}^+ \setminus [0, m^j], \quad \text{and} \quad \Delta_j(-\infty) = \mathbb{R}^- \setminus [-m^j, 0],$$

where $\mathbb{R}^+ = [0, +\infty)$ and $\mathbb{R}^- = (-\infty, 0]$. If \mathcal{B} is a family of sets then for some set E we use the following notation $\mathcal{B} \cap E = \{G \cap E : G \in \mathcal{B}\}$.

3.1.1. Maximal function. Let

$$(6) \quad M_{\mathcal{M}}f(x) = \sup_{x \in \Delta, \Delta \in \mathcal{M}} \frac{1}{|\Delta|} \int_{\Delta} |f(t)| dt, \quad f \in L^1_{loc}(\mathbb{R}).$$

Let us consider also a maximal function with respect to a weight function ω defined by the following equation

$$(7) \quad M_{\mathcal{M},\omega}f(x) = \sup_{x \in \Delta, \Delta \in \mathcal{M}} \frac{1}{\omega(\Delta)} \int_{\Delta} |f(t)|\omega(t) dt, \quad f \in L^1_{loc}(\mathbb{R}, \omega),$$

where $\omega(\Delta) = \int_{\Delta} \omega(t) dt$.

PROPOSITION 3.1. *Let $\omega(x) \geq 0$ for $x \in \mathbb{R}$, $\omega \in L^1_{loc}(\mathbb{R})$ and let $f \in L^1_{loc}(\mathbb{R}, \omega)$. Then for any $\lambda > 0$*

$$(8) \quad \omega(\{t \in \mathbb{R} : M_{\mathcal{M},\omega}f(t) > \lambda\}) \leq \frac{1}{\lambda} \int_{\mathbb{R}} |f(t)|\omega(t) dt.$$

PROOF. If $x \in \Omega_{\lambda}(f) := \{t \in \mathbb{R} : M_{\mathcal{M},\omega}f(t) > \lambda\}$ then for some $\Delta \in \mathcal{M}$ such that $x \in \Delta$

$$\frac{1}{\omega(\Delta)} \int_{\Delta} |f(t)|\omega(t) dt > \lambda.$$

Observe that among all intervals which have the above properties there exists $\Delta_x \in \mathcal{M}$ with maximal ω -measure. Thus, having in mind that \mathcal{M} is numerable we can find a sequence of mutually disjoint intervals $\{\Delta_v\} \subset \mathcal{M}$ so that $\Omega_{\lambda} = \bigcup_{v=1}^{\infty} \Delta_v$. □

We also have

PROPOSITION 3.2. *Let $\omega(x) \geq 0$ for $x \in \mathbb{R}$, $\omega \in L^1_{loc}(\mathbb{R})$. Then for any $f \in L^p_{loc}(\mathbb{R}, \omega)$, $p > 1$*

$$(9) \quad \int_{\mathbb{R}} M_{\mathcal{M},\omega}f(t)^p \omega(t) dt \leq \frac{2^p p}{p-1} \int_{\mathbb{R}} |f(t)|^p \omega(t) dt.$$

PROOF. Following the proof of the corresponding result for the Lebesgue measure (see [42], p.7) we split f into two parts, $f = f_1 + f_2$, where $f_1(x) = f(x)$ if $|f(x)| \geq \frac{\lambda}{2}$ and $f_1(x) = 0$ otherwise. Then we have $|f(x)| \leq |f_1(x)| + \frac{\lambda}{2}$. Hence, $M_{\mathcal{M},\omega}f(x) \leq M_{\mathcal{M},\omega}f_1(x) + \frac{\lambda}{2}$ and $\Omega_{\lambda}(f) \subseteq \Omega_{\frac{\lambda}{2}}(f_1)$. Thus by Proposition 3.1 we have that

$$(10) \quad \omega(\Omega_{\lambda}(f)) \leq \frac{2}{\lambda} \int_{\mathbb{R}} |f_1(t)|\omega(t) dt = \frac{2}{\lambda} \int_{\Omega_{\frac{\lambda}{2}}(f_1)} |f(t)|\omega(t) dt.$$

Afterwards we have to use the following equality for any measurable function $g : \mathbb{R} \rightarrow \mathbb{R}$

$$\begin{aligned} \int_{\mathbb{R}} |g(t)|^p \omega(t) dt &= p \int_{\mathbb{R}} \int_{[0, |g(t)|]} \lambda^{p-1} d\lambda \omega(t) dt \\ &= p \int_0^{+\infty} \lambda^{p-1} \omega(\{t : |g(t)| > \lambda\}) d\lambda. \end{aligned}$$

Hence, by (10) we obtain

$$\begin{aligned} \int_{\mathbb{R}} M_{\mathcal{M}, \omega} f(t)^p \omega(t) dt &\leq 2p \int_0^{+\infty} \lambda^{p-2} \int_{\Omega_{\frac{\lambda}{2}}(f)} |f(t)| \omega(t) dt d\lambda \\ &= 2p \int_{\mathbb{R}} |f(t)| \omega(t) \int_0^{2|f(t)|} \lambda^{p-2} d\lambda dt = \frac{2^p p}{p-1} \int_{\mathbb{R}} |f(t)|^p \omega(t) dt \end{aligned}$$

□

3.1.2. Calderón-Zygmund decomposition for m -adic intervals. We need a modified version for the Calderón-Zygmund decomposition (see [43]) for the m -adic intervals. Let $f \in L^1[0, 1]$, $f \geq 0$ and let $\lambda > 0$ is such that

$$\int_{[0,1]} f(t) dt < \lambda.$$

At the first step we take m intervals $\{I_k\}_{k=1}^m \subset \mathcal{M} \cap [0, 1]$ such that $|I_k| = \frac{1}{m}$, $1 \leq k \leq m$ and $\bigcup_{k=1}^m I_k = [0, 1]$. Let $\{I_{l_i}\}_{i=1}^{m_1} \subset \{I_k\}_{k=1}^m$ be all those intervals for which $\eta_{I_{l_i}} > \lambda$, $1 \leq i \leq m_1 < m$, where

$$\frac{1}{|I|} \int_I f(t) dt := \eta_I.$$

Those intervals are renamed G_1, \dots, G_{m_1} . Clearly,

$$(11) \quad \lambda < \frac{1}{|G_l|} \int_{G_l} f(t) dt \leq m\lambda.$$

If $\eta_{I_k} \leq \lambda$ for all $1 \leq k \leq m$ then we put $m_1 = 0$. On the next step we repeat the same procedure on any of those intervals that were not renamed. The collection of all m -adic intervals which are separated on the second step are renamed $G_{m_1+1}, \dots, G_{m_2}$. For those intervals the condition (11) holds again. On the v th step all m -adic intervals which are separated are renamed $G_{m_{v-1}+1}, \dots, G_{m_v}$. If no any interval is separated then we put $m_{v+1} = m_v$. This procedure produces a collection of disjoint m -adic intervals $\{G_l\}$ for which the condition (11) holds and

$$(12) \quad |\Omega| = \sum_l |G_l| < \frac{1}{\lambda} \sum_l \int_{G_l} f(t) dt \leq \frac{1}{\lambda} \int_{[0,1]} f(t) dt,$$

where $\Omega = \bigcup_l G_l$ and for any $I \in \mathcal{M} \cap [0, 1]$, $I \subset \Omega^c := [0, 1] \setminus \Omega$ we have that $\eta_I \leq \lambda$. The collection $\{G_l\}$ will be called Calderón-Zygmund m -adic decomposition at level λ . Let

$$(13) \quad g(x) = f(x) \chi_{\Omega^c}(x) + \sum_l \eta_{G_l} \chi_{G_l}(x) \quad \text{and} \quad b(x) = f(x) - g(x).$$

We skip the details of the proof of the following

PROPOSITION 3.3. *Let $f \in L^1[0, 1]$, and let $\lambda > 0$ be such that $\int_{[0,1]} |f(t)|dt < \lambda$. Then there exists a family of disjoint sets $\{G_l\}_{l \in \Upsilon} \subset \mathcal{M}$ such that*

$$(14) \quad |f(x)| \leq \lambda \quad \text{a.e. on } \Omega^c, \quad \text{where } \Omega = \bigcup_{l \in \Upsilon} G_l,$$

(12) is true and for any $l \in \Upsilon$ holds (11). Moreover, $f(x) = g(x) + b(x)$, where g is defined by (13) and the following conditions hold:

$$(15) \quad |g(x)| \leq m\lambda \quad \text{a.e. on } [0, 1];$$

$$(16) \quad \|g(x)\|_p^p \leq (m\lambda)^{p-1} \|f\|_1 \quad \text{for all } 1 \leq p < \infty;$$

$$(17) \quad \int_{G_l} b(t)dt = 0 \quad \text{for all } l \in \Upsilon.$$

For dyadic intervals a similar result was obtained by C. Watari [44].

3.1.3. Classes of $\mathcal{M}_p, p \geq 1$ weights.

DEFINITION 3.4. We say that a non negative locally integrable function ω satisfies the condition $\mathcal{M}_p, p \geq 1$ if

$$(18) \quad \omega(\Delta) \left[\int_{\Delta} \omega^{-\frac{1}{p-1}}(t)dt \right]^{p-1} \leq C_p |\Delta|^p \quad \forall \Delta \in \mathcal{M},$$

where $C_p > 0$ is independent of $\Delta \in \mathcal{M}$. For $p = 1$ it is understood that

$$\left[\int_{\Delta} \omega^{-\frac{1}{p-1}}(t)dt \right]^{p-1} := \|\omega^{-1}\|_{L^\infty(\Delta)}.$$

We say that ω satisfies the condition $\mathcal{M}_p(G)$, where $G \subset \mathbb{R}$ if (18) holds for all $\Delta \in \mathcal{M} \cap G$. The reader should observe that the conditions $\mathcal{M}_p([0, 1])$ and $\mathcal{M}_p((0, 1])$ are distinct. In the second case the intervals $[0, 2^{-j}]$, $j \in \mathbb{N}$ should be excluded when one checks the inequality (18). The following lemma is obvious.

LEMMA 3.5. *Let ω satisfy the condition $\mathcal{M}_p, p > 1$ then $\psi = \omega^{-\frac{1}{p-1}}$ satisfies the condition $\mathcal{M}_{p'}$, where $\frac{1}{p} + \frac{1}{p'} = 1$.*

We follow the ideas given in [7] to prove the following result.

PROPOSITION 3.6. *Let $\omega(x) \geq 0$ for $x \in \mathbb{R}$, $\omega \in L^1_{loc}(\mathbb{R})$. Then for any $f \in L^p_{loc}(\mathbb{R}, \omega), p > 1$*

$$(19) \quad \int_{\mathbb{R}} M_{\mathcal{M}} f(t)^p \omega(t)dt \leq B_p \int_{\mathbb{R}} |f(t)|^p \omega(t)dt,$$

for some $B_p > 0$ independent of f if and only if ω satisfies the condition \mathcal{M}_p .

PROOF. Suppose that (19) is true. For any $\Delta \in \mathcal{M}$ we have by (6) that

$$\frac{1}{|\Delta|} \int_{\Delta} |f(t)| dt \chi_{\Delta}(x) \leq M_{\mathcal{M}} f(x).$$

Hence, by (19) we have that

$$\left(\frac{1}{|\Delta|} \int_{\Delta} |f(t)| dt \right)^p \omega(\Delta) \leq B_p \int_{\mathbb{R}} |f(t)|^p \omega(t) dt.$$

Letting $f(t) = \omega^{-\frac{1}{p-1}}(t) \chi_{\Delta}(t)$ we obtain (18) with $C_p = B_p$. To prove the opposite assertion one has to use Proposition 3.1 and the following

LEMMA 3.7. *Let ω satisfy the condition \mathcal{M}_p , $p > 1$ then there exists $\varepsilon > 0$ such that ω satisfies the condition $\mathcal{M}_{p-\varepsilon}$.*

We skip the rest of the proof because the proof in [7] works with small changes. □

DEFINITION 3.8. We say that a non negative locally integrable function ω satisfies the condition \mathcal{M}_{∞} if there exists $C > 0$ and $\delta > 0$ such that for any $\Delta \in \mathcal{M}$ and any measurable subset $E \subseteq \Delta$

$$(20) \quad \frac{\omega(E)}{\omega(\Delta)} \leq C \left(\frac{|E|}{|\Delta|} \right)^{\delta}.$$

We skip the detailed proofs of the following two lemmas because the corresponding proofs in [7] for A_p weights work with obvious changes.

LEMMA 3.9. *Let ω satisfy the condition \mathcal{M}_p , $p > 1$ then there exist $r > 0$ and $C > 0$ such that*

$$(21) \quad \left(\frac{1}{|\Delta|} \int_{\Delta} \omega(x)^{1+r} \right)^{\frac{1}{1+r}} \leq C \frac{\omega(\Delta)}{|\Delta|} \quad \forall \Delta \in \mathcal{M}.$$

LEMMA 3.10. *Let ω satisfy the condition \mathcal{M}_p for some $p > 1$ then ω satisfies the condition \mathcal{M}_{∞} .*

We need also the following related with \mathcal{M} -neighborhoods of $+\infty$ and $-\infty$ (see (5)).

LEMMA 3.11. *Let ω satisfy the condition \mathcal{M}_p for some $p > 1$ then*

$$\omega \notin L(\Delta_j(+\infty)), \quad \omega \notin L(\Delta_j(-\infty)), \quad \forall j \in \mathbb{N}_0.$$

PROOF. We prove the assertion for the neighborhoods of $+\infty$. For this purpose we observe that there exists $C_p > 0$ such that for any $j \in \mathbb{N}_0$

$$(22) \quad \omega(\Delta_{-j-1}) \leq C_p \omega(\Delta_{-j-1} \setminus \Delta_{-j}).$$

For any $j \in \mathbb{N}_0$ and any locally integrable function $f \geq 0$ we have that $M_{\mathcal{M}} f(x) \geq m^{-j} \int_{[0, m^j]} f(t) dt$ if $x \in [0, m^j]$. By Proposition 3.6 we obtain that

$$\left(m^{-j} \int_{[0, m^j]} f(t) dt \right)^p \omega([0, m^j]) \leq B_p \int_{[0, m^j]} f(t)^p \omega(t) dt.$$

Putting $j + 1$ instead of j in the above inequality and letting f be the characteristic function of the set $\Delta_{-j-1} \setminus \Delta_{-j}$ we obtain the inequality (22). If $\omega \in L(\Delta_{j_0}(+\infty))$ then for any $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that

$$\int_{[m^N, +\infty)} \omega(t)dt < \varepsilon .$$

Which leads to a contradiction with the condition (22). Evidently the proof for the neighborhoods of $-\infty$ is similar. □

DEFINITION 3.12. Let $\omega \geq 0$ be a weight function defined on \mathbb{R}^+ . We will say that ω satisfies the condition $\mathcal{M}_p^y(\mathbb{R}^+)$, $p \geq 1$ for some $y \in \mathbb{R}^+$ if

$$(23) \quad \omega(\Delta_j(y)) \left[\int_{\mathbb{R}^+ \setminus \Delta_j(y)} \omega^{-\frac{1}{p-1}}(t)dt \right]^{p-1} \leq C_p |\Delta_j(y)|^p \quad \forall j \in \mathbb{Z},$$

where $C_p > 0$ is independent of $j \in \mathbb{Z}$.

We will not formulate the definition of the condition $\mathcal{M}_p^y(\mathbb{R}^-)$ because it is clear from the context.

DEFINITION 3.13. Let $\omega \geq 0$ be a weight function defined on Δ , where $\Delta \in \mathcal{M}$, $|\Delta| = m^l$. We will say that ω satisfies the condition $\mathcal{M}_p^y(\Delta)$, $p \geq 1$ for some $y \in \Delta$ if

$$(24) \quad \omega(\Delta_j(y)) \left[\int_{\Delta \setminus \Delta_j(y)} \omega^{-\frac{1}{p-1}}(t)dt \right]^{p-1} \leq C_p |\Delta_j(y)|^p \quad \forall j > l,$$

where $C_p > 0$ is independent of j .

LEMMA 3.14. Let $w \geq 0$ be a weight function defined on $[0, 1]$ such that w satisfies the condition $\mathcal{M}_p^y([0, 1])$ for some $y \in [0, 1]$ and $1 < p < \infty$. Then there exists $q_p > 1$ such that

$$\int_{[0,1] \setminus \Delta_{j+1}(y)} w^{-\frac{1}{p-1}}(t)dt \left(\int_{[0,1] \setminus \Delta_j(y)} w^{-\frac{1}{p-1}}(t)dt \right)^{-1} \geq q_p$$

for all $j \in \mathbb{N}$.

PROOF. For any $j \in \mathbb{N}$ we have that

$$\begin{aligned} & \int_{[0,1] \setminus \Delta_{j+1}(y)} w^{-\frac{1}{p-1}}(t)dt \left(\int_{[0,1] \setminus \Delta_j(y)} w^{-\frac{1}{p-1}}(t)dt \right)^{-1} \\ & \geq 1 + \int_{\Delta_j(y) \setminus \Delta_{j+1}(y)} w^{-\frac{1}{p-1}}(t)dt \left(\int_{[0,1] \setminus \Delta_j(y)} w^{-\frac{1}{p-1}}(t)dt \right)^{-1} \\ & \geq 1 + \int_{\Delta_j(y) \setminus \Delta_{j+1}(y)} w^{-\frac{1}{p-1}}(t)dt \left(w(\Delta_j(y)) \right)^{\frac{1}{p-1}} C_p^{-\frac{1}{p-1}} |\Delta_j(y)|^{-\frac{p}{p-1}} \\ & \geq 1 + |\Delta_j(y) \setminus \Delta_{j+1}(y)|^{\frac{p}{p-1}} C_p^{-\frac{1}{p-1}} |\Delta_j(y)|^{-\frac{p}{p-1}} \geq 1 + C_p^{-\frac{1}{p-1}} \left(\frac{m-1}{m} \right)^{\frac{p}{p-1}} . \end{aligned}$$

Putting $q_p = 1 + C_p^{-\frac{1}{p-1}} \left(\frac{m-1}{m}\right)^{\frac{p}{p-1}}$ we finish the proof. □

Following three lemmas will be used in the last section. In the proofs we will use the following notation: $aE = \{at : t \in E\}$.

LEMMA 3.15. *Let $\omega \geq 0$ satisfy the condition $\mathcal{M}_p(\mathbb{R}^+)$, $p > 1$ with a constant $C_p > 0$. Then for any $N \in \mathbb{N}$ the weight function $\omega_N(x) := \omega(m^N x)$ satisfies the condition $\mathcal{M}_p([0, 1])$ with the same constant C_p .*

PROOF. For any $E \in \mathcal{M} \cap [0, 1]$ we have

$$\begin{aligned} \omega_N(E) \left[\int_E \omega_N^{-\frac{1}{p-1}}(t) dt \right]^{p-1} &= m^{-N} \int_{m^N E} \omega(x) dx \left[m^{-N} \int_{m^N E} \omega^{-\frac{1}{p-1}}(x) dx \right]^{p-1} \\ &\leq m^{-pN} C_p |m^N E|^p = C_p |E|^p. \end{aligned}$$

We have used that $m^N E \in \mathcal{M}$. □

LEMMA 3.16. *Let $N \in \mathbb{N}$ and let $y \in [0, m^N]$. Suppose that $\omega \geq 0$ satisfies the condition $\mathcal{M}_p([0, m^N] \setminus \{y\})$, $p > 1$ with a constant $C_p > 0$. Then the weight function $\omega_N(x) := \omega(m^N x)$ satisfies the condition $\mathcal{M}_p([0, 1] \setminus \{y_N\})$ with the same constant C_p , where $y_N = m^{-N} y$.*

PROOF. For any $E \in \mathcal{M} \cap ([0, 1] \setminus \{y_N\})$ we observe that the interval $m^N E \in \mathcal{M} \cap ([0, m^N] \setminus \{y\})$. The rest of the proof is the same as above. □

LEMMA 3.17. *Let $N \in \mathbb{N}$ and let $y \in [0, m^N]$. Suppose that $\omega \geq 0$ satisfies the condition $\mathcal{M}_p^y([0, m^N])$, $p > 1$ with a constant $C_p > 0$. Then the weight function $\omega_N(x) := \omega(m^N x)$ satisfies the condition $\mathcal{M}_p^{y_N}([0, 1])$ with the same constant C_p , where $y_N = m^{-N} y$.*

PROOF. For any $j \in \mathbb{N}$ we have

$$\begin{aligned} \omega_N(\Delta_j(y_N)) \left[\int_{[0,1] \setminus \Delta_j(y_N)} \omega_N^{-\frac{1}{p-1}}(t) dt \right]^{p-1} \\ = m^{-N} \int_{\Delta_{j-N}(y)} \omega(x) dx \left[m^{-N} \int_{[0,m^N] \setminus \Delta_{j-N}(y)} \omega^{-\frac{1}{p-1}}(x) dx \right]^{p-1} \\ \leq m^{-pN} C_p |\Delta_{j-N}(y)|^p = C_p |\Delta_j(y)|^p. \end{aligned}$$

□

3.2. Higher rank Haar wavelets. We bring the definition of higher rank Haar wavelets without recalling the general theory of multiresolution analysis. For relations of these type of wavelets with p -adic analysis see [33]. Let $\varphi(x) = \chi_{[0,1]}(x)$ and let

$$(25) \quad V(m) = \text{span}\{\varphi_{1,j,m}(x) : 0 \leq j \leq m - 1\}$$

for any $m = 2, 3, \dots$. Afterwards, let $\{h^{(\nu)}(x) : 0 \leq \nu \leq m - 1\}$ be an orthonormal basis in $V(m)$ such that $h^{(0)}(x) = \varphi(x)$. The system

$$(26) \quad H(m) = \{h_{k,j,m}^{(\nu)}(x) : k \in \mathbb{Z}; j \in \mathbb{Z}; 1 \leq \nu \leq m - 1\},$$

where

$$(27) \quad h_{k,j,m}^{(\nu)}(x) = m^{k/2} h^{(\nu)}(m^k x - j)$$

will be called m -th rank Haar system. Sometimes we will use also the following notation

$$(28) \quad h_{\Delta}^{(\nu)} := h_{k,j,m}^{(\nu)}(x) \quad \text{when} \quad \Delta = \left[\frac{j}{m^k}, \frac{j+1}{m^k} \right].$$

The orthogonality of the system (26) is obvious.

THEOREM 3.18. *The system $H(m)$ is complete in $L^p(\mathbb{R})$, $1 < p < \infty$.*

PROOF. Let p , $1 < p < \infty$ be fixed. It is easy to observe that the proof will be finished if we show that for any $\varphi_{1,j,m}(x)$, $0 \leq j \leq m - 1$ and any $\varepsilon > 0$ there exists a finite linear combination P_j of functions $\{h_{k,l,m}^{(\nu)}(x) : k \in \mathbb{Z} \setminus \mathbb{N}_0; l \in \mathbb{Z}; 1 \leq \nu \leq m - 1\}$ such that $\|\varphi_{1,j,m} - P_j\|_{L^p(\mathbb{R})} < \varepsilon$.

Let $l \in \mathbb{N}$ be such that $m^{1/2} m^{l(1/p-1)} < \varepsilon$. We set

$$V^{(l)}(m) = \text{span}\{\varphi_{1,j,m}(x) : 0 \leq j \leq m^{l+1} - 1\}.$$

It is clear that $\dim V^{(l)}(m) = m^{l+1}$. Let us show by induction that there are exactly $m^{l+1} - 1$ functions from the system $\{h_{k,j,m}^{(\nu)}(x) : 1 \geq k \geq -l + 1; j \in \mathbb{Z}; 1 \leq \nu \leq m - 1\}$ with supports in $[0, m^l]$.

If $l = 0$ then it is obvious. Suppose that for some $\mu \in \mathbb{N}$ we have that the number of functions from the system

$$(29) \quad \{h_{k,j,m}^{(\nu)}(x) : 1 \geq k \geq -\mu + 1; j \in \mathbb{Z}; 1 \leq \nu \leq m - 1\}$$

with supports in $[0, m^\mu]$ is equal to $m^{\mu+1} - 1$. Then it is clear that there are $(m^{\mu+1} - 1)m$ functions from the system (29) that have their supports in $[0, m^{\mu+1}]$. Note that the functions $\{h_{-\mu,0,m}^{(\nu)}(x) : 1 \leq \nu \leq m - 1\}$ vanish outside the closed interval $[0, m^{\mu+1}]$. Thus we have $(m^{\mu+1} - 1)m + m - 1 = m^{\mu+2} - 1$ mutually orthogonal functions in (29) which have their supports in $[0, m^{\mu+1}]$.

Let $\{g_i\}_{i=1}^{m^{l+1}-1}$ be all functions from the system

$$\{h_{k,j,m}^{(\nu)}(x) : 1 \geq k \geq -l + 1; j \in \mathbb{Z}; 1 \leq \nu \leq m - 1\}$$

that have their supports in $[0, m^l]$. Evidently $\{g_i\}_{i=1}^{m^{l+1}-1} \subset V^{(l)}(m)$.

Let $g_0(x) = \chi_{[0,m^l]}(x)$. Then $g_0 \in V^{(l)}(m)$ and g_0 is orthogonal to all elements of $\{g_i\}_{i=1}^{m^{l+1}-1}$. Hence, $\{g_i\}_{i=0}^{m^{l+1}-1}$ is a basis in $V^{(l)}(m)$ and

$$\varphi_{1,j,m} = \sum_{i=0}^{m^{l+1}-1} a_i^{(j)} g_i, \quad \text{where} \quad a_0^{(j)} = m^{-l} \int_{[0,m^l]} \varphi_{1,j,m}(t) dt.$$

Thus we obtain that for any $0 \leq j \leq m - 1$

$$\left\| \varphi_{1,j,m} - \sum_{i=1}^{m^{l+1}-1} a_i^{(j)} g_i \right\|_{L^p(\mathbb{R})} = \left\| a_0^{(j)} g_0 \right\|_{L^p(\mathbb{R})} = m^{1/2} m^{l(1/p-1)} < \varepsilon.$$

□

4. m -th rank Haar system on $[0, 1]$. Let $h_0(x) \equiv 1$ for $x \in [0, 1]$. For any $n \in \mathbb{N}$ we have a unique representation

$$(30) \quad n = m_k + j - 1, \quad \text{where } k \in \mathbb{N}, 1 \leq j \leq m^k,$$

and

$$(31) \quad m_k = 1 + m + m^2 + \dots + m^{k-1}, \quad m_1 = 1.$$

For any $1 \leq \nu \leq m - 1$ we put

$$h_n^{(\nu)}(x) = h_{k,j-1,m}^{(\nu)}(x) \quad \text{for } x \in [0, 1].$$

Afterwards we enumerate the functions in the following way

$$(32) \quad h_l(x) = h^{(l)}(x) \quad \text{for } 1 \leq l \leq m - 1;$$

$$(33) \quad h_l(x) = h_n^{(\nu)}(x) \quad \text{for } l = \nu + n(m - 1), n \in \mathbb{N}.$$

We denote the m th rank Haar system by $\mathcal{H}(m) = \{h_l(x)\}_{l=0}^\infty$. We also let

$$(34) \quad \mu_0 = 0, \mu_1 = \mu_0 + m - 1, \dots, \mu_{k+1} = \mu_k + (m - 1)m^k, \dots$$

The following lemma is the analogue of Schauder’s lemma for the classical Haar system (see [41]). For any $f \in L^1[0, 1]$ and for any $1 \leq j \leq m^k, k \in \mathbb{N}$ we put

$$\Theta_{\mu_k+j(m-1)}(f, x) = \sum_{l=0}^{\mu_k} a_l(f) h_l(x) + \sum_{s=0}^{j-1} \sum_{\nu=1}^{m-1} a_{k,s,m}^{(\nu)}(f) h_{k,s,m}^{(\nu)}(x),$$

where

$$a_l(f) = \int_{[0,1]} f(t) h_l(t) dt; \quad a_{k,s,m}^{(\nu)}(f) = \int_{[0,1]} f(t) h_{k,s,m}^{(\nu)}(t) dt.$$

LEMMA 4.1. *Let $f \in L^1[0, 1]$ and let $1 \leq j \leq m^k, k \in \mathbb{N}_0$. Then the partial sum $\Theta_{\mu_k+j(m-1)}(f, x)$ is constant on any interval from the collection of sets*

$$(35) \quad \left\{ \left[\frac{s}{m^{k+1}}, \frac{s+1}{m^{k+1}} \right] : 0 \leq s \leq jm - 1 \right\},$$

$$(36) \quad \left\{ \left[\frac{l}{m^k}, \frac{l+1}{m^k} \right] : j \leq l \leq m^k - 1 \right\}.$$

Moreover, for any Δ from (35) or from (36)

$$(37) \quad \Theta_{\mu_k+j(m-1)}(f, x) = \frac{1}{|\Delta|} \int_{\Delta} f(t) dt \quad \text{for } x \in \Delta.$$

PROOF. At first we show that the assertion of the lemma is true for $\Theta_{\mu_k}(f, x)$, $k \in \mathbb{N}_0$.
Indeed,

$$\text{span}\{h_l, 0 \leq l \leq m^k - 1\} = V(m),$$

where $V(m)$ is defined by (25). Hence, for any Δ from (36) with $j = 1$ we have that for $x \in \Delta$

$$\Theta_{\mu_k}(f, x) = \sum_{l=0}^{m^k-1} \int_{[0,1]} f(t)\varphi_{k,l,m}(t)dt \varphi_{k,l,m}(x) = \frac{1}{|\Delta|} \int_{\Delta} f(t)dt.$$

Afterwards we observe that in the general case

$$\Theta_{\mu_k+j(m-1)}(f, x) = \Theta_{\mu_{k+1}}(f, x) \quad \text{if } x \in \left[0, \frac{j}{m^k}\right]$$

and

$$\Theta_{\mu_k+j(m-1)}(f, x) = \Theta_{\mu_k}(f, x) \quad \text{if } x \in \left[\frac{j}{m^k}, 1\right].$$

Which finishes the proof. □

By Lemma 4.1 we obtain the following corollaries.

COROLLARY 4.2. *For any $m = 2, 3, \dots$ the system $\mathcal{H}(m)$ is a basis in any space $L^p[0, 1]$, $1 \leq p < \infty$.*

PROOF. By Lemma 4.1 as in the case of the classical Haar system we have that

$$\|\Theta_{\mu_k+j(m-1)}\|_{L^p \rightarrow L^p} \leq 1 \quad \text{for all } 1 \leq j \leq m^k, k \in \mathbb{N}.$$

To finish the proof we have to check that $\lim_{l \rightarrow \infty} |a_l(f)| \|h_l\|_{L^p[0,1]} = 0$. We skip the technical details because afterwards we are going to return to the similar question in the weighted norm case. □

COROLLARY 4.3. *For any $f \in L^1[0, 1]$ the Fourier series of f with respect to the system $\mathcal{H}(m)$, $m = 2, 3, \dots$ converges almost everywhere to f on $[0, 1]$.*

PROOF. For every $x \in [0, 1]$ which is a Lebesgue point of f we have that

$$\lim_{k \rightarrow \infty} \Theta_{\mu_k}(f, x) = f(x).$$

Afterwards we observe that $|a_l(f)| \|h_l(x)\| \leq CM_{\mathcal{M}}(f, x)$ which finishes the proof. □

For any $k \in \mathbb{N}_0$ and any $1 \leq j \leq m^k$ consider the kernel

$$(38) \quad K_{kj}(t, x) = \sum_{l=0}^{\mu_k} h_l(t)h_l(x) + \sum_{s=0}^{j-1} \sum_{v=1}^{m-1} h_{k,s,m}^{(v)}(t)h_{k,s,m}^{(v)}(x).$$

Let $\{G_i : 1 \leq i \leq m^k + j(m - 1)\}$ be mutually disjoint sets from (35) and (36). Further in the paper we will need the following result.

LEMMA 4.4. *Let $k \in \mathbb{N}_0$ and $1 \leq j \leq m^k$. Then the kernel*

$$(39) \quad K_{kj}(t, x) = \frac{1}{|G_i|} \quad \text{for } (t, x) \in G_i^2, 1 \leq i \leq m^k + j(m - 1);$$

and

$$K_{kj}(t, x) = 0, \quad \text{if } (t, x) \in [0, 1]^2 \setminus \bigcup_{i=1}^{m^k+j(m-1)} G_i^2.$$

PROOF. We have that $\{|G_i|^{-1/2}\chi_{G_i}(x)\}_{i=1}^{m^k+j(m-1)}$ is an orthonormal system of functions. From Lemma 4.1 it follows that the orthonormal system of functions $\{h_l(x)\}_{l=0}^{\mu_k} \cup \{h_{k,s,m}^{(v)}(x) : 0 \leq s \leq j - 1, 1 \leq v \leq m - 1\}$ can be obtained from the set of functions $\{|G_i|^{-1/2}\chi_{G_i}(x)\}_{i=1}^{m^k+j(m-1)}$ by an orthogonal transformation. Hence,

$$\begin{aligned} K_{kj}(t, x) &= \sum_{i=1}^{m^k+j(m-1)} |G_i|^{-1/2}\chi_{G_i}(t)|G_i|^{-1/2}\chi_{G_i}(x) \\ &= \sum_{i=1}^{m^k+j(m-1)} |G_i|^{-1}\chi_{G_i}(t)\chi_{G_i}(x). \end{aligned}$$

□

DEFINITION 4.5. We say that a system of functions $\{\phi_k\}_{k=1}^\infty \subset L^\infty[0, 1]$ is total with respect to $L^1[0, 1]$ if

$$(40) \quad \int_{[0,1]} f(t)\phi_k(t)dt = 0 \quad \text{for all } k \in \mathbb{N} \quad \text{for some } f \in L^1[0, 1]$$

if and only if $f = 0$ a.e. on $[0, 1]$.

By Lemma 4.1 it follows immediately

COROLLARY 4.6. *The system $\mathcal{H}(m)$, $m = 2, 3, \dots$ is total with respect to $L^1[0, 1]$.*

THEOREM 4.7. *For any $m = 2, 3, \dots$ the system $\mathcal{H}(m)$ is an unconditional basis in any space $L^p[0, 1]$, $1 < p < \infty$.*

The reader can find well known facts about unconditional bases in [36]. For any sequence $\varepsilon = \{\varepsilon_l\}_{l=0}^\infty$, where $\varepsilon = \pm 1$ we consider an operator $I_\varepsilon : L^1[0, 1] \rightarrow L^0[0, 1]$ defined as follows $I_\varepsilon(f, x) = \sum_{l=0}^\infty \varepsilon_l a_l(f)h_l$.

PROPOSITION 4.8. *The operator I_ε is of weak-(1, 1) type.*

PROOF. We adopt the idea of the proof given in [44]. Let $f \in L^\infty[0, 1]$ and suppose that $\lambda > \|f\|_1$. Without loss in generality we can suppose that $f \geq 0$ (see [42], pp. 21–22). By Proposition 3.3 we write $f(x) = g(x) + b(x)$, where g satisfies the condition (15).

The system $\mathcal{H}(m)$ is a complete orthonormal system. Hence, $I_\varepsilon : L^2[0, 1] \rightarrow L^2[0, 1]$ is an isometry. Thus by the Tchebychev inequality we will have

$$(41) \quad |\{x \in [0, 1] : |I_\varepsilon(g, x)| > \lambda\}| \leq \frac{1}{\lambda^2} \int_{[0,1]} g^2(x)dx \leq \frac{m}{\lambda} \|g\|_{L^1[0,1]} \leq \frac{m}{\lambda} \|f\|_{L^1[0,1]},$$

where the last inequality follows by (16). Afterwards, we apply the following property of m -adic intervals. If $\Delta_1, \Delta_2 \subset \mathcal{M}$ then only two relations are possible or $\Delta_1 \cap \Delta_2 = \emptyset$ or one of those intervals is a subset of another interval. By the definition of the system $\mathcal{H}(m)$ and by (17) it is easy to deduce that $I_\varepsilon(b, x) = 0$ for $x \in \Omega^c$. Thus by (12) and (41) we obtain

$$(42) \quad |\{x \in [0, 1] : |I_\varepsilon(f, x)| > \lambda\}| \leq \frac{m+1}{\lambda} \|f\|_{L^1[0,1]}.$$

From the last inequality readily follows that for any $f \in L^1[0, 1]$ the series $\sum_{l=0}^\infty \varepsilon_l a_l(f) h_l$ converges in measure on $[0, 1]$. Observe that in the proof of the inequality (42) the condition $f \in L^\infty[0, 1]$ was used only to claim the existence of $I_\varepsilon(f, x)$. Hence, the proof is complete. \square

The analogue of Proposition 4.8 for the Haar system was obtained by S. Yano [47].

PROOF. By Proposition 4.8 and the Marcinkiewicz interpolation theorem (see [48]) we obtain that $\mathcal{H}(m)$ is an unconditional basis in $L^p[0, 1]$, $1 < p \leq 2$. Afterwards by duality we finish the proof of Theorem 4.7. \square

For the system $\mathcal{H}(m)$ we put

$$G_m(f, x) = \left(\sum_{l=0}^\infty |a_l(f) h_l(x)|^2 \right)^{\frac{1}{2}}, \quad \text{where } f \in L^1[0, 1].$$

For the operator $G_m : L^1[0, 1] \rightarrow L^0[0, 1]$ the following proposition holds.

PROPOSITION 4.9. *The operator G_m is of weak-(1, 1) type.*

PROOF. Let $f = \sum_{l=0}^N a_l h_l$ be any polynomial with respect to the system $\mathcal{H}(m)$ and let $\varepsilon = \{\varepsilon_l\}_{l=0}^N$ be any Rademacher sequence. By a well known inequality (see [25],p.8) we have that for any $0 < \alpha < 1$ and any $x \in [0, 1]$

$$\mathcal{P}(I_\varepsilon(f, x) > \alpha G_m(f, x)) > \frac{1}{3}(1 - \alpha)^2.$$

Observe that $G_m(I_\varepsilon(f, \cdot), x) = G_m(f, x)$ for any Rademacher sequence ε . For any $\lambda > 0$ we have that if for some $x \in [0, 1]$ $I_\varepsilon(f, x) > \alpha G_m(f, x)$ and $G_m(f, x) > \frac{\lambda}{\alpha}$ then $I_\varepsilon(f, x) > \lambda$. Hence, by Proposition 4.8 we finish the proof for the polynomials with respect to the system $\mathcal{H}(m)$.

For arbitrary $f \in L^1[0, 1]$ we have that the sequence $G_m(\Theta_n(f, \cdot), x)$ is an increasing sequence which a.e. converges to $G_m(f, x)$. Hence,

$$|\{G_m(f, x) > \lambda\}| = \lim_{n \rightarrow +\infty} |\{G_m(\Theta_n(f, \cdot), x) > \lambda\}|$$

$$< \frac{C}{\lambda} \sup_n \|\Theta_n(f, \cdot)\|_{L^1[0,1]} \leq \frac{C_1}{\lambda} \|f\|_{L^1[0,1]}.$$

□

By standard arguments (see [46]) one can derive from Theorem 4.7 that for all $1 < p < \infty$

$$(43) \quad B_p \|f\|_{L^p[0,1]} \leq \|G_m(f, \cdot)\|_{L^p[0,1]} \leq C_p \|f\|_{L^p[0,1]},$$

where $B_p > 0, C_p > 0$ are independent of $f \in L^p[0, 1]$.

4.1. Haar wavelet systems as unconditional bases in $L^p(\mathbb{R}), 1 < p < \infty$. From Theorem 4.7 we easily derive

THEOREM 4.10. *For any $m = 2, 3, \dots$ the system $H(m)$ is an unconditional basis in any space $L^p(\mathbb{R}), 1 < p < \infty$.*

Further we will use the following notations: $\mathbb{R}^+ = [0, +\infty), \mathbb{R}^- = (-\infty, 0]$ and $\mathbb{Z}^+ = \mathbb{R}^+ \cap \mathbb{Z}$ having in mind the agreement introduced in Subsection 3.1. For technical reasons we divide the system $H(m)$ into two parts:

$$(44) \quad H^+(m) = \{h_{k,j,m}^{(v)}(x) : k \in \mathbb{Z}; j \geq 0; 1 \leq v \leq m - 1\},$$

$$(45) \quad H^-(m) = \{h_{k,j,m}^{(v)}(x) : k \in \mathbb{Z}; j \leq -1; 1 \leq v \leq m - 1\}.$$

We are going to show that the systems $H^+(m), H^-(m)$ are unconditional bases respectively in the spaces $L^p(\mathbb{R}^+)$ and $L^p(\mathbb{R}^-), 1 < p < \infty$. Let us prove the following

THEOREM 4.11. *For any $m = 2, 3, \dots$ the system $H^+(m)$ is an unconditional basis in any space $L^p(\mathbb{R}^+), 1 < p < \infty$.*

PROOF. Let $f \in L^p(\mathbb{R}^+)$ and let $\Omega \subset \mathbb{Z} \times \mathbb{Z}^+$ be a finite set. For any $1 \leq v \leq m - 1$ consider the sum

$$S_\Omega^{(v)}(f, x) = \sum_{(k,j) \in \Omega} c_{kj}^{(v)}(f) h_{k,j,m}^{(v)}(x),$$

where

$$c_{kj}^{(v)}(f) = \int_{\mathbb{R}^+} f(t) h_{k,j,m}^{(v)}(t) dt.$$

Let $N \in \mathbb{N}$ be such that for all $(k, j) \in \Omega$ $h_{k,j,m}^{(v)}(x) = 0$ if $x \in [m^N, +\infty)$. Consider the dilation operator $D_N(\phi)(x) = m^{\frac{N}{2}} \phi(m^N x)$. It is clear that $D_N(h_{k,j,m}^{(v)}) \in \mathcal{H}(m)$ for any $h_{k,j,m}^{(v)}$ which satisfies to the above conditions if we consider the restriction on $[0, 1]$ of the image of the operator. Thus

$$D_N(S_\Omega^{(v)}(f, \cdot))(x) = \sum_{(k,j) \in \Omega} c_{kj}^{(v)}(f) D_N(h_{k,j,m}^{(v)})(x)$$

on $[0, 1]$ is a finite linear combination of elements from $\mathcal{H}(m)$. We also have that if $(k, j) \in \Omega$

$$c_{kj}^{(v)}(f) = \int_{[0, m^N]} f(t)h_{k,j,m}^{(v)}(t)dt = \int_{[0,1]} D_N(f)(t)D_N(h_{k,j,m}^{(v)})(t)dt.$$

Hence, $D_N(S_\Omega^{(v)}(f, \cdot))(x)$ on $[0, 1]$ coincides with the sum of a subsequence of the expansion of the function $D_N(f)$ with respect to the system $\mathcal{H}(m)$. By Theorem 4.10 we obtain that there exists $C_p > 0$ which depends only on p such that

$$\int_{[0,1]} |D_N(S_\Omega^{(v)}(f, \cdot))(t)|^p dt \leq C_p^p \int_{[0,1]} |D_N(f)(t)|^p dt$$

which yields

$$\|S_\Omega^{(v)}(f, \cdot)\|_{L^p(\mathbb{R}^+)} \leq C_p \|f\|_{L^p(\mathbb{R}^+)}.$$

□

It is clear that in a similar way we can check that $H^-(m)$ is an unconditional basis in any space $L^p(\mathbb{R}^-)$, $1 < p < \infty$. Thus we the proof of Theorem 4.10 is finished.

4.2. m -th rank Haar system in $L^p([0, 1], w)$, $1 \leq p < \infty$.

THEOREM 4.12. *For any $m = 2, 3, \dots$ the system $\mathcal{H}(m)$ is a basis in the weighted norm space $L^p([0, 1], w)$, $1 \leq p < \infty$ if and only if w satisfies the condition $\mathcal{M}_p([0, 1])$.*

PROOF. By Corollary 4.6 we easily obtain that the system $\mathcal{H}(m)$ is complete in $L^p([0, 1], w)$, $1 \leq p < \infty$. Suppose that w satisfies the condition $\mathcal{M}_p([0, 1])$. Then it is evident that

$$\frac{\chi_{[0,1]}}{w} \in L^{p'}([0, 1], w), \quad \text{where} \quad \frac{1}{p} + \frac{1}{p'} = 1.$$

Hence, the system $\mathcal{H}(m) = \{h_l(x)\}_{l=0}^\infty$ is minimal in $L^p([0, 1], w)$ and its conjugate system is the system $\mathcal{H}^*(m) = \{\frac{1}{w(x)}h_l(x)\}_{l=0}^\infty$. Thus for any $f \in L^p([0, 1], w)$, the coefficients of its expansion with respect to the system $\mathcal{H}(m)$ are equal to

$$b_l(f) = \int_{[0,1]} f(t)\frac{1}{w(t)}h_l(t)w(t)dt = \int_{[0,1]} f(t)h_l(t)dt = a_l(f).$$

Hence, for any $k \in \mathbb{N}$ and $1 \leq j \leq m^k$ the partial sums of the mentioned expansion with indices $\mu_k + j(m - 1)$ coincide with $\Theta_{\mu_k+j(m-1)}(f, x)$ (see the beginning of Section 4). By Lemma 4.1 it follows easily that

$$(46) \quad \|\Theta_{\mu_k+j(m-1)}(f, \cdot)\|_{L^p([0,1],w)} \leq C \|f\|_{L^p([0,1],w)},$$

where $C = C(w, p, m)$ is independent of f . If we prove that

$$(47) \quad \lim_{l \rightarrow +\infty} |b_l(f)| \|h_l\|_{L^p([0,1],w)} = 0$$

then it will follow that (46) holds for all $n \in \mathbb{N}$. Which yields that the system $\mathcal{H}(m)$ is a basis $L^p([0, 1], w)$. We have that $b_l(f) = \int_{[0,1]} [f(t) - P(t)]h_l(t)dt$ for any $P(t) = \sum_{k=0}^{l-1} d_k h_k(t)$. If $h_l(x) = h_{k,j,m}^{(v)}(x)$ for $x \in [0, 1]$ and $\Delta = [\frac{j}{m^k}, \frac{j+1}{m^k}]$ then

$$\begin{aligned} |b_l(f)| \|h_l\|_{L^p([0,1],w)} &\leq \|f - P\|_{L^p([0,1],w)} \left\| \frac{h_{k,j,m}^{(v)}}{w^{\frac{1}{p}}} \right\|_{L^{p'}(\Delta)} \|h^{(v)}\|_{L^\infty([0,1])} m^{\frac{k}{2}} [w(\Delta)]^{\frac{1}{p}} \\ &\leq \|f - P\|_{L^p([0,1],w)} \|h^{(v)}\|_{L^\infty([0,1])}^2 m^k [w(\Delta)]^{\frac{1}{p}} \left[\int_{\Delta} w^{-\frac{1}{p-1}} \right]^{\frac{1}{p'}} \\ &\leq \|f - P\|_{L^p([0,1],w)} \|h^{(v)}\|_{L^\infty([0,1])}^2 C_p^{\frac{1}{p}}. \end{aligned}$$

The last inequality yields (47) because the system $\mathcal{H}(m)$ is complete in $L^p([0, 1], w)$. To prove the necessity suppose that the system $\mathcal{H}(m)$ is a basis in the weighted norm space $L^p([0, 1], w)$, where $1 \leq p < \infty$.

Let $\mathcal{H}^*(m) = \{h_l^*(x)\}_{l=0}^\infty$ be the conjugate system of the basis $\mathcal{H}(m)$. Then we have that

$$\int_{[0,1]} [h_0^*(t)w(t) - 1]h_l(t)dt = 0 \quad \text{for all } l \in \mathbb{N}_0.$$

Hence, $h_0^*(t) = \frac{1}{w(t)} \in L^{p'}([0, 1], w)$. Thus we obtain that

$$h_l^*(x) = \frac{h_l(x)}{w(x)} \quad \text{for all } l \in \mathbb{N}_0.$$

Thus for any $f \in L^p([0, 1], w)$ n -th partial sums of its expansion with respect to the basis $\mathcal{H}(m)$ coincide with $\Theta_n(f, x)$. By Lemma 4.1 it follows that for some $C \geq 1$ such that for any $\Delta \in \mathcal{M}$

$$\sup \frac{1}{|\Delta|^p} \left| \int_{\Delta} f(t)dt \right|^p w(\Delta) \leq C^p,$$

where the supremum is taken over all $\|f\|_{L^p([0,1],w)} \leq 1$. The last inequality easily yields (18) with $C_p = C^p$. □

The prove of the following result technically is much more complicated. The main line of our proof is close to the one given in [16](see also [7] and [28], [10]).

THEOREM 4.13. *For any $m = 2, 3, \dots$ the system $\mathcal{H}(m)$ is an unconditional basis in the weighted norm space $L^p([0, 1], w)$, $1 < p < \infty$ if and only if w satisfies the condition $\mathcal{M}_p([0, 1])$.*

LEMMA 4.14. *Let w be a weight function which satisfies the condition $\mathcal{M}_\infty([0, 1])$. Then for any $\lambda > 0$, any $0 < \gamma < 1$ and for any $f \in L^1[0, 1]$*

$$\begin{aligned} w(\{x \in [0, 1] : G_m(f, x) > 2\lambda \text{ and } M_{\mathcal{M}}f(x) \leq \gamma\lambda\}) \\ \leq C\gamma^\delta w(\{x \in [0, 1] : G_m(f, x) > \lambda\}), \end{aligned}$$

where $C > 0$ is independent of f , $\lambda > 0$ and $\gamma > 0$, while $\delta > 0$ is the corresponding constant from Definition 3.8.

In the formulation of the following assertion we use the agreement formulated in Subsection 3.1.

LEMMA 4.15. *For any $f \in L^1[0, 1]$ and any $\lambda > 0$ there exists a finite or denumerable set of disjoint closed intervals $\{\Delta_k\}_{k \in \Upsilon}$ such that the set*

$$(48) \quad E_\lambda(f) = \{x \in [0, 1] : G_m(f, x) > \lambda\} = \bigcup_{k \in \Upsilon} \Delta_k .$$

PROOF. According to our agreement for any $x \in [0, 1]$ there exists a unique sequence of closed intervals $\Gamma_k(x) \subset \mathcal{M}$ such that $\Gamma_k(x) \subset \Gamma_{k-1}(x)$ for all $k \in \mathbb{N}$ and $|\Gamma_k(x)| = m^{-k}$, $\bigcap_{k=0}^\infty \Gamma_k(x) = x$. For any $x_0 \in E_\lambda(f)$ there exists $k(x_0) \in \mathbb{N}_0$ so that $\Gamma_{k(x_0)}(x_0) \subseteq E_\lambda(f)$ and $\Gamma_{k(x_0)-1}(x_0)$ at least contains a point which does not belong to $E_\lambda(f)$. Indeed, if $G_m(f, x_0) > \lambda$ then there exists $N \in \mathbb{N}$ such that

$$[G_m(f, x_0)]^2 \geq \sum_{l=0}^N |a_l(f)h_l(x_0)|^2 > \lambda^2 .$$

Hence, for some $\nu \in \mathbb{N}$ the sum $\sum_{l=0}^N |a_l(f)h_l(x_0)|^2$ is constant on $\Gamma_\nu(x_0)$. Which means that $\Gamma_\nu(x_0) \subseteq E_\lambda(f)$. The number $k(x_0)$ will be the smallest index for which the last relation holds. Afterwards, one observes that $\max_{x \in E_\lambda(f)} |\Gamma_{k(x)}(x)| := \mu_0$ exists. There exist only finitely many disjoint intervals in the set $\{\Gamma_{k(x)}(x) : x \in E_\lambda(f)\}$ with length equal to μ_0 . Let Δ_j ($1 \leq j \leq n_1$) be all such intervals. Let $E_2 = E_\lambda(f) \setminus (\bigcup_{j=1}^{n_1} \Delta_j)$ and repeat the same procedure taking E_2 instead of $E_\lambda(f)$. Thus step by step we construct the finite or denumerable set of disjoint closed intervals $\{\Delta_k\}_{k \in \Upsilon}$ which satisfy the conditions of lemma. □

PROOF OF LEMMA 4.14. Let Δ_l be an arbitrary closed interval from (48). At the first step we have to prove the following relation

$$(49) \quad \left| \left\{ x \in \Delta_l : G_m(f, x) > 2\lambda \quad \text{and} \quad M_{\mathcal{M}}f(x) \leq \gamma\lambda \right\} \right| \leq C\gamma|\Delta_l| ,$$

where $C > 0$ is independent of f, λ, γ and Δ_l . Suppose that there is at least a point $y_l \in \Delta_l$ such that $M_{\mathcal{M}}f(y_l) \leq \gamma\lambda$. Otherwise there is nothing to prove. Let $\Delta_l^* \in \mathcal{M}$ be the interval which satisfies the following conditions: $\Delta_l^* \supset \Delta_l$, $|\Delta_l^*| = m|\Delta_l|$. Let $f(x) = f_1(x) + f_2(x)$, where

$$f_1(x) = (f(x) - f_{\Delta_l^*})\chi_{\Delta_l^*}(x), \quad f_{\Delta} = \frac{1}{|\Delta|} \int_{\Delta} f(t)dt, \quad \Delta \in \mathcal{M}$$

and $f_2(x) = f(x) - f_1(x)$. By Proposition 4.9 we have that

$$(50) \quad \left| \left\{ G_m(f_1, x) > \frac{\lambda}{2} \right\} \right| \leq \frac{2C_1}{\lambda} \|f_1\|_{L^1[0,1]} = \frac{4C_1}{\lambda} \int_{\Delta_l^*} |f(t)|dt \\ \leq \frac{4mC_1}{\lambda} |\Delta_l| M_{\mathcal{M}}f(y_l) \leq 4mC_1\gamma|\Delta_l| .$$

On the other hand we have that $|\Delta_l^*| = m^{-\kappa}$ for some $\kappa \in \mathbb{N}_0$. Thus for all $0 \leq l \leq \mu_\kappa$ we have that $a_l(f) = a_l(f_2)$, which yields $G_m(f_2, x) = G_m(\Theta_{\mu_\kappa}(f, \cdot), x)$ if $x \in \Delta_l^*$. There exists at least one point $z_l \in \Delta_l^*$ such that $G_m(f, z_l) \leq \lambda$. Thus if $x \in \Delta_l^*$ then

$$G_m(f_2, x) = G_m(\Theta_{\mu_\kappa}(f, \cdot), x) \leq G_m(f, z_l) \leq \lambda.$$

We have that if $x \in \Delta_l^*$ then

$$G_m(f, x) \leq G_m(f_1, x) + G_m(f_2, x) \leq G_m(f_1, x) + \lambda.$$

Hence, by (50) we finish the proof of (49).

We have that the weight function w satisfies the condition $\mathcal{M}_\infty([0, 1])$. Thus we obtain that

$$w(\{x \in \Delta_l : G_m(f, x) > 2\lambda \text{ and } M_{\mathcal{M}}f(x) \leq \gamma\lambda\}) \leq C\gamma^\delta w(\Delta_l)$$

where $C > 0$ is independent of $f, \lambda > 0, \gamma > 0$ and Δ_l . Hence, by Lemma 48 we finish the proof. □

PROOF OF THEOREM 4.13. The necessity follows from Theorem 4.12. Suppose that w satisfies the condition $\mathcal{M}_p([0, 1])$. By Lemma 4.14 we derive

$$\begin{aligned} \int_{[0,1]} G_m^p(f, x)w(x)dx &= p2^p \int_0^{+\infty} \lambda^{p-1}w(\{x \in [0, 1] : G_m(f, x) > 2\lambda\})d\lambda \\ &\leq K_p \int_0^{+\infty} \lambda^{p-1}w(\{x \in [0, 1] : M_{\mathcal{M}}f(x) > \gamma\lambda\})d\lambda \\ &\quad + K_p C \gamma^\delta \int_0^{+\infty} \lambda^{p-1}w(\{x \in [0, 1] : G_m(f, x) > \lambda\})d\lambda. \end{aligned}$$

Let $\gamma_0 > 0$ be such that $K_p C \gamma_0^\delta < \frac{1}{2}$. Then we obtain that

$$\int_{[0,1]} G_m^p(f, x)w(x)dx \leq 2K_p \gamma_0^{-p} \int_{[0,1]} M_{\mathcal{M}}^p f(x)w(x)dx.$$

By Proposition 3.6 we finish the proof. □

5. The system $\mathcal{H}_0(m) = \{h_l(x)\}_{l=1}^\infty$ in $L^p([0, 1], w)$. In this section we will use the following result (see [26]–[30])

THEOREM 5.1. *Let $\{f_n\}_{n=1}^\infty \subseteq L^\infty(E)$ be an orthonormal system of real-valued functions defined on a measurable set $E, 0 < |E| < +\infty$ and suppose that $\{f_n\}_{n=1}^\infty$ is total with respect to $L^1(E)$. Let, furthermore, $N \in \mathbb{N}$ and $w \in L^1(E)$ be a weight function. For the system $\{f_n\}_{n=N+1}^\infty$ to be closed and/or minimal it is necessary and sufficient that the following conditions 1) and/or 2), respectively, are satisfied:*

1) *any function of the form $(w)^{-1} \sum_{n=1}^N c_n f_n$, where $c_n (1 \leq n \leq N)$ are real numbers, belongs to $L^{p'}(E, w)$ if and only if every c_n is zero;*

2) for every k ($k = N + 1, N + 2, \dots$) there exist uniquely determined real numbers $b_n^{(k)}$ ($1 \leq n \leq N$) such that the function

$$g_k = \frac{1}{w} \left[\sum_{n=1}^N b_n^{(k)} f_n + f_k \right]$$

belongs to $L^{p'}(E, w)$ ($\frac{1}{p} + \frac{1}{p'} = 1$).

The following two lemmas are easy consequences of Theorem 5.1. We skip the details of the proofs because they are similar to the case of the Haar system [31].

LEMMA 5.2. For any $m = 2, 3, \dots$ the system $\mathcal{H}_0(m)$ is complete in a weighted norm space $L^p([0, 1], w)$, $1 \leq p < \infty$ if and only if there exists at least one point $y \in [0, 1]$ such that

$$(51) \quad \frac{1}{w} \notin L^{\frac{1}{p-1}}(\Delta_j(y)) \quad \text{for all } j \in \mathbb{N}.$$

LEMMA 5.3. For any $m = 2, 3, \dots$ the system $\mathcal{H}_0(m)$ is minimal in a weighted norm space $L^p([0, 1], w)$, $1 \leq p < \infty$ if and only if $\frac{1}{w} \in L^1([0, 1])$ or for a point $y \in [0, 1]$

$$(52) \quad \frac{1}{w} \in L^{\frac{1}{p-1}}([0, 1] \setminus \Delta_j(y)) \quad \text{for all } j \in \mathbb{N}.$$

By Lemmas 5.2 and 5.3 it follows easily

LEMMA 5.4. For any $m = 2, 3, \dots$ the system $\mathcal{H}_0(m)$ is complete and minimal in a weighted norm space $L^p([0, 1], w)$, $1 \leq p < \infty$ if and only if there exists only one point $y \in [0, 1]$ such that the conditions (51), (52) hold.

Further in this section we will suppose that the weight function w satisfies the conditions (51) and (52). Hence, the system $\mathcal{H}_0(m)$ is complete and minimal in the weighted norm space $L^p([0, 1], w)$, $1 \leq p < \infty$ with the unique conjugate system $\mathcal{H}_0^*(m)$. By Theorem 5.1 applied for our case it is easy to see that the system $\mathcal{H}_0^*(m) = \{h_l^*(x)\}_{l=1}^\infty$ is defined by the following equations:

$$(53) \quad h_l^*(x) = \frac{h_l(x) - h_l(y)}{w(x)}.$$

For any $f \in L^p([0, 1], w)$ and for any $1 \leq j \leq m^k$, $k \in \mathbb{N}$ we put

$$(54) \quad \Theta_{\mu_k + j(m-1)}^{(0)}(f, x) = \sum_{l=1}^{\mu_k} c_l(f) h_l(x) + \sum_{s=0}^{j-1} \sum_{v=1}^{m-1} c_{k,s,m}^{(v)}(f) h_{k,s,m}^{(v)}(x),$$

where

$$c_l(f) = \int_{[0,1]} f(t) h_l^*(t) dt; \quad c_{k,s,m}^{(v)}(f) = \int_{[0,1]} f(t) [h_{k,s,m}^{(v)}(t) - h_{k,s,m}^{(v)}(y)] dt.$$

Let $\Delta_{kj}(y)$ be the interval from the collection of sets (35), (36) such that $y \in \Delta_{kj}(y)$.

LEMMA 5.5. For any $f \in L^p([0, 1], w)$ and for any $1 \leq j \leq m^k, k \in \mathbb{N}$ we have that $\Theta_{\mu_k+j(m-1)}^{(0)}(f, x)$ is constant on any interval from the collection of sets (35), (36). Moreover,

$$(55) \quad \Theta_{\mu_k+j(m-1)}^{(0)}(f, x) = -\frac{1}{|\Delta_{kj}(y)|} \int_{[0,1] \setminus \Delta_{kj}(y)} f(t)dt \quad \text{for } x \in \Delta_{kj}(y),$$

and for any Δ from (35) or from (36) which does not coincide with $\Delta_{kj}(y)$

$$(56) \quad \Theta_{\mu_k+j(m-1)}^{(0)}(f, x) = \frac{1}{|\Delta|} \int_{\Delta} f(t)dt \quad \text{for } x \in \Delta.$$

PROOF. In the proof we use the notation of Lemma 4.4. Let Δ be any interval from the collection of sets (35), (36) such that $\Delta \cap \Delta_{kj}(y) = \emptyset$. Then we have that

$$\begin{aligned} \Theta_{\mu_k+j(m-1)}^{(0)}(f, x) &= \int_{[0,1]} f(t)[\tilde{K}_{kj}(t, x) - \tilde{K}_{kj}(y, x)]dt \\ &= \sum_{i=1}^{m^k+j(m-1)} \int_{G_i} f(t)[K_{kj}(t, x) - K_{kj}(y, x)]dt, \end{aligned}$$

where $\tilde{K}_{kj}(t, x) = K_{kj}(t, x) - 1$, hence, $\tilde{K}_{kj}(t, x) - \tilde{K}_{kj}(y, x) = K_{kj}(t, x) - K_{kj}(y, x)$. Suppose that $G_v = \Delta_{kj}(y)$ and take any $i_0 \neq v, 1 \leq i_0 \leq m^k + j(m - 1)$. Then by Lemma 4.4 we obtain that for $x \in G_{i_0}$ we obtain that

$$\begin{aligned} \sum_{i=1}^{m^k+j(m-1)} \int_{G_i} f(t)[K_{kj}(t, x) - K_{kj}(y, x)]dt &= \int_{G_{i_0}} f(t)K_{kj}(t, x)dt \\ &= \frac{1}{|G_{i_0}|} \int_{G_{i_0}} f(t)dt. \end{aligned}$$

On the other hand if $x \in G_v$ by Lemma 4.4 we will have that

$$\begin{aligned} \Theta_{\mu_k+j(m-1)}^{(0)}(f, x) &= \sum_{i=1, i \neq v}^{m^k+j(m-1)} \int_{G_i} f(t)[K_{kj}(t, x) - K_{kj}(y, x)]dt \\ &= -\frac{1}{|G_v|} \sum_{i=1, i \neq v}^{m^k+j(m-1)} \int_{G_i} f(t)dt = -\frac{1}{|G_v|} \int_{[0,1] \setminus G_v} f(t)dt. \end{aligned}$$

□

Lemma 3.14 and Lemma 5.4 easily yield

LEMMA 5.6. Let $w \geq 0$ be a weight function defined on $[0, 1]$ such that w satisfies the condition $\mathcal{M}_p^y([0, 1])$ for some $y \in [0, 1]$ and $1 < p < \infty$. Then the conditions (51), (52) hold and the system $\mathcal{H}_0(m)$ is complete and minimal in a weighted norm space $L^p([0, 1], w)$.

THEOREM 5.7. For any $m = 2, 3, \dots$ the system $\mathcal{H}_0(m)$ is a basis in the weighted norm space $L^p([0, 1], w), 1 < p < \infty$ if and only if there exists a point $y \in [0, 1]$ such that w satisfies the conditions $\mathcal{M}_p([0, 1] \setminus \{y\})$ and $\mathcal{M}_p^y([0, 1])$.

PROOF. Necessity. If the system $\mathcal{H}_0(m)$ is a basis in the weighted norm space $L^p([0, 1], w)$ then it is a complete minimal system in $L^p([0, 1], w)$. Then we will have that for any $f \in L^p([0, 1], w)$ and for any $1 \leq j \leq m^k, k \in \mathbb{N}$ the partial sum operators are uniformly bounded

$$\sup_{1 \leq j \leq m^k, k \in \mathbb{N}} \|\Theta_{\mu_k+j(m-1)}^{(0)}\|_{L^p([0,1],w) \rightarrow L^p([0,1],w)} := M_0 \leq B_p,$$

where $B_p > 0$. We have that

$$M_0 \geq \max_{1 \leq i \leq m^k+j(m-1)} \sup_{\|f\|_{L^p(G_i,w)} \leq 1} \|\Theta_{\mu_k+j(m-1)}^{(0)}\|_{L^p([0,1],w)}.$$

If $G_i \cap \Delta_{kj}(y) = \emptyset$ then by Lemma 5.5 we will have that

$$\begin{aligned} \sup_{\|f\|_{L^p(G_i,w)} \leq 1} \|\Theta_{\mu_k+j(m-1)}^{(0)}\|_{L^p(G_i,w)} &= \frac{1}{|G_i|} \sup_{\|f\|_{L^p(G_i,w)} \leq 1} \left| \int_{G_i} f(t) dt \right| w(G_i)^{\frac{1}{p}} \\ &= |G_i|^{-1} w(G_i)^{\frac{1}{p}} \|w^{-\frac{1}{p}}\|_{L^{p'}(G_i)}. \end{aligned}$$

If $G_v = \Delta_{kj}(y)$ then in the same way as above we obtain that

$$|\Delta_{kj}(y)|^{-p} w(\Delta_{kj}(y)) \left(\int_{\Delta_{kj}(y)} w^{-\frac{1}{p-1}} dt \right)^{p-1} \leq B_p^p.$$

Sufficiency. By Lemma 5.6 we have that the system $\mathcal{H}_0(m)$ is complete and minimal in a weighted norm space $L^p([0, 1], w)$. Hence, by Lemma 5.5 we obtain that for any $f \in L^p([0, 1], w)$ and for any $1 \leq j \leq m^k, k \in \mathbb{N}$

$$\begin{aligned} \int_{[0,1]} |\Theta_{\mu_k+j(m-1)}^{(0)}(f, t)|^p w(t) dt &= \sum_{\substack{1 \leq i \leq m^k+j(m-1) \\ i \neq v}} \left| \frac{1}{|G_i|} \int_{G_i} f(t) dt \right|^p \int_{G_i} w(t) dt \\ &+ \left| \frac{1}{|G_v|} \int_{[0,1] \setminus G_v} f(t) dt \right|^p \int_{G_v} w(t) dt \leq B_p^p \int_{[0,1]} |f(t)|^p w(t) dt. \end{aligned}$$

The last inequality follows because w satisfies the conditions $\mathcal{M}_p([0, 1] \setminus \{y\})$ and $\mathcal{M}_p^y([0, 1])$. To finish the proof we have to show that $\lim_{l \rightarrow \infty} |c_l(f)| \|\mathbf{h}_l(\cdot)\|_{L^p([0,1],w)} = 0$. We skip the details because a similar result we have proved for the proof of Theorem 4.12. \square

In the case $p = 1$ we have the following result.

THEOREM 5.8. *For any $m = 2, 3, \dots$ the system $\mathcal{H}_0(m)$ is a basis in the weighted norm space $L^1([0, 1], w)$, if and only if $\frac{1}{w} \notin L^\infty([0, 1])$ and there exists a point $y \in [0, 1]$ such that w satisfies the conditions $\mathcal{M}_1([0, 1] \setminus \{y\})$ and $\mathcal{M}_1^y([0, 1])$.*

We will not give the details of the proof because it is similar to the proof of Theorem 5.7. The main theorem of this section is the following

THEOREM 5.9. *If the system $\mathcal{H}_0(m), m = 2, 3, \dots$ is a basis in the weighted norm space $L^p([0, 1], w), 1 < p < \infty$ then $\mathcal{H}_0(m)$ is an unconditional basis in the same space.*

PROOF. By Theorem 5.7 we have that there exists $y \in [0, 1]$ such that the weight function w satisfies the conditions $\mathcal{M}_p([0, 1] \setminus \{y\})$ and $\mathcal{M}_p^y([0, 1])$. For any $f \in L^p([0, 1], w)$ there exists a unique sequence $\{a_l(f)\}_{l=1}^\infty$ such that

$$(57) \quad f = \sum_{l=1}^\infty a_l(f)h_l.$$

The coefficients which correspond to the functions $h_{\Delta_j(y)}^{(v)}$ in the series (57) we denote by b_{jv} . We split formally the series (57) into two parts

$$(58) \quad \sum_{l=1}^\infty a_l(f)h_l = \sum_{j=0}^\infty \sum_{v=1}^{m-1} b_{jv}(f)h_{\Delta_j(y)}^{(v)} + \sum' a_l(f)h_l,$$

where by \sum' we have denoted the series obtained after excluding the terms which are present in the first series.

For any $k \in \mathbb{N}_0$ let $G_{kl} \subset \mathcal{M}$, $1 \leq l \leq m - 1$ be mutually disjoint intervals such that $|G_{kl}| = m^{-k-1}$, $1 \leq l \leq m - 1$ and $\Delta_k(y) = \Delta_{k+1}(y) \cup \bigcup_{l=1}^{m-1} G_{kl}$.

By Theorem 4.13 we easily obtain that the series $\sum' a_l(f)h_l$ converges unconditionally in $L^p(G_{kl}, w)$ for any $k \in \mathbb{N}$ and for all $1 \leq l \leq m - 1$. Hence, if we check that the series $\sum' a_l(f)h_l$ converges in $L^p([0, 1], w)$ we will have that it converges unconditionally in $L^p([0, 1], w)$. Thus the proof of theorem will be finished if we prove that the first series on the right hand side of the equality (58) converges unconditionally in $L^p([0, 1], w)$.

Recall that we are using the notation introduced in (28). Let

$$(59) \quad F(x) = \sum_{j=0}^\infty \sum_{v=1}^{m-1} b_{jv}(f)h_{\Delta_j(y)}^{(v)}(x) = d_k^l \quad \text{if } x \in G_{kl}$$

for all $k \in \mathbb{N}_0$ and $1 \leq l \leq m - 1$. By Lemma 5.5 we have that

$$d_k^l = \frac{1}{|G_{kl}|} \int_{G_{kl}} f(t)dt, \quad \text{for } x \in G_{kl}.$$

The weight function w satisfies the condition $\mathcal{M}_p([0, 1] \setminus \{y\})$. Hence,

$$\begin{aligned} \int_{[0,1]} |F(x)|^p w(x)dx &= \sum_{k=0}^\infty \sum_{l=1}^{m-1} |d_k^l|^p w(G_{kl}) \\ &\leq \sum_{k=0}^\infty \sum_{l=1}^{m-1} |G_{kl}|^{-p} \int_{G_{kl}} |f(t)|^p w(t)dt \left(\int_{G_{kl}} w(t)^{-\frac{1}{p-1}} dt \right)^{p-1} w(G_{kl}) \\ &\leq C_p \int_{[0,1]} |f(t)|^p w(t)dt. \end{aligned}$$

The system $\mathcal{H}_0(m)$ is a basis in the weighted norm space $L^p([0, 1], w)$. Hence, the first series in the right hand side of the equality (58) converges in $L^p([0, 1], w)$. Thus to finish the

proof of Theorem 5.9 we have to prove that the series in (59) converges unconditionally in $L^p([0, 1], w)$.

For any $j \in \mathbb{N}_0$ we have that

$$\sum_{v=1}^{m-1} b_{jv}(f)h_{\Delta_j(y)}^{(v)}(x) = d_j^l = \frac{1}{|G_{jl}|} \int_{G_{jl}} f(t)dt,$$

for $x \in G_{jl}, 1 \leq l \leq m - 1$ and

$$\sum_{v=1}^{m-1} b_{jv}(f)h_{\Delta_j(y)}^{(v)}(x) := -c_j = -\sum_{l=1}^{m-1} d_j^l, \quad \text{for } x \in \Delta_{j+1}(y).$$

Let $\{\gamma_l\}_{l=0}^{m-1}$ be a collection of numbers such that

$$(60) \quad \sum_{l=0}^{m-1} \gamma_l^2 = 1 \quad \text{and} \quad \sum_{l=0}^{m-1} \gamma_l = 0.$$

We put

$$(61) \quad \xi_j(x) = |\Delta_{j+1}(y)|^{-\frac{1}{2}} \left[\gamma_0 \chi_{\Delta_{j+1}(y)}(x) + \sum_{l=1}^{m-1} \gamma_l \chi_{G_{jl}}(x) \right]$$

and

$$(62) \quad \begin{aligned} \alpha_j(f) &= \int_{[0,1]} f(t)[\xi_j(t) - |\Delta_{j+1}(y)|^{-\frac{1}{2}}\gamma_0]dt \\ &= -\frac{\gamma_0}{\sqrt{|\Delta_{j+1}(y)|}} \int_{[0,1] \setminus \Delta_{j+1}(y)} f(t)dt + \int_{[0,1] \setminus \Delta_{j+1}(y)} f(t)\xi_j(t)dt \\ &= |\Delta_{j+1}(y)|^{\frac{1}{2}} \sum_{l=1}^{m-1} (\gamma_l - \gamma_0)d_j^l - \frac{\gamma_0}{\sqrt{|\Delta_{j+1}(y)|}} \int_{[0,1] \setminus \Delta_j(y)} f(t)dt \\ &= |\Delta_{j+1}(y)|^{\frac{1}{2}} \left(\sum_{l=1}^{m-1} (\gamma_l - \gamma_0)d_j^l - \gamma_0 \sum_{s=0}^{j-1} m^{j-s-1}c_s \right). \end{aligned}$$

LEMMA 5.10. For any $\varepsilon = \{\varepsilon_j\}_{j=0}^\infty$ let

$$(63) \quad F_\varepsilon^*(x) = \sum_{j=0}^\infty \varepsilon_j \alpha_j \xi_j(x).$$

Then for all $k \in \mathbb{N}$ and $x \in \Delta_k(y) \setminus \Delta_{k+1}(y)$

$$|F_\varepsilon^*(x)| \leq 2 \sum_{s=0}^k \sum_{l=1}^{m-1} |d_s^l| + \frac{1}{m-1} \sum_{s=0}^{k-1} m^{k-s} |c_s|.$$

PROOF. By (61) and (62) we obtain that for $x \in G_{k\nu}$, $1 \leq \nu \leq m - 1$

$$\begin{aligned} |F_\varepsilon^*(x)| &\leq \sum_{j=0}^k |\alpha_j \xi_j(x)| = |\gamma_0| \sum_{s=0}^{k-1} \sum_{l=1}^{m-1} |\gamma_l - \gamma_0| d_s^l + |\gamma_\nu| \sum_{l=1}^{m-1} |\gamma_l - \gamma_0| d_k^l \\ &+ \sum_{j=1}^{k-1} \sum_{s=0}^{j-1} m^{j-s-1} |c_s| \leq 2 \sum_{s=0}^k \sum_{l=1}^{m-1} |d_s^l| + \sum_{j=1}^{k-1} \sum_{s=0}^{j-1} m^{j-s-1} |c_s| \\ &\leq 2 \sum_{s=0}^k \sum_{l=1}^{m-1} |d_s^l| + \frac{1}{m-1} \sum_{s=0}^{k-1} m^{k-s} |c_s|. \end{aligned}$$

□

LEMMA 5.11. For any $f \in L^p([0, 1], w)$, $1 < p < \infty$ and any $\varepsilon = \{\varepsilon_j\}_{j=0}^\infty$ the function $F_\varepsilon^* \in L^p([0, 1], w)$ and

$$\|F_\varepsilon^*\|_{L^p([0,1],w)} \leq C'_p \|f\|_{L^p([0,1],w)},$$

where $C'_p > 0$ is independent of f and ε .

PROOF. By Lemma 5.10 we have that

$$\begin{aligned} \int_{\Delta_k(y) \setminus \Delta_{k+1}(y)} |F^*(x)|^p w(t) dt &\leq 4^p \left(\sum_{s=0}^k \sum_{l=1}^{m-1} |d_s^l| \right)^p w(\Delta_k(y)) \\ &+ \frac{2^p}{(m-1)^p} \left([w(\Delta_k(y))]^{\frac{1}{p}} \sum_{s=0}^{k-1} m^{k-s} |c_s| \right)^p. \end{aligned}$$

Afterwards write

$$\begin{aligned} &|\Delta_{s+1}(y)| \sum_{l=1}^{m-1} |d_s^l| [w(\Delta_k(y))]^{\frac{1}{p}} \\ &\leq \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} |f(t)|^p w(t) dt \right)^{\frac{1}{p}} \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} w(t)^{-\frac{1}{p-1}} dt \right)^{\frac{1}{p'}} [w(\Delta_k(y))]^{\frac{1}{p}} \\ &\leq C_p^{\frac{1}{p}} \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} |f(t)|^p w(t) dt \right)^{\frac{1}{p}} |\Delta_k(y)|. \end{aligned}$$

Hence, we obtain that

$$\sum_{s=0}^k \sum_{l=1}^{m-1} |d_s^l| [w(\Delta_k(y))]^{\frac{1}{p}} \leq C_p^{\frac{1}{p}} \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} |f(t)|^p w(t) dt \right)^{\frac{1}{p}} \frac{1}{m^{k-s}}.$$

Now we apply the following lemma which is a consequence of Theorem 274 from [18].

LEMMA 5.12. *Let $u = \{u_j\}_{j=0}^\infty$ and $v = \{v_j\}_{j=0}^\infty$ be numerical sequences such that $u \in l^1$ and $v \in l^p, p > 1$. Then the Cauchy product $w = \{w_n\}_{n=0}^\infty, w_n = \sum_{j=0}^n u_{n-j}v_j$ of the sequences u and v belongs to l^p . Moreover $\|w\|_{l^p} \leq \|u\|_{l^1} \|v\|_{l^p}$.*

Which gives us the convergence of the series

$$\sum_{k=1}^\infty \left(\sum_{s=0}^k \sum_{l=1}^{m-1} |d_s^l| \right)^p w(\Delta_k(y)) \leq 2^p C_p \int_{[0,1]} |f(t)|^p w(t) dt .$$

To finish the proof of Lemma 5.11 we have to show that

$$(64) \quad \sum_{k=1}^\infty \left([w(\Delta_k(y))]^{\frac{1}{p}} \sum_{s=0}^{k-1} m^{k-s} |c_s| \right)^p < +\infty .$$

We have that

$$\begin{aligned} & \left(\sum_{s=0}^{k-1} m^{-s-1} |c_s| \right)^p \\ & \leq \left(\sum_{s=0}^{k-1} \sum_{l=1}^{m-1} \left(\int_{G_{sl}} |f(t)|^p w(t) dt \right)^{\frac{1}{p}} \left(\int_{G_{sl}} w(t)^{-\frac{1}{p-1}} dt \right)^{\frac{1}{p'}} \right)^p \\ & \leq \left(\sum_{s=0}^{k-1} \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} |f(t)|^p w(t) dt \right)^{\frac{1}{p}} \right. \\ & \quad \left. \times \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} w(t)^{-\frac{1}{p-1}} dt \right)^{\frac{1}{p'}} \right)^p . \end{aligned}$$

Recall that w satisfies $\mathcal{M}_p^y([0, 1])$. By Lemma 3.14 we obtain that

$$\begin{aligned} & m^k \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} w(t)^{-\frac{1}{p-1}} dt \right)^{\frac{1}{p'}} w(G_{kl})^{\frac{1}{p}} \\ & \leq C_p^{\frac{1}{p}} \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} w(t)^{-\frac{1}{p-1}} dt \right)^{\frac{1}{p'}} \left[\int_{[0,1] \setminus \Delta_k(y)} \omega^{-\frac{1}{p-1}}(t) dt \right]^{-\frac{1}{p'}} \\ & \leq C_p^{\frac{1}{p}} q_p^{-\frac{k-s}{p}} . \end{aligned}$$

If we write

$$\begin{aligned} & \sum_{s=0}^{k-1} m^{k-s-1} |c_s| w(G_{kl})^{\frac{1}{p}} \\ & \leq C_p^{\frac{1}{p}} \sum_{s=0}^{k-1} \left(\int_{\Delta_s(y) \setminus \Delta_{s+1}(y)} |f(t)|^p w(t) dt \right)^{\frac{1}{p}} q_p^{-\frac{k-s}{p}} \end{aligned}$$

and put $v_j = (\int_{\Delta_j(y) \setminus \Delta_{j+1}(y)} |f(t)|^p w(t) dt)^{\frac{1}{p}}$, $u_j = q_p^{-\frac{j}{p}}$ then by Lemma 5.12 we will obtain

$$\sum_{k=1}^{\infty} \left(\sum_{s=0}^{k-1} m^{k-s-1} |c_s| \right)^p w(G_{kl}) \leq C_p B_p \int_{[0,1]} |f(t)|^p w(t) dt .$$

□

Lemma 5.11 yields the convergence of the series

$$\sum_{j=0}^{\infty} \varepsilon_j^{(v)} b_{jv}(f) h_{\Delta_j(y)}^{(v)}$$

for any $1 \leq v \leq m - 1$ and for all $\varepsilon_j^{(v)} = \{\varepsilon_j^{(v)}\}_{j=0}^{\infty}$, where $\varepsilon_j^{(v)} = \pm 1$. Moreover, we obtain that for some $B_p > 0$

$$\left\| \sum_{j=0}^{\infty} \sum_{v=1}^{m-1} \varepsilon_j^{(v)} b_{jv}(f) h_{\Delta_j(y)}^{(v)} \right\|_{L^p([0,1],w)} \leq B_p \|f\|_{L^p([0,1],w)} .$$

□

6. Higher rank Haar wavelets in $L^p(\mathbb{R}, \omega)$. Let $\omega \geq 0$ be a locally integrable function defined on \mathbb{R} . In this section we study the phenomenon described in the introduction with respect to the higher rank Haar wavelet systems $H(m)$, $m = 2, 3, \dots$. Let $\chi^-(x) = \chi_{\mathbb{R}^-}(x)$ and $\chi^+(x) = \chi_{\mathbb{R}^+}(x)$. The following result is the first step in that direction.

LEMMA 6.1. *For any $m = 2, 3, \dots$ let $H(m)$ be the wavelet system defined by (26) and (27). Let U_m be the linear subspace of locally integrable functions ξ on \mathbb{R} such that*

$$(65) \quad \int_{\mathbb{R}} \xi(t) h_{k,j,m}^{(v)}(x)(t) dt = 0 \quad \forall j, k \in \mathbb{Z}, 1 \leq v \leq m - 1 .$$

Then $\dim U_m = 2$ and χ^-, χ^+ as vectors constitute a basis in U_m .

PROOF. It is clear that if we prove that a locally integrable function ξ such that $\xi(x) = 0$ if $x \in \mathbb{R}^-$ and holds (65) if and only if $\xi = c\chi^+$ for some $c \in \mathbb{R}$ then the proof will be finished. By Corollary 4.6 we have that the system $\mathcal{H}(m)$, $m = 2, 3, \dots$ is total with respect to $L^1[0, 1]$. Hence, by definition of the system $\mathcal{H}(m)$ and by (65) it follows that

$$\int_{[0,1]} \xi(t) h_l(t) dt = 0 \quad \text{for all } l \in \mathbb{N} .$$

Which yields that $\xi(x) = c h_0(x)$ for $x \in [0, 1]$. We finish the proof by induction. Suppose that for some $N \in \mathbb{N}$ it is true that if ξ is a locally integrable function such that $\xi(x) = 0$ if $x \in \mathbb{R}^-$ and (65) is true then $\xi(x) = c_0$ if $x \in [0, m^N]$, where $c_0 \in \mathbb{R}$. If ξ is a function which satisfies to all mentioned conditions then by definition of the system $H(m)$ it follows that the functions $\xi_\nu(x) = \xi(x - \nu m^N)$, $1 \leq \nu \leq m - 1$ also satisfy to those conditions. Thus by our supposition it follows that $\xi_\nu(x) = c_\nu$ if $x \in [0, m^N]$, where $c_\nu \in \mathbb{R}$. Hence, $\xi(x) =$

c_ν if $x \in [\nu m^N, (\nu + 1)m^N], 0 \leq \nu \leq m - 1$. Afterwards we observe that the functions $h^{(\nu)}(m^{N+1}x), 1 \leq \nu \leq m - 1$ belong to the system $H(m)$, which yields

$$\int_0^{m^{N+1}} \xi(x)h^{(\nu)}(m^{N+1}x)dx = 0 \quad \text{for all } 1 \leq \nu \leq m - 1.$$

After a change of the variable we have that

$$\int_0^1 \xi(m^{-N-1}t)h^{(\nu)}(t)dt = 0 \quad \text{for all } 1 \leq \nu \leq m - 1.$$

By definition of the functions $h^{(\nu)}, 1 \leq \nu \leq m - 1$ we obtain that $\xi(m^{-N-1}x) = c$ if $x \in [0, 1]$. □

It is convenient to continue our study considering the systems $H^+(m), H^-(m)$ respectively in the spaces $L^p(\mathbb{R}^+, \omega)$ and $L^p(\mathbb{R}^-, \omega)$. It is easy to see some sort of symmetry between those systems. Thus it would be sufficient to study the system $H^+(m)$ in the space $L^p(\mathbb{R}^+)$. In fact we have proved the analogue of the above lemma for the system $H^+(m)$ which is formulated as follows.

LEMMA 6.2. *Let U_m^+ be the linear subspace of locally integrable functions ξ on \mathbb{R}^+ such that*

$$(66) \quad \int_{\mathbb{R}^+} \xi(t)h_{k,j,m}^{(\nu)}(x)(t)dt = 0 \quad \forall k \in \mathbb{Z}, \forall j \in \mathbb{Z}^+, 1 \leq \nu \leq m - 1.$$

Then $\dim U_m^+ = 1$ and $\chi^+ \in U_m^+$.

We need the analogues of Lemmas 5.2, 5.3 for this case.

LEMMA 6.3. *The system $H^+(m)$ is complete in $L^p(\mathbb{R}^+, \omega), 1 \leq p < \infty$ if and only if*

$$(67) \quad \frac{\chi^+}{\omega} \notin L^{\frac{1}{p-1}}(\mathbb{R}^+).$$

PROOF. Suppose that $H^+(m)$ is complete in $L^p(\mathbb{R}^+, \omega)$. If $g = \frac{\chi^+}{\omega} \in L^{\frac{1}{p-1}}(\mathbb{R}^+)$ then $g \in L^{p'}(\mathbb{R}^+, \omega)$, where $\frac{1}{p} + \frac{1}{p'} = 1$. Thus

$$(68) \quad \int_{\mathbb{R}^+} g(t)h_{k,j,m}^{(\nu)}(x)(t)\omega(t)dt = 0 \quad \forall k \in \mathbb{Z}, \forall j \in \mathbb{Z}^+, 1 \leq \nu \leq m - 1.$$

Which yields that $H^+(m)$ is not complete in $L^p(\mathbb{R}^+, \omega)$. Which is a contradiction.

Suppose that $\frac{\chi^+}{\omega} \notin L^{\frac{1}{p-1}}(\mathbb{R}^+)$. If $H^+(m)$ is not complete in $L^p(\mathbb{R}^+, \omega)$ then there exists $g \in L^{p'}(\mathbb{R}^+, \omega)$ such that (68) holds. By Lemma 6.3 it follows that $g(t)\omega(t) = c\chi^+(t)$ a.e. on \mathbb{R}^+ , where $c \in \mathbb{R}$. We came to a contradiction which finishes the proof. □

From Lemma 6.3 it follows that

LEMMA 6.4. *The system $H^+(m)$ is complete in $L^p(\mathbb{R}^+, \omega)$, $1 \leq p < \infty$ if and only if there exists at least one point $y \in [0, +\infty]$ such that*

$$(69) \quad \frac{1}{w} \notin L^{\frac{1}{p-1}}(\Delta_j(y)) \quad \text{for all } j \in \mathbb{N}.$$

We also have

LEMMA 6.5. *The system $H^+(m)$ is minimal in $L^p(\mathbb{R}^+, \omega)$, $1 \leq p < \infty$ if and only if*

For any $h_{k,j,m}^{(v)}(x) \in H^+(m)$ there exists a coefficient $a_{k,j,m}^{(v)}$ such that

$$(M) \quad g_{k,j,m}^{(v)} = \frac{a_{k,j,m}^{(v)}\chi^+ + h_{k,j,m}^{(v)}}{\omega} \in L^{\frac{1}{p-1}}(\mathbb{R}^+).$$

PROOF. Suppose that the system $H^+(m)$ is minimal in $L^p(\mathbb{R}^+, \omega)$. Then there exists a system $\{g_{k,j,m}^{(v)} : k \in \mathbb{Z}, j \in \mathbb{Z}^+, 1 \leq v \leq m-1\}$ byorthogonal to $H^+(m)$. Hence, if for some $v_0, 1 \leq v_0 \leq m-1$ we fix any $l \in \mathbb{Z}$ and any $\mu \in \mathbb{Z}^+$ then for all $k \in \mathbb{Z}, j \in \mathbb{Z}^+$ and $1 \leq v \leq m-1$

$$(70) \quad \int_{\mathbb{R}^+} [g_{l,\mu,m}^{(v_0)}(x)\omega(x) - h_{l,\mu,m}^{(v_0)}(x)]h_{k,j,m}^{(v)}(x)dx = 0.$$

By Lemma 6.2 we obtain that

$$g_{l,\mu,m}^{(v_0)} = \frac{a_{l,\mu,m}^{(v_0)}\chi^+ + h_{l,\mu,m}^{(v_0)}}{\omega} \in L^{\frac{1}{p-1}}(\mathbb{R}^+).$$

The proof of sufficiency is direct. We easily check that the system $\{g_{k,j,m}^{(v)} : k \in \mathbb{Z}, j \in \mathbb{Z}^+, 1 \leq v \leq m-1\}$ is biorthogonal to $H^+(m)$. □

From Lemma 6.5 it follows easily that

LEMMA 6.6. *The system $H^+(m)$ is minimal in $L^p(\mathbb{R}^+, \omega)$, $1 \leq p < \infty$ if and only if there exists at most one point $y \in [0, +\infty]$ such that (69) holds.*

By Lemmas 6.3 and 6.5 we obtain immediately

LEMMA 6.7. *The system $H^+(m)$ is complete and minimal in $L^p(\mathbb{R}^+, \omega)$, $1 \leq p < \infty$ if and only if conditions (67) and (M) hold.*

Lemmas 6.4 and 6.6 yield

LEMMA 6.8. *The system $H^+(m)$ is complete and minimal in $L^p(\mathbb{R}^+, \omega)$, $1 \leq p < \infty$ if and only if there exists a unique point $y \in [0, +\infty]$ such that the condition (69) holds.*

If we analyze the proofs of results which brought us the last lemma then it is not hard to see that the following result also holds.

LEMMA 6.9. *The system $H^-(m)$ is complete minimal in $L^p(\mathbb{R}^-, \omega)$, $1 \leq p < \infty$ if and only if there exists a unique point $y \in [-\infty, 0]$ such that the condition (69) holds.*

Lemma 6.8 and Lemma 6.9 easily yield

LEMMA 6.10. *The system $H(m)$ is complete and minimal in $L^p(\mathbb{R}, \omega)$, $1 \leq p < \infty$ if and only if there exists a unique point $y^+ \in [0, +\infty]$ and a unique point $y^- \in [-\infty, 0]$ such that the condition (69) holds for both of those points.*

The following lemma will be used in the proof of the main result of the present section.

LEMMA 6.11. *Let $\{\phi_j\}_{j=1}^\mu, \{\psi_j\}_{j=1}^\mu$ be some measurable functions defined on a measurable set E and let*

$$K(x, t) = \sum_{j=1}^\mu \phi_j(x)\psi_j(t) \quad (x, t) \in E \times E.$$

Furthermore, for a given real valued orthogonal matrix

$$A = \left(a_{ij} \right)_{\substack{1 \leq i \leq \mu \\ 1 \leq j \leq \mu}}$$

let $f_k(x) = \sum_{i=1}^\mu a_{ik}\phi_i(x)$, $g_k(x) = \sum_{i=1}^\mu a_{ik}\psi_i(t)$. If we consider a new kernel $\Phi(x, t) = \sum_{k=1}^\mu f_k(x)g_k(t)$ then

$$K(x, t) = \Phi(x, t) \quad \text{for } (x, t) \in E \times E.$$

PROOF. We have

$$\begin{aligned} \Phi(x, t) &= \sum_{k=1}^\mu \sum_{i=1}^\mu a_{ik}\phi_i(x) \sum_{v=1}^\mu a_{vk}\psi_v(t) \\ &= \sum_{i=1}^\mu \sum_{v=1}^\mu \phi_i(x)\psi_v(t) \sum_{k=1}^\mu a_{vk}a_{ik} \\ &= \sum_{v=1}^\mu \phi_v(x)\psi_v(t) = K(x, t). \end{aligned}$$

□

We are going to apply Lemma 6.11 in the proof of the next theorem. As $\{\phi_j\}_{j=1}^\mu$ and $\{\psi_k\}_{k=1}^\mu$ we will take two orthonormal bases in $V(m)$ considered in Subsection 3.2. Concretely we will consider the following orthonormal bases of $V(m)$: $\{h^{(v)}(x) : 0 \leq v \leq m - 1\}$ and $\{\varphi_{1,j,m}(x) : 0 \leq j \leq m - 1\}$.

THEOREM 6.12. *For any $m = 2, 3, \dots$ the system $H^+(m)$ is an unconditional basis in the weighted norm space $L^p(\mathbb{R}^+, \omega)$, $1 < p < \infty$ if and only if there exists a point $y \in [0, +\infty]$ such that:*

If $y \neq +\infty$ then ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^+ \setminus \{y\})$ and the condition $\mathcal{M}_p^y(\mathbb{R}^+)$;

If $y = +\infty$ then ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^+)$.

PROOF. Suppose that $H^+(m)$ is an unconditional basis in the weighted norm space $L^p(\mathbb{R}^+, \omega)$, $1 < p < \infty$. Then $H^+(m)$ is complete minimal in $L^p(\mathbb{R}^+, \omega)$ and there exists a unique point $y \in [0, +\infty]$ such that the condition (69) holds. First consider the case $y = +\infty$. In this case the uniqueness of the point y means that for any $h_{k,j,m}^{(v)}(x) \in H^+(m)$

$$(71) \quad g_{k,j,m}^{(v)} = \frac{h_{k,j,m}^{(v)}}{\omega} \in L^{\frac{1}{p-1}}(\mathbb{R}^+).$$

The proof of the necessity can be easily completed following the scheme of the proof of Theorem 4.12.

If $y \in [0, +\infty)$ then by Lemma 6.5 the system byorthogonal to $H^+(m)$ is defined by the following equations:

$$(72) \quad g_{k,j,m}^{(v)}(t) = \frac{h_{k,j,m}^{(v)}(t) - h_{k,j,m}^{(v)}(y)\chi^+(t)}{\omega(t)}$$

for all $k \in \mathbb{Z}$, $j \in \mathbb{Z}^+$ and $1 \leq v \leq m-1$.

Let $\Delta = \Delta_{l+1}(y)$ and let $\Delta_l(y) = \Delta_l(0) + j_y m^{-l}$, where $j_y \in \mathbb{N}_0$. For $f \in L^p(\mathbb{R}^+, \omega)$, consider the sum

$$\begin{aligned} & \sum_{v=1}^{m-1} c_{l,j_y,m}^{(v)}(f) h_{l,j_y,m}^{(v)}(x) \\ &= \sum_{v=0}^{m-1} \int_{\mathbb{R}^+} f(t) g_{k,j_y,m}^{(v)}(t) \omega(t) dt h_{l,j_y,m}^{(v)}(x) - c_{l,j_y,m}^{(0)}(f) h_{l,j_y,m}^{(0)}(x) \\ &= \int_{\mathbb{R}^+} f(t) \sum_{v=0}^{m-1} g_{k,j_y,m}^{(v)}(t) h_{l,j_y,m}^{(v)}(x) \omega(t) dt - c_{l,j_y,m}^{(0)}(f) h_{l,j_y,m}^{(0)}(x), \end{aligned}$$

where $h_{l,j_y,m}^{(0)}(x) = \chi_{\Delta_l(y)}(x)$ and

$$g_{k,j,m}^{(0)}(t) = \frac{h_{k,j,m}^{(0)}(t) - h_{k,j,m}^{(0)}(y)\chi^+(t)}{\omega(t)}.$$

By Lemma 6.11 it follows that

$$(73) \quad \int_{\mathbb{R}^+} f(t) \sum_{v=0}^{m-1} g_{k,j_y,m}^{(v)}(t) h_{l,j_y,m}^{(v)}(x) \omega(t) dt$$

$$(74) \quad = \int_{\mathbb{R}^+} f(t) \sum_{j=0}^{m-1} \varphi_{l+1,j_y+j,m}(x) \psi_{l,j,m}(t) \omega(t) dt,$$

where

$$\psi_{l,j,m}(t) = \frac{\varphi_{l+1,j_y+j,m}(t) - \varphi_{l+1,j_y+j,m}(y)\chi^+(t)}{\omega(t)}.$$

Thus we obtain that if $x \in \Delta_{l+1}(y)$ then

$$\sum_{\nu=1}^{m-1} c_{l,j_y,m}^{(\nu)}(f) h_{l,j_y,m}^{(\nu)}(x) = -m^{l+1} \int_{\mathbb{R}^+ \setminus \Delta_{l+1}(y)} f(t) dt - m^l \int_{\mathbb{R}^+ \setminus \Delta_l(y)} f(t) dt .$$

If $f(t) \geq 0$ for $t \in \mathbb{R}^+$ then it follows that for $x \in \Delta_{l+1}(y)$

$$\left| \sum_{\nu=1}^{m-1} c_{l,j_y,m}^{(\nu)}(f) h_{l,j_y,m}^{(\nu)}(x) \right| \geq |\Delta_{l+1}(y)|^{-1} \left| \int_{\mathbb{R}^+ \setminus \Delta_{l+1}(y)} f(t) dt \right| .$$

Afterwards in the same way as in the proof of Theorem 5.7 we obtain that for some $B_p > 0$ and for all $l \in \mathbb{Z}$

$$|\Delta_l(y)|^{-p} \omega(\Delta_l(y)) \left(\int_{\mathbb{R}^+ \setminus \Delta_l(y)} \omega^{-\frac{1}{p-1}} dt \right)^{p-1} \leq B_p^p .$$

Let $E_{lj} \in \mathcal{M}$, $1 \leq j \leq m-1$ be mutually disjoint intervals such that $|E_{lj}| = m^{-1} |\Delta_l(y)|$, $1 \leq l \leq m-1$ and $\Delta_l(y) = \Delta_{l+1}(y) \cup \bigcup_{j=1}^{m-1} E_{lj}$. By (73) we obtain that if $f(t) = 0$ when $x \in \mathbb{R}^+ \setminus \Delta_l(y)$ then

$$\sum_{\nu=1}^{m-1} c_{l,j_y,m}^{(\nu)}(f) h_{l,j_y,m}^{(\nu)}(x) = |E_{lj}|^{-1} \int_{E_{lj}} f(t) dt \quad \text{if } x \in E_{lj} (1 \leq j \leq m-1) .$$

Which yields

$$|E_{lj}|^{-p} \omega(E_{lj}) \left(\int_{E_{lj}} \omega^{-\frac{1}{p-1}} dt \right)^{p-1} \leq B_p^p \quad \text{for any } l \in \mathbb{Z} \quad \text{and } 1 \leq j \leq m-1 .$$

Let $\Delta \in \mathcal{M}$, $|\Delta| = m^{-l-1}$ be such that $\Delta \cap \Delta_l(y) = \emptyset$. We consider the interval $\Delta^* \in \mathcal{M}$, $|\Delta^*| = m^{-l}$ be such that $\Delta \subset \Delta^*$. Let $\Delta = \Delta_l(0) + k^* m^{-l}$, where $k^* \in \mathbb{N}_0$.

For $f \in L^p(\mathbb{R}^+, \omega)$ consider the sum

$$\sum_{\nu=1}^{m-1} c_{l,k^*,m}^{(\nu)}(f) h_{l,k^*,m}^{(\nu)}(x) .$$

Using the same idea as above we show that if $x \in \Delta$ then

$$\sum_{\nu=1}^{m-1} c_{l,k^*,m}^{(\nu)}(f) h_{l,k^*,m}^{(\nu)}(x) = |\Delta|^{-1} \int_{\Delta} f(t) dt - |\Delta^*|^{-1} \int_{\Delta^*} f(t) dt .$$

Thus if $f(t) = 0$ for $t \in \mathbb{R}^+ \setminus \Delta$ it follows that

$$\sum_{\nu=1}^{m-1} c_{l,k^*,m}^{(\nu)}(f) h_{l,k^*,m}^{(\nu)}(x) = \frac{m-1}{m} |\Delta|^{-1} \int_{\Delta} f(t) dt$$

and the proof of the necessity is completed easily.

The proof of the sufficiency will be given following the same idea as in proof of Theorem 4.10. By Lemma 6.8 we have that the system $H^+(m)$ is complete and minimal in $L^p(\mathbb{R}^+, \omega)$. Let $G^+(m) = \{g_{k,j,m}^{(\nu)} : k \in \mathbb{Z}, j \in \mathbb{Z}^+, 1 \leq \nu \leq m-1\}$ be the conjugate sys-

tem of the basis $H^+(m)$. Suppose that ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^+)$. Then the system $G^+(m)$ is defined by the equations (71). Let $f \in L^p(\mathbb{R}^+, \omega)$ and let $\Omega \subset \mathbb{Z} \times \mathbb{Z}^+$ be a finite set. Moreover, let $N \in \mathbb{N}$ be such that $h_{k,j,m}^{(v)}(x) = 0$ if $x \in [m^N, +\infty)$ for all $(k, j) \in \Omega$. For any $1 \leq v \leq m - 1$ consider the sum

$$(75) \quad S_{\Omega}^{(v)}(f, x) = \sum_{(k,j) \in \Omega} c_{kj}^{(v)}(f) h_{k,j,m}^{(v)}(x),$$

where

$$c_{kj}^{(v)}(f) = \int_{\mathbb{R}^+} f(t) h_{k,j,m}^{(v)}(t) dt.$$

Applying Lemma 3.15 as in the proof of Theorem 4.10 we obtain that

$$\|S_{\Omega}^{(v)}(f, \cdot)\|_{L^p(\mathbb{R}^+, \omega)} \leq B_p \|f\|_{L^p(\mathbb{R}^+, \omega)},$$

where $B_p > 0$ is independent of f and Ω .

If $y \in [0, +\infty)$ then we take $N \in \mathbb{N}$ so that $h_{k,j,m}^{(v)}(x) = 0$ if $x \in [m^N, +\infty)$ for all $(k, j) \in \Omega$ and $y \in [0, m^N]$. By Lemma 6.5 the system $G^+(m)$ is defined by the equations (72). In this case the coefficients of the sum (75) are defined as follows:

$$c_{kj}^{(v)}(f) = \int_{\mathbb{R}^+} f(t) [h_{k,j,m}^{(v)}(t) - h_{k,j,m}^{(v)}(y)] dt.$$

We write $f(t) = f_1(t) + f_2(t)$, where $f_1(t) = f(t)\chi_{[0, m^N]}(t)$. By Lemma 3.16 and Lemma 3.17 in the same way as above we obtain that

$$\|S_{\Omega}^{(v)}(f_1, \cdot)\|_{L^p(\mathbb{R}^+, \omega)} \leq B_p \|f_1\|_{L^p(\mathbb{R}^+, \omega)},$$

where $B_p > 0$ is independent of f_1 and Ω . The proof will be complete if we show that

$$\|S_{\Omega}^{(v)}(f_2, \cdot)\|_{L^p(\mathbb{R}^+, \omega)} \leq B_p^* \|f_2\|_{L^p(\mathbb{R}^+, \omega)},$$

where $B_p^* > 0$ is independent of f_2 and Ω . We have that

$$\begin{aligned} S_{\Omega}^{(v)}(f_2, x) &= \sum_{(k,j) \in \Omega} c_{kj}^{(v)}(f_2) h_{k,j,m}^{(v)}(x) \\ &= - \int_{[m^N, +\infty)} f(t) dt \sum_{(k,j) \in \Omega} h_{k,j,m}^{(v)}(y) h_{k,j,m}^{(v)}(x). \end{aligned}$$

Let $f_N^*(t) = m^{-N} \chi_{[0, m^N]}(t)$ then

$$\begin{aligned} \sum_{(k,j) \in \Omega} h_{k,j,m}^{(v)}(y) h_{k,j,m}^{(v)}(x) &= \sum_{(k,j) \in \Omega} \int_{\mathbb{R}^+} f_N^*(t) [h_{k,j,m}^{(v)}(t) - h_{k,j,m}^{(v)}(y)] dt h_{k,j,m}^{(v)}(x) \\ &= S_{\Omega}^{(v)}(f_N^*, x). \end{aligned}$$

Thus we have that

$$\|S_{\Omega}^{(v)}(f_2, \cdot)\|_{L^p(\mathbb{R}^+, \omega)} = \left| \int_{[m^N, +\infty)} f(t) dt \right| \|S_{\Omega}^{(v)}(f_N^*, \cdot)\|_{L^p(\mathbb{R}^+, \omega)}$$

$$\begin{aligned} &\leq \|f_2\|_{L^p(\mathbb{R}^+, \omega)} \left[\int_{[m^N, +\infty)} \omega(t)^{-\frac{1}{p-1}} dt \right]^{\frac{1}{p'}} B_p \|f_N^*\|_{L^p(\mathbb{R}^+, \omega)} \\ &= B_p m^{-N} \left[\int_{[m^N, +\infty)} \omega(t)^{-\frac{1}{p-1}} dt \right]^{\frac{1}{p'}} \left[\int_{[0, m^N]} \omega(t) dt \right]^{\frac{1}{p}} \|f_2\|_{L^p(\mathbb{R}^+, \omega)}. \end{aligned}$$

Using that ω satisfies the condition $\mathcal{M}_p^y(\mathbb{R}^+)$ we complete the proof. □

It is easy to check that any function $\omega_r(x) = x^r$ if $r > p - 1$ satisfies the condition $\mathcal{M}_p^0(\mathbb{R}^+)$ and the condition $\mathcal{M}_p((0, +\infty))$.

COROLLARY 6.13. *Let $1 < p < \infty$ and let $\omega_r(x) = |x|^r$ if $r > p - 1$. Then the system $H^+(m)$ is an unconditional basis in the weighted norm space $L^p(\mathbb{R}^+, \omega_r)$.*

THEOREM 6.14. *For any $m = 2, 3, \dots$ the system $H(m)$ is an unconditional basis in the weighted norm space $L^p(\mathbb{R}, \omega)$, $1 < p < \infty$ if and only if there exist two points $y_1 \in [0, +\infty]$, $y_2 \in [-\infty, 0]$ such that:*

- If $y_1 \neq +\infty$ then ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^+ \setminus \{y_1\})$ and the condition $\mathcal{M}_p^{y_1}(\mathbb{R}^+)$;*
- If $y_1 = +\infty$ then ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^+)$;*
- If $y_2 \neq -\infty$ then ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^- \setminus \{y_2\})$ and the condition $\mathcal{M}_p^{y_2}(\mathbb{R}^-)$;*
- If $y_2 = -\infty$ then ω satisfies the condition $\mathcal{M}_p(\mathbb{R}^-)$;*

COROLLARY 6.15. *Let $1 < p < \infty$ and let $\omega_r(x) = |x|^r$ if $r > p - 1$. Then the system $H(m)$ is an unconditional basis in the weighted norm space $L^p(\mathbb{R}, \omega_r)$.*

REFERENCES

- [1] H. A. AIMAR, A. L. BERNARDIS AND F. J. MARTÍN-REYES, Multiresolution approximations and wavelet bases of weighted L^p spaces, *J. Fourier Anal. Appl.* 9 (2003), 497–510.
- [2] H. A. AIMAR, A. L. BERNARDIS AND B. IAFFEI, Multiresolution approximations and unconditional bases on weighted Lebesgue spaces on spaces of homogeneous type, *J. Approx. Theory* 148 (2007), 12–34.
- [3] K. I. BABENKO, On conjugate functions (in Russian), *Dokl. Akad. Nauk (N.S.)* 62 (2) (1948), 157–160.
- [4] Y. M. CHEN, Theorems of asymptotic approximation, *Math. Ann.* 140 (1960), 360–407.
- [5] P. CIFUENTES, K. S. KAZARIAN AND A. SAN-ANTOLÍN, Characterization of scaling functions in a Multiresolution Analysis, *Proc. Amer. Math. Soc.* 133 (2005), 1013–1022.
- [6] P. CIFUENTES, K. S. KAZARIAN AND A. SAN-ANTOLÍN, Characterization of scaling functions, *Wavelets and splines: Athens 2005*, 152–163, *Mod. Methods Math.*, Nashboro Press, Brentwood, TN, 2006.
- [7] R. R. COIFMAN AND C. FEFFERMAN, Weighted norm inequalities for maximal functions and singular integrals, *Studia Math.* 51 (1974), 241–250.
- [8] I. DAUBECHIES, *Ten lectures on wavelets*, SIAM, Philadelphia, 1992.
- [9] R. E. EDWARDS, The stability of weighted Lebesgue spaces, *Trans. Amer. Math. Soc.* 93 (1959), 369–394.
- [10] J. GARCÍA-CUERVA AND K. S. KAZARIAN, Calderón-Zygmund operators and unconditional bases of weighted Hardy spaces, *Studia Math.* 109 (1994), 255–276.
- [11] J. GARCÍA-CUERVA AND K. S. KAZARIAN, Spline wavelet bases of weighted L^p spaces, *Proc. Amer. Math. Soc.* 123 (1995), 433–439.
- [12] J. GARCÍA-CUERVA AND K. S. KAZARIAN, Spline wavelet bases of weighted spaces, *Fourier Analysis and Part. Diff. Eq., Studies in Advanced Mathematics*, CRC Press (1995), 169–184.

- [13] V. F. GAPOSHKIN, A generalization of a theorem of M. Riesz on conjugate functions (in Russian), *Mat. Sb.* 46 (88) (1958), 359–372.
- [14] B. I. GOLUBOV, Series in the Haar system, *Itogi Nauki, Ser. Matematika, Mat. Anal.* 1970 (1971), 109–146; trans. *J. Soviet Math.* 1 (1973), no. 6, 704–726.
- [15] G. GRIPENBERG, Necessary and sufficient conditions for the existence of a father wavelet, *Studia Math.* 114 (1995), 207–226.
- [16] R. F. GUNDY AND R. L. WHEEDEN, Weighted integral inequalities for the non-tangential maximal function, Lusin area integral and Walsh-Paley series, *Studia Math.* 49 (1974), 107–124.
- [17] G. H. HARDY AND J. E. LITTLEWOOD, Some more theorems concerning Fourier series and Fourier power series, *Duke Math. J.* 2 (1936), 354–382.
- [18] G. H. HARDY, J. E. LITTLEWOOD AND G. PÓLYA, *Inequalities*, Cambridge University Press, 1952.
- [19] H. HELSON AND G. SZEGÖ, A problem in prediction theory, *Ann. Mat. Pura Appl.* 51 (1960), 107–138.
- [20] E. HERNÁNDEZ AND G. WEISS, *A first course on Wavelets*, CRC Press, Inc. 1996.
- [21] E. HERNÁNDEZ, X. WANG AND G. WEISS, Characterization of wavelets, scaling functions and wavelets associated with multiresolution analyses, *Function spaces, interpolation spaces, and related topics (Haifa, 1995)*, 51–87, *Israel Math. Conf. Proc.*, 13, Bar-Ilan Univ., Ramat Gan, 1999.
- [22] I. I. HIRSCHMAN, The decomposition of Walsh and Fourier series, *Mem. Amer. Math. Soc. No. 15* (1955), 65pp.
- [23] M. IZUKI, The Haar wavelets and the Haar scaling function in weighted L^p spaces with $A_p^{dy:m}$ weights, *Hokkaido Math. J.* 36 (2007), no. 2, 417–444.
- [24] M. IZUKI AND Y. SAWANO, The Haar wavelet characterization of weighted Herz spaces and greediness of the Haar wavelet basis, *J. Math. Anal. Appl.* 362 (2010), no. 1, 140–155.
- [25] J.-P. KAHANE, *Some random series of functions*, Cambridge University Press, 1993.
- [26] K. S. KAZARIAN, The multiplicative completion of basic sequences in L^p , $1 \leq p < \infty$ to bases in L^p , *Dokl. Akad. Nauk Arm. SSR*, 62 (1976), 203–209, (Russian).
- [27] K. S. KAZARIAN, On the multiplicative completion of some incomplete orthonormal systems to bases in L^p , $1 \leq p < \infty$, (Russian), *Analysis Math.* 4 (1978), 37–52.
- [28] K. S. KAZARIAN, On bases and unconditional bases in the spaces $L^p(d\mu)$, $1 \leq p < \infty$, *Studia Math.* 71 (1982), 227–249.
- [29] K. S. KAZARIAN, Uniform continuity in weight spaces L^p , $1 \leq p < \infty$ of families of operators generated by truncated kernels, *Doklady AN USSR*, 272 (1983), 1048–1052.
- [30] K. S. KAZARIAN, Summability of generalized Fourier series and Dirichlet’s problem in $L^p(d\mu)$ and weighted H^p -spaces ($p > 1$), *Analysis. Math.* 13 (1987), no. 3, 173–197.
- [31] K. S. KAZARIAN, The multiplicative completion of certain systems, (Russian), *Izv. Akad. Nauk Armyan. SSR Ser. Mat.* 13 (1978), 315–351.
- [32] T. KOPALIANI, Higher rank Haar wavelet bases in spaces $L^p_w(\mathbb{R})$, *Georgian Math. J.* 18 (2011), 517–532.
- [33] S. V. KOZYREV, Wavelet theory as p -adic spectral analysis, *Izv. Math.* 66 (2002), 367–376.
- [34] A. S. KRANTSBERG, On the basisity of the Haar system in weighted spaces, *Mosk. Inst. Elektr. Mat.* 24 (1971).
- [35] P. G. LEMARIÉ-RIEUSSET, Ondelettes et poids de Muckenhoupt, *Studia Mathematica* 108 (1994), 127–147.
- [36] J. LINDENSTRAUSS AND L. TZAFRIRI, *Classical Banach spaces*, v.1, Springer, 1977.
- [37] S. MALLAT, Multiresolution approximations and wavelet orthonormal bases for $L^2(\mathbb{R})$, *Trans. Amer. Math. Soc.* 315 (1989), 69–87.
- [38] B. MUCKENHOUP, Weighted norm inequalities for the Hardy maximal function, *Trans. Amer. Math. Soc.* 165 (1972), 207–226.
- [39] M. C. PEREYRA, Lecture notes on dyadic harmonic analysis. Second Summer School in Analysis and Mathematical Physics (Cuernavaca, 2000), 1–60, *Contemp. Math.* 289, Amer. Math. Soc., Providence, RI, 2001.
- [40] M. ROSENBLUM, Summability of Fourier series in $L^p(d\mu)$, *Trans. Amer. Math. Soc.* 105 (1962), 32–42.

- [41] J. SCHAUDER, Eine eigenschaft des Haarschen orthogonalsystems, Math. Z. 28 (1928), 317–320.
- [42] E. M. STEIN, Singular integrals and differentiability properties of functions, Princeton, 1970.
- [43] A. TORCHINSKY, Real-variable methods in harmonic analysis, Acad. Press, 1986.
- [44] C. WATARI, Mean convergence of Walsh Fourier series, Tôhoku Math. J. 16 (1964), 183–188.
- [45] A. WOJCIECHOWSKA, A remark on wavelet bases in weighted L_p spaces, J. Funct. Spaces, 2012, Art. ID 328310, 10 pp.
- [46] P. WOJTASZCZYK, A mathematical introduction to wavelets, London Mathematical Society Student Texts 37, Cambridge University Press, Cambridge, 1997.
- [47] S. YANO, On a lemma of Marcinkiewicz and its applications to Fourier series, Tôhoku Math. J. 11 (1959), 191–215.
- [48] A. ZYGMUND, Trigonometric series, v. 1-2, Cambridge Univ. Press, 1959.

DEPARTMENT OF MATHEMATICS
UNIVERSIDAD AUTÓNOMA DE MADRID
28049, MADRID
SPAIN

E-mail address: kazaros.kazarian@uam.es

INSTITUTE OF MATHEMATICS
ARMENIAN ACADEMY OF SCIENCES
375019 YEREVAN
ARMENIA

E-mail address: kazarian@instmath.sci.am

DEPARTMENT OF MATHEMATICAL ANALYSIS
UNIVERSIDAD DE ALICANTE
03690 ALICANTE
SPAIN

E-mail address: angel.sanantolin@ua.es