

PRODUCTS OF RANDOM WALKS ON FINITE GROUPS WITH MODERATE GROWTH

GUAN-YU CHEN AND TAKASHI KUMAGAI

(Received March 21, 2017, revised May 1, 2017)

Abstract. In this article, we consider products of random walks on finite groups with moderate growth and discuss their cutoffs in the total variation. Based on several comparison techniques, we are able to identify the total variation cutoff of discrete time lazy random walks with the Hellinger distance cutoff of continuous time random walks. Along with the cutoff criterion for Laplace transforms, we derive a series of equivalent conditions on the existence of cutoffs, including the existence of pre-cutoffs, Peres' product condition and a formula generated by the graph diameters. For illustration, we consider products of Heisenberg groups and randomized products of finite cycles.

1. Introduction. Let G be a finite group equipped with a probability Q . A random walk on G driven by Q is a discrete time Markov chain with state space G and transition matrix K given by $K(x, y) = Q(x^{-1}y)$. If K is irreducible, then the stationary distribution U is uniform on G . For simplicity, we write the triple (G, Q, U) for such a random walk. Here, Q is called symmetric if $Q(x) = Q(x^{-1})$ for all $x \in G$ and, in this case, (G, Q, U) is named a symmetric random walk. Note that if Q is symmetric, then K is reversible. To study the convergence of (G, Q, U) , we consider the total variation and its corresponding mixing time, which are defined respectively by

$$(1.1) \quad d_{\text{TV}}(x, m) := \max_{E \subset G} \{Q^{(m)}(x^{-1}E) - U(E)\},$$

and

$$(1.2) \quad T_{\text{TV}}(x, \varepsilon) := \min\{m \geq 0 \mid d_{\text{TV}}(x, m) \leq \varepsilon\},$$

where $x^{-1}E = \{x^{-1}y \mid y \in E\}$ and $Q^{(m)}$ is the m -fold convolution product $Q * \cdots * Q$ with

$$f * g(x) = \sum_{z \in G} f(z)g(z^{-1}x).$$

As the total variation and, thus, its mixing time are constant in x , we shall write $d_{\text{TV}}(m)$ and $T_{\text{TV}}(\varepsilon)$ for short.

A subset $E \subset G$ is called symmetric, if $x \in E$ implies $x^{-1} \in E$, and is called a generating set of G , if $E^n := \{x_1 x_2 \cdots x_n \mid x_i \in E, \forall 1 \leq i \leq n\} = G$ for some $n > 0$. We write (G, E) for

2010 *Mathematics Subject Classification.* Primary 60J10; Secondary 60J27.

Key words and phrases. Product chains, random walks, moderate growth

The first author is partially supported by grant MOST 104-2115-M-009-013-MY3 and NCTS. The second author is partially supported by the Grant-in-Aid for Scientific Research (A) 25247007 and 17H01093.

the Cayley graph with vertex set G and edge set $\{(x, xy) | x \in G, y \in E\}$ and define its volume growth function and diameter by

$$V(m) = |E^m|, \quad \rho = \min\{m \geq 1 | V(m) = |G|\}.$$

A group G is said to have (A, d) -moderate growth with respect to a generating set E if

$$\frac{V(m)}{V(\rho)} \geq \frac{1}{A} \left(\frac{m}{\rho}\right)^d, \quad \forall 1 \leq m \leq \rho.$$

The following are some typical groups with moderate growth.

Example 1: When $G = \mathbb{Z}_n$ and $E = \{0, \pm 1\}$, the graph (G, E) has diameter $\rho = \lfloor n/2 \rfloor$ and G has $(1, 1)$ -moderate growth w.r.t. E for $n \geq 2$.

Example 2: When $G = \mathbb{Z}_n$ and $E = \{0, \pm 1, \pm \lfloor \sqrt{n} \rfloor\}$, the diameter ρ is of order \sqrt{n} and G has $(1, 2)$ -moderate growth w.r.t. E for $n \geq 2$.

Example 3: When G is the Heisenberg group mod $n + 2$, which is the set of 3×3 matrices of the form

$$(1.3) \quad \begin{pmatrix} 1 & i & k \\ 0 & 1 & j \\ 0 & 0 & 1 \end{pmatrix}, \quad \forall i, j, k \in \mathbb{Z}_{n+2},$$

and E contains the following matrices

$$(1.4) \quad I, \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix},$$

it was proved in [7, Lemma 4.1] that (G, E) has diameter $n + 1 \leq \rho \leq n + 4$ and G has $(48, 3)$ -moderate growth w.r.t. E for $n \geq 1$.

Throughout this article, we will simply write id for the identity of any group. In [7], Diaconis and Saloff-Coste considered random walks on finite groups with moderate growth and achieved the following proposition.

PROPOSITION 1.1 (Theorem 3.1 in [7]). *Let (G, Q, U) be a symmetric random walk on a finite group and E be a symmetric generating set of G containing id . Assume that G has (A, d) -moderate growth with respect to E and $\eta = \min\{Q(x) | x \in E\} > 0$. Then, there is $C_1 = C_1(A, d) > 0$ such that*

$$(1.5) \quad d_{TV}(m) \leq C_1 e^{-\eta m / \rho^2}, \quad \forall m \geq 0,$$

where ρ is the diameter of (G, E) . If it is assumed further that Q is supported on E and that $\rho \geq A2^{2d+2}$, then there is $C_2 = C_2(A, d) > 0$ such that

$$(1.6) \quad d_{TV}(m) \geq \frac{1}{2} e^{-C_2 m / \rho^2}, \quad \forall m \geq 0.$$

In fact, the authors of [7] obtain $C_1 = A^{1/2} 2^{d(d+3)/4}$ and $C_2 = A^2 2^{4d+2}$. This means that the bounds in (1.5)-(1.6) are far from comparable when A or d is large.

We now consider product chains. Let $(G_i, Q_i, U_i)_{i=1}^n$ be irreducible random walks on finite groups and (p_1, \dots, p_n) be a probability vector. Define

$$(1.7) \quad G = G_1 \times \dots \times G_n, \quad U = U_1 \times \dots \times U_n, \quad Q(x) = \sum_{i=1}^n p_i Q_i(x_i),$$

for $x = (x_1, \dots, x_n) \in G$. Here, (G, Q, U) is called the product of $(G_i, Q_i, U_i)_{i=1}^n$ with respect to the probability vector (p_1, \dots, p_n) . Note that if E_i is the support of Q_i and contains id , then Q is supported on $E = \check{E}_1 \cup \dots \cup \check{E}_n$, where $\check{E}_i = \{x = (x_1, \dots, x_n) \in G \mid x_i \in E_i, x_j = id, \forall j \neq i\}$. Further, if E_i is a symmetric generating set of G_i and ρ_i is the diameter of (G_i, E_i) , then E is a symmetric generating set of G and the diameter ρ of (G, E) satisfies $\rho = \rho_1 + \dots + \rho_n$. To see the moderate growth of direct products of groups, let E_i, E be as before and assume that G_i has (A_i, d_i) -moderate growth w.r.t. E_i . As G is a finite group and E generates G , there are always positive constants A, d such that G has (A, d) -moderate growth w.r.t. E . However, the relation between (A, d) and $(A_i, d_i)_{i=1}^n$ could be complicated and, in general, A or d can be very large when n grows. (For instance, consider $G_i = \mathbb{Z}_N$ and $E_i = \{0, \pm 1\}$ for $1 \leq i \leq n$. It is mentioned earlier that G_i has $(1, 1)$ moderate growth w.r.t. E_i . In some combinatoric computations, one may show that, under the assumption of $N \geq n^2$, G has (A, d) moderate growth with $A = (1 - 1/N)^{-2}$ and $d = n$.) Consequently, (1.5) and (1.6) might not be sharp enough to provide efficient bounds on the total variation even if the prerequisites, $\eta > 0$ and $\rho \leq A2^{2d+2}$, are fulfilled. To proceed the analysis of product chains, as the total variation mixing times are comparable between (G, Q, U) and its associated continuous time walk, see e.g. [5], it is more convenient, as is discussed below, to consider the continuous time chain rather than the discrete time one.

Given a random walk (G, Q, U) , we associate it with a continuous time random walk (G, H_t, U) , where $H_t = e^{t(K-U)}$ and K is the transition matrix given by Q . One realization of (G, H_t, U) is to change the constant waiting times of (G, Q, U) into an i.i.d. sequence of exponential random variables. Note that, if (G, Q, U) is the product of $(G_i, Q_i, U_i)_{i=1}^n$ with respect to the probability vector (p_1, \dots, p_n) and $(G_i, H_{i,t}, U_i)$ is the continuous time random walk associated with (G_i, Q_i, U_i) , then

$$(1.8) \quad H_t = H_{1,p_1 t} \otimes \dots \otimes H_{n,p_n t},$$

where $A \otimes B$ denotes the tensor product of matrices A and B . In general, K^m does not have the form of (1.8). Through (1.8), one may study H_t via $(H_{i,t})_{i=1}^n$ but, unfortunately, there lacks an efficient expression of the total variation of (G, H_t, U) in terms of the total variations of $(G_i, H_{i,t}, U_i)_{i=1}^n$.

In [2], two inequalities were used to compare the total variation and the Hellinger distance and this leads to a different way to analyze their mixing times. In detail, the Hellinger distance of (G, Q, U) is defined by

$$(1.9) \quad d_H(x, m) := \left(\frac{1}{2} \sum_{y \in G} (\sqrt{K^m(x, y)} - \sqrt{U(y)})^2 \right)^{1/2},$$

while the Hellinger distance of (G, H_t, U) is defined by replacing K^m with H_t in (1.9) and denoted by $d_H^{(c)}(x, t)$ in avoidance of confusion. As before, we will write $d_H(m)$ (resp. $d_H^{(c)}(t)$) for short since $d_H(x, m)$ (resp. $d_H^{(c)}(x, t)$) is constant in x . In the above setting, Equation (1.3) in [2] says that

$$(1.10) \quad 1 - \sqrt{1 - d_{TV}^2(m)} \leq d_H^2(m) \leq d_{TV}(m),$$

and also hold in the continuous time case. In the Hellinger distance, if (G, Q, U) is the product of $(G_i, Q_i, U_i)_{i=1}^n$ with respect to the probability vector (p_1, \dots, p_n) , then the Hellinger distances, $d_H^{(c)}$ and $d_{i,H}^{(c)}$, of (G, H_t, U) and $(G_i, H_{i,t}, U_i)$ satisfy

$$(1.11) \quad d_H^{(c)}(t)^2 = 1 - \prod_{i=1}^n (1 - d_{i,H}^{(c)}(p_i t)^2).$$

Such an equality is derived from (1.8) but not applicable to the discrete time case. See p.365 in [12] or Lemma 2.4 in [2] for a proof of (1.10) and see [2] for more comparisons of mixing times of product chains.

In this article, we focus on the cutoff phenomenon, or briefly cutoff, for products of random walks on finite groups with moderate growth. The cutoff of Markov chains was introduced by Aldous and Diaconis in early 1980s in order to catch up the phase transition of the mixing time. To see a definition, let $\mathcal{F} = (G_n, Q_n, U_n)_{n=1}^\infty$ be a family of random walks on finite groups. For $n \geq 1$, let $d_{n,TV}$ and $T_{n,TV}$ be the total variation and corresponding mixing time of the n th chain in \mathcal{F} . Assume that $T_{n,TV}(\varepsilon_0) \rightarrow \infty$ for some $\varepsilon_0 \in (0, 1)$. The family \mathcal{F} is said to present a cutoff in the total variation if

$$(1.12) \quad \lim_{n \rightarrow \infty} \frac{T_{n,TV}(\varepsilon)}{T_{n,TV}(\delta)} = 1, \quad \forall \varepsilon, \delta \in (0, 1),$$

or, equivalently (see [3, Proposition 2.4]), there is a sequence of positive reals $(t_n)_{n=1}^\infty$ such that

$$(1.13) \quad \lim_{n \rightarrow \infty} d_{n,TV}(\lceil at_n \rceil) = 0 \quad \forall a > 1, \quad \lim_{n \rightarrow \infty} d_{n,TV}(\lfloor at_n \rfloor) = 1, \quad \forall a \in (0, 1).$$

When a cutoff exists, the sequence $(t_n)_{n=1}^\infty$, or briefly t_n , in (1.13) is called a cutoff time. By (1.12), it is easy to see that $T_{n,TV}(\varepsilon)$ can be selected as cutoff time for any $\varepsilon \in (0, 1)$. In the continuous time case, we write $\mathcal{F}_c = (G_n, H_{n,t}, U_n)_{n=1}^\infty$ for the family of continuous time Markov chains associated with \mathcal{F} and use $d_{n,TV}^{(c)}$ and $T_{n,TV}^{(c)}$ to denote the total variation and its mixing time of the n th chain in \mathcal{F}_c . The total variation cutoff of \mathcal{F}_c is defined in the same way through (1.12) or (1.13) under the replacement of $T_{n,TV}, d_{n,TV}$ with $T_{n,TV}^{(c)}, d_{n,TV}^{(c)}$ and the removal of $\lceil \cdot \rceil, \lfloor \cdot \rfloor$ but without the prerequisite of $T_{n,TV}^{(c)}(\varepsilon_0) \rightarrow \infty$. All above is also applicable to the Hellinger distance. We refer readers to [6, 11] for more discussions on cutoffs for random walks on finite groups.

Cutoffs in the total variation and in the Hellinger distance were proved to be equivalent in [2] via (1.10). Since no similar formula to (1.11) is available for the total variation or for the discrete time case, it is straightforward to consider the cutoff in the Hellinger distance for

Total variation cutoffs for discrete time Markov chains
⇕ (Proposition 1.1 in [5] holds for lazy random walks)
Total variation cutoffs for continuous time Markov chains
⇕ (Proposition 1.1 in [2] holds for any Markov chain)
Hellinger distance cutoffs for continuous time Markov chains
⇕ (Proposition 3.3 holds for groups with moderate growth)
Cutoffs for Laplace transforms
⇕ (Theorem 2.4 in [1] holds for reversible Markov chains)
Precise condition on cutoffs

TABLE 1. A scheme to analyze cutoffs.

families of continuous time product chains. For finite groups with moderate growth, we obtain a continuous time variant of Proposition 1.1 in Proposition 3.3 with a refined assumption on the lower bound (from $\rho \geq A2^{2d+2}$ to $\rho \geq 4$). Through (1.11), the Hellinger distances of product chains can be expressed in a form related to sums of exponential functions. By regarding those sums as Laplace transforms, a criterion in [1] was proposed to determine the cutoff and to characterize the cutoff time. Table 1 is the conclusive scheme of all above discussions.

The aim of this paper is to establish necessary and sufficient conditions for cutoff for products of random walks on finite groups with moderate growth, and apply them to stimulating examples. In the first main theorem (Theorem 2.3), we give various equivalent conditions for cutoff in our framework. It should be noted that in this framework the cutoff is equivalent to a weaker concept, called the pre-cutoff (note that such an equivalence generally fails; see [8]). Moreover, one equivalent condition in Theorem 2.3 is consistent with Peres' conjecture (see Remark 2.4 (3)), while another is simply determined by the graph diameters and a sequence \mathcal{P} given below. In the second main theorem (Theorem 2.5), we apply Theorem 2.3 to the specific type of products introduced in [2] and derive more concrete conditions on their cutoffs. To illustrate our results, let us consider products of random walks on Heisenberg groups and randomized products of random walks on finite cycles. Let $\mathcal{G} = (G_n, Q_n, U_n)_{n=1}^\infty$ be a family of random walks on finite groups and $\mathcal{P} = (p_n)_{n=1}^\infty$ be a sequence of positive reals. For $n \geq 1$, let $q_n = \sum_{i=1}^n p_i$ and write $\mathcal{G}^{\mathcal{P}}$ for the family of which n th random walk is the product of $(G_i, Q_i, U_i)_{i=1}^n$ according to the probability vector $(p_1/q_n, \dots, p_n/q_n)$. Then the following hold.

PROPOSITION 1.2. *Let $\mathcal{G} = (G_n, Q_n, U_n)_{n=1}^\infty$, where G_n is the Heisenberg group in (1.3), Q_n is the probability uniformly supported on the set E_n in (1.4) and $p_n = n^2 \exp\{-n^\gamma\}$ with $\gamma > 0$. Then, $\mathcal{G}^{\mathcal{P}}$ has a total variation cutoff if and only if $0 < \gamma < 1$.*

PROPOSITION 1.3. *Consider the family $\mathcal{G} = (\mathbb{Z}_{n+2}, Q_n, U_n)_{n=1}^\infty$, where $Q_n(0) = 1/2$ and $Q_n(1) = Q_n(-1) = 1/4$. Let $(X_n)_{n=1}^\infty$ be i.i.d. positive random variables and $\mathcal{P} = (p_n)_{n=1}^\infty$ be a random sequence given by $(X_n)_{n=1}^\infty$.*

- (1) Suppose $p_n = (X_1 + \dots + X_n)^\gamma$ with $\gamma > 0$ and X_1 has a finite expectation. For $\gamma \in (0, 2]$, $\mathcal{G}^{\mathcal{P}}$ has a total variation cutoff with probability 1. For $\gamma > 2$, $\mathcal{G}^{\mathcal{P}}$ has no total variation cutoff with probability 1.
- (2) Suppose $p_n = X_1 \times \dots \times X_n$ and $\log X_1$ has a positive finite expectation. Then, $\mathcal{G}^{\mathcal{P}}$ has no total variation cutoff with probability 1.

Proposition 1.2 is an immediate result of Corollary 2.9, which shows a phase transition of cutoffs at $\gamma = 1$. Proposition 1.3 is of its own interest and discussed in detail in Subsection 2.3.

The remaining of this paper is organized in the following way. In Section 2, we introduce the core results of the paper. The main theorems are given in Subsections 2.1 and 2.2. As an example, we consider the randomized product and discuss its cutoff in Subsection 2.3. Section 3 is dedicated to the construction of framework in Section 2. In Subsection 3.1, we review and develop some theoretical results that are crucial to the equivalences in Table 1, while in Subsection 3.2, Theorem 2.3 is proved in detail. To make this paper more readable, we address those minor and involved results in the appendix.

We end the introduction by quoting the following notations. Let $x, y \in \mathbb{R}$ and a_n, b_n be sequences of positive reals. We write $x \vee y$ and $x \wedge y$ for the maximum and minimum of x, y . When a_n/b_n is bounded, we write $a_n = O(b_n)$; when $a_n/b_n \rightarrow 0$, we write $a_n = o(b_n)$. In the case of $a_n = O(b_n)$ and $b_n = O(a_n)$, we simply say $a_n \asymp b_n$. If $a_n/b_n \rightarrow 1$, we write $a_n \sim b_n$. In computations, $O(a_n)$ and $o(b_n)$ denote two sequences c_n and d_n satisfying $|c_n/a_n| = O(1)$ and $|d_n/b_n| = o(1)$.

2. Main theorems and applications. In this section, we will introduce our main results in the general setting and discuss their applications, including Proposition 1.2.

2.1. Framework and main theorem. In this subsection, we introduce the theoretical framework and one of the main theorems in this article. First, let us consider a concept weaker than cutoff.

DEFINITION 2.1. Let $\mathcal{F} = (G_n, Q_n, U_n)_{n=1}^\infty$ be a family of random walks on finite groups and $d_{n,TV}$ be the total variation of (G_n, Q_n, U_n) . \mathcal{F} is said to present a pre-cutoff in the total variation if there are $B > A > 0$ and a sequence $t_n > 0$ such that

$$(2.1) \quad \lim_{n \rightarrow \infty} d_{n,TV}(\lceil Bt_n \rceil) = 0, \quad \liminf_{n \rightarrow \infty} d_{n,TV}(\lfloor At_n \rfloor) > 0.$$

REMARK 2.2. (1) The removal of $\lceil \cdot \rceil, \lfloor \cdot \rfloor$ provides the pre-cutoff for \mathcal{F}_c and the replacement of $d_{n,TV}$ with $d_{n,H}$ yields the pre-cutoff in the Hellinger distance. When $t_n \rightarrow \infty$, the pre-cutoff in Definition 2.1 is equivalent to

$$(2.2) \quad \lim_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{T_{n,TV}(\varepsilon)}{T_{n,TV}(\varepsilon_0 - \varepsilon)} < \infty,$$

for some $\varepsilon_0 \in (0, 1)$. Such an equivalence also holds for \mathcal{F}_c without the prerequisite of $t_n \rightarrow \infty$.

(2) Definition 2.1 was introduced for the purpose of studying the mixing times and cutoffs for families of Markov chains. Readers are referred to [10, 3] for more discussions on this

subject. It is worthwhile to note that there is indeed another (stronger) variant of pre-cutoff in [9, Chapter 18], of which definition is similar to (2.1) except the replacement of the second limit by

$$\lim_{n \rightarrow \infty} d_{n,TV}(\lfloor At_n \rfloor) = 1.$$

When $t_n \rightarrow \infty$, the equivalence of the pre-cutoff and (2.2) also holds for such a variant with $\varepsilon_0 = 1$. We would like to emphasize that Theorem 2.3 (discussed later) remains true when the pre-cutoff refers to the stronger one.

In the following, we consider a rather general setting than Proposition 1.2. Let $(k_n)_{n=1}^\infty$ be a sequence of positive integers and

$$(2.3) \quad \mathcal{F} = \{(G_{n,i}, Q_{n,i}, U_{n,i}) | 1 \leq i \leq k_n, n \geq 1\}, \quad \mathcal{P} = \{p_{n,i} | 1 \leq i \leq k_n, n \geq 1\},$$

where $(G_{n,i}, Q_{n,i}, U_{n,i})$ is a random walk on a finite group and $p_{n,i} > 0$. We write $\mathcal{F}^{\mathcal{P}}$ for the family $(G_n, Q_n, U_n)_{n=1}^\infty$, where (G_n, Q_n, U_n) is the product of $(G_{n,i}, Q_{n,i}, U_{n,i})_{i=1}^{k_n}$ according to the probability vector $(p_{n,i}/q_n)_{i=1}^{k_n}$ and $q_n = p_{n,1} + \dots + p_{n,k_n}$. As before, we use $d_{n,TV}^{(c)}, T_{n,TV}^{(c)}$ to denote the total variation and its mixing time of the n th chain in $\mathcal{F}_c^{\mathcal{P}}$. Along with these notations, we are ready to state the first main theorem of this article.

THEOREM 2.3. *Refer to the triangular arrays in (2.3). Let $E_{n,i}$ be the support of $Q_{n,i}$ and $\rho_{n,i}$ be the diameter of $(G_{n,i}, E_{n,i})$. Assume that $Q_{n,i}$ is symmetric, $\inf\{Q_{n,i}(x) | x \in E_{n,i}, 1 \leq i \leq k_n, n \geq 1\} > 0$ and $G_{n,i}$ has (A, d) -moderate growth with respect to $E_{n,i}$ for all n, i . Assume further that $\rho_{n,1} \geq 4$ for n large enough and there are $C > 1$ and $\ell_{n,i} > 0$ satisfying $\ell_{n,i} \leq \ell_{n,i+1}$ such that $\ell_{n,i}/C \leq p_{n,i}/(q_n \rho_{n,i}^2) \leq C \ell_{n,i}$ for all n, i . By setting $t_n = \max\{\log(i+1)/\ell_{n,i} | 1 \leq i \leq k_n\}$, one has:*

- (1) If $k_n = O(1)$, then there are $A > 0$ and $\sigma_2 > \sigma_1 > 0$ such that

$$1 - \exp\{-e^{-a\sigma_2}\} \leq d_{n,TV}^{(c)}(at_n) \leq 1 - \exp\{-e^{-a\sigma_1}\}, \quad \forall a > A, n \geq 1.$$

In particular, $\mathcal{F}_c^{\mathcal{P}}$ has no pre-cutoff in the total variation.

- (2) If $k_n \rightarrow \infty$ and $\min\{\rho_{n,i} | i \geq m, n \geq 1\} \geq 4$ for m large enough, then the following are equivalent.

- (i) $\mathcal{F}_c^{\mathcal{P}}$ has a total variation cutoff.
- (ii) $\mathcal{F}_c^{\mathcal{P}}$ has a total variation pre-cutoff.
- (iii) $t_n \ell_{n,1} \rightarrow \infty$.
- (iv) $T_{n,TV}^{(c)}(\varepsilon) \ell_{n,1} \rightarrow \infty$ for all $\varepsilon \in (0, 1)$.
- (v) $T_{n,TV}^{(c)}(\varepsilon) \ell_{n,1} \rightarrow \infty$ for some $\varepsilon \in (0, 1)$.

Moreover, if $\mathcal{F}_c^{\mathcal{P}}$ has a total variation cutoff, then $T_{n,TV}^{(c)}(\varepsilon) \asymp t_n$ and $T_{n,TV}^{(c)}(\varepsilon) = T_{n,TV}^{(c)}(\delta) + O(1/\ell_{n,1})$ for all $0 < \varepsilon < \delta < 1$.

When $E_{n,i}$ contains id , (2) also holds under the replacement of $\mathcal{F}_c^{\mathcal{P}}$ and $T_{n,TV}^{(c)}$ with $\mathcal{F}^{\mathcal{P}}$ and $T_{n,TV}$.

Theorem 2.3 is built on a list of theoretical results in Subsection 3.1. As its proof is a little complicated, we leave it to Subsection 3.2. In the following, we provide some remarks to comment the importance of Theorem 2.3.

REMARK 2.4. (1) Note that Theorem 2.3 also holds in the Hellinger distance due to (1.10) and Proposition 3.2 and this is exactly what is done in the proof of the continuous time case. See Subsection 3.2 for details.

(2) In the proof of Theorem 2.3, we obtain the order of the cutoff times, which is the same as t_n , but could not determine its asymptotic value, which relies on a more precise estimation of the convergence rate in the Helinger distance in Proposition 3.3.

(3) A conjectured condition for the existence of cutoffs was introduced by Peres in 2004, which says that

$$(2.4) \quad \text{A cutoff exists} \iff \text{Mixing time} \times \text{Spectral gap} \rightarrow \infty .$$

In the setting of Theorem 2.3, the spectral gaps of the n th random walks in $\mathcal{F}^{\mathcal{P}}$ and $\mathcal{F}_c^{\mathcal{P}}$ are of the same order as $\ell_{n,1}$ due to the assumption of $\inf\{Q_{n,i}(x)|x \in E_{n,i}, 1 \leq i \leq k_n, n \geq 1\} > 0$. Consequently, the equivalence of (i) and (v) in (2) confirms the conjecture in (2.4) for products of random walks on finite groups with moderate growth.

2.2. Applications. In this subsection, we apply Theorem 2.3 to the specific type of products introduced in [2] and derive conditions on their cutoffs. Let $\mathcal{G} = (G_n, Q_n, U_n)_{n=1}^{\infty}$ be a family of random walks on finite groups driven by symmetric probabilities and $\mathcal{P} = (p_n)_{n=1}^{\infty}$ be a sequence of positive reals. Throughout this subsection, we write $\mathcal{G}^{\mathcal{P}}$ for the family, of which n th random walk refers to the product of $(G_i, Q_i, U_i)_{i=1}^n$ with respect to the probability vector $(p_1/q_n, \dots, p_n/q_n)$, where $q_n = p_1 + \dots + p_n$. We now state the second main theorem of this article.

THEOREM 2.5. *Consider the family $\mathcal{G}^{\mathcal{P}}$ introduced above. Let E_n be the support of Q_n and ρ_n be the diameter of (G_n, E_n) . Assume that G_n has (A, d) -moderate growth with respect to E_n , $\inf\{Q_n(x)|x \in E_n, n \geq 1\} > 0$, $\rho_n \geq 4$ for n large enough and $p_n/\rho_n^2 \asymp \ell_n$ for some sequence $(\ell_n)_{n=1}^{\infty}$.*

- (1) *If $\ell_n \leq \ell_{n+1}$ and $u_n = \max\{\log(i+1)/\ell_i|1 \leq i \leq n\}$, then $\mathcal{G}_c^{\mathcal{P}}$ has a total variation cutoff if and only if $u_n \rightarrow \infty$.*
- (2) *If $\ell_n \geq \ell_{n+1}$ and $u_n = \max\{\log(i+1)/\ell_{n-i+1}|1 \leq i \leq n\}$, then $\mathcal{G}_c^{\mathcal{P}}$ has a total variation cutoff if and only if $u_n \ell_n \rightarrow \infty$.*

In either case of (1) and (2), if $\mathcal{G}_c^{\mathcal{P}}$ has a total variation cutoff, then the cutoff time is of order $u_n q_n$. Further, if E_n contains id for all $n \geq 1$, then all above also holds for $\mathcal{G}^{\mathcal{P}}$.

REMARK 2.6. The lower bound of the graph diameter (at least 4) in Theorem 2.5 is due to the requirement in Proposition 3.3. As the product of finitely many random walks has negligible contribution to the total variation (see e.g. Theorem 2.3(1) for an illustration), one may suitably relax such a restriction on graph diameters as in Theorem 2.5.

PROOF OF THEOREM 2.5. By Propositions 3.1 and 3.2 in the next section, it suffices to prove this theorem for $\mathcal{G}_c^{\mathcal{P}}$ in the Hellinger distance. In the following, we will discuss (2), while (1) can be treated in a similar way.

Let $n_0 > 0$ be an integer such that $\rho_n \geq 4$ for $n \geq n_0$. For $n > n_0$, let (G, Q, U) and $(G_n^{\mathcal{R}}, Q_n^{\mathcal{R}}, U_n^{\mathcal{R}})$ be products of $(G_i, Q_i, U_i)_{i=1}^{n_0}$ and $(G_i, Q_i, U_i)_{i=n_0+1}^n$ with respect to the probability vectors $(p_i/q)_{i=1}^{n_0}$ and $(p_i/q_n^{\mathcal{R}})_{i=n_0+1}^n$, where $q = p_1 + \dots + p_{n_0}$ and $q_n^{\mathcal{R}} = p_{n_0+1} + \dots + p_n$. Clearly, the n th random walk in $\mathcal{G}^{\mathcal{P}}$ is the product of (G, Q, U) and $(G_n^{\mathcal{R}}, Q_n^{\mathcal{R}}, U_n^{\mathcal{R}})$ with respect to the probability vector $(q/q_n, q_n^{\mathcal{R}}/q_n)$. Let $d_H, d_{n,H}^{\mathcal{R}}$ and $d_{n,H}^{(c)}$ be the Hellinger distances of the continuous time random walks associated with $(G, Q, U), (G_n^{\mathcal{R}}, Q_n^{\mathcal{R}}, U_n^{\mathcal{R}})$ and the n th random walk in $\mathcal{G}^{\mathcal{P}}$. By (1.11), one has

$$(2.5) \quad d_{n,H}^{(c)}(t)^2 = 1 - \left(1 - d_H(qt/q_n)^2\right) \left(1 - d_{n,H}^{\mathcal{R}}(q_n^{\mathcal{R}}t/q_n)^2\right).$$

For the family $\mathcal{H} := (G_n^{\mathcal{R}}, Q_n^{\mathcal{R}}, U_n^{\mathcal{R}})_{n=n_0+1}^{\infty}$ and the random walk (G, Q, U) , we set

$$v_n = q_n^{\mathcal{R}} \max_{1 \leq i \leq n-n_0} \frac{\log(i+1)}{\ell_{n-i+1}}, \quad v = q \max_{1 \leq i \leq n_0} \frac{\log(i+1)}{\ell_{n_0-i+1}}.$$

By Theorem 2.3(1), there are constants $A > 0$ and $\sigma_2 > \sigma_1 > 0$ such that

$$(2.6) \quad 1 - \exp\{-e^{-a\sigma_2}\} \leq d_H(av) \leq 1 - \exp\{-e^{-a\sigma_1}\}, \quad \forall a > A,$$

and, by Theorem 2.3(2), \mathcal{H}_c has a cutoff in the Hellinger distance if and only if $v_n \ell_n / q_n^{\mathcal{R}} \rightarrow \infty$. Observe that, for $n > n_0$,

$$\frac{v_n}{q_n^{\mathcal{R}}} \leq u_n \leq \frac{v_n}{q_n^{\mathcal{R}}} + \max_{n-n_0 < i \leq n} \frac{\log(i+1)}{\ell_{n-i+1}} \leq \frac{v_n}{q_n^{\mathcal{R}}} \left(1 + \frac{\log(n+1)}{\log(n-n_0+1)}\right).$$

This implies $v_n/q_n^{\mathcal{R}} \asymp u_n$.

We are now ready to derive (2). Assume that $u_n \ell_n \rightarrow \infty$ or equivalently $v_n \ell_n / q_n^{\mathcal{R}} \rightarrow \infty$. By Theorem 2.3(2), \mathcal{H}_c has a cutoff in the Hellinger distance and the cutoff time w_n satisfies $w_n \asymp v_n$. Immediately, this implies

$$\lim_{n \rightarrow \infty} d_{n,H}^{\mathcal{R}}(aw_n) = \begin{cases} 0 & \text{for } a > 1, \\ 1 & \text{for } 0 < a < 1, \end{cases} \quad \lim_{n \rightarrow \infty} d_H(bw_n/q_n^{\mathcal{R}}) = 0, \quad \forall b > 0,$$

where the second limit results from the second inequality of (2.6) and the observation of $u_n \geq \log(n+1)/\ell_1 \rightarrow \infty$. Applying the above computations to (2.5) yields

$$\lim_{n \rightarrow \infty} d_{n,H}^{(c)}(aq_n w_n / q_n^{\mathcal{R}}) = \begin{cases} 0 & \text{for } a > 1, \\ 1 & \text{for } 0 < a < 1. \end{cases}$$

This proves that $\mathcal{G}_c^{\mathcal{P}}$ has a cutoff in the Hellinger distance and the cutoff time is of order $u_n q_n$. Conversely, assume that $\mathcal{G}_c^{\mathcal{P}}$ has a cutoff in the Hellinger distance with cutoff time w'_n . Then, for $a > 1$,

$$\lim_{n \rightarrow \infty} d_{n,H}^{\mathcal{R}}(aq_n^{\mathcal{R}} w'_n / q_n) = 0, \quad \lim_{n \rightarrow \infty} d_H(aq w'_n / q_n) = 0.$$

By the first inequality of (2.6), the latter limit implies $w'_n/q_n \rightarrow \infty$, which yields $d_H(bw'_n/q_n) \rightarrow 0$ for all $b > 0$. In addition with the cutoff for $\mathcal{G}_c^{\mathcal{P}}$, we may derive from (2.5) that, for $0 < a < 1$,

$$1 = \lim_{n \rightarrow \infty} d_{n,H}^{(c)}(aw'_n) = \lim_{n \rightarrow \infty} d_{n,H}^{\mathcal{R}}(aq_n^{\mathcal{R}}w'_n/q_n).$$

As a consequence, \mathcal{H}_c has a cutoff in the Hellinger distance or equivalently $u_n \ell_n \asymp v_n \ell_n / q_n^{\mathcal{R}} \rightarrow \infty$, as desired. □

REMARK 2.7. Theorem 2.5(1) can be in fact proved in a more direct way. Consider the exchange of the first random walk in \mathcal{G} and the first random walk of which graph diameter is at least 4, say the N th random walk. For the new family, all assumptions in Theorem 2.3 are fulfilled except the monotonicity of the sequence $\{\ell_N, \ell_2, \dots, \ell_{N-1}, \ell_1, \ell_{N+1}, \dots\}$. Such a concern can be eliminated by using the original sequence $(\ell_n)_{n=1}^{\infty}$ along with a larger multiplicative constant and its reciprocal to bound the sequence $\{p_N/\rho_N^2, p_2/\rho_2^2, \dots, p_{N-1}/\rho_{N-1}^2, p_1/\rho_1^2, p_{N+1}/\rho_{N+1}^2, \dots\}$. Under the above construction, Theorem 2.5 follows immediately from Theorem 2.3(2).

The following lemma is auxiliary to Theorem 2.5, which provides conditions on the boundedness of u_n and $u_n \ell_n$.

LEMMA 2.8. *Let ℓ_n, u_n be constants in Theorem 2.5.*

(1) *If $\ell_n \leq \ell_{n+1}$, then*

$$u_n \rightarrow \infty \iff \sup_{n \geq 1} \frac{\log n}{\ell_n} = \infty.$$

(2) *Assume $\ell_n \geq \ell_{n+1}$. If $\ell_n/\ell_{n+1} \rightarrow 1$, then $u_n \ell_n \rightarrow \infty$. If $\liminf_{n \rightarrow \infty} \ell_n/\ell_{n+1} > 1$, then $u_n \ell_n = O(1)$.*

PROOF. (1) is obvious from the definition of u_n . For (2), we first consider the case $\ell_n/\ell_{n+1} \rightarrow 1$. Note that, for $m \geq 1$,

$$\liminf_{n \rightarrow \infty} u_n \ell_n \geq \liminf_{n \rightarrow \infty} \frac{\log(m+1)}{\ell_{n-m+1}/\ell_n} = \log(m+1).$$

Letting m tend to infinity gives the desired limit. Next, we consider the case $\liminf_{n \rightarrow \infty} \ell_n/\ell_{n+1} > 1$ and choose $N > 0$ and $M > 1$ such that $\ell_n/\ell_{n+1} \geq M$ for $n \geq N$. Immediately, this implies that $\ell_{n-m}/\ell_n \geq M^{m-N+1}$ for all $0 \leq m < n$ and $n \geq 1$. As a result, one has

$$u_n \ell_n = \max_{1 \leq i \leq n} \frac{\log(i+1)}{\ell_{n-i+1}/\ell_n} \leq M^N \sup_{i \geq 1} \frac{\log(i+1)}{M^i} < \infty.$$

□

The following corollary is a combination of Theorem 2.5 and Lemma 2.8, of which proof is obvious and skipped.

COROLLARY 2.9. *Let $\mathcal{G}^{\mathcal{P}}, E_n, \rho_n$ be as in Theorem 2.5. Assume that G_n has (A, d) -moderate growth with respect to E_n , $\inf\{Q_n(x) | x \in E_n, n \geq 1\} > 0$, $\rho_n \rightarrow \infty$ and $p_n \asymp \rho_n^2 \ell_n$ for some monotonic sequence $(\ell_n)_{n=1}^{\infty}$.*

- (1) If $\sup\{\log n/\ell_n | n \geq 1\} < \infty$ or $\liminf_{n \rightarrow \infty} \ell_n/\ell_{n+1} > 1$, then $\mathcal{G}_c^{\mathcal{P}}$ has no total variation cutoff.
- (2) If $\sup\{\log n/\ell_n | n \geq 1\} = \infty$ and $\lim_{n \rightarrow \infty} \ell_n/\ell_{n+1} = 1$, then $\mathcal{G}_c^{\mathcal{P}}$ has a total variation cutoff.

In particular, if $\ell_n = \exp\{-n^\gamma\}$ with $\gamma > 0$, then $\mathcal{G}_c^{\mathcal{P}}$ has a total variation cutoff if and only if $0 < \gamma < 1$. When E_n contains id for all $n \geq 1$, all above also holds for $\mathcal{G}^{\mathcal{P}}$.

2.3. Examples. In this subsection, we consider the randomized product in Proposition 1.3 for illustration of the results developed in Subsections 2.1-2.2. Recall that $\mathcal{G} = (\mathbb{Z}_{n+2}, Q_n, U_n)_{n=1}^\infty$, where $Q_n(0) = 1/2$ and $Q_n(1) = Q_n(-1) = 1/4$. It has been stated in the introduction that the diameter ρ_n of G_n w.r.t. $\{0, \pm 1\}$ is $\lfloor n/2 + 1 \rfloor$ and \mathbb{Z}_{n+2} has $(1, 1)$ -moderate growth w.r.t. $\{0, \pm 1\}$. As the randomness refers to the case that $\mathcal{P} = (p_n)_{n=1}^\infty$ is a sequence of positive random variables, we treat the specified cases separately in the following.

EXAMPLE 2.10 (Polynomial random sequences). Let X_1, X_2, \dots be i.i.d. positive random variables, $\gamma > 0$ and set $p_n = (X_1 + \dots + X_n)^\gamma$. Assume that the expectation μ of X_1 is finite. By the strong law of large numbers, one has

$$\frac{X_1 + \dots + X_n}{n} \sim \mu, \quad \text{almost surely,}$$

which implies

$$\frac{p_n}{\rho_n^2} \sim \ell_n := \mu^\gamma n^{\gamma-2}, \quad \text{almost surely.}$$

Clearly, $(\ell_n)_{n=1}^\infty$ is monotonic, $\ell_n/\ell_{n+1} \rightarrow 1$ and

$$\sup_{n \geq 1} \frac{\log n}{\ell_n} \begin{cases} < \infty & \text{for } \gamma > 2, \\ = \infty, & \text{for } 0 < \gamma \leq 2. \end{cases}$$

As a consequence of Corollary 2.9, if $0 < \gamma \leq 2$, then $\mathcal{G}^{\mathcal{P}}$ has a total variation cutoff almost surely; if $\gamma > 2$, then $\mathcal{G}^{\mathcal{P}}$ has no total variation cutoff with probability 1.

EXAMPLE 2.11 (Exponential random sequences). Let Y_1, Y_2, \dots be i.i.d. positive random variables and $p_n = Y_1 \times \dots \times Y_n$. For $n \geq 1$, let $(\ell_{n,i})_{i=1}^n$ be a non-decreasing arrangement of $(p_i/\rho_i^2)_{i=1}^n$ and set

$$t_n = \sup_{1 \leq i \leq n} \frac{\log(i+1)}{\ell_{n,i}}.$$

Using a similar reasoning as in the proof of Theorem 2.5, one may conclude using Theorem 2.3 that $\mathcal{G}^{\mathcal{P}}$ has a total variation cutoff if and only if $t_n \ell_{n,1} \rightarrow \infty$.

To analyze the product $t_n \ell_{n,1}$, we assume that the expectation ν of $\log Y_1$ is finite. By the strong law of large numbers, there is an event E with probability 1 such that

$$\nu_n := \frac{\log(p_n/\rho_n)}{n} \rightarrow \nu \quad \text{on } E.$$

In the following, we focus on the case $\nu > 0$. By writing $p_n/\rho_n^2 = e^{\nu n}$, one may select, for each $\omega \in E$, a constant $C(\omega) \in (0, 1)$ such that $p_n(\omega)/\rho_n^2 \geq C(\omega)e^{C(\omega)\nu n}$ for all $n \geq 1$. This implies that, on the event E ,

$$\ell_{n,1} \asymp 1, \quad \ell_{n,i} \geq Ce^{C\nu i}, \quad \forall 1 \leq i \leq n, n \geq 1.$$

Consequently, we obtain $t_n \ell_{n,1} = O(1)$ on E , which is equivalent to say that $\mathcal{G}^{\mathcal{P}}$ has no total variation cutoff with probability 1.

The results in the above discussion are summarized in Proposition 1.3.

3. Constructions of theoretical frameworks. This section is dedicated to proving Theorem 2.3. In the first subsection, we review those required but developed results in the introduction. In the second subsection, we treat the discrete time and continuous time cases separately and provide proofs in detail.

3.1. Review of technical supports. In this subsection, we survey those equivalences in Table 1 and state them by following the setting in the introduction. The first two propositions are supportive to the first two equivalences in Table 1 and, in fact, hold under more general assumptions.

PROPOSITION 3.1 (Theorems 3.1 and 3.3 in [5]). *Let $\mathcal{F} = (G_n, Q_n, U_n)_{n=1}^\infty$ be a family of random walks on finite groups and $\delta = \inf_n Q_n(id)$. Assume that $\delta > 0$ and, for some $\varepsilon_0 \in (0, 1)$, $T_{n,\text{TV}}(\varepsilon_0) \rightarrow \infty$ or $T_{n,\text{TV}}^{(c)}(\varepsilon_0) \rightarrow \infty$. Then, in the total variation, \mathcal{F} has a cutoff (resp. pre-cutoff) if and only if \mathcal{F}_c has a cutoff (resp. pre-cutoff). Furthermore, if \mathcal{F} or \mathcal{F}_c presents a total variation cutoff, then $T_{n,\text{TV}}(\varepsilon) \sim T_{n,\text{TV}}^{(c)}(\varepsilon)$ for all $\varepsilon \in (0, 1)$ and, for sequences of positive reals, $(t_n)_{n=1}^\infty$ and $(b_n)_{n=1}^\infty$, satisfying $b_n = o(t_n)$,*

$$|T_{n,\text{TV}}(\varepsilon) - t_n| = O(b_n), \quad \forall \varepsilon \in (0, 1) \quad \Leftrightarrow \quad |T_{n,\text{TV}}^{(c)}(\varepsilon) - t_n| = O(b_n), \quad \forall \varepsilon \in (0, 1).$$

PROPOSITION 3.2. *Let $\mathcal{F} = (G_n, Q_n, U_n)_{n=1}^\infty$ be a family of random walks on finite groups and let $T_{n,\text{TV}}^{(c)}$ and $T_{n,H}^{(c)}$ be the total variation and the Hellinger distance of the n th chain in \mathcal{F}_c . Then, \mathcal{F}_c has a total variation cutoff (resp. pre-cutoff) if and only if \mathcal{F}_c has a Hellinger distance cutoff (resp. pre-cutoff). Furthermore, if \mathcal{F}_c presents a cutoff in either measurement, then $T_{n,\text{TV}}^{(c)}(\varepsilon) \sim T_{n,H}^{(c)}(\varepsilon)$ for all $\varepsilon \in (0, 1)$.*

PROOF. The equivalence of cutoffs is already discussed in Proposition 1.1 of [2], while the equivalence of pre-cutoffs is given by (1.10). □

To analyze products of random walks, we need a variant of Proposition 1.1 in the Hellinger distance and, particularly, in the continuous time case.

PROPOSITION 3.3. *Let (G, Q, U) be a symmetric random walk on a finite group and $d_{\text{TV}}^{(c)}, d_H^{(c)}$ be the total variation and the Hellinger distance of its associated continuous time random walk. If G has (A, d) -moderate growth with respect to the support E of Q , then there is $C > 0$ depending only on A, d such that*

$$\frac{1}{2}e^{-Ct/\rho^2} \leq d_{\text{TV}}^{(c)}(t) \leq Ce^{-\eta t/(2\rho^2)}, \quad \forall t \geq 0,$$

and

$$\frac{1}{4}e^{-Ct/\rho^2} \leq d_H^{(c)}(t) \leq Ce^{-\eta t/(4\rho^2)}, \quad \forall t \geq 0,$$

where ρ is the diameter of (G, E) , $\eta = \min\{Q(x)|x \in E\}$ and both lower bounds require $\rho \geq 4$ in addition.

REMARK 3.4. (1) Compared with Proposition 1.1, the generating set E in Proposition 3.3 need not contain id and this means that the laziness of (G, Q, U) is not required at all. In fact, the laziness of a continuous time walk can be seen from the identity $\frac{1}{2}(K - I) = \frac{1}{2}(K + I) - I$, where K refers to the transition matrix determined by Q .

(2) The prerequisite of the lower bound on the graph diameter (at least 4) is due to the development of an upper bound on the spectral gap. See the proof of (3.2) in [7] for details.

PROOF OF PROPOSITION 3.3. First, we set $Q' = (Q + \mathbf{1}_{\{id\}})/2$ and $E' = E \cup \{id\}$. Let K, K' be the transition matrices determined by Q, Q' , set ρ, ρ' for the diameters of $(G, E), (G, E')$ and define $\eta = \min\{Q(x)|x \in E\}$ and $\eta' = \min\{Q'(x)|x \in E'\}$. Obviously, one has $K' = (K + I)/2$ and $H_t = H'_{2t}$, where $H_t = e^{-t(I-K)}$ and $H'_t = e^{-t(I-K')}$. Let $d'_{TV}, d_{TV}^{(c)}$ be the total variations of $(G, Q', U), (G, H_t, U)$. By applying Proposition 1.1 to (G, Q', U) , since G has (A, d) -moderate growth with respect to E (and, hence, with respect to E'), there is $C_2 > 0$ depending only on A, d such that

$$d'_{TV}(m) \leq C_2 e^{-\eta' m/(\rho')^2}, \quad \forall m \geq 0.$$

By the triangle inequality, this implies

$$d_{TV}^{(c)}(t) \leq e^{-2t} \sum_{m=0}^{\infty} \frac{(2t)^m}{m!} d'_{TV}(m) \leq C_2 \exp\{2t(e^{-\eta'/(\rho')^2} - 1)\} \leq C_2 e^{-\eta t/(2\rho^2)},$$

where the last inequality comes from $\rho' \leq \rho, \eta' \geq \eta/2$ and the fact that $e^{-u} \leq 1 - u/2$ for $u \in [0, 1]$. To see a lower bound of the total variation, let λ be the smallest nonzero eigenvalue of $I - K$. Note that $2d_{TV}^{(c)}(t) = \|H_t - \Pi\|_{\infty \rightarrow \infty}$, where $\Pi f := \pi(f)\mathbf{1}, \|L\|_{\infty \rightarrow \infty} := \sup\{\|Lf\|_{\infty} : \|f\|_{\infty} \leq 1\}$ and $\|f\|_{\infty} := \max_x |f(x)|$. By the symmetry of Q , this implies

$$d_{TV}^{(c)}(t) \geq \frac{1}{2}e^{-\lambda t}, \quad \forall t \geq 0.$$

Based on the (A, d) -moderate growth of (G, E) , Diaconis and Saloff-Coste showed in [7, Equation (3.2)] that if $\rho \geq 4$, then there is a constant $C_1 > 0$ depending only on A, d such that $\lambda \leq C_1/\rho^2$, where the assumption of $id \in E$ is in fact not required. This proves the desired bounds for the total variation with $C = \max\{C_1, C_2\}$, while the combination of (1.10) with such total variation bounds leads to bounds for the Hellinger distance. \square

Finally, we introduce the fourth equivalence in Table 1. Let $\mathcal{A} = \{a_{n,i} | 1 \leq i \leq k_n, n \geq 1\}$ and $\Lambda = \{\lambda_{n,i} | 1 \leq i \leq k_n, n \geq 1\}$ be triangular arrays of positive reals and set

$$(3.1) \quad \mathcal{F}(\mathcal{A}, \Lambda) = (f_n)_{n=1}^{\infty}, \quad f_n(t) = \sum_{i=1}^{k_n} a_{n,i} e^{-\lambda_{n,i} t}.$$

As f_n is nonnegative and decreasing, we define the mixing time of f_n by $T_n(\varepsilon) = \min\{t \geq 0 \mid f_n(t) \leq \varepsilon\}$ for $\varepsilon > 0$ and define the cutoff for $\mathcal{F}(\mathcal{A}, \Lambda)$ as follows.

DEFINITION 3.5. The family $\mathcal{F}(\mathcal{A}, \Lambda)$ is said to present a cutoff if there is a sequence $(t_n)_{n=1}^\infty$ of positive reals such that

$$\lim_{n \rightarrow \infty} f_n(at_n) = \begin{cases} 0 & \text{if } a > 1, \\ \infty & \text{if } 0 < a < 1. \end{cases}$$

In the above, $(t_n)_{n=1}^\infty$ or briefly t_n is called a cutoff time.

REMARK 3.6. It is easy to check from Definition 3.5 that $\mathcal{F}(\mathcal{A}, \Lambda)$ has a cutoff if and only if $T_n(\varepsilon) \sim T_n(\delta)$ for all $\varepsilon > 0$ and $\delta > 0$. In particular, if $\mathcal{F}(\mathcal{A}, \Lambda)$ has a cutoff, then $T_n(\varepsilon)$ is a cutoff time for all $\varepsilon > 0$.

By expressing f_n as a Laplace transform of some positive measure, the authors of [4] provided a criterion (Theorems 3.5 and 3.8 in [4]) to determine the cutoff for $\mathcal{F}(\mathcal{A}, \Lambda)$. Later, such a method was refined in [1, Theorem 2.4]. To see the details, we set, for $c > 0$,

$$(3.2) \quad \lambda_n(c) = \lambda_{n, j_n(c)}, \quad \tau_n(c) = \max_{i \geq j_n(c)} \frac{\log(1 + a_{n,1} + \cdots + a_{n,i})}{\lambda_{n,i}},$$

where $j_n(c) := \min\{i \geq 1 \mid a_{n,1} + \cdots + a_{n,i} > c\}$.

PROPOSITION 3.7 (Theorem 2.4 in [1]). *Let \mathcal{F} be the family in (3.1), $T_n(\varepsilon)$ be the mixing time of f_n and λ_n, τ_n be the quantities in (3.2). Then, the following are equivalent.*

- (1) $\mathcal{F}(\mathcal{A}, \Lambda)$ has a cutoff.
- (2) There is $\varepsilon > 0$ such that $T_n(\varepsilon)\lambda_n(c) \rightarrow \infty$ for all $c > 0$.
- (3) $\tau_n(c)\lambda_n(c) \rightarrow \infty$ for all $c > 0$.

In particular, $\tau_n(c)$ is a cutoff time for all $c > 0$.

REMARK 3.8. It was shown in [1, Lemma 2.5] that, if $\tau_n(c)\lambda_n(c) \rightarrow \infty$, then $\tau_n(c')\lambda_n(c') \rightarrow \infty$ for all $c' > c$.

3.2. Proof of Theorem 2.3. The proof of Theorem 2.3 is based on some crucial techniques, of which proofs are either developed or involved and are addressed in the appendix for reference. See Lemmas A.1, A.2 and A.3 for details.

PROOF OF THEOREM 2.3 (The continuous time case). To prove this proposition, it suffices to consider, by Proposition 3.2 and (1.10), the Hellinger distance. Let $d_{n,H}^{(c)}$ and $d_{n,i,H}^{(c)}$ be the Hellinger distances of the n th and (n, i) th random walks in $\mathcal{F}_c^{\mathcal{P}}$ and \mathcal{F}_c , and set $\eta = \inf\{Q_{n,i}(x) \mid x \in E_{n,i}, 1 \leq i \leq k_n, n \geq 1\}$. By Proposition 3.3, there is $C_1 > 1$ such that, for all $1 \leq i \leq k_n$ and $n \geq 1$,

$$(3.3) \quad \frac{1}{4}e^{-C_1 t / \rho_{n,i}^2} \leq d_{n,i,H}^{(c)}(t) \leq C_1 e^{-\eta t / (4\rho_{n,i}^2)}, \quad \forall t \geq 0,$$

where $\rho_{n,i} \geq 4$ is required for the first inequality.

For (1), set $M = \sup_n k_n$. By (3.3), one has

$$d_{n,i,H}^{(c)}(ap_{n,i}t_n/q_n) \leq C_1 e^{-a\eta\ell_{n,i}t_n/(4C)} \leq C_1(i+1)^{-a\eta/(4C)} \leq C_1 2^{-a\eta/(4C)},$$

and, in addition with the fact $t_n \leq \log(M+1)/\ell_{n,1}$,

$$d_{n,1,H}^{(c)}(ap_{n,1}t_n/q_n) \geq \frac{1}{4} e^{-aC_1 C \ell_{n,1} t_n} \geq \frac{1}{4} (M+1)^{-aC_1 C}.$$

Consequently, the replacement of A, p_i with $1/\sqrt{2}, p_{n,i}/q_n$ in Lemma A.1 yields

$$1 - \exp\left\{-\frac{1}{16}(M+1)^{-2aC_1 C}\right\} \leq d_{n,H}^{(c)}(at_n)^2 \leq 1 - \exp\left\{-MC_1^2 2^{1-a\eta/(2C)}\right\},$$

for all $a > A := (4C/\eta)(\log_2 C_1 + 1/2)$ and $n \geq 1$. This proves (1).

For (2), note that (i) \Rightarrow (ii) is clear from the definition of cutoffs and pre-cutoffs, and (iv) \Rightarrow (v) is trivial. To prove the other equivalences, we first make some analysis on $d_{n,H}^{(c)}$. Let C_1 be the constant in (3.3), A be the constant defined as above and N be a positive integer such that $\rho_{n,i} \geq 4$ for $i \geq N$ and $n \geq 1$. In a similar reasoning as before, one can show that

$$(3.4) \quad d_{n,H}^{(c)}(t)^2 \geq 1 - \exp\left\{-\sum_{i=1}^{k_n} d_{n,i,H}^{(c)}(p_{n,i}t/q_n)^2\right\}, \quad \forall t > 0,$$

and

$$(3.5) \quad d_{n,H}^{(c)}(t)^2 \leq 1 - \exp\left\{-2\sum_{i=1}^{k_n} d_{n,i,H}^{(c)}(p_{n,i}t/q_n)^2\right\}, \quad \forall t > At_n.$$

By (3.3), we have

$$\sum_{i=1}^{k_n} d_{n,i,H}^{(c)}(p_{n,i}t/q_n)^2 \leq C_1^2 \sum_{i=1}^{k_n} e^{-\eta p_{n,i}t/(2q_n \rho_{n,i}^2)} \leq C_1^2 \sum_{i=1}^{k_n} e^{-\eta t \ell_{n,i}/(2C)}$$

and

$$\sum_{i \in I_n} d_{n,i,H}^{(c)}(p_{n,i}t/q_n)^2 \geq \frac{1}{16} \sum_{i \in I_n} e^{-2C_1 p_{n,i}t/\rho_{n,i}^2} \geq \frac{1}{16} \sum_{i \in I_n} e^{-2CC_1 t \ell_{n,i}},$$

where $I_n = \{1 \leq i \leq k_n | \rho_{n,i} \geq 4\}$. Putting the last terms in the above computations back to (3.4) and (3.5) yields

$$(3.6) \quad 1 - \exp\left\{-\frac{1}{16}g_n(2CC_1t)\right\} \leq d_{n,H}^{(c)}(t)^2 \leq 1 - \exp\left\{-2C_1^2 f_n(\eta t/(2C))\right\},$$

where $f_n(t) = \sum_{i=1}^{k_n} e^{-\ell_{n,i}t}$, $g_n(t) = \sum_{i \in I_n} e^{-\ell_{n,i}t}$ and the second inequality holds for $t > At_n$. We are now ready to proceed the proof of (ii) \Rightarrow (iii) \Rightarrow (iv) and (v) \Rightarrow (i).

To see (ii) \Rightarrow (iii), assume that $\mathcal{F}_c^{\mathcal{P}}$ presents a pre-cutoff in the Hellinger distance and let $s_n > 0$ and $B_2 > B_1 > 0$ be such that

$$(3.7) \quad \lim_{n \rightarrow \infty} d_{n,H}^{(c)}(B_2 s_n) = 0, \quad \liminf_{n \rightarrow \infty} d_{n,H}^{(c)}(B_1 s_n) = \alpha > 0.$$

Note that

$$f_n(at_n) \leq \sum_{i=1}^{k_n} (i+1)^{-a} \leq \int_1^\infty u^{-a} du = \frac{1}{a-1}, \quad \forall a > 1.$$

By the second inequality of (3.6) and the fact $A > 2C/\eta$, we have

$$(3.8) \quad d_{n,H}^{(c)}(at_n)^2 \leq 1 - \exp\left\{-\frac{2C_1^2}{a\eta/(2C) - 1}\right\}, \quad \forall a > A, n \geq 1.$$

Next, let's fix $a > (2C_1^2/\alpha^2 + 1)A$. By the fact of $A > 2C/\eta$, one may derive $a\eta/(2C) - 1 > 2C_1^2/\alpha^2$ and, by (3.8), this leads to $d_{n,H}^{(c)}(at_n)^2 \leq 1 - e^{-\alpha^2} < \alpha^2$. As a result of the second limit in (3.7), we obtain that $at_n \geq B_1s_n$ for n large enough. In addition with the fact $1 \in I_n$ for all $n \geq 1$, we may conclude from the first inequality of (3.6) that

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} d_{n,H}^{(c)}(B_2s_n)^2 \geq \lim_{n \rightarrow \infty} d_{n,H}^{(c)}(aB_2t_n/B_1)^2 \\ &\geq 1 - \exp\left\{-\frac{1}{16} \limsup_{n \rightarrow \infty} e^{-2aB_2CC_1t_n\ell_{n,1}/B_1}\right\}, \end{aligned}$$

which leads to (iii).

For (iii) \Rightarrow (iv), assume that $t_n\ell_{n,1} \rightarrow \infty$. By Proposition 3.7 and Remark 3.8, the family $(f_n)_{n=1}^\infty$ has a cutoff with cutoff time t_n . By (3.6), this implies

$$\lim_{n \rightarrow \infty} d_{n,H}^{(c)}(at_n) = 0, \quad \forall a > A, \quad \lim_{n \rightarrow \infty} d_{n,H}^{(c)}(at_n) = 1, \quad \forall 0 < a < 1/(2CC_1),$$

where the second limit also uses the fact

$$g_n(t) = f_n(t) - (f_n(t) - g_n(t)) \geq f_n(t) - Ne^{-\ell_{n,1}t}.$$

As a consequence, when $\varepsilon \in (0, 1)$, one has $t_n/(4CC_1) \leq T_{n,H}^{(c)}(\varepsilon) \leq 2t_n$ for n large enough, which gives (iv) and the order of the mixing time.

To show (v) \Rightarrow (i), it suffices to prove $T_{n,H}^{(c)}(\varepsilon) - T_{n,H}^{(c)}(\delta) = O(1/\ell_{n,1})$ for all $0 < \varepsilon < \delta < 1$, which is exactly the specific conclusion in (2). First, we need a refinement of (3.5). Let $\delta \in (0, 1)$. In Lemma A.1, the first inequality implies $T_{n,H}^{(c)}(\delta) \geq \max\{T_{n,i,H}^{(c)}(\delta)q_n/p_{n,i} | 1 \leq i \leq k_n\}$, while the third inequality yields

$$d_{n,H}^{(c)}(t)^2 \leq 1 - \exp\left\{-\frac{1}{1-\delta^2} \sum_{i=1}^{k_n} d_{n,i,H}^{(c)}\left(\frac{p_{n,i}t}{q_n}\right)^2\right\}, \quad \forall t \geq T_{n,H}^{(c)}(\delta).$$

Let $t = T_{n,H}^{(c)}(\delta) + a/\ell_{n,1}$ with $a > 0$. By the quasi-submultiplicativity in Lemma A.3, one has

$$d_{n,i,H}^{(c)}\left(\frac{p_{n,i}t}{q_n}\right) \leq 4d_{n,i,H}^{(c)}\left(\frac{p_{n,i}T_{n,H}^{(c)}(\delta)}{q_n}\right) d_{n,i,H}^{(c)}\left(\frac{ap_{n,i}}{\ell_{n,1}q_n}\right).$$

Putting this back to the upper bound for $d_{n,H}^{(c)}(t)$ yields

$$d_{n,H}^{(c)}(t)^2 \leq 1 - \exp \left\{ -\frac{16}{1-\delta^2} \sum_{i=1}^{k_n} d_{n,i,H}^{(c)} \left(\frac{p_{n,i} T_{n,H}^{(c)}(\delta)}{q_n} \right)^2 d_{n,i,H}^{(c)} \left(\frac{ap_{n,i}}{\ell_{n,1} q_n} \right)^2 \right\}.$$

Observe that the second inequality in (3.3) and (3.4) give

$$d_{n,i,H}^{(c)} \left(\frac{ap_{n,i}}{\ell_{n,1} q_n} \right)^2 \leq C_1^2 e^{-a\eta/(2C)}, \quad \sum_{i=1}^{k_n} d_{n,i,H}^{(c)} \left(\frac{p_{n,i} T_{n,H}^{(c)}(\delta)}{q_n} \right)^2 \leq \log \frac{1}{1-\delta^2}.$$

Combining the last three inequalities leads to

$$d_{n,H}^{(c)} \left(T_{n,H}^{(c)}(\delta) + a/\ell_{n,1} \right)^2 \leq 1 - \exp \left\{ -\frac{C_1^2 e^{-a\eta/(2C)}}{1-\delta^2} \log \frac{1}{1-\delta^2} \right\}, \quad \forall a > 0.$$

As the right hand side tends to 0 as a tends to infinity, we obtain $T_{n,H}^{(c)}(\varepsilon) - T_{n,H}^{(c)}(\delta) = O(1/\ell_{n,1})$ for $\varepsilon \in (0, \delta)$, as desired. This finishes the proof of (2). \square

PROOF OF THEOREM 2.3 (The discrete time case). We shall prove the discrete time case by identifying the items in (2) with the continuous time case. First, we show that $T_{n,TV}(\varepsilon) \rightarrow \infty$ and $T_{n,TV}^{(c)}(\varepsilon) \rightarrow \infty$ for some $\varepsilon \in (0, 1)$. Let $d_{n,i,TV}^{(c)}$ be the total variation of the (n, i) th random walk in \mathcal{F}_c and let $N > 0$ be a positive integer such that $\rho_{n,i} \geq 4$ for $i \geq N$ and $n \geq 1$. By Proposition 3.3, there is $C_2 > 0$ such that

$$d_{n,i,TV}^{(c)}(t) \geq \frac{1}{2} e^{-C_2 t / \rho_{n,i}^2} \geq \frac{1}{2} e^{-C_2 t / 16}, \quad \forall t \geq 0, i \geq N, n \geq 1.$$

Note that the first inequality in Lemma A.1 also holds for the total variation (see [2, Proposition 3.2]) and this implies that, for $k_n > N$,

$$\begin{aligned} d_{n,TV}^{(c)}(t) &\geq \max_{N \leq i \leq k_n} d_{n,i,TV}^{(c)} \left(\frac{p_{n,i} t}{q_n} \right) \geq \frac{1}{2} \left\{ -\frac{C_2 t}{16} \min_{N \leq i \leq k_n} \frac{p_{n,i}}{q_n} \right\} \\ &\geq \frac{1}{2} \exp \left\{ -\frac{C_2 t}{16(k_n - N)} \sum_{i=N}^{k_n} \frac{p_{n,i}}{q_n} \right\} \geq \frac{1}{2} \exp \left\{ -\frac{C_2 t}{16(k_n - N)} \right\}, \end{aligned}$$

where the last inequality uses the fact that $(p_{n,i}/q_n)_{i=1}^{k_n}$ is a probability vector. As $k_n \rightarrow \infty$, one has $T_{n,TV}^{(c)}(1/4) \geq 8(k_n - N)/C_2$ for n large enough, which yields $T_{n,TV}^{(c)}(1/4) \rightarrow \infty$. In the discrete time case, observe that, by the triangle inequality,

$$d_{n,TV}^{(c)}(t) \leq \sum_{m=0}^{\infty} e^{-t} \frac{t^m}{m!} d_{n,TV}(m) \leq \sum_{m=0}^{\ell} e^{-t} \frac{t^m}{m!} + \left(\sum_{m=\ell+1}^{\infty} e^{-t} \frac{t^m}{m!} \right) d_{n,TV}(\ell).$$

When $t = T_{n,TV}^{(c)}(1/4)$ and $\ell = \lceil t/2 \rceil$, it's easy to check (or to see from [3, Lemma A.1]) that

$$\lim_{n \rightarrow \infty} e^{-t} \sum_{m=0}^{\ell} \frac{t^m}{m!} = 0,$$

which leads to

$$\lim_{n \rightarrow \infty} d_{n,\text{TV}} \left(\left\lceil \frac{1}{2} T_{n,\text{TV}}^{(c)}(1/4) \right\rceil \right) \geq \frac{1}{4}.$$

Consequently, we obtain $T_{n,\text{TV}}(1/5) \geq \frac{1}{2} T_{n,\text{TV}}^{(c)}(1/4)$ for n large enough and, thus, $T_{n,\text{TV}}(1/5) \rightarrow \infty$. Now, we are ready to prove (2) for the discrete time case.

Let $(*)'$ with $*$ \in {i, ii, iv, v} be respectively the corresponding statements for $\mathcal{F}^{\mathcal{P}}$ in Theorem 2.3(2). Immediately, the equivalence of $(*)$ and $(*)'$ with $*$ \in {i, ii} is given by Proposition 3.1. Let μ_n and $\mu_{n,1}$ be the second largest eigenvalues of the transition matrices determined by Q_n and $Q_{n,1}$. It is easy to check that $1 - \mu_n \leq (p_{n,1}/q_n)(1 - \mu_{n,1})$. A similarly reasoning as in the proof of Proposition 3.3 implies

$$d_{n,\text{TV}}(m) \geq \mu_n^m, \quad 1 - \mu_{n,1} \leq C_3/\rho_{n,1}^2, \quad \forall m \geq 0, n \geq 1,$$

where C_3 is a positive constant depending on A, d . As a result, this yields

$$d_{n,\text{TV}}(m) \geq e^{-m(1-\mu_n)} \geq e^{-C_3 m p_{n,1}/(q_n \rho_{n,1}^2)} \geq e^{-CC_3 m \ell_{n,1}}.$$

When $\mathcal{F}^{\mathcal{P}}$ has a total variation cutoff, one has

$$\exp \left\{ -2CC_3 \liminf_{n \rightarrow \infty} T_{n,\text{TV}}(\varepsilon) \ell_{n,1} \right\} \leq \lim_{n \rightarrow \infty} d_{n,\text{TV}}(2T_{n,\text{TV}}(\varepsilon)) = 0, \quad \forall \varepsilon \in (0, 1).$$

This proves (i)' \Rightarrow (iv)'.

Based on the above discussions, it remains to show (v)' \Rightarrow (v). Assume that $T_{n,\text{TV}}(\varepsilon_1) \ell_{n,1} \rightarrow \infty$ for some $\varepsilon_1 \in (0, 1)$. We will prove (v) by contradiction and thus assume the inverse that $T_{\xi_n,\text{TV}}^{(c)}(\varepsilon_2) \ell_{\xi_n,1} = O(1)$, where $\varepsilon_2 \in (0, 1)$ and $(\xi_n)_{n=1}^\infty$ is an increasing sequence of positive integers. Note that we may restrict ourselves to the case of $\varepsilon_2 \leq 1/4$. Set $r_n = \sqrt{T_{\xi_n,\text{TV}}^{(c)}(\varepsilon_1)/\ell_{\xi_n,1}}$. Obviously, $T_{\xi_n,\text{TV}}^{(c)}(\varepsilon_2) = o(r_n)$ and $r_n = o(T_{\xi_n,\text{TV}}(\varepsilon_1))$. By the quasi-submultiplicativity of the total variation, one has

$$\begin{aligned} \limsup_{n \rightarrow \infty} 2d_{\xi_n,\text{TV}}^{(c)}(r_n) &\leq \limsup_{n \rightarrow \infty} \left(2d_{\xi_n,\text{TV}} \left(T_{\xi_n,\text{TV}}^{(c)}(\varepsilon_2) \right) \right)^{\lceil r_n/T_{\xi_n,\text{TV}}^{(c)}(\varepsilon_2) \rceil} \\ &\leq \limsup_{n \rightarrow \infty} 2^{-\lceil r_n/T_{\xi_n,\text{TV}}^{(c)}(\varepsilon_2) \rceil} = 0. \end{aligned}$$

As a consequence of Lemma A.2(3), this implies

$$\lim_{n \rightarrow \infty} d_{\xi_n,\text{TV}}(\lceil ar_n \rceil) = 0, \quad \forall a > 1.$$

But, however, as $ar_n = o(T_{\xi_n,\text{TV}}(\varepsilon_1))$, we have

$$\liminf_{n \rightarrow \infty} d_{\xi_n,\text{TV}}(\lceil ar_n \rceil) \geq \liminf_{n \rightarrow \infty} d_{\xi_n,\text{TV}} \left(T_{\xi_n,\text{TV}}(\varepsilon_1) - 1 \right) \geq \varepsilon_1 > 0.$$

This makes a contradiction and, hence, $T_{n,\text{TV}}(\varepsilon_2) \ell_{n,1} \rightarrow \infty$, as desired. □

Appendix A. Techniques and auxiliary results.

LEMMA A.1 ([2, Proposition 3.2]). *Let $(G_i, Q_i, U_i)_{i=1}^n$ be random walks on finite groups and (G, Q, U) be their product according to the probability vector (p_1, \dots, p_n) . Let $d_{i,H}^{(c)}$ and $d_H^{(c)}$ be the Hellinger distances of the continuous time random walks associated with (G_i, Q_i, U_i) and (G, Q, U) . Then, one has*

$$d_H^{(c)}(t) \geq \max \left\{ d_{i,H}^{(c)}(p_i t) \mid 1 \leq i \leq n \right\},$$

and

$$1 - \exp \left\{ - \sum_{i=1}^n d_{i,H}^{(c)}(p_i t)^2 \right\} \leq d_H^{(c)}(t)^2 \leq 1 - \exp \left\{ - \frac{1}{1 - A^2} \sum_{i=1}^n d_{i,H}^{(c)}(p_i t)^2 \right\},$$

where the second inequality requires $t \geq \max \{ T_{i,H}^{(c)}(A)/p_i \mid 1 \leq i \leq n \}$ with $A \in (0, 1)$.

LEMMA A.2 ([5, Proposition 3.1]). *Let $\mathcal{F} = (G_n, Q_n, U_n)_{n=1}^\infty$ be a family of random walks on finite groups and \mathcal{F}_c be the family of continuous time random walks associated with \mathcal{F} . For $n \geq 1$, let $d_{n,TV}$, $d_{n,TV}^{(c)}$ be the total variations of the n th random walks in \mathcal{F} , \mathcal{F}_c . Suppose $\inf_{n \geq 1} Q_n(id) > 0$. Then, for any sequence $(t_n)_{n=1}^\infty$ tending to infinity,*

- (1) $d_{n,TV}(\lfloor t_n \rfloor) = 1$ if and only if $d_{n,TV}^{(c)}(t_n) = 1$.
- (2) If $d_{n,TV}(\lceil t_n \rceil) = 0$, then $d_{n,TV}^{(c)}(at_n) \rightarrow 0$ for all $a > 1$.
- (3) If $d_{n,TV}^{(c)}(t_n) \rightarrow 0$, then $d_{n,TV}(\lceil at_n \rceil) \rightarrow 0$ for all $a > 1$.

LEMMA A.3. *Consider an irreducible Markov chain on a finite or countable set \mathcal{X} with transition matrix K and stationary distribution π . Set $H_t = e^{-t(I-K)}$ and let $d_H, d_H^{(c)}$ be the maximum Hellinger distances defined by*

$$d_H(m) = \sup_{x \in \mathcal{X}} \left(\frac{1}{2} \sum_{y \in \mathcal{X}} (\sqrt{K^m(x, y)} - \sqrt{\pi(y)})^2 \right)^{1/2},$$

and

$$d_H^{(c)}(t) = \sup_{x \in \mathcal{X}} \left(\frac{1}{2} \sum_{y \in \mathcal{X}} (\sqrt{H_t(x, y)} - \sqrt{\pi(y)})^2 \right)^{1/2}.$$

Then, the mappings

$$m \mapsto 4d_H(m), \quad t \mapsto 4d_H^{(c)}(t),$$

are non-increasing and submultiplicative.

PROOF. We deal with the discrete time case, while the continuous time case can be shown in a similar way. Let n, m be positive integers and $x, y \in \mathcal{X}$. Note that

$$\begin{aligned} \sqrt{K^{n+m}(x, y)} - \sqrt{\pi(y)} &= \frac{\sum_{z \in \mathcal{X}} [K^n(x, z) - \pi(z)][K^m(z, y) - \pi(y)]}{\sqrt{K^{n+m}(x, y)} + \sqrt{\pi(y)}} \\ (A.1) \quad &= \sum_{z \in \mathcal{X}} A_z B_z, \end{aligned}$$

where

$$A_z = \left(\sqrt{K^n(x, z)} - \sqrt{\pi(z)} \right) \left(\sqrt{K^m(z, y)} - \sqrt{\pi(y)} \right)$$

and

$$B_z = \frac{\left(\sqrt{K^n(x, z)} + \sqrt{\pi(z)} \right) \left(\sqrt{K^m(z, y)} + \sqrt{\pi(y)} \right)}{\sqrt{K^{n+m}(x, y)} + \sqrt{\pi(y)}}.$$

Consider the following two cases.

Case 1: $K^{n+m}(x, y) \geq \pi(y)$. In this case, one may apply the Cauchy-Schwarz inequality to obtain

$$\left| \sqrt{K^{n+m}(x, y)} - \sqrt{\pi(y)} \right|^2 \leq \left(\sum_{z \in \mathcal{X}} A_z^2 \right) \left(\sum_{z \in \mathcal{X}} B_z^2 \right).$$

Note that $(\sqrt{a} + \sqrt{b})^2 \leq 2(a + b)$ for all $a, b \geq 0$. As a result, this implies

$$\begin{aligned} \sum_z B_z^2 &\leq \frac{4 \sum_{z \in \mathcal{X}} (K^n(x, z) + \pi(z))(K^m(z, y) + \pi(y))}{\left(\sqrt{K^{n+m}(x, y)} + \sqrt{\pi(y)} \right)^2} \\ &= \frac{4[K^{n+m}(x, y) + 3\pi(y)]}{K^{n+m}(x, y) + \pi(y) + 2\sqrt{K^{n+m}(x, y)\pi(y)}} \leq 4, \end{aligned}$$

and, hence,

$$\left| \sqrt{K^{n+m}(x, y)} - \sqrt{\pi(y)} \right|^2 \leq 4 \left(\sum_{z \in \mathcal{X}} A_z^2 \right).$$

Case 2: $K^{n+m}(x, y) < \pi(y)$. For this case, set

$$E_1 = \{z | \pi(z) > K^n(x, z), K^m(z, y) > \pi(y)\}$$

and

$$E_2 = \{z | \pi(z) < K^n(x, z), K^m(z, y) < \pi(y)\}.$$

By (A.1), one has

$$\begin{aligned} 0 < \sqrt{\pi(y)} - \sqrt{K^{n+m}(x, y)} &\leq \frac{\sum_{z \in E_1 \cup E_2} [\pi(z) - K^n(x, z)][K^m(z, y) - \pi(y)]}{\sqrt{K^{n+m}(x, y)} + \sqrt{\pi(y)}} \\ &= \sum_{z \in E_1 \cup E_2} A_z B_z. \end{aligned}$$

As before, we may apply the Cauchy-Schwarz inequality to get

$$\left| \sqrt{K^{n+m}(x, y)} - \sqrt{\pi(y)} \right|^2 \leq \left(\sum_{z \in E_1 \cup E_2} A_z^2 \right) \left(\sum_{z \in E_1 \cup E_2} B_z^2 \right).$$

Note that, for $z \in E_1$,

$$\begin{aligned} & (\sqrt{K^n(x, z)} + \sqrt{\pi(z)})^2 (\sqrt{K^m(z, y)} + \sqrt{\pi(y)})^2 \\ & \leq 4 (\sqrt{K^n(x, z)} + \sqrt{\pi(z)})^2 K^m(z, y) = 4 \left(\sqrt{\frac{K^n(x, z)}{\pi(z)}} + 1 \right)^2 \pi(z) K^m(z, y), \end{aligned}$$

and, for $z \in E_2$,

$$\begin{aligned} & (\sqrt{K^n(x, z)} + \sqrt{\pi(z)})^2 (\sqrt{K^m(z, y)} + \sqrt{\pi(y)})^2 \\ & \leq 4K^n(x, z) (\sqrt{K^m(z, y)} + \sqrt{\pi(y)})^2. \end{aligned}$$

By the Jensen inequality, this implies

$$\begin{aligned} & \sum_{z \in E_1} (\sqrt{K^n(x, z)} + \sqrt{\pi(z)})^2 (\sqrt{K^m(z, y)} + \sqrt{\pi(y)})^2 \\ \text{(A.2)} \quad & \leq 4 \left(\sqrt{\sum_{z \in E_1} K^n(x, z) K^m(z, y)} + \sqrt{c_1 \pi(y)} \right)^2, \end{aligned}$$

where $c_1 = \sum_{z \in E_1} \pi(z) K^m(z, y) / \pi(y)$, and

$$\begin{aligned} & \sum_{z \in E_2} (\sqrt{K^n(x, z)} + \sqrt{\pi(z)})^2 (\sqrt{K^m(z, y)} + \sqrt{\pi(y)})^2 \\ \text{(A.3)} \quad & \leq 4 \left(\sqrt{\sum_{z \in E_2} K^n(x, z) K^m(z, y)} + \sqrt{c_2 \pi(y)} \right)^2, \end{aligned}$$

where $c_2 = \sum_{z \in E_2} K^n(x, z)$. Summing up (A.2) and (A.3) yields

$$\sum_{z \in E_1 \cup E_2} B_z^2 \leq \frac{4 \left(\sqrt{\sum_{z \in E_1 \cup E_2} K^n(x, z) K^m(z, y)} + \sqrt{(c_1 + c_2) \pi(y)} \right)^2}{(\sqrt{K^{n+m}(x, y)} + \sqrt{\pi(y)})^2} \leq 8,$$

while the last inequality uses the fact of $c_1 \leq 1$ and $c_2 \leq 1$. As a consequence, this leads to

$$\left| \sqrt{K^{n+m}(x, y)} - \sqrt{\pi(y)} \right|^2 \leq 8 \left(\sum_{z \in E_1 \cup E_2} A_z^2 \right) \leq 8 \sum_{z \in \mathcal{X}} A_z^2,$$

which is also applicable for Case 1.

Based on the result in the above discussions, we obtain

$$\begin{aligned} 4d_H(n + m) &= \sup_{x \in \mathcal{X}} \left\{ 8 \sum_{y \in \mathcal{X}} (\sqrt{K^{n+m}(x, y)} - \sqrt{\pi(y)})^2 \right\}^{1/2} \\ &\leq \sup_{x \in \mathcal{X}} \left\{ \sum_{y, z \in \mathcal{X}} 8 (\sqrt{K^n(x, z)} - \sqrt{\pi(z)})^2 \times 8 (\sqrt{K^m(z, y)} - \sqrt{\pi(y)})^2 \right\}^{1/2} \end{aligned}$$

$$\leq \sup_{x \in \mathcal{X}} \left\{ 8 \left(\sqrt{K^n(x, z)} - \sqrt{\pi(z)} \right)^2 \right\}^{1/2} \times 4d_H(m) = 4d_H(n) \times 4d_H(m).$$

□

Acknowledgments. The authors thank an anonymous referee for a careful reading of the manuscript and helpful comments.

REFERENCES

- [1] G.-Y. CHEN, J.-M. HSU AND Y.-C. SHEU, The L^2 -cutoff for reversible Markov chains, *Ann. Appl. Probab.* 27 (2017), no. 4, 2305–2341.
- [2] G.-Y. CHEN AND T. KUMAGAI, Cutoffs for product chains, *Stochastic Process. Appl.* 128 (2018), no. 11, 3840–3879.
- [3] G.-Y. CHEN AND L. SALOFF-COSTE, The cutoff phenomenon for ergodic Markov processes, *Electron. J. Probab.* 13 (2008), no. 3, 26–78.
- [4] G.-Y. CHEN AND L. SALOFF-COSTE, The L^2 -cutoff for reversible Markov processes, *J. Funct. Anal.* 258 (2010), no. 7, 2246–2315.
- [5] G.-Y. CHEN AND L. SALOFF-COSTE, Comparison of cutoffs between lazy walks and Markovian semigroups, *J. Appl. Probab.* 50 (2013), no. 4, 943–959.
- [6] P. DIACONIS, Group representations in probability and statistics, Institute of Mathematical Statistics Lecture Notes—Monograph Series, 11, Institute of Mathematical Statistics, Hayward, CA, 1988.
- [7] P. DIACONIS AND L. SALOFF-COSTE, Moderate growth and random walk on finite groups, *Geom. Funct. Anal.* 4 (1994), no. 1, 1–36.
- [8] H. LACONIN, A product chain without cutoff, *Electron. Commun. Probab.* 20 (2015), no. 19, 9.
- [9] D.A. LEVIN, Y. PERES AND E.L. WILMER, Markov chains and mixing times, American Mathematical Society, Providence, RI, 2009.
- [10] L. SALOFF-COSTE, Lectures on finite Markov chains, In *Lectures on probability theory and statistics* (Saint-Flour, 1996), *Lecture Notes in Math.* 1665, pp. 301–413, Springer, Berlin, 1997.
- [11] L. SALOFF-COSTE, Random walks on finite groups, In *Probability on discrete structures*, *Encyclopaedia Math. Sci.* 110, pp. 263–346, Springer, Berlin, 2004.
- [12] A.N. SHIRYAEV, *Probability*, Graduate Texts in Mathematics 95, Springer-Verlag, New York, second edition, 1996. Translated from the first (1980) Russian edition by R.P. Boas.

DEPARTMENT OF APPLIED MATHEMATICS
NATIONAL CHIAO TUNG UNIVERSITY
HSINCHU 300
TAIWAN

E-mail address: gychen@math.nctu.edu.tw

RIMS
KYOTO UNIVERSITY
KYOTO 606–8502
JAPAN

E-mail address: kumagai@kurims.kyoto-u.ac.jp