

Pseudo-Anosovs on closed surfaces having small entropy and the Whitehead sister link exterior

By Eiko KIN and Mitsuhiro TAKASAWA

(Received Aug. 15, 2011)

Abstract. We denote by δ_g (resp. δ_g^+), the minimal dilatation for pseudo-Anosovs (resp. pseudo-Anosovs with orientable invariant foliations) on a closed surface of genus g . This paper concerns the pseudo-Anosovs which occur as monodromies of fibrations on manifolds obtained from the Whitehead sister link exterior W by Dehn filling two cusps, where the fillings are on the boundary slopes of fibers of W . We give upper bounds of δ_g for $g \equiv 0, 1, 5, 6, 7, 9 \pmod{10}$, δ_g^+ for $g \equiv 1, 5, 7, 9 \pmod{10}$. Our bounds improve the previous one given by Hironaka. We note that the monodromies of fibrations on W were also studied by Aaber and Dunfield independently.

1. Introduction.

Let $\text{Mod}(\Sigma)$ be the mapping class group on an orientable surface Σ . An element $\phi \in \text{Mod}(\Sigma)$ which contains a pseudo-Anosov homeomorphism $\Phi : \Sigma \rightarrow \Sigma$ as a representative is called a pseudo-Anosov mapping class. There are two numerical invariants for pseudo-Anosov mapping classes. One is the dilatation $\lambda(\phi) > 1$ (or the entropy $\text{ent}(\phi) = \log \lambda(\phi)$) which is defined to be the dilatation $\lambda(\Phi)$ of Φ , and the other is the hyperbolic volume $\text{vol}(\phi) = \text{vol}(\mathbb{T}(\phi))$ of the mapping torus $\mathbb{T}(\phi)$. It is natural to ask whether there is a relation between $\text{ent}(\phi)$ and $\text{vol}(\phi)$. Computer experiments in [13] tell us that if we fix a surface Σ , then pseudo-Anosovs with small dilatation have small volume. This is true in a sense. In fact it is proved in [6] that pseudo-Anosovs on any surfaces with small dilatation have bounded volume, see Theorem 1.4.

We denote by δ_g , the minimal dilatation for pseudo-Anosov elements $\phi \in \text{Mod}(\Sigma_g)$ on a closed surface Σ_g of genus g . A natural question is: what is the value δ_g ? To discuss the minimal dilatations, we introduce the polynomial

$$f_{(k,\ell)}(t) = t^{2k} - t^{k+\ell} - t^k - t^{k-\ell} + 1 \quad \text{for } k > 0, -k < \ell < k.$$

2010 *Mathematics Subject Classification.* Primary 57M27, 37E30; Secondary 37B40.

Key Words and Phrases. mapping class group, pseudo-Anosov, dilatation, entropy, fibered 3-manifold.

The first author was partially supported by Grant-in-Aid for Young Scientists (B) (No. 20740031), MEXT, Japan.

This polynomial has the largest real root $\lambda_{(k,\ell)}$ which is greater than 1 (Theorem 3.2 and Lemma 4.4). For any fixed $\ell > 0$, it follows that $k \log \lambda_{(k,\ell)}$ converges to $\log((3 + \sqrt{5})/2)$ if k goes to ∞ (Lemma 4.16). It is easy to show that $\delta_1 = \lambda_{(1,0)} = (3 + \sqrt{5})/2$. It was proved by Cho-Ham that $\delta_2 = \lambda_{(2,1)} \approx 1.72208$ [4]. It is open to determine the values δ_g for $g \geq 3$. Questions on properties of δ_g were posed by McMullen and Farb:

QUESTION 1.1 ([23] for (1), [5] for (2)).

- (1) Does $\lim_{g \rightarrow \infty} g \log \delta_g$ exist? What is its value?
- (2) Is the sequence $\{\delta_g\}_{g \geq 2}$ (strictly) monotone decreasing?

Related questions are ones for orientable pseudo-Anosovs. A pseudo-Anosov mapping class ϕ is said to be *orientable* if the invariant (un)stable foliation of a pseudo-Anosov homeomorphism $\Phi \in \phi$ is orientable. We denote by δ_g^+ , the minimal dilatation for orientable pseudo-Anosov elements of $\text{Mod}(\Sigma_g)$. The minima δ_g^+ were determined for $g = 2$ by Zhurov [31], for $3 \leq g \leq 5$ by Lanneau-Thiffeault [17], and for $g = 8$ by Lanneau-Thiffeault and Hironaka [17], [9]. Those values are given by $\delta_2^+ = \lambda_{(2,1)}$, $\delta_3^+ = \lambda_{(3,1)} = \lambda_{(4,3)} \approx 1.40127$, $\delta_4^+ = \lambda_{(4,1)} \approx 1.28064$, $\delta_5^+ = \lambda_{(6,1)} = \lambda_{(7,4)} \approx 1.17628$ and $\delta_8^+ = \lambda_{(8,1)} \approx 1.12876$.

Lanneau-Thiffeault obtained the inequality $\delta_5^+ \leq \delta_6^+$ ([17]) which implies that $\{\delta_g^+\}_{g \geq 2}$ is not strictly monotone decreasing. This leads us to ask an alternative question related to Question 1.1 (2): is the sequence $\{\delta_g^+\}_{g \geq 2}$ monotone decreasing? Also, one can ask: which g does the inequality $\delta_g < \delta_g^+$ hold? It is easy to see that $\delta_1 = \delta_1^+$. The equality $\delta_g = \delta_g^+$ holds for $g = 2$ [4], [31]. We do not know whether $\delta_3 = \delta_3^+$ holds or not. By work of Lanneau-Thiffeault and Hironaka, it follows that $\delta_g < \delta_g^+$ for $g = 4, 6, 8$ [17], [9].

To discuss Question 1.1 (1), we recall the previous upper bound of δ_g given by Hironaka.

THEOREM 1.2 ([9]). (1) $\delta_g \leq \lambda_{(g+1,3)}$ if $g \equiv 0, 1, 3, 4 \pmod{6}$ and $g \geq 3$.
 (2) $\delta_g \leq \lambda_{(g+1,1)}$ if $g \equiv 2, 5 \pmod{6}$ and $g \geq 5$.

By using Lemma 4.16 and Theorem 1.2, the following asymptotic inequality holds.

THEOREM 1.3 ([9]).

$$\limsup_{g \rightarrow \infty} g \log \delta_g \leq \log \left(\frac{3 + \sqrt{5}}{2} \right).$$

This improves the upper bound $g \log \delta_g \leq g \log \delta_g^+ \leq \log(2 + \sqrt{3})$ for any $g \geq 2$ by Minakawa [22] and Hironaka-Kin [10]. Since $\log \delta_g$ tends to 0 as g tends to ∞ ,




Figure 1. (left) 3 chain link C_3 . (center) $(-2, 3, 8)$ -pretzel link or Whitehead sister link. (right) link 6_2^2 .

Theorem 1.3 implies that

$$\limsup_{g \rightarrow \infty} |\chi(\Sigma_g)| \log \delta_g \leq 2 \log \left(\frac{3 + \sqrt{5}}{2} \right),$$

where $\chi(\Sigma)$ is the Euler characteristic of a surface Σ .

Let N be the *magic manifold*, which is the exterior of the 3 chain link C_3 illustrated in Figure 1 (left). This manifold has the smallest known volume among orientable hyperbolic 3-manifolds having 3 cusps. Many manifolds having at most 2 cusps with small volume are obtained from N by Dehn fillings, see [20]. In this paper, we study the *small dilatation pseudo-Anosov homeomorphisms* which occur as monodromies of fibrations on manifolds obtained from N by Dehn filling all three cusps. In [6], Farb, Leininger and Margalit introduced small dilatation pseudo-Anosov homeomorphisms which we recall below.

For any number $P > 1$, define the set of pseudo-Anosov homeomorphisms

$$\Psi_P = \{\text{pseudo-Anosov } \Phi : \Sigma \rightarrow \Sigma \mid \chi(\Sigma) < 0, |\chi(\Sigma)| \log \lambda(\Phi) \leq \log P\}.$$

They call elements $\Phi \in \Psi_P$ small dilatation pseudo-Anosov homeomorphisms. Theorem 1.3 says that if one takes P sufficiently large, then Ψ_P contains a pseudo-Anosov homeomorphism $\Phi_g : \Sigma_g \rightarrow \Sigma_g$ for each $g \geq 2$. By a result by Hironaka-Kin [10], Ψ_P also contains a pseudo-Anosov homeomorphism $\Phi_n : D_n \rightarrow D_n$ on an n -punctured disk D_n for each $n \geq 3$. Let $\Sigma^\circ \subset \Sigma$ be the surface obtained by removing the singularities of the (un)stable foliation for Φ and $\Phi|_{\Sigma^\circ} : \Sigma^\circ \rightarrow \Sigma^\circ$ denotes the restriction. Observe that $\lambda(\Phi) = \lambda(\Phi|_{\Sigma^\circ})$. The set

$$\Psi_P^\circ = \{\Phi|_{\Sigma^\circ} : \Sigma^\circ \rightarrow \Sigma^\circ \mid (\Phi : \Sigma \rightarrow \Sigma) \in \Psi_P\}$$

is infinite. Let $\mathcal{T}(\Psi_P^\circ)$ be the set of homeomorphism classes of mapping tori by elements of Ψ_P° .

THEOREM 1.4 ([6]). *The set $\mathcal{T}(\Psi_P^\circ)$ is finite. Namely, for each $P > 1$, there exist finite many complete, non compact hyperbolic 3-manifolds M_1, M_2, \dots, M_r fibering over S^1 so that the following holds. Any pseudo-Anosov $\Phi \in \Psi_P$ occurs as the monodromy of a Dehn filling of one of the M_k . In particular, there exists a constant $V = V(P)$ such that $\text{vol}(\Phi) \leq V$ holds for any $\Phi \in \Psi_P$.*

Agol also proved Theorem 1.4 by using periodic splitting sequences of pseudo-Anosov mapping tori [3]. By Theorem 1.4, one sees that the following sets \mathcal{U} , \mathcal{U}^+ and \mathcal{V} are finite.

$$\begin{aligned} \mathcal{U} &= \{ \mathbb{T}(\Phi|_{\Sigma^\circ}) \mid g \geq 2, \Phi \text{ is a pseudo-Anosov homeomorphism} \\ &\quad \text{on } \Sigma = \Sigma_g \text{ such that } \lambda(\Phi) = \delta_g \}, \\ \mathcal{U}^+ &= \{ \mathbb{T}(\Phi|_{\Sigma^\circ}) \mid g \geq 2, \Phi \text{ is an orientable pseudo-Anosov homeomorphism} \\ &\quad \text{on } \Sigma = \Sigma_g \text{ such that } \lambda(\Phi) = \delta_g^+ \}, \\ \mathcal{V} &= \{ \mathbb{T}(\Phi|_{\Sigma^\circ}) \mid n \geq 3, \Phi \text{ is a pseudo-Anosov homeomorphism} \\ &\quad \text{on } \Sigma = D_n \text{ such that } \lambda(\Phi) = \delta(D_n) \}, \end{aligned}$$

where $\delta(D_n)$ denotes the minimal dilatation for pseudo-Anosov elements of $\text{Mod}(D_n)$ on an n -punctured disk D_n .

The previous study [15] by the authors implies that $N \in \mathcal{V}$. In fact, the mapping class $\phi \in \text{Mod}(D_4)$ represented by the 4-braid $\sigma_1\sigma_2\sigma_3^{-1}$ has the minimal dilatation $\delta(D_4)$ [16]. For the pseudo-Anosov representative Φ of this mapping class ϕ , the mapping torus $\mathbb{T}(\Phi|_{\Sigma^\circ})$ is homeomorphic to N [15]. Moreover for each $n \geq 6$ (resp. $n = 3, 4, 5$), a pseudo-Anosov homeomorphism $\Phi_n : D_n \rightarrow D_n$ having the smallest known dilatation (resp. smallest dilatation) occurs as the monodromy on a particular fibration on a manifold obtained from N by Dehn filling [15]. See also work of Venzke [29].

Hironaka obtained Theorems 1.2 and 1.3 by viewing the monodromies of fibrations on manifolds obtained from the 6_2^2 link exterior $S^3 \setminus 6_2^2$ by Dehn filling two cusps. (For the link 6_2^2 , see Figure 1 (right) or Rolfsen’s table [24, Appendix C].) There exists an orientable monodromy: $\Sigma_2 \rightarrow \Sigma_2$, with dilatation $\delta_2 = \delta_2^+$ of a fibration on a manifold obtained from $S^3 \setminus 6_2^2$ by Dehn filling two cusps. This implies that $S^3 \setminus 6_2^2 \in \mathcal{U} \cap \mathcal{U}^+$ (Lemma 4.24 or [9]). We see that $S^3 \setminus 6_2^2$ is homeomorphic to $N(-1/2)$ (see [20, Table A.1] for example), where $N(r)$ is the manifold obtained from N by Dehn filling one cusp along the slope r . As mentioned, computer experiments say that the pseudo-Anosovs with small dilatation have small volume, and N is the candidate having the smallest volume among orientable 3-manifolds with 3 cusps. These results led us to see monodromies of fibrations on manifold obtained from N by Dehn filling.

In this paper, we investigate the fibrations on manifolds obtained from the three 2-cusped manifolds $N(-1/2)$, $N(-3/2)$ and $N(2)$ by Dehn filling 2 cusps. The second one $N(-3/2)$ is homeomorphic to $N(-4)$ and this is the Whitehead sister link exterior, i.e, the $(-2, 3, 8)$ -pretzel link exterior (see [20, Table A.1]), see Figure 1 (center). The manifold $N(-3/2)$ and the Whitehead link exterior have the smallest volume among orientable 2-cusped hyperbolic 3-manifolds [2]. We shall see that $N(-3/2)$ and $N(2)$ are elements of \mathcal{U}^+ (Lemmas 4.21, 4.36). Our main result is that $N(-3/2)$ (resp. $N(2)$) also admits Dehn fillings giving a sequence of fibers over the circle, with closed fibers Σ_g of genus g for each $g \geq 3$ such that the monodromies associated to the fibrations satisfy the same asymptotic inequality as Theorem 1.3. More precisely, we shall prove the following.

THEOREM 1.5. *Let $r \in \{3/-2, 1/-2, 2\}$. For each $g \geq 3$, there exist Σ_g -bundles over the circle obtained from $N(r)$ by Dehn filling all two cusps along the boundary slopes of fibers of $N(r)$. Among them, there exist the monodromies $\Phi_g(r) : \Sigma_g \rightarrow \Sigma_g$ of the fibrations such that*

- (1) $\lim_{g \rightarrow \infty} g \log \lambda(\Phi_g(r)) = \log((3 + \sqrt{5})/2)$,
- (2) $\lim_{g \rightarrow \infty} \text{vol}(\Phi_g(r)) = \text{vol}(N(r))$.

Independently, Aaber and Dunfield have investigated Σ_g -bundles over the circle obtained from $N(-3/2)$ by Dehn filling two cusps, see [1] and Remark 4.33. They have obtained similar results on the dilatation to those given in this paper. Theorem 1.5 in the case $r = -3/2$ was also established by [1].

By using monodromies on closed fibers coming from $N(-3/2)$, we find an upper bound of δ_g .

- THEOREM 1.6.** (1) $\delta_g \leq \lambda_{(g+2,1)}$ if $g \equiv 0, 1, 5, 6 \pmod{10}$ and $g \geq 5$.
 (2) $\delta_g \leq \lambda_{(g+2,2)}$ if $g \equiv 7, 9 \pmod{10}$ and $g \geq 7$.

THEOREM 1.7. *Let $g \equiv 2, 4 \pmod{10}$. Suppose that $g + 2 \not\equiv 0 \pmod{4641}$ ($= 3 \cdot 7 \cdot 13 \cdot 17$).*

- (1) $\delta_g \leq \lambda_{(g+2,3)}$ if $\text{gcd}(g + 2, 3) = 1$.
- (2) $\delta_g \leq \lambda_{(g+2,7)}$ if 3 divides $g + 2$ and $\text{gcd}(g + 2, 7) = 1$.
- (3) $\delta_g \leq \lambda_{(g+2,13)}$ if 21 ($= 3 \cdot 7$) divides $g + 2$ and $\text{gcd}(g + 2, 13) = 1$.
- (4) $\delta_g \leq \lambda_{(g+2,17)}$ if 273 ($= 3 \cdot 7 \cdot 13$) divides $g + 2$ and $\text{gcd}(g + 2, 17) = 1$.

We will verify the bounds in Theorems 1.6, 1.7 are sharper than the ones in Theorem 1.2 (see Propositions 4.26 (1), (2) and 4.28). Theorems 1.6, 1.7 do not include the case $g \equiv 3, 8 \pmod{10}$. This is because in this case, $N(-3/2)$ can not give rise to the monodromy on a closed fiber of genus g whose dilatation is strictly smaller than the one obtained from $N(-1/2)$, see Proposition 4.26 (3),

(4). However in the case $g = 8, 13$, we find a sharper upper bound than the one in Theorem 1.2. Let $\lambda_{(x,y,z)}$ be the largest real root of the polynomial

$$f_{(x,y,z)}(t) = t^{x+y-z} - t^x - t^y - t^{x-z} - t^{y-z} + 1.$$

- PROPOSITION 1.8. (1) $\delta_8 \leq \lambda_{(18,17,7)} (\approx 1.10403) < \lambda_{(9,1)} (\approx 1.11350)$.
 (2) $\delta_{13} \leq \lambda_{(27,21,8)} (\approx 1.07169) < \lambda_{(14,3)} (\approx 1.07266)$.

We turn to the study on δ_g^+ . We record results by Lanneau-Thiffeault.

THEOREM 1.9 ([17]). *The minimal dilatation δ_g^+ for $g = 6, 7$ is not less than the largest real root of the following polynomial.*

- (1) $f_{(6,1)}(t) = t^{12} - t^7 - t^6 - t^5 + 1$ if $g = 6$. ($\delta_6^+ \geq \lambda_{(6,1)} \approx 1.17628$.)
 (2) $f_{(9,2)}(t) = (t^4 - t^3 + t^2 - t + 1)(t^{14} + t^{13} - t^9 - t^8 - t^7 - t^6 - t^5 + t + 1)$ if $g = 7$.
 ($\delta_7^+ \geq \lambda_{(9,2)} \approx 1.11548$.)

Lanneau-Thiffeault asked the following.

QUESTION 1.10 ([17]). For g even, is δ_g^+ equal to the largest real root of the polynomial

$$f_{(g,1)}(t) = t^{2g} - t^{g+1} - t^g - t^{g-1} + 1?$$

Namely, is δ_g^+ equal to $\lambda_{(g,1)}$ for g even?

An upper bound of δ_g^+ given by Hironaka is as follows.

- THEOREM 1.11 ([9]). (1) $\delta_g^+ \leq \lambda_{(g+1,3)}$ if $g \equiv 1, 3 \pmod{6}$.
 (2) $\delta_g^+ \leq \lambda_{(g,1)}$ if $g \equiv 2, 4 \pmod{6}$.
 (3) $\delta_g^+ \leq \lambda_{(g+1,1)}$ if $g \equiv 5 \pmod{6}$.

We do not know whether there exists an orientable pseudo-Anosov homeomorphism of genus g having the dilatation $\lambda_{(g,1)}$ (appeared in Question 1.10) or not for each $g \equiv 0 \pmod{6}$. (For the recent work on the upper bound of δ_g^+ for $g \equiv 0 \pmod{6}$, see [14].) Under the assumption that Question 1.10 is true, the inequality $\delta_g^+ \leq \delta_{g+1}^+$ holds whenever $g \equiv 5 \pmod{6}$ and $\delta_g < \delta_g^+$ holds for all even g , see [9].

We give an upper bound of δ_g^+ in the case $g \equiv 1, 5, 7, 9 \pmod{10}$ using orientable pseudo-Anosov monodromies coming from $N(-3/2)$.

- THEOREM 1.12. (1) $\delta_g^+ \leq \lambda_{(g+2,2)}$ if $g \equiv 7, 9 \pmod{10}$ and $g \geq 7$.
 (2) $\delta_g^+ \leq \lambda_{(g+2,4)}$ if $g \equiv 1, 5 \pmod{10}$ and $g \geq 5$.

We shall see that the bound in Theorem 1.12 improves the one in Theorem 1.11 (see Proposition 4.34). Theorem 1.12 (1) together with Theorem 1.9 (2) gives:

COROLLARY 1.13. $\delta_7^+ = \lambda_{(9,2)}$.

Independently, Corollary 1.13 was established by Aaber and Dunfield [1].

The following tells us that the sequence $\{\delta_g^+\}_{g \geq 2}$ is not monotone decreasing.

PROPOSITION 1.14. *If Question 1.10 is true, then $\delta_g^+ < \delta_{g+1}^+$ whenever $g \equiv 1, 5, 7, 9 \pmod{10}$ and $g \geq 7$. In particular the inequality $\delta_7^+ < \delta_8^+$ holds.*

Our pseudo-Anosov homeomorphisms providing the upper bound of δ_g in Theorem 1.6 (1) are not orientable (Remark 4.27). This together with the inequality $\lambda_{(7,1)} < \lambda_{(6,1)} = \delta_5^+$ implies:

COROLLARY 1.15. $\delta_5 < \delta_5^+$.

We have a question:

QUESTION 1.16. Does the magic manifold N satisfy the following (1), (2) and (3)?

- (1) There exist Dehn fillings of N giving an infinite sequence of fiberings over the circle, with closed fibers Σ_{g_i} of genus $g_i \geq 2$ with $g_i \rightarrow \infty$, and with monodromy Φ_i so that $\delta_{g_i} = \lambda(\Phi_i)$.
- (2) There exist Dehn fillings of N giving an infinite sequence of fiberings over the circle, with closed fibers Σ_{g_i} of genus $g_i \geq 2$ with $g_i \rightarrow \infty$, and with monodromy Φ_i having the orientable (un)stable foliation so that $\delta_{g_i}^+ = \lambda(\Phi_i)$.
- (3) There exist Dehn fillings of N giving an infinite sequence of fiberings over the circle, with fibers D_{n_i} having n_i punctures with $n_i \rightarrow \infty$, and with monodromy Φ_i so that $\delta(D_{n_i}) = \lambda(\Phi_i)$.

The existence of the manifold satisfying each of (1), (2) and (3) is guaranteed from Theorem 1.4. Question 1.16 asks whether the magic manifold enjoys all (1), (2) and (3) or not.

The paper is organized as follows. We review basic facts in Section 2. The fibered faces and the entropy function for N are described in Section 3. The (un)stable foliation for the monodromy of the fibration on N is discussed in the section. We prove theorems in Section 4.

2. Notation and basic facts.

2.1. Pseudo-Anosov.

The *mapping class group* $\text{Mod}(\Sigma)$ is the group of isotopy classes of orientation preserving homeomorphisms of an orientable surface Σ , where the group operation is induced by composition of homeomorphisms. An element of this group is called a *mapping class*.

A homeomorphism $\Phi : \Sigma \rightarrow \Sigma$ is *pseudo-Anosov* if there exists a constant $\lambda = \lambda(\Phi) > 1$ called the *dilatation of Φ* and there exists a pair of transverse measured foliations \mathcal{F}^s and \mathcal{F}^u such that

$$\Phi(\mathcal{F}^s) = \frac{1}{\lambda}\mathcal{F}^s \text{ and } \Phi(\mathcal{F}^u) = \lambda\mathcal{F}^u.$$

Measured foliations \mathcal{F}^s and \mathcal{F}^u are called the *stable* and *unstable foliations* or *invariant foliations* for Φ . In this case the mapping class $\phi = [\Phi]$ is called pseudo-Anosov. We define the dilatation of ϕ , denoted by $\lambda(\phi)$, to be the dilatation of Φ .

The (topological) entropy $\text{ent}(f)$ is a measure of the complexity of a continuous self-map f on a compact manifold, see for instance [30]. The inequality

$$\log \text{sp}(f_*) \leq \text{ent}(f)$$

holds (see [19]), where $\text{sp}(f_*)$ is the spectral radius of the induced map $f_* : H_1(S; \mathbb{R}) \rightarrow H_1(S; \mathbb{R})$ on the first homology group. For any pseudo-Anosov homeomorphism $\Phi : \Sigma \rightarrow \Sigma$, the equality

$$\text{ent}(\Phi) = \log(\lambda(\Phi))$$

holds and $\text{ent}(\Phi)$ attains the minimal entropy among all homeomorphisms which are isotopic to Φ , see [7, Exposé 10]. We denote by $\text{ent}(\phi)$, this characteristic number. If Φ has orientable invariant foliations, then the equality

$$\log \text{sp}(\Phi_*) = \text{ent}(\Phi)$$

holds, see [25]. The converse is true:

THEOREM 2.1 ([17]). *A pseudo-Anosov homeomorphism Φ is orientable if and only if $\text{sp}(\Phi_*) = \lambda(\Phi)$.*

If we fix a surface Σ and take a constant $c > 1$, then the set of dilatations $\lambda(\Phi) < c$ for pseudo-Anosov homeomorphisms $\Phi : \Sigma \rightarrow \Sigma$ is finite, see [11]. In

particular the set

$$\text{Dil}(\Sigma) = \{\lambda(\phi) \mid \text{pseudo-Anosov } \phi \in \text{Mod}(\Sigma)\}$$

achieves a minimum $\delta(\Sigma)$.

Thurston’s hyperbolization theorem [27] asserts that ϕ is pseudo-Anosov if and only if its mapping torus

$$\mathbb{T}(\phi) = \Sigma \times [0, 1] / \sim,$$

where \sim identifies $(x, 1)$ with $(f(x), 0)$ for a representative $f \in \phi$, is hyperbolic. We denote the hyperbolic volume of $\mathbb{T}(\phi)$ by $\text{vol}(\phi)$.

Let us suppose that Σ is a compact orientable surface of genus g and we consider a pseudo-Anosov homeomorphism $\Phi : \Sigma \rightarrow \Sigma$. The stable foliation for Φ is denote by \mathcal{F} . Let x_1, \dots, x_m be all the singularities for \mathcal{F} in the interior $\text{int}(\Sigma)$, and $p(x_i) \geq 3$ denotes the number of prongs of \mathcal{F} at x_i . Let y_1, \dots, y_n be all the singularities for \mathcal{F} on the boundary $\partial\Sigma$, and $p(y_j) \geq 1$ denotes the number of prongs of \mathcal{F} at y_j . The following Euler-Poincaré formula holds:

$$\sum_{i=1}^m (p(x_i) - 2) + \sum_{j=1}^n (p(y_j) - 2) = -2\chi(\Sigma_g) = 4g - 4$$

(see [7, Exposé 5] for example). The pair of integers

$$(p(x_1) - 2, p(x_2) - 2, \dots, p(x_m) - 2, p(y_1) - 2, p(y_2) - 2, \dots, p(y_n) - 2)$$

is called the *singularity data* of Φ .

2.2. Thurston norm.

Let M be an irreducible, atoroidal and oriented 3-manifold with boundary ∂M (possibly $\partial M = \emptyset$). We recall the Thurston norm $\|\cdot\| : H_2(M, \partial M; \mathbb{R}) \rightarrow \mathbb{R}$ (see [26]). The norm $\|\cdot\|$ has the property that for any integral class $a \in H_2(M, \partial M; \mathbb{R})$,

$$\|a\| = \min_F \{-\chi(F)\},$$

where the minimum is taken over all oriented surface F embedded in M , satisfying $a = [F]$, with no components of non-negative Euler characteristic. The surface F which realizes this minimum is called a *minimal representative* of a . For a rational class $a \in H_2(M, \partial M; \mathbb{R})$, take a rational number r so that ra is an integral class. Then $\|a\|$ is defined to be

$$\|a\| = \frac{1}{|r|} \|ra\|.$$

The function $\|\cdot\|$ defined on rational classes admits a unique continuous extension to $H_2(M, \partial M; \mathbb{R})$ which is linear on the ray through the origin. The unit ball $U = \{a \in H_2(M, \partial M; \mathbb{R}) \mid \|a\| \leq 1\}$ is a compact, convex polyhedron [26].

The following notations are needed to describe how fibrations of M are related to $\|\cdot\|$.

- A top dimensional face in the boundary ∂U is denoted by Δ , and its open face is denoted by $\text{int}(\Delta)$.
- C_Δ is the cone over Δ with the origin and $\text{int}(C_\Delta)$ is its interior.
- The set of integral classes (resp. rational classes) of $\text{int}(C_\Delta)$ is denoted by $\text{int}(C_\Delta(\mathbb{Z}))$ (resp. $\text{int}(C_\Delta(\mathbb{Q}))$).

THEOREM 2.2 ([26]). *Suppose that M is a surface bundle over the circle and let F be a fiber. Then there exists a top dimensional face Δ satisfying the following.*

- (1) $[F] \in \text{int}(C_\Delta(\mathbb{Z}))$.
- (2) For any $a \in \text{int}(C_\Delta(\mathbb{Z}))$, a minimal representative of a is a fiber of fibrations on M .

The face Δ in Theorem 2.2 is called a *fibred face* and an integral class $a \in \text{int}(C_\Delta)$ is called a *fibred class*.

2.3. Entropy function.

Let M be a hyperbolic surface bundle over the circle. We fix a fibred face Δ for M . The *entropy function* $\text{ent} : \text{int}(C_\Delta) \rightarrow \mathbb{R}$ introduced by Fried [8] is defined as follows. The minimal representative F_a for a primitive fibred class $a \in \text{int}(C_\Delta)$ is connected and is a fiber of fibrations on M . Let $\Phi_a : F_a \rightarrow F_a$ be the monodromy. Since M is hyperbolic, $\phi_a = [\Phi_a]$ is pseudo-Anosov. The entropy $\text{ent}(a)$ and dilatation $\lambda(a)$ are defined to be the entropy and dilatation of ϕ_a . For $r \in \mathbb{Q}$, the entropy $\text{ent}(ra)$ is defined by $1/|r| \text{ent}(a)$. Fried proved that $1/\text{ent} : \text{int}(C_\Delta(\mathbb{Q})) \rightarrow \mathbb{R}$ is concave [8], and in particular it admits a unique continuous extension $\text{ent} : \text{int}(C_\Delta) \rightarrow \mathbb{R}$. Moreover, he proved that the restriction of ent to $\text{int}(\Delta)$ is proper, that is $\text{ent}(a)$ goes to ∞ as a goes to a boundary point of $\partial\Delta$. Note that $1/\text{ent} : \text{int}(C_\Delta) \rightarrow \mathbb{R}$ is linear along each ray through the origin by definition and cannot be strictly concave for this direction. However Matsumoto and later McMullen proved that it is strictly concave for other directions.

THEOREM 2.3 ([21], [23]). $1/\text{ent} : \text{int}(\Delta) \rightarrow \mathbb{R}$ is strictly concave.

By definition of ent , $\text{Ent} = \|\cdot\| \text{ent} : \text{int}(C_\Delta) \rightarrow \mathbb{R}$ is constant on each ray in $\text{int}(C_\Delta)$ through the origin. We call $\text{Ent}(a)$ the *normalized entropy* of $a \in \text{int}(C_\Delta)$. By Theorem 2.3 together with the properness of ent by Fried, Ent admits a unique minimum at a unique ray through the origin. In other words, if we regard Ent as a function defined on $\text{int}(\Delta)$, then it has the minimum at a unique point in $\text{int}(\Delta)$.

The following question was posed by McMullen.

QUESTION 2.4 ([23]). On which ray in $\text{int}(C_\Delta)$ does Ent attain the minimum? Is the minimum attained on a rational class of $\text{int}(\Delta)$?

We consider Question 2.4 for $N(-3/2)$, $N(-1/2)$ and $N(2)$, see Proposition 4.13.

3. Magic manifold.

3.1. Fibered face and entropy function.

The magic manifold N is a surface bundle over the circle ([15] for instance). In this section, we recall the entropy function on a fibered face for N .

Let K_α, K_β and K_γ be the components of the 3 chain link \mathcal{C}_3 . They bound the oriented disks F_α, F_β and F_γ with 2 holes, see Figure 2 (right). Let $\alpha = [F_\alpha]$, $\beta = [F_\beta]$, and $\gamma = [F_\gamma]$. The Thurston unit ball U for N is the parallelepiped with vertices $\pm\alpha = (\pm 1, 0, 0)$, $\pm\beta = (0, \pm 1, 0)$, $\pm\gamma = (0, 0, \pm 1)$, $\pm(\alpha + \beta + \gamma)$, see Figure 2 (left). The set $\{\alpha, \beta, \gamma\}$ is a basis of $H_2(N, \partial N; \mathbb{Z})$. The symmetry of \mathcal{C}_3 tells us that every top dimensional face is a fibered face. We fix a face Δ with vertices $\alpha = (1, 0, 0)$, $\alpha + \beta + \gamma = (1, 1, 1)$, $\beta = (0, 1, 0)$ and $-\gamma = (0, 0, -1)$. Then

$$\text{int}(\Delta) = \{x\alpha + y\beta + z\gamma \mid x + y - z = 1, x > 0, y > 0, x > z, y > z\}.$$

Hence if $x\alpha + y\beta + z\gamma \in \text{int}(C_\Delta)$, then

$$\|x\alpha + y\beta + z\gamma\| = x + y - z. \tag{3.1}$$




Figure 2. (left) Thurston norm ball for N . (right) $F_\alpha, F_\beta, F_\gamma$ [arrows indicate the normal direction of oriented surfaces].

Let $\mathcal{N}(L)$ be the regular neighborhood of a link L in S^3 . We denote the tori $\partial\mathcal{N}(K_\alpha)$, $\partial\mathcal{N}(K_\beta)$, $\partial\mathcal{N}(K_\gamma)$ by T_α , T_β , T_γ respectively. Let $x\alpha + y\beta + z\gamma$ be a primitive fibered class in $\text{int}(C_\Delta)$. The minimal representative of this class is denoted by $F_{x\alpha+y\beta+z\gamma}$ or $F_{(x,y,z)}$. Let us put $\partial_\alpha F_{(x,y,z)} = \partial F_{(x,y,z)} \cap T_\alpha$ which consists of parallel simple closed curves on T_α . We define $\partial_\beta F_{(x,y,z)}$, $\partial_\gamma F_{(x,y,z)} \subset \partial F_{(x,y,z)}$ in the same manner.

LEMMA 3.1. *Let $x\alpha + y\beta + z\gamma$ be a primitive fibered class in $\text{int}(C_\Delta)$. The number of boundary components $\sharp(\partial F_{(x,y,z)})$ is equal to $\gcd(x, y + z) + \gcd(y, z + x) + \gcd(z, x + y)$, where $\gcd(0, w)$ is defined by $|w|$. More precisely*

- (1) $\sharp(\partial_\alpha F_{(x,y,z)}) = \gcd(x, y + z)$,
- (2) $\sharp(\partial_\beta F_{(x,y,z)}) = \gcd(y, z + x)$,
- (3) $\sharp(\partial_\gamma F_{(x,y,z)}) = \gcd(z, x + y)$.

PROOF. We prove (1). The proof of (2), (3) is similar. We have the meridian and longitude bases $\{m_\alpha, \ell_\alpha\}$ for T_α , $\{m_\beta, \ell_\beta\}$ for T_β and $\{m_\gamma, \ell_\gamma\}$ for T_γ . We consider the long exact sequence of the homology groups of the pair $(N, \partial N)$. The boundary map is given by

$$\begin{aligned} \partial_* : H_2(N, \partial N; \mathbb{R}) &\rightarrow H_1(\partial N; \mathbb{R}), \\ \alpha &\mapsto \ell_\alpha - m_\beta - m_\gamma, \\ \beta &\mapsto \ell_\beta - m_\gamma - m_\alpha, \\ \gamma &\mapsto \ell_\gamma - m_\alpha - m_\beta. \end{aligned}$$

Hence

$$\partial_*(x\alpha + y\beta + z\gamma) = x\ell_\alpha - (y + z)m_\alpha + y\ell_\beta - (z + x)m_\beta + z\ell_\gamma - (x + y)m_\gamma. \tag{3.2}$$

Since $F_{(x,y,z)}$ is the minimal representative, $\partial_\alpha F_{(x,y,z)}$ is a union of oriented parallel simple closed curves on T_α whose homology class equals $x\ell_\alpha - (y + z)m_\alpha \in H_1(T_\alpha; \mathbb{R})$, see (3.2). Thus the number of components of $\partial_\alpha F_{(x,y,z)}$ equals $\gcd(x, y + z)$. This completes the proof. \square

From the proof of Lemma 3.1, one sees that the boundary slope of each simple closed curve of $\partial_\alpha F_{(x,y,z)}$ equals $-(y + z)/x$. Similarly the boundary slope of each component of $\partial_\beta F_{(x,y,z)}$ (resp. $\partial_\gamma F_{(x,y,z)}$) is given by $-(z + x)/y$ (resp. $-(x + y)/z$). Let us define

$$\text{slope}(x\alpha + y\beta + z\gamma) = \left(\frac{-(y + z)}{x}, \frac{-(z + x)}{y}, \frac{-(x + y)}{z} \right). \tag{3.3}$$

This notation $\text{slope}(\cdot)$ is needed for the study of Dehn fillings of N in Section 4.

One can compute the entropy for any element of $\text{int}(C_\Delta(\mathbb{Z}))$ by using the next theorem.

THEOREM 3.2 ([15]). *The dilatation $\lambda_{(x,y,z)}$ of $x\alpha + y\beta + z\gamma \in \text{int}(C_\Delta(\mathbb{Z}))$ is the largest real root of the polynomial*

$$P(t_1, t_2, t_3) = -t_1 - t_2 + t_3 + t_1t_2 - t_1t_3 - t_2t_3.$$

Since $P(t^x, t^y, t^z) = t^z(t^{x+y-z} - t^x - t^y - t^{x-z} - t^{y-z} + 1)$, $\lambda_{(x,y,z)}$ is the largest real root of

$$f_{(x,y,z)}(t) = t^{x+y-z} - t^x - t^y - t^{x-z} - t^{y-z} + 1.$$

The minimum of $\text{Ent} : \text{int}(C_\Delta) \rightarrow \mathbb{R}$ is equals to $2\log(2 + \sqrt{3})$ and it is attained by $\alpha + \beta$ [15]. Since the Thurston norm ball of N has a symmetry, $\min \text{Ent}$ does not depend on fibered faces of N .

3.2. Invariant foliations.

Let $\Phi_{(x,y,z)}$ be the monodromy of the fibration of N associated to a primitive fibered class $x\alpha + y\beta + z\gamma \in \text{int}(C_\Delta)$ and let $F_{(x,y,z)}$ be its fiber. We denote the stable foliation for $\Phi_{(x,y,z)}$ by $\mathcal{F}_{(x,y,z)}$. We shall compute the number of prongs at the singularities of $\mathcal{F}_{(x,y,z)}$.

PROPOSITION 3.3. *The singularity data of $\Phi_{(x,y,z)}$ is given by*

$$\left(\underbrace{\left(\frac{x}{\gcd(x, y + z)} - 2, \dots, \frac{x}{\gcd(x, y + z)} - 2 \right)}_{\gcd(x, y + z)}, \right. \\ \left. \underbrace{\left(\frac{y}{\gcd(y, x + z)} - 2, \dots, \frac{y}{\gcd(y, x + z)} - 2 \right)}_{\gcd(y, x + z)}, \right. \\ \left. \underbrace{\left(\frac{x + y - 2z}{\gcd(z, x + y)} - 2, \dots, \frac{x + y - 2z}{\gcd(z, x + y)} - 2 \right)}_{\gcd(z, x + y)} \right).$$

More precisely $\mathcal{F}_{(x,y,z)}$ is

- (1) $x/\gcd(x, y + z)$ -pronged at each component of $\partial_\alpha F_{(x,y,z)}$,
- (2) $y/\gcd(y, x + z)$ -pronged at each component of $\partial_\beta F_{(x,y,z)}$,

- (3) $(x + y - 2z)/\gcd(z, x + y)$ -pronged at each component of $\partial_\gamma F_{(x,y,z)}$, and
- (4) $\mathcal{F}_{(x,y,z)}$ has no singularities in the interior of $F_{(x,y,z)}$.

Here we recall the formula of the intersection number $i([c], [c'])$ between isotopy classes of essential simple closed curves c, c' on a torus T . Let $p/q, r/s$ be rational numbers or $1/0$ with irreducible forms and suppose that $p/q, r/s$ are slopes on T which represent isotopy classes $[c], [c']$ respectively. Then $i([c], [c']) = |ps - qr|$.

PROOF OF PROPOSITION 3.3. Observe that a fiber $F = F_{(1,1,0)}$ associated to the fibered class $\alpha + \beta$ is a sphere with 4 boundary components. The monodromy $\Phi = \Phi_{(1,1,0)} : F \rightarrow F$ of the fibration on N is represented by the 3-braid $b = \sigma_2 \sigma_1^{-1} \sigma_2$. In particular $S^3 \setminus \bar{b}$ is homeomorphic to $S^3 \setminus \mathcal{C}_3 = N$, where \bar{b} is a union of the closed braid of b and the braid axis, see Figure 3 (right). We define a homeomorphism $H : S^3 \setminus \mathcal{N}(\mathcal{C}_3) \rightarrow S^3 \setminus \mathcal{N}(\bar{b})$ as follows. Notice that the link illustrated in Figure 3 (center) is isotopic to \mathcal{C}_3 . We cut the twice-punctured disk F_α bounded by the component K_α . Let F'_α and F''_α be the resulting twice-punctured disks after cutting F_α . Reglue F'_α and F''_α after twisting the neighborhood of F'_α by 360 degrees in the clockwise direction. Then we obtain the link \bar{b} whose exterior $S^3 \setminus \bar{b}$ is homeomorphic to $S^3 \setminus \mathcal{C}_3$, see Figure 3. The inverse H^{-1} is denoted by h . We set $T_\alpha^H = H(T_\alpha)$, $T_\beta^H = H(T_\beta)$ and $T_\gamma^H = H(T_\gamma)$, see Figure 5. (Then $\partial\mathcal{N}(\bar{b}) = T_\alpha^H \cup T_\beta^H \cup T_\gamma^H$.)




Figure 3. (left, center) \mathcal{C}_3 . (right) \bar{b} . (This figure explains how to obtain H .)

The invariant train track τ which carries the stable lamination ℓ^s for Φ is illustrated in Figure 4 (left). The stable foliation \mathcal{F} for Φ has 1 prong at each component of ∂F and it has no singularity in the interior of F . We consider the suspension flow induced on the mapping torus $N = F \times [0, 1] / \sim$, where \sim identifies $(x, 1)$ with $(\Phi(x), 0)$. One obtains the simple closed curve $c_\alpha \subset T_\alpha^H$ which is the closed orbit of the singularity of \mathcal{F} on $\partial F \cap T_\alpha^H$. Similarly one has the closed orbits $c_\beta \subset T_\beta^H$, $c_\gamma \subset T_\gamma^H$, see Figure 5 (right). (One can draw these closed orbits by using the orbit of each cusp of $F \setminus \tau$.) Let $\mathcal{L}^s \subset N$ be the suspended stable lamination constructed from $\ell^s \times I \subset F \times I$ by gluing $\ell^s \times \{1\}$ to $\ell^s \times \{0\}$ using Φ . By construction, \mathcal{L}^s is carried by the branched surface B_τ which is obtained from $\tau \times I$ by gluing $\tau \times \{1\}$ to $\tau \times \{0\}$ using Φ . Notice that c_α, c_β and c_γ correspond to

the branched loci of B_τ . By work of Fried [8] (see also work of Long-Oertel [18]), we may assume that the fiber $F_{(x,y,z)}$ is transverse to \mathcal{L}^s . The stable lamination $\ell_{(x,y,z)}^s$ for $\Phi_{(x,y,z)}$ is given by the intersection $\mathcal{L}^s \cap F_{(x,y,z)}$ and $\ell_{(x,y,z)}^s$ is carried by the train track $B_\tau \cap F_{(x,y,z)}$. This implies that $\mathcal{F}_{(x,y,z)}$ has no singularity in the interior of $F_{(x,y,z)}$ and we finish the proof of (4).

We consider the number of prongs of $\mathcal{F}_{(x,y,z)}$ at each component of $\partial_\alpha F_{(x,y,z)}$. The boundary slope of each simple closed curve of $\partial_\alpha F_{(x,y,z)}$ is given by $-(y+z)/x$. The desired number is equal to the intersection number

$$i([c_{-(y+z)/x}], [h(c_\alpha)]) = i([H(c_{-(y+z)/x})], [c_\alpha]),$$

where c_r is a simple closed curve with slope $r \in \mathbb{Q} \cup \{1/0\}$ on T_α . Observe that $h(c_\alpha)$ has the slope $1/0$ (see Figure 5). Hence

$$i([c_{-(y+z)/x}], [h(c_\alpha)]) = \left| 1 \cdot \frac{x}{\gcd(x, y+z)} + 0 \cdot \frac{y+z}{\gcd(x, y+z)} \right| = \frac{x}{\gcd(x, y+z)}.$$

This completes the proof of (1).

One verifies that $h(c_\beta)$ and $h(c_\gamma)$ have slopes $1/0$ and $-2/1$ respectively. By using a similar argument, one can prove (2), (3). □

We consider the orientability of $\mathcal{F}_{(x,y,z)}$ using Theorem 2.1. Alexander polynomial of \mathcal{C}_3 is

$$A(t_1, t_2, t_3) = t_1 t_2 + t_2 t_3 + t_3 t_1 - t_1 - t_2 - t_3.$$

The following is a consequence of Proposition 7.3.10 in [12] which tells us the relation between the Alexander polynomial of links and the characteristic polynomial of $\Phi_* : H_1(\Sigma; \mathbb{R}) \rightarrow H_1(\Sigma; \mathbb{R})$ on fibers Σ in the link exteriors.




Figure 4. (left) Invariant train track τ for $\Phi_{(1,1,0)}$. (right) 1-pronged singularity.




Figure 5. (left) $h(c_\alpha), h(c_\beta), h(c_\gamma)$. (right) $c_\alpha, c_\beta, c_\gamma$.

LEMMA 3.4. *The spectral radius of $(\Phi_{(x,y,z)})_*$ is the largest absolute value among roots of*

$$A(t^x, t^y, t^z) = t^{x+y} + t^{y+z} + t^{z+x} - t^x - t^y - t^z.$$

PROPOSITION 3.5. *The pseudo-Anosov homeomorphism $\Phi_{(x,y,z)}$ is orientable if and only if x and y are even and z is odd.*

PROOF. (If part.) Suppose that x and y are even and z is odd. Then

$$P(t^x, t^y, t^z) = A((-t)^x, (-t)^y, (-t)^z).$$

This implies that $\lambda(\Phi_{(x,y,z)}) = \text{sp}((\Phi_{(x,y,z)})_*)$. By Theorem 2.1 $\mathcal{F}_{(x,y,z)}$ is orientable.

(Only if part.) Suppose that x or y is odd. We may assume that x is odd. The number of prongs of $\mathcal{F}_{(x,y,z)}$ at each component of $\partial_\alpha F_{(x,y,z)}$ equals $x/\text{gcd}(x, y+z)$ which is odd. Thus $\mathcal{F}_{(x,y,z)}$ can not be orientable. \square

3.3. Non-hyperbolic Dehn fillings.

Let M be a 3-manifold with boundary tori T_0, \dots, T_j and let $r_i \in \mathbb{Q} \cup \infty$ be a slope on T_i . Then $M(r_0, r_1, \dots, r_j)$ denotes the manifold obtained from M by Dehn filling along the slope r_i for each i , that is $M(r_0, r_1, \dots, r_j)$ is the manifold attaching a solid torus \tilde{T}_i to M along T_i in such a way that r_i bounds a disk in \tilde{T}_i .

Martelli and Petronio classified all the non-hyperbolic fillings of the magic manifold [20, Theorems 1.1, 1.2, 1.3]. We denote by T_0, T_1, T_2 , the boundary tori of $N = S^3 \setminus \mathcal{N}(\mathcal{C}_3)$.

THEOREM 3.6 ([20]). (1) $N(p/q)$ is hyperbolic if and only if

$$\frac{p}{q} \notin \{\infty, -3, -2, -1, 0\}.$$

(2) $N(p/q, r/s)$ is hyperbolic if and only if

$$\frac{p}{q}, \frac{r}{s} \notin \{\infty, -3, -2, -1, 0\} \text{ and } \left(\frac{p}{q}, \frac{r}{s}\right) \notin \left\{ (1, 1), \left(-4, \frac{-1}{2}\right), \left(\frac{-3}{2}, \frac{-5}{2}\right) \right\}.$$

As a corollary of Theorem 3.6 one has:

COROLLARY 3.7. *If $N(p/q, r/s, t/u)$ is hyperbolic, then*

$$\frac{p}{q}, \frac{r}{s} \notin \{\infty, -3, -2, -1, 0\} \text{ and } \left(\frac{p}{q}, \frac{r}{s}\right) \notin \left\{ (1, 1), \left(-4, \frac{-1}{2}\right), \left(\frac{-3}{2}, \frac{-5}{2}\right) \right\}.$$

Let us consider the monodromy $\Phi_{(x,y,z)} : F_{(x,y,z)} \rightarrow F_{(x,y,z)}$ of the fibration on N associated to a primitive fibered class $x\alpha + y\beta + z\gamma \in \text{int}(C_\Delta)$. Recall that $\text{slope}(x\alpha + y\beta + z\gamma) = (-(y+z)/x, -(z+x)/y, -(x+y)/z)$, see (3.3). By capping each boundary component of $F_{(x,y,z)}$, $\Phi_{(x,y,z)}$ extends to the monodromy $\bar{\Phi}_{(x,y,z)}$ with a closed fiber $\bar{F}_{(x,y,z)}$ of the fibration on $N(-(y+z)/x, -(z+x)/y, -(x+y)/z)$. If the stable foliation $\mathcal{F}_{(x,y,z)}$ is not 1-pronged at each component of $\partial F_{(x,y,z)}$, then $\bar{\Phi}_{(x,y,z)}$ is pseudo-Anosov and $\lambda(\bar{\Phi}_{(x,y,z)}) = \lambda(\Phi_{(x,y,z)})$. If $\mathcal{F}_{(x,y,z)}$ is 1-pronged at a component of $\partial F_{(x,y,z)}$, then $\bar{\Phi}_{(x,y,z)}$ may not be pseudo-Anosov.

4. Hyperbolic Dehn fillings $N(-3/2)$, $N(-1/2)$ and $N(2)$.

4.1. Thurston norm balls of $N(-3/2)$, $N(-1/2)$ and $N(2)$.

Let $N(r)$ be the manifold obtained from N by Dehn filling the cusp specified by T_β along the slope $r \in \mathbb{Q}$. Then there exists a natural injection $\iota_\beta : H_2(N(r), \partial N(r)) \rightarrow H_2(N, \partial N)$ whose image equals

$$S_\beta(r) = \{(x, y, z) \in H_2(N, \partial N) \mid -ry = z + x\},$$

see [14]. By Theorem 3.6(1), $N(r)$ is hyperbolic if and only if $r \in \mathbb{Q} \setminus \{-3, -2, -1, 0\}$. Choose $r \in \mathbb{Q} \setminus \{-3, -2, -1, 0\}$, and assume that $a \in S_\beta(r) = \text{Im } \iota_\beta$ is a fibered class of $H_2(N, \partial N)$. Then $\bar{a} = \iota_\beta^{-1}(a) \in H_2(N(r), \partial N(r))$ is also a fibered class of $N(r)$.

Similarly, when $N(r)$ is the manifold obtained from N by Dehn filling the cusp specified by T_α or T_γ along the slope r , one has natural injections,

$$\iota_\alpha : H_2(N(r), \partial N(r)) \rightarrow H_2(N, \partial N),$$

$$\iota_\gamma : H_2(N(r), \partial N(r)) \rightarrow H_2(N, \partial N)$$

such that their images are

$$\begin{aligned} S_\alpha(r) &= \{(x, y, z) \in H_2(N, \partial N) \mid -rx = y + z\}, \\ S_\gamma(r) &= \{(x, y, z) \in H_2(N, \partial N) \mid -rz = x + y\}. \end{aligned}$$

We set

$$\begin{aligned} \mathbf{a} &= 2\alpha + 2\beta + \gamma, & \mathbf{b} &= \alpha + 2\beta + 2\gamma, \\ \mathbf{p} &= \alpha + 2\beta, & \mathbf{q} &= 2\beta + \gamma, \\ \mathbf{r} &= \alpha + \beta - \gamma, & \mathbf{s} &= \alpha - \beta. \end{aligned}$$

For $k, \ell \in \mathbb{Z}$, we have

$$\begin{aligned} \text{slope}(k\mathbf{a} + \ell\mathbf{b}) &= \left(\frac{-3k - 4\ell}{2k + \ell}, \frac{-3}{2}, \frac{-4k - 3\ell}{k + 2\ell} \right), \\ \text{slope}(k\mathbf{p} + \ell\mathbf{q}) &= \left(\frac{-2k - 3\ell}{k}, \frac{-1}{2}, \frac{-3k - 2\ell}{\ell} \right), \\ \text{slope}(k\mathbf{r} + \ell\mathbf{s}) &= \left(\frac{\ell}{k + \ell}, \frac{-\ell}{k - \ell}, 2 \right), \text{ and} \end{aligned} \tag{4.1}$$

$$\begin{aligned} k\mathbf{a} + \ell\mathbf{b} &= (2k + \ell)\alpha + (2k + 2\ell)\beta + (k + 2\ell)\gamma \in \iota_\beta \left(H_2 \left(N \left(\frac{-3}{2} \right), \partial N \left(\frac{-3}{2} \right) \right) \right), \\ k\mathbf{p} + \ell\mathbf{q} &= k\alpha + (2k + 2\ell)\beta + \ell\gamma \in \iota_\beta \left(H_2 \left(N \left(\frac{-1}{2} \right), \partial N \left(\frac{-1}{2} \right) \right) \right), \\ k\mathbf{r} + \ell\mathbf{s} &= (k + \ell)\alpha + (k - \ell)\beta - k\gamma \in \iota_\gamma (H_2(N(2), \partial N(2))). \end{aligned} \tag{4.2}$$

It is easy to check that $\{\bar{\mathbf{a}}, \bar{\mathbf{b}}\}$, $\{\bar{\mathbf{p}}, \bar{\mathbf{q}}\}$ and $\{\bar{\mathbf{r}}, \bar{\mathbf{s}}\}$ are bases of $H_2(N(-3/2), \partial N(-3/2); \mathbb{Z})$, $H_2(N(-1/2), \partial N(-1/2); \mathbb{Z})$ and $H_2(N(2), \partial N(2); \mathbb{Z})$ respectively.

Note that $\gcd(k, \ell) = 1$ if and only if $k\mathbf{a} + \ell\mathbf{b}$, $k\mathbf{p} + \ell\mathbf{q}$ and $k\mathbf{r} + \ell\mathbf{s}$ are primitive integral classes of $H_2(N, \partial N; \mathbb{R})$. All $k\mathbf{a} + \ell\mathbf{b}$, $k\mathbf{p} + \ell\mathbf{q}$, $k\mathbf{r} + \ell\mathbf{s}$ are fibered classes in $\text{int}(C_\Delta)$ for $k > 0$ and $-k < \ell < k$.

We first focus on the topological types of fibers for primitive fibered classes in $\text{int}(C_\Delta)$. Let $\Sigma_{g,p}$ be a compact orientable surface of genus g with p boundary components.

LEMMA 4.1. *Suppose that $k > 0$, $-k < \ell < k$ and $\gcd(k, \ell) = 1$.*

(1) $F_{k\mathbf{a} + \ell\mathbf{b}} = \Sigma_{k-2, k + \ell + 6}$ if $\gcd(2k + \ell, 5) = 5$ or $\gcd(5, k + 2\ell) = 5$. Otherwise

- $$F_{k\mathbf{a}+\ell\mathbf{b}} = \Sigma_{k,k+\ell+2}.$$
- (2) $F_{k\mathbf{p}+\ell\mathbf{q}} = \Sigma_{k-1,k+\ell+4}$ if $\gcd(k, 3) = 3$ or $\gcd(3, \ell) = 3$. Otherwise $F_{k\mathbf{p}+\ell\mathbf{q}} = \Sigma_{k,k+\ell+2}$.
- (3) $F_{k\mathbf{r}+\ell\mathbf{s}} = \Sigma_{k,k+2}$.

PROOF OF (1). By Lemma 3.1,

$$\begin{aligned} \sharp(\partial F_{k\mathbf{a}+\ell\mathbf{b}}) &= \gcd(2k + \ell, 3k + 4\ell) + \gcd(2k + 2\ell, 3k + 3\ell) + \gcd(4k + 3\ell, k + 2\ell) \\ &= \gcd(2k + \ell, 5k) + k + \ell + \gcd(5\ell, k + 2\ell) \\ &= \gcd(2k + \ell, 5) + k + \ell + \gcd(5, k + 2\ell). \end{aligned}$$

The last equality holds since $\gcd(k, \ell) = 1$. The following 3 cases can occur.

- (1) $\gcd(2k + \ell, 5) = 1$ and $\gcd(5, k + 2\ell) = 1$.
- (2) $\gcd(2k + \ell, 5) = 5$ and $\gcd(5, k + 2\ell) = 1$.
- (3) $\gcd(2k + \ell, 5) = 1$ and $\gcd(5, k + 2\ell) = 5$.

In the case (1), the genus g of $F_{k\mathbf{a}+\ell\mathbf{b}}$ must satisfy

$$-(2 - 2g - k - \ell - 2) = \|k\mathbf{a} + \ell\mathbf{b}\| = 3k + \ell$$

(see (3.1)). Thus $g = k$ and $F_{k\mathbf{a}+\ell\mathbf{b}} = \Sigma_{k,k+\ell+2}$. In the cases (2) and (3), $F_{k\mathbf{a}+\ell\mathbf{b}} = \Sigma_{k-2,k+\ell+6}$.

The proof of claims (2), (3) of the lemma is similar. □

LEMMA 4.2. *Suppose that $k > 0$ and $-k < \ell < k$.*

- (1) $F_{k\mathbf{a}+\ell\mathbf{b}} = F_{\ell\mathbf{a}+k\mathbf{b}}$ and $\lambda(k\mathbf{a} + \ell\mathbf{b}) = \lambda(\ell\mathbf{a} + k\mathbf{b})$.
- (2) $F_{k\mathbf{p}+\ell\mathbf{q}} = F_{\ell\mathbf{p}+k\mathbf{q}}$ and $\lambda(k\mathbf{p} + \ell\mathbf{q}) = \lambda(\ell\mathbf{p} + k\mathbf{q})$.

PROOF. (1) By the symmetry of the Thurston norm ball of N , it is not hard to see that the topological type of the minimal representative (resp. the dilatation) for $\ell\mathbf{a} + k\mathbf{b} = (2\ell + k)\alpha + (2\ell + 2k)\beta + (\ell + 2k)\gamma$ is the same as the one for $k\mathbf{a} + \ell\mathbf{b} = (2k + \ell)\alpha + (2k + 2\ell)\beta + (k + 2\ell)\gamma$.

The proof of (2) is similar. □

We make a remark that it is not true in general that $F_{k\mathbf{r}+\ell\mathbf{s}} = F_{\ell\mathbf{r}+k\mathbf{s}}$ and $\lambda(k\mathbf{r} + \ell\mathbf{s}) = \lambda(\ell\mathbf{r} + k\mathbf{s})$ for $k > 0$ and $-k < \ell < k$. We do not use this remark in the rest of the paper.

LEMMA 4.3. *Suppose that $0 < \ell < k$ and $\gcd(k, \ell) = 1$.*

- (1) *The genus of $F_{k\mathbf{a}+\ell\mathbf{b}}$ equals the one of $F_{k\mathbf{a}-\ell\mathbf{b}}$.*

- (2) *The genus of $F_{k\mathfrak{p}+\ell\mathfrak{q}}$ equals the one of $F_{k\mathfrak{p}-\ell\mathfrak{q}}$.*
- (3) *The genera of $F_{k\mathfrak{r}+\ell\mathfrak{s}}$ and $F_{k\mathfrak{r}-\ell\mathfrak{s}}$ equal k .*

PROOF. (1) By Lemma 4.1 (1), the genus of $F_{k\mathfrak{a}\pm\ell\mathfrak{b}}$ equals $k-2$ if $\gcd(2k\pm\ell, 5) = 5$ or $\gcd(5, k\pm 2\ell) = 5$. Otherwise its genus equals k . It is easy to check that

- $\gcd(2k + \ell, 5) = 5$ if and only if $\gcd(5, k - 2\ell) = 5$, and
- $\gcd(2k - \ell, 5) = 5$ if and only if $\gcd(5, k + 2\ell) = 5$.

This implies the desired claim (1).

By using a similar argument, one can prove (2). The claim (3) is obvious from Lemma 4.1 (3). □

LEMMA 4.4. *Suppose that $0 < \ell < k$. Then*

$$\lambda(k\mathfrak{a} \pm \ell\mathfrak{b}) = \lambda(k\mathfrak{p} \pm \ell\mathfrak{q}) = \lambda(k\mathfrak{r} \pm \ell\mathfrak{s}) = \lambda_{(k,\ell)}.$$

PROOF. We use Theorem 3.2. The dilatations of $k\mathfrak{a} \pm \ell\mathfrak{b}$ and $k\mathfrak{p} \pm \ell\mathfrak{q}$ are the largest real root of

$$f_{(2k\pm\ell, 2k\pm 2\ell, k\pm 2\ell)}(t) = f_{(k, 2k\pm 2\ell, \pm\ell)}(t) = (t^{k\pm\ell} + 1)(t^{2k} - t^{k+\ell} - t^k - t^{k-\ell} + 1).$$

The dilatation of $k\mathfrak{r} \pm \ell\mathfrak{s}$ is the largest real root of

$$f_{(k\pm\ell, k\mp\ell, -k)}(t) = (t^k + 1)(t^{2k} - t^{k+\ell} - t^k - t^{k-\ell} + 1).$$

Since the absolute values of all roots of $t^{k\pm\ell} + 1$ and $t^k + 1$ are equal to 1, one finishes the proof. □

By Proposition 3.5 and (4.2), we immediately obtain the following.

COROLLARY 4.5. *Suppose that $k > 0$, $-k < \ell < k$ and $\gcd(k, \ell) = 1$.*

- (1) *The monodromy of the fibration associated to $k\mathfrak{a} + \ell\mathfrak{b}$ on N is orientable if and only if k is odd and ℓ is even.*
- (2) *The monodromy of the fibration associated to $k\mathfrak{p} + \ell\mathfrak{q}$ on N is orientable if and only if k is even and ℓ is odd.*
- (3) *The monodromy of the fibration associated to $k\mathfrak{r} + \ell\mathfrak{s}$ on N is orientable if and only if both k and ℓ are odd.*

The following can be obtained from Proposition 3.3

COROLLARY 4.6. *Suppose that $k > 0$, $-k < \ell < k$ and $\gcd(k, \ell) = 1$.*

- (1) *The singularity data of the monodromy of the fibration associated to $k\mathbf{a} + \ell\mathbf{b}$ on N is given by*

$$\left(\underbrace{\left(\frac{2k + \ell}{\gcd(2k + \ell, 5)} - 2, \dots, \frac{2k + \ell}{\gcd(2k + \ell, 5)} - 2 \right)}_{\gcd(2k + \ell, 5)}, \underbrace{\left(\frac{2k - \ell}{\gcd(5, k + 2\ell)} - 2, \dots, \frac{2k - \ell}{\gcd(5, k + 2\ell)} - 2 \right)}_{\gcd(5, k + 2\ell)} \right).$$

- (2) *The singularity data of the monodromy of the fibration associated to $k\mathbf{p} + \ell\mathbf{q}$ on N is given by*

$$\left(\underbrace{\left(\frac{k}{\gcd(k, 3)} - 2, \dots, \frac{k}{\gcd(k, 3)} - 2 \right)}_{\gcd(k, 3)}, \underbrace{\left(\frac{3k}{\gcd(3, \ell)} - 2, \dots, \frac{3k}{\gcd(3, \ell)} - 2 \right)}_{\gcd(3, \ell)} \right).$$

- (3) *The singularity data of the monodromy of the fibration associated to $k\mathbf{r} + \ell\mathbf{s}$ on N is given by*

$$(k + \ell - 2, k - \ell - 2, \underbrace{2, \dots, 2}_k).$$

REMARK 4.7. (1) The stable foliation for the monodromy of the fibration associated to $k\mathbf{a} + \ell\mathbf{b}$ (resp. $k\mathbf{p} + \ell\mathbf{q}$) is 2-pronged at each boundary component on T_β . (Hence there is no singular leaf on $\partial_\beta F_{k\mathbf{a} + \ell\mathbf{b}}$ (resp. $\partial_\beta F_{k\mathbf{p} + \ell\mathbf{q}}$).

- (2) The stable foliation for the monodromy of the fibration associated to $k\mathbf{r} + \ell\mathbf{s}$ is 4-pronged at each boundary component on T_γ .

LEMMA 4.8. *Suppose that $k > 0$, $-k < \ell < k$ and $\gcd(k, \ell) = 1$.*

- (1) *The stable foliation for the monodromy of the fibration associated to $k\mathbf{a} + \ell\mathbf{b}$ is 1-pronged at a boundary component if and only if $(k, \ell) \in \{(2, \pm 1), (3, \pm 1), (4, \pm 3)\}$.*
- (2) *The stable foliation for the monodromy of the fibration associated to $k\mathbf{p} + \ell\mathbf{q}$ is 1-pronged at a boundary component if and only if $(k, \ell) \in \{(1, 0), (3, \pm 1), (3, \pm 2)\}$.*
- (3) *The stable foliation for the monodromy of the fibration associated to $k\mathbf{r} + \ell\mathbf{s}$ is 1-pronged at a boundary component if and only if $k + \ell = 1$ or $k - \ell = 1$.*




Figure 6. (left) Thurston norm ball. (right) Open cone $\text{int}(C_{\Delta(r)})$ [shaded region].

PROOF. (1) By Corollary 4.6, the stable foliation of the monodromy for $k\alpha + \ell\mathbf{b}$ is 1-pronged at a boundary component if and only if $(2k + \ell)/\gcd(2k + \ell, 5) = 1$ or $(2k - \ell)/\gcd(5, k + 2\ell) = 1$.

Suppose that $(2k + \ell)/\gcd(2k + \ell, 5) = 1$. Clearly $\gcd(2k + \ell, 5) = 1$ or 5 . If $\gcd(2k + \ell, 5) = 1$, then $2k + \ell = 1$. Since $-k < \ell < k$, one has $-k < -2k + 1 < k$ which implies that $1/3 < k < 1$. This does not occur since k is an integer. If $\gcd(2k + \ell, 5) = 5$, then $2k + \ell = 5$. Since $-k < \ell < k$, one has $-k < 5 - 2k < k$ which implies $5/3 < k < 5$. Hence $(k, \ell) \in \{(2, 1), (3, -1), (4, -3)\}$.

Suppose that $(2k - \ell)/\gcd(5, k + 2\ell) = 1$. In this case one sees that $(k, \ell) \in \{(2, -1), (3, 1), (4, 3)\}$. This completes the proof of (1).

By using the same argument one can prove (2), (3). □

REMARK 4.9. Suppose that $k > 0$, $-k < \ell < k$ and $\gcd(k, \ell) = 1$. By Lemma 4.8 and Corollary 3.7, we see that:

- (1) $N((-3k - 4\ell)/(2k + \ell), -3/2, (-4k - 3\ell)/(k + 2\ell))$ is non-hyperbolic for $(k, \ell) \in \{(2, \pm 1), (3, \pm 1), (4, \pm 3)\}$. Otherwise it is hyperbolic.
- (2) $N((-2k - 3\ell)/k, -1/2, (-3k - 2\ell)/\ell)$ is non-hyperbolic for $(k, \ell) \in \{(1, 0), (3, \pm 1), (3, \pm 2)\}$. Otherwise it is hyperbolic.
- (3) Suppose that $k + \ell = 1$ or $k - \ell = 1$. Then we have the following. $N(\ell/(k + \ell), -\ell/(k - \ell), 2)$ is non-hyperbolic if $(k, \ell) \in \{(2, \pm 1), (3, \pm 2), (4, \pm 3)\}$ and it is hyperbolic if $(k, \ell) \notin \{(2, \pm 1), (3, \pm 2), (4, \pm 3)\}$. Suppose that $k + \ell \neq 1$ and $k - \ell \neq 1$. Then $N(\ell/(k + \ell), -\ell/(k - \ell), 2)$ is hyperbolic.

For each of $N(-3/2)$, $N(-1/2)$ and $N(2)$, its Thurston norm ball of radius 2 is a rectangle with vertices $(k, \ell) = (\pm 1, \pm 1)$ illustrated in Figure 6. By using (3.1) we see that for $k, \ell \in \mathbb{R}$,

$$\|k\bar{\mathbf{a}} + \ell\bar{\mathbf{b}}\| = \|k\bar{\mathbf{p}} + \ell\bar{\mathbf{q}}\| = \|k\bar{\mathbf{r}} + \ell\bar{\mathbf{s}}\| = \max\{2|k|, 2|\ell|\}. \tag{4.3}$$

The following lemma asserts that fibered faces for $N(-3/2)$ and $N(-1/2)$ has

a symmetry. Thus, for the study of monodromies on fibrations on $N(-3/2)$ (resp. $N(-1/2)$), it is enough to consider the open cone over an arbitrary picked fibered face.

LEMMA 4.10. *Suppose that $k > 0$ and $-k < \ell < k$.*

- (1) $F_{k\bar{a}+\ell\bar{b}} = F_{\ell\bar{a}+k\bar{b}}$ and $\lambda(k\bar{a} + \ell\bar{b}) = \lambda(\ell\bar{a} + k\bar{b})$.
- (2) $F_{k\bar{p}+\ell\bar{q}} = F_{\ell\bar{p}+k\bar{q}}$ and $\lambda(k\bar{p} + \ell\bar{q}) = \lambda(\ell\bar{p} + k\bar{q})$.

PROOF. See Lemma 4.2 and Remark 4.7 (1). □

Let us fix an open cones

$$\text{int}(C_{\Delta(-3/2)}) = \{k\bar{a} + \ell\bar{b} \mid k > 0, -k < \ell < k\} \subset H_2\left(N\left(\frac{-3}{2}\right), \partial N\left(\frac{-3}{2}\right); \mathbb{R}\right),$$

$$\text{int}(C_{\Delta(-1/2)}) = \{k\bar{p} + \ell\bar{q} \mid k > 0, -k < \ell < k\} \subset H_2\left(N\left(\frac{-1}{2}\right), \partial N\left(\frac{-1}{2}\right); \mathbb{R}\right),$$

$$\text{int}(C_{\Delta(2)}) = \{k\bar{r} + \ell\bar{s} \mid k > 0, -k < \ell < k\} \subset H_2(N(2), \partial N(2); \mathbb{R}).$$

Lemmas 4.11 and 4.12 tell us that it is enough to consider the fibered classes of $\text{int}(C_{\Delta(r)})$ for $0 < \ell < k$.

LEMMA 4.11. *Suppose that $0 < \ell < k$ and $\text{gcd}(k, \ell) = 1$.*

- (1) $F_{k\bar{a}\pm\ell\bar{b}} = \Sigma_{k-2,6}$ if $\text{gcd}(2k + \ell, 5) = 5$ or $\text{gcd}(5, k + 2\ell) = 5$. Otherwise $F_{k\bar{a}+\ell\bar{b}} = \Sigma_{k,2}$.
- (2) $F_{k\bar{p}\pm\ell\bar{q}} = \Sigma_{k-1,4}$ if $\text{gcd}(k, 3) = 3$ or $\text{gcd}(3, \ell) = 3$. Otherwise $F_{k\bar{p}+\ell\bar{q}} = \Sigma_{k,2}$.
- (3) $F_{k\bar{r}\pm\ell\bar{s}} = \Sigma_{k,2}$.

PROOF. The number of components of $\partial_\beta F_{k\bar{a}+\ell\bar{b}}$ equals $k + \ell$. By Lemma 4.1 (1), we have the desired claim (1). One can prove (2), (3) by using Lemma 4.1 (2), (3) respectively. □

LEMMA 4.12. *Suppose that $0 < \ell < k$. Then*

$$\lambda(k\bar{a} \pm \ell\bar{b}) = \lambda(k\bar{p} \pm \ell\bar{q}) = \lambda(k\bar{r} \pm \ell\bar{s}) = \lambda_{(k,\ell)}.$$

PROOF. See Lemma 4.4 and Remark 4.7. □

PROPOSITION 4.13. *Let $r \in \{-3/2, -1/2, 2\}$. The minimum of $\text{Ent} : \text{int}(C_{\Delta(r)}) \rightarrow \mathbb{R}$ equals $2 \log \lambda_{(1,0)} = 2 \log((3 + \sqrt{5})/2)$. The minimizer is given by \bar{a} if $r = -3/2$, \bar{p} if $r = -1/2$ and \bar{r} if $r = 2$.*

PROOF. Recall that Ent is constant on each ray thorough the origin and it attains its minimum at a unique ray. By Lemma 4.12 and (4.3), this ray must satisfy $\ell = 0$. Using the representative $(k, \ell) = (1, 0)$, Theorem 3.2 implies that $\min \text{Ent} = 2 \log \lambda_{(1,0)} = 2 \log((3 + \sqrt{5})/2)$. \square

REMARK 4.14. The monodromies of the fibrations associated to $\bar{\mathbf{a}}$ (resp. $\bar{\mathbf{p}}$) on $N(-3/2)$ (resp. $N(-1/2)$) are intriguing examples.

- (1) $N(-3/2)$ admits a fiber of genus 1 with 2 boundary components corresponding to $\bar{\mathbf{a}}$. The stable foliation of its monodromy $\Phi : \Sigma_{1,2} \rightarrow \Sigma_{1,2}$ is 2-pronged at each boundary component. Thus Φ extends to the monodromy $\bar{\Phi} : \Sigma_{1,1} \rightarrow \Sigma_{1,1}$ of the fibration on $N(-3/2, -3/2)$ (which is the figure-8 knot sister manifold, see [20, Table A.2]) by capping the boundary component of $\Sigma_{1,2}$ on T_α . It is well-known that $\bar{\Phi}$ realizes the minimal dilatation $(3 + \sqrt{5})/2$ among pseudo-Anosovs on $\Sigma_{1,1}$.
- (2) $N(-1/2)$ admits a fiber of genus 0 with 4 boundary components corresponding to $\bar{\mathbf{p}}$. The stable foliation of its monodromy Φ fixes a boundary component, and hence it can be considered that Φ is a pseudo-Anosov homeomorphism on a 3-punctured disk D_3 . This monodromy Φ realizes the minimal dilatation $\delta(D_3) = (3 + \sqrt{5})/2$ among pseudo-Anosovs on D_3 .

4.2. Property of algebraic integers $\lambda_{(k,\ell)}$.

LEMMA 4.15. *Suppose that $1 < \ell + 1 < k$ and $\gcd(k, \ell) = 1$. Then $\lambda_{(k+1,\ell)} < \lambda_{(k,\ell)} < \lambda_{(k,\ell+1)}$.*

PROOF. The ray that attains the minimum of $\text{Ent} : \text{int}(C_{\Delta(r)}) \rightarrow \mathbb{R}$ satisfies $\ell = 0$. Recall that the function $1/\text{ent}(\cdot) : \text{int}(C_{\Delta(r)}(\mathbb{Q})) \rightarrow \mathbb{R}$ is strictly concave. Thus one has

$$\log \lambda_{(k,k\ell/(k+1))} < \log \lambda_{(k,\ell)} < \log \lambda_{(k,\ell+1)}.$$

The inequality $\log \lambda_{(k+1,\ell)} < \log \lambda_{(k,\ell)}$ holds since

$$\begin{aligned} \log \lambda_{(k+1,\ell)} &= \text{ent}((k+1)\bar{\mathbf{a}} + \ell\bar{\mathbf{b}}) = \frac{k}{k+1} \text{ent}\left(k\bar{\mathbf{a}} + \frac{k\ell}{k+1}\bar{\mathbf{b}}\right) \\ &= \frac{k}{k+1} \log \lambda_{(k,k\ell/(k+1))} < \log \lambda_{(k,\ell+1)}. \end{aligned}$$

Hence $\log \lambda_{(k+1,\ell)} < \log \lambda_{(k,\ell)} < \log \lambda_{(k,\ell+1)}$. \square

LEMMA 4.16. *For any fixed $\ell > 0$,*

$$\lim_{k \rightarrow \infty} k \log \lambda_{(k,\ell)} = \log \lambda_{(1,0)} = \log \left(\frac{3 + \sqrt{5}}{2} \right).$$

PROOF. The ray through $k\bar{\mathbf{a}} + \ell\bar{\mathbf{b}}$ from the origin goes to the ray through $\bar{\mathbf{a}}$ if k goes to ∞ . Hence

$$\lim_{k \rightarrow \infty} \text{Ent}(k\bar{\mathbf{a}} + \ell\bar{\mathbf{b}}) = \lim_{k \rightarrow \infty} 2k \log \lambda_{(k,\ell)} = \text{Ent}(\bar{\mathbf{a}}) = 2 \log \lambda_{(1,0)}.$$

This completes the proof. □

PROPOSITION 4.17. *If $\lambda_{(k+1,\ell)} < \lambda_{(k,1)}$ for some $k \geq \ell \geq 2$, then $\lambda_{(k+2,\ell)} < \lambda_{(k+1,1)}$.*

PROOF. We denote the homology class $k\bar{\mathbf{a}} + \ell\bar{\mathbf{b}}$ by (k, ℓ) . Since Ent is constant on each ray thorough the origin, $k \text{ent}(k, \ell) = \text{ent}(1, \ell/k)$. One takes 4 points

$$p_1 = \left(1, \frac{1}{k+1} \right), p_2 = \left(1, \frac{1}{k} \right), p_3 = \left(1, \frac{\ell}{k+2} \right), p_4 = \left(1, \frac{\ell}{k+1} \right),$$

see Figure 7. We have $1/(k+1) < 1/k < 2/(k+2) \leq \ell/(k+2) < \ell/(k+1)$. Let us set t, t' and c as follows.

$$\begin{aligned} 0 < t &= \frac{|p_3 - p_2|}{|p_4 - p_2|} = \frac{(k+1)(k\ell - k - 2)}{(k+2)(k\ell - k - 1)} < 1, \\ 0 < t' &= \frac{|p_3 - p_2|}{|p_3 - p_1|} = \frac{|p_4 - p_2|}{|p_3 - p_1|} t = \frac{(k+2)(k\ell - k - 1)}{k(k\ell - k + \ell - 2)} t < 1, \\ 1 < c &= \frac{(k+2)(k\ell - k - 1)}{k(k\ell - k + \ell - 2)}. \end{aligned} \tag{4.4}$$

(Hence $t' = ct$.) Then

$$\begin{aligned} |p_3 - p_2| : |p_4 - p_3| &= t : 1 - t, \\ |p_3 - p_2| : |p_2 - p_1| &= ct : 1 - ct. \end{aligned}$$

These ratios together with Theorem 2.3 imply that

$$\frac{1}{(k+2) \text{ent}(k+2, \ell)} > (1-t) \frac{1}{k \text{ent}(k, 1)} + t \frac{1}{(k+1) \text{ent}(k+1, \ell)}, \quad \text{and} \tag{4.5}$$

$$\frac{1}{k \operatorname{ent}(k, 1)} > ct \frac{1}{(k+1) \operatorname{ent}(k+1, 1)} + (1-ct) \frac{1}{(k+2) \operatorname{ent}(k+2, \ell)}. \tag{4.6}$$

By (4.5) and by the assumption $\operatorname{ent}(k, 1) > \operatorname{ent}(k+1, \ell)$,

$$\begin{aligned} & \frac{1}{(k+2) \operatorname{ent}(k+2, \ell)} \\ & > (1-t) \frac{1}{k \operatorname{ent}(k, 1)} + t \frac{1}{(k+1) \operatorname{ent}(k+1, \ell)} \\ & > (1-t) \frac{1}{k \operatorname{ent}(k, 1)} + t \frac{1}{(k+1) \operatorname{ent}(k, 1)} \\ & = \frac{k+1-t}{k+1} \frac{1}{k \operatorname{ent}(k, 1)} \\ & > \frac{k+1-t}{k+1} \left\{ ct \frac{1}{(k+1) \operatorname{ent}(k+1, 1)} + (1-ct) \frac{1}{(k+2) \operatorname{ent}(k+2, \ell)} \right\}. \end{aligned}$$

The last inequality is given by (4.6). Hence

$$\left\{ \frac{1}{k+2} - \frac{(k+1-t)(1-ct)}{(k+1)(k+2)} \right\} \frac{1}{\operatorname{ent}(k+2, \ell)} > \frac{(k+1-t)ct}{(k+1)^2} \frac{1}{\operatorname{ent}(k+1, 1)},$$

which gives, by calculation,

$$\frac{(k+1-t)c+1}{k+2} \frac{1}{\operatorname{ent}(k+2, \ell)} > \frac{(k+1-t)c}{k+1} \frac{1}{\operatorname{ent}(k+1, 1)}.$$

Thus




Figure 7. Four boxes \square (from the bottom to the top) on the line $k=1$ indicate p_1, p_2, p_3 and p_4 .

$$\text{ent}(k + 2, \ell) < \left\{ \frac{k + 1}{(k + 1 - t)c} \right\} \left\{ \frac{(k + 1 - t)c + 1}{k + 2} \right\} \text{ent}(k + 1, 1).$$

For the proof of the claim it is enough to verify the equality $\{(k + 1)/(k + 1 - t)c\}\{(k + 1 - t)c + 1/(k + 2)\} = 1$. Clearly,

$$\begin{aligned} & \left\{ \frac{k + 1}{(k + 1 - t)c} \right\} \left\{ \frac{(k + 1 - t)c + 1}{k + 2} \right\} = 1 \\ & \Leftrightarrow (k + 1)\{(k + 1 - t)c + 1\} = (k + 2)(k + 1 - t)c \\ & \Leftrightarrow (k + 1)(k + 1 - t)c + k + 1 = (k + 2)(k + 1 - t)c \\ & \Leftrightarrow k + 1 = (k + 1 - t)c \\ & \Leftrightarrow c = \frac{k + 1}{k + 1 - t}. \end{aligned}$$

One can verify the last equality $c = (k + 1)/(k + 1 - t)$ by substituting the constants t and c given by (4.4). □

4.3. Fibrations of manifolds obtained from $N(-1/2), N(-3/2)$ and $N(2)$ by Dehn filling two cusps.

As a consequence of Lemma 4.8, we see the following.

REMARK 4.18. If $(k, \ell) \notin \{(2, \pm 1), (3, \pm 1), (4, \pm 3)\}$, then the monodromy $\Phi_{k\mathbf{a}+\ell\mathbf{b}} : F_{k\mathbf{a}+\ell\mathbf{b}} \rightarrow F_{k\mathbf{a}+\ell\mathbf{b}}$ of the fibration associated to $k\mathbf{a} + \ell\mathbf{b}$ on N extends to the monodromy $\bar{\Phi}_{k\mathbf{a}+\ell\mathbf{b}} : \bar{F}_{k\mathbf{a}+\ell\mathbf{b}} \rightarrow \bar{F}_{k\mathbf{a}+\ell\mathbf{b}}$ of the fibration on $N((-3k - 4\ell)/(2k + \ell), -3/2, (-4k - 3\ell)/(k + 2\ell))$ with the dilatation $\lambda_{(k,\ell)} (= \lambda(\Phi_{k\mathbf{a}+\ell\mathbf{b}}))$. Similarly, if $(k, \ell) \notin \{(1, 0), (3, \pm 1), (3, \pm 2)\}$ (resp. if $k + \ell \neq 1$ and $k - \ell \neq 1$), then the monodromy $\Phi_{k\mathbf{p}+\ell\mathbf{q}}$ (resp. $\Phi_{k\mathbf{r}+\ell\mathbf{s}}$) of the fibration associated to $k\mathbf{p} + \ell\mathbf{q}$ (resp. $k\mathbf{r} + \ell\mathbf{s}$) on N extends to the monodromy $\bar{\Phi}_{k\mathbf{p}+\ell\mathbf{q}}$ (resp. $\bar{\Phi}_{k\mathbf{r}+\ell\mathbf{s}}$) of the fibration on $N((-2k - 3\ell)/k, -1/2, (-3k - 2\ell)/\ell)$ (resp. $N(\ell/(k + \ell), -\ell/(k - \ell), 2)$) with the dilatation $\lambda_{(k,\ell)}$.

Let $\bar{\phi}_{k\mathbf{a}+\ell\mathbf{b}}, \bar{\phi}_{k\mathbf{p}+\ell\mathbf{q}}$ and $\bar{\phi}_{k\mathbf{r}+\ell\mathbf{s}}$ be elements of $\text{Mod}(\Sigma_g)$ containing $\bar{\Phi}_{k\mathbf{a}+\ell\mathbf{b}}, \bar{\Phi}_{k\mathbf{p}+\ell\mathbf{q}}$ and $\bar{\Phi}_{k\mathbf{r}+\ell\mathbf{s}}$ as a representative.

PROPOSITION 4.19. For any fixed integer $\ell > 0$, we have the following.

- (1) $\lim_{\substack{k \rightarrow \infty \\ \gcd(k,\ell)=1}} \text{vol}(\bar{\phi}_{k\mathbf{a}+\ell\mathbf{b}}) = \text{vol}\left(N\left(\frac{-3}{2}\right)\right) \approx 3.66386.$
- (2) $\lim_{\substack{k \rightarrow \infty \\ \gcd(k,\ell)=1}} \text{vol}(\bar{\phi}_{k\mathbf{p}+\ell\mathbf{q}}) = \text{vol}\left(N\left(\frac{-1}{2}\right)\right) \approx 4.05977.$

$$(3) \quad \lim_{\substack{k \rightarrow \infty \\ \gcd(k, \ell) = 1}} \text{vol}(\overline{\phi}_{k\tau + \ell\epsilon}) = \text{vol}(N(2)) \approx 4.41533.$$

PROOF. We will prove the claim (1). The proof of claims (2), (3) is similar. The mapping torus $\mathbb{T}(\overline{\phi}_{k\mathfrak{a} + \ell\mathfrak{b}})$ is homeomorphic to $N((-3k - 4\ell)/(2k + \ell), -3/2, (-4k - 3\ell)/(k + 2\ell))$. Since $\gcd(-3k - 4\ell, 2k + \ell)$ (resp. $\gcd(-4k - 3\ell, k + 2\ell)$) is either 1 or 5, the two points

$$\begin{aligned} & \left(\frac{-3k - 4\ell}{\gcd(-3k - 4\ell, 2k + \ell)}, \frac{2k + \ell}{\gcd(-3k - 4\ell, 2k + \ell)} \right) \in \mathbb{R}^2, \\ & \left(\frac{-4k - 3\ell}{\gcd(-4k - 3\ell, k + 2\ell)}, \frac{k + 2\ell}{\gcd(-4k - 3\ell, k + 2\ell)} \right) \in \mathbb{R}^2 \end{aligned}$$

tend to ∞ as k tends to ∞ . Thurston’s hyperbolic Dehn surgery theorem (see [28]) implies the volume of $N((-3k - 4\ell)/(2k + \ell), -3/2, (-4k - 3\ell)/(k + 2\ell))$ converges to $\text{vol}(N(-3/2))$ as k tends to ∞ . \square

PROOF OF THEOREM 1.5 (Case $r = -3/2$). For the proof of (1), first of all we find a pair $(k(g), \ell(g)) = (g + \tilde{k}(g), \ell(g))$ for each $g \geq 3$ satisfying the following: the both $\tilde{k}(g) > 0$, $\ell(g) > 0$ are bounded, and the genus of $F_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}$ equals g . Next we check that the stable foliation of $\Phi_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}$ has no 1 prong at each boundary component of $F_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}$. Then one can extend $\Phi_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}$ to the pseudo-Anosov homeomorphism $\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}$ on a closed surface of genus g . This finishes the proof of (1). In fact, by Lemma 4.16

$$\lim_{g \rightarrow \infty} k(g) \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) = \lim_{g \rightarrow \infty} k(g) \log \lambda_{(k(g), \ell(g))} = \log \left(\frac{3 + \sqrt{5}}{2} \right).$$

On the other hand

$$\lim_{g \rightarrow \infty} \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) = \lim_{g \rightarrow \infty} \frac{1}{g + \tilde{k}(g)} \log \left(\frac{3 + \sqrt{5}}{2} \right) = 0.$$

Thus one obtains

$$\begin{aligned} \log \left(\frac{3 + \sqrt{5}}{2} \right) &= \lim_{g \rightarrow \infty} k(g) \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) \\ &= \lim_{g \rightarrow \infty} (g + \tilde{k}(g)) \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) \\ &= \lim_{g \rightarrow \infty} g \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) + \lim_{g \rightarrow \infty} \tilde{k}(g) \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) \\ &= \lim_{g \rightarrow \infty} g \log \lambda(\overline{\Phi}_{k(g)\mathfrak{a} + \ell(g)\mathfrak{b}}) + 0, \end{aligned}$$

which implies (1).

One sees that the genera of F_{3a+2b} and F_{4a+b} equal 3 and 4 respectively. If $g \not\equiv 0 \pmod{5}$ and $g \geq 6$, the genus of $F_{g_{a+5b}}$ equals g . In the case $g \equiv 2 \pmod{5}$ and $g \geq 7$, the genus of $F_{g_{a+b}}$ equals $g - 2 \equiv 0 \pmod{5}$. By Lemma 4.8, one has the desired equality (1).

The claim (2) on the volume holds by Proposition 4.19 (1).

(Case $r = -1/2$.) If $g \equiv 0, 1 \pmod{3}$ and $g \geq 3$, the genus of $F_{(g+1)p+3q}$ equals g . If $g \equiv 2 \pmod{3}$ and $g \geq 3$, the genus of $F_{(g+1)p+q}$ equals g . By Lemma 4.8 and Proposition 4.19 (2), one obtains the claims (1), (2).

(Case $r = 2$.) The genus of $F_{g_{r+s}}$ equals g . By Lemma 4.8 and Proposition 4.19 (3), one obtains the claims (1), (2). \square

REMARK 4.20. For $g \geq 4$ even, there exists a Σ_g -bundle over the circle with the dilatation $\lambda_{(g,1)}$, which is obtained from the extension of the monodromy of the fibration associated to $g\tau + s$ on N . However the invariant foliation associated to this Σ_g -bundle over the circle is non-orientable, see Corollary 4.5 and Question 1.10. (In the case $g = 2$, the monodromy of the fibration associated to $2\tau + s$ cannot extend to the *pseudo-Anosov* monodromy on a closed fiber, see Remark 4.9 (3).)

For $r \in \mathbb{Q}$, $\Lambda_g(r)$ (resp. $\Lambda_g^+(r)$) is defined to be the set of dilatations of all Σ_g -bundles (resp. all Σ_g -bundles with orientable invariant foliations) which are obtained from $N(r)$ by Dehn filling two cusps along the boundary slopes of the fibers of $N(r)$. Recall that \mathcal{U} and \mathcal{U}^+ are finite sets of fibered hyperbolic 3-manifolds defined in the introduction.

LEMMA 4.21. $N(2) \in \mathcal{U}^+$.

PROOF. One sees that the pseudo-Anosov $\bar{\phi}_{3\tau+s} \in \text{Mod}(\Sigma_3)$ is orientable and has the dilatation $\lambda_{(3,1)} (= \delta_3^+)$. Hence $\delta_3^+ \in \Lambda_3^+(2)$. \square

In the rest of this section, we mainly consider the sets $\Lambda_g^{(+)}(-1/2)$ and $\Lambda_g^{(+)}(-3/2)$. We first recall the number $\min \Lambda_g^{(+)}(-1/2)$.

PROPOSITION 4.22 ([9]). Let $g \geq 3$.

- (1) $\lambda_{(g+1,3)} = \min \Lambda_g(-1/2)$ if $g \equiv 0, 1, 3, 4 \pmod{6}$.
- (2) $\lambda_{(g+1,1)} = \min \Lambda_g(-1/2)$ if $g \equiv 2, 5 \pmod{6}$.

PROPOSITION 4.23 ([9]). Let $g \geq 3$.

- (1) $\lambda_{(g+1,3)} = \min \Lambda_g^+(-1/2)$ if $g \equiv 1, 3 \pmod{6}$.

- (2) $\lambda_{(g,1)} = \min \Lambda_g^+(-1/2)$ if $g \equiv 2, 4 \pmod{6}$.
- (3) $\lambda_{(g+1,1)} = \min \Lambda_g^+(-1/2)$ if $g \equiv 5 \pmod{6}$.

LEMMA 4.24 ([9]). $N(-1/2) \in \mathcal{U} \cap \mathcal{U}^+$.

PROOF. One sees that $\bar{\phi}_{2p+q} \in \text{Mod}(\Sigma_2)$ is an orientable pseudo-Anosov mapping class having dilatation $\lambda_{(2,1)} (= \delta_2 = \delta_2^+)$. Hence $\delta_2 = \delta_2^+ \in \Lambda_2(-1/2) \cap \Lambda_2^+(-1/2)$. □

We turn to $N(-3/2)$. By Lemma 4.1 (1), if $\lambda \in \Lambda_g(-3/2)$, then $\lambda = \lambda_{(g+2,\ell)}$ for some $1 \leq \ell < g+2$ or $\lambda = \lambda_{(g,\ell')}$ for some $1 \leq \ell' < g$.

It is easy to verify the following by a direct computation.

LEMMA 4.25. *For integers k and ℓ , $\gcd(2k + \ell, 5) = 5$ or $\gcd(5, k + 2\ell) = 5$ if and only if k and ℓ are either (1), (2), (3), (4) or (5) in the following table.*

	$k \pmod{5}$	$\ell \pmod{5}$
(1)	0	0
(2)	2, 3	1
(3)	1, 4	2
(4)	1, 4	3
(5)	2, 3	4

We compute $\min \Lambda_g(-3/2)$ for $g \equiv 0, 1, 3, 5, 6, 7, 8, 9 \pmod{10}$.

- PROPOSITION 4.26. (1) $\lambda_{(g+2,1)} = \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ if $g \equiv 0, 1, 5, 6 \pmod{10}$ and $g \geq 5$.
- (2) $\lambda_{(g+2,2)} = \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ if $g \equiv 7, 9 \pmod{10}$ and $g \geq 7$.
- (3) $\lambda_{(g,2)} = \min \Lambda_g(-3/2) > \min \Lambda_g(-1/2)$ if $g \equiv 3 \pmod{10}$ and $g \geq 3$.
- (4) Let $g \equiv 8 \pmod{10}$ and $g \geq 8$.
- (i) $\lambda_{(g,3)} = \min \Lambda_g(-3/2) > \min \Lambda_g(-1/2)$ if $g \equiv 8, 28 \pmod{30}$,
 - (ii) $\lambda_{(g,5)} = \min \Lambda_g(-3/2) > \min \Lambda_g(-1/2)$ if $g \equiv 18 \pmod{30}$.

PROOF. (1) If $k \equiv 2, 3 \pmod{5}$, then $\gcd(2k+1, 5) = 5$ or $\gcd(5, k+2) = 5$. We set $k = g+2$. (Hence $g \equiv 0, 1 \pmod{5}$ or equivalently $g \equiv 0, 1, 5, 6 \pmod{10}$.) The genus of $F_{(g+2)a+b}$ is equal to g by Lemma 4.1 (1), and hence $\lambda_{(g+2,1)} \in \Lambda_g(-3/2)$ by Remark 4.18. One can check that $\lambda_{(g+2,1)}$ attains $\min \Lambda_g(-3/2)$ by Lemma 4.15. In fact for any $g > 1$, $1 \leq \ell' < g$ and $1 \leq \ell < g+2$, it follows that

$$\lambda_{(g+2,1)} < \lambda_{(g+1,1)} < \lambda_{(g,1)} \leq \lambda_{(g,\ell')} \quad \text{and} \quad \lambda_{(g+2,1)} < \lambda_{(g+2,\ell)}.$$

Thus $\lambda_{(g,1)} = \min \Lambda_g(-3/2)$.

By Proposition 4.22, the lower and upper bound of $\min \Lambda_g(-1/2)$ is given by

$$\lambda_{(g+1,1)} \leq \min \Lambda_g\left(\frac{-1}{2}\right) \leq \lambda_{(g+1,3)} \text{ for any } g. \tag{4.7}$$

Since $\lambda_{(g+2,1)} < \lambda_{(g+1,1)}$, one obtains the inequality $\min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$.

(2) If $k \equiv 1, 4 \pmod{5}$, then $\gcd(2k + 2, 5) = 5$ or $\gcd(5, k + 4) = 5$. We set $k = g + 2$. (Hence $g \equiv 2, 4 \pmod{5}$.) Suppose that $\gcd(g + 2, 2) = 1$. Then $\lambda_{(g+2,2)} \in \Lambda_g(-3/2)$ and $g \equiv 7, 9 \pmod{10}$ since g must be odd. One sees that $\lambda_{(g+2,1)} \notin \Lambda_g(-3/2)$ since $\gcd(2k + 1, 5) = 1$ and $\gcd(5, k + 2) = 1$. For any $g > 1$ and $1 \leq \ell < g$, it follows that $\lambda_{(g+1,1)} < \lambda_{(g,1)} \leq \lambda_{(g,\ell)}$. On the other hand

$$\lambda_{(5,2)} \approx 1.23039 < \lambda_{(4,1)} \approx 1.28064$$

and by Proposition 4.17, one has $\lambda_{(g+2,2)} < \lambda_{(g+1,1)}$ holds for any $g \geq 3$. Thus $\lambda_{(g+2,2)}$ attains $\min \Lambda_g(-3/2)$. The inequality $\min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ holds by (4.7).

(3), (4) Suppose that $g \equiv 3 \pmod{5}$, that is $g \equiv 3, 8 \pmod{10}$. One observes that the genus of $F_{(g+2)\mathfrak{a}+\ell\mathfrak{b}}$ equals $g + 2$ whenever $\gcd(g + 2, \ell) = 1$. Hence if $\lambda \in \Lambda_g(-3/2)$, then $\lambda = \lambda_{(g,\ell)}$ for some $1 \leq \ell < g$. Suppose that $g \equiv 3 \pmod{10}$. By Lemmas 4.1 (1) and 4.25, the genera of $F_{g\mathfrak{a}+\mathfrak{b}}$ and $F_{g\mathfrak{a}+2\mathfrak{b}}$ are $g - 2$ and g respectively. Hence $\lambda_{(g,2)} = \min \Lambda_g(-3/2)$.

One has

$$\lambda_{(3,2)} \approx 1.50614 > \lambda_{(3,1)} = \lambda_{(4,3)} \approx 1.40127,$$

and hence $\min \Lambda_3(-3/2) > \min \Lambda_3(-1/2)$. By Proposition 4.17 together with the inequality

$$\lambda_{(4,1)} \approx 1.28064 > \lambda_{(5,3)} \approx 1.26123,$$

one obtains $\lambda_{(k,1)} > \lambda_{(k+1,3)}$ for any $k \geq 4$. The inequality $\min \Lambda_g(-3/2) > \min \Lambda_g(-1/2)$ holds for $g \equiv 3 \pmod{10}$ and $g > 3$ since

$$\min \Lambda_g\left(\frac{-3}{2}\right) = \lambda_{(g,2)} > \lambda_{(g,1)} > \lambda_{(g+1,3)} \geq \min \Lambda_g\left(\frac{-1}{2}\right).$$

One completes the proof of (3). Similarly one can prove (4). □

REMARK 4.27. The pseudo-Anosov homeomorphism whose dilatation equals $\min \Lambda_g(-3/2)$ in the proof of Proposition 4.26 (1) (resp. (2)) is non-orientable (resp. orientable), see Corollary 4.5.

PROOF OF THEOREM 1.6. See Proposition 4.26 (1), (2). □

In the case $g \equiv 2, 4 \pmod{10}$, we compute $\min \Lambda_g(-3/2)$ under certain conditions of g .

PROPOSITION 4.28. *Let $g \equiv 2, 4 \pmod{10}$ and $g \geq 12$. Suppose that $g+2 \not\equiv 0 \pmod{4641(=3 \cdot 7 \cdot 13 \cdot 17)}$.*

- (1) $\lambda_{(g+2,3)} = \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ if $\gcd(g+2, 3) = 1$.
- (2) $\lambda_{(g+2,7)} = \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ if 3 divides $g+2$ and $\gcd(g+2, 7) = 1$.
- (3) $\lambda_{(g+2,13)} = \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ if $21(=3 \cdot 7)$ divides $g+2$ and $\gcd(g+2, 13) = 1$.
- (4) $\lambda_{(g+2,17)} = \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ if $273(=3 \cdot 7 \cdot 13)$ divides $g+2$ and $\gcd(g+2, 17) = 1$.

The following will be used for proving Proposition 4.28. Its proof is similar to the one for Proposition 4.26 (3).

LEMMA 4.29. (1) *Let $g \equiv 2 \pmod{10}$ and $g \geq 12$.*

(i) *Suppose that $g \equiv 2, 22 \pmod{30}$. If $\lambda_{(g,\ell)} \in \Lambda_g(-3/2)$, then $\ell \geq 3$.*

(ii) *Suppose that $g \equiv 12 \pmod{30}$. If $\lambda_{(g,\ell)} \in \Lambda_g(-3/2)$, then $\ell \geq 5$.*

(2) *Let $g \equiv 4 \pmod{10}$ and $g \geq 14$. Then $\lambda_{(g,1)} \in \Lambda_g(-3/2)$.*

LEMMA 4.30. *Suppose that $g \equiv 2, 4 \pmod{10}$ and $g \geq 12$. If $\gcd(g+2, \ell) = 1$, $\ell \equiv 2, 3 \pmod{5}$ and $0 < \ell < g+2$, then $\lambda_{(g+2,\ell)} \in \Lambda_g(-3/2)$.*

PROOF. We use Lemma 4.1 (1). We set $k = g+2$ ($k \equiv 1, 4 \pmod{5}$). If $\ell \equiv 2, 3 \pmod{5}$, then $\gcd(2k + \ell, 5) = 5$ or $\gcd(5, k + 2\ell) = 5$. Thus if ℓ satisfies that $\gcd(k, \ell) = \gcd(g+2, \ell) = 1$ and $0 < \ell < g+2$, then one obtains the desired claim $\lambda_{(g+2,\ell)} \in \Lambda_g(-3/2)$. □

One can check the following inequalities.

LEMMA 4.31. (1) $\lambda_{(9,7)} \approx 1.16873 < \lambda_{(8,1)} \approx 1.12876$.

(2) $\lambda_{(73,13)} \approx 1.013457447 < \lambda_{(72,1)} \approx 1.013457858$.

(3) $\lambda_{(125,17)} \approx 1.007791640 < \lambda_{(124,1)} \approx 1.007791898$.

PROOF OF PROPOSITION 4.28. (1) By Lemma 4.30, $\lambda_{(g+2,3)} \in \Lambda_g(-3/2)$. We have shown that $\lambda_{(k+1,3)} < \lambda_{(k,1)}$ for any $k \geq 4$ in the proof of Proposi-

tion 4.26 (3), (4). Hence $\lambda_{(g+2,3)} < \lambda_{(g+1,1)}$ for any $g \geq 3$. By (4.7), we have $\min \Lambda_g(-3/2) \leq \lambda_{(g+2,3)} < \lambda_{(g+1,1)} \leq \min \Lambda_g(-1/2)$. We can prove that $\lambda_{(g+2,3)}$ attains $\min \Lambda_g(-3/2)$ by using the foregoing argument together with Lemma 4.29 (1).

The claims (2), (3), (4) can be verified by using Lemmas 4.29, 4.30 and 4.31. □

PROOF OF THEOREM 1.7. See Proposition 4.28. □

QUESTION 4.32. Is it true that $\delta_g \leq \min \Lambda_g(-3/2) < \min \Lambda_g(-1/2)$ for all $g \equiv 2, 4 \pmod{10}$ and $g \geq 12$?

REMARK 4.33. Independently, Aaber and Dunfield identified the pair $(k(g), \ell(g))$ such that the pseudo-Anosov homeomorphism $\bar{\Phi}_{k(g)\mathfrak{a}+\ell(g)\mathfrak{b}} : \Sigma_g \rightarrow \Sigma_g$ which attains $\min \Lambda_g(-3/2)$ for large g . They proved that under a plausible assumption, the mapping class $\bar{\phi}_{k(g)\mathfrak{a}+\ell(g)\mathfrak{b}} = [\bar{\Phi}_{k(g)\mathfrak{a}+\ell(g)\mathfrak{b}}]$ has the least volume among pseudo-Anosov elements of $\text{Mod}(\Sigma_g)$ for large g , see [1].

We turn to $\min \Lambda_g^+(-3/2)$. By Corollary 4.5 (1) and Lemma 4.11 (1), one sees that if g is even, then there exist no orientable pseudo-Anosov monodromies with a closed fiber of genus g of fibrations on $N(-3/2)$. Hence in this case $\Lambda_g^+(-3/2) = \emptyset$. We compute $\min \Lambda_g^+(-3/2)$ for g odd.

PROPOSITION 4.34. *Let $g \geq 5$.*

- (1) $\lambda_{(g+2,2)} = \min \Lambda_g^+(-3/2) < \min \Lambda_g^+(-1/2)$ if $g \equiv 7, 9 \pmod{10}$.
- (2) $\lambda_{(g+2,4)} = \min \Lambda_g^+(-3/2) \leq \min \Lambda_g^+(-1/2)$ if $g \equiv 1, 5 \pmod{10}$. *The equality holds if and only if $g = 5$.*
- (3) $\lambda_{(g,2)} = \min \Lambda_g^+(-3/2) > \min \Lambda_g^+(-1/2)$ if $g \equiv 3 \pmod{10}$.

PROOF. We use Corollary 4.5 to see whether $\lambda_{(k,\ell)} \in \Lambda_g(-3/2)$ is an element of $\Lambda_g^+(-3/2)$ or not.

(1) We see that $\lambda_{(g+2,2)} \in \Lambda_g^+(-3/2)$, see Remark 4.27. By Proposition 4.26 (2), we have

$$\lambda_{(g+2,2)} = \min \Lambda_g \left(\frac{-3}{2} \right) = \min \Lambda_g^+ \left(\frac{-3}{2} \right) < \min \Lambda_g \left(\frac{-1}{2} \right) \leq \min \Lambda_g^+ \left(\frac{-1}{2} \right).$$

(2) It can be shown that $\lambda_{(g+2,4)} = \min \Lambda_g^+(-3/2)$. Since $\lambda_{(7,4)} = \lambda_{(6,1)}$, the equality $\min \Lambda_5^+(-3/2) = \min \Lambda_5^+(-1/2)$ holds. Suppose that $g \neq 5$. By Proposition 4.17 together with

$$\lambda_{(8,4)} \approx 1.14555 < \lambda_{(7,1)} \approx 1.14879,$$

we obtain the inequality $\lambda_{(k,4)} < \lambda_{(k-1,1)}$ for any $k \geq 8$. Thus $\min \Lambda_g^+(-3/2) < \min \Lambda_g^+(-1/2)$.

One can prove (3) by using a similar argument together with Proposition 4.26 (3). □

PROOF OF THEOREM 1.12. See Proposition 4.34 (1), (2). □

PROOF OF PROPOSITION 1.14. We have proved the inequality $(\lambda_{(g+2,2)} <)\lambda_{(g+2,4)} < \lambda_{(g+1,1)}$ for any $g \geq 6$ in the proof of Proposition 4.34 (2). By Theorem 1.12 and by the assumption $\delta_{g+1}^+ = \lambda_{(g+1,1)}$, one has

$$\delta_g^+ \leq \max\{\lambda_{(g+2,2)}, \lambda_{(g+2,4)}\} \leq \lambda_{(g+2,4)} < \lambda_{(g+1,1)} = \delta_{g+1}^+.$$

This completes the proof. □

- REMARK 4.35. (1) The $(-2, 3, 7)$ -pretzel knot complement is homeomorphic to $N(-3/2, -8/3)$, see [20, Table A.4]. On the other hand, $\text{slope}(7\mathbf{a} + 4\mathbf{b}) = (-37/18, -3/2, -8/3)$. The monodromy $\Phi_{7\mathbf{a}+4\mathbf{b}} : \Sigma_{5,17} \rightarrow \Sigma_{5,17}$ of the fibration associated to $7\mathbf{a} + 4\mathbf{b}$ on N is orientable (see Corollary 4.5 (1)) and its singularity data is given by (16) (see Corollary 4.6 (1)). Thus $\Phi_{7\mathbf{a}+4\mathbf{b}} : \Sigma_{5,17} \rightarrow \Sigma_{5,17}$ extends to the pseudo-Anosov monodromy $\bar{\Phi}_{7\mathbf{a}+4\mathbf{b}} : \Sigma_{5,1} \rightarrow \Sigma_{5,1}$ of the fibration on $N(-3/2, -8/3)$ (with the dilatation $\lambda_{(7,4)}$) by capping all the boundary components on $T_\beta \cup T_\gamma$.
- (2) $\bar{\Phi}_{7\mathbf{a}+4\mathbf{b}} : \Sigma_{5,1} \rightarrow \Sigma_{5,1}$ extends to the monodromy: $\Sigma_5 \rightarrow \Sigma_5$ of the fibration on $N(-37/18, -3/2, -8/3)$ with dilatation $\delta_5^+ = \lambda_{(7,4)}$. Since this extended monodromy is orientable, we have $\delta_5^+ \in \Lambda_5^+(-3/2)$.

By Remark 4.35 (2), we have:

LEMMA 4.36. $N(-3/2) \in \mathcal{U}^+$.

4.4. Fibers of genera 8 and 13.

By using the foregoing discussion one can prove the following which implies Proposition 1.8.

- PROPOSITION 4.37. (1) $N(-4/3, -25/17, -5)$ is a Σ_8 -bundle over the circle with dilatation $\lambda_{(18,17,7)} \approx 1.10403$ and with singularity data $(\underbrace{1, \dots, 1}_6, 15, \underbrace{1, \dots, 1}_7)$.
- (2) $N(-29/27, -5/3, -6)$ is a Σ_{13} -bundle over the circle with dilatation $\lambda_{(27,21,8)} \approx 1.07169$ and with singularity data $(25, \underbrace{1, \dots, 1}_7, \underbrace{2, \dots, 2}_8)$.

References

- [1] J. W. Aaber and N. M. Dunfield, Closed surface bundles of least volume, *Algebr. Geom. Topol.*, **10** (2010), 2315–2342.
- [2] I. Agol, The minimal volume orientable hyperbolic 2-cusped 3-manifolds, *Proc. Amer. Math. Soc.*, **138** (2010), 3723–3732.
- [3] I. Agol, Ideal triangulations of pseudo-Anosov mapping tori, *Contemp. Math.*, **560** (2011), 1–17.
- [4] J.-H. Cho and J.-Y. Ham, The minimal dilatation of a genus-two surface, *Experiment. Math.*, **17** (2008), 257–267.
- [5] B. Farb, Some problems on mapping class groups and moduli space, In: Problems on Mapping Class Groups and Related Topics, (ed. B. Farb), Proc. Sympos. Pure Math., **74**, Amer. Math. Soc., Providence, RI, 2006, pp. 11–55.
- [6] B. Farb, C. J. Leininger and D. Margalit, Small dilatation pseudo-Anosov homeomorphisms and 3-manifolds, *Adv. Math.*, **228** (2011), 1466–1502.
- [7] A. Fathi, F. Laudenbach and V. Poenaru, Travaux de Thurston sur les surfaces, *Asterisque*, **66–67**, Soc. Math. France, Paris, 1979.
- [8] D. Fried, Flow equivalence, hyperbolic systems and a new zeta function for flows, *Comment. Math. Helv.*, **57** (1982), 237–259.
- [9] E. Hironaka, Small dilatation mapping classes coming from the simplest hyperbolic braid, *Algebr. Geom. Topol.*, **10** (2010), 2041–2060.
- [10] E. Hironaka and E. Kin, A family of pseudo-Anosov braids with small dilatation, *Algebr. Geom. Topol.*, **6** (2006), 699–738.
- [11] N. V. Ivanov, Stretching factors of pseudo-Anosov homeomorphisms, *J. Soviet Math.*, **52** (1990), 2819–2822, which is translated from Zap. Nauch. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), **167** (1988), 111–116.
- [12] A. Kawachi, A Survey of Knot Theory, Birkhäuser Verlag, Basel, 1996.
- [13] E. Kin, S. Kojima and M. Takasawa, Entropy versus volume for pseudo-Anosovs, *Experiment. Math.*, **18** (2009), 397–407.
- [14] E. Kin, S. Kojima and M. Takasawa, Minimal dilatations of pseudo-Anosovs generated by the magic 3-manifold and their asymptotic behavior, to appear in *Algebraic & Geometric Topology*.
- [15] E. Kin and M. Takasawa, Pseudo-Anosov braids with small entropy and the magic 3-manifold, *Comm. Anal. Geom.*, **19** (2011), 705–758.
- [16] K. H. Ko, J. E. Los and W. T. Song, Entropies of braids, *J. Knot Theory Ramifications*, **11** (2002), 647–666.
- [17] E. Lanneau and J.-L. Thiffeault, On the minimum dilatation of pseudo-Anosov homeomorphisms on surfaces of small genus, *Ann. Inst. Fourier (Grenoble)*, **61** (2011), 105–144.
- [18] D. D. Long and U. Oertel, Hyperbolic surface bundles over the circle, In: Progress in Knot Theory and Related Topics, (Eds. M. Boileau, M. Domergue, Y. Mathieu and K. Millett), Travaux en Cours, **56**, Hermann, Paris, 1997, pp. 121–142.
- [19] A. Manning, Topological entropy and the first homology group, In: Dynamical Systems–Warwick 1974, (Eds. A. Dold and B. Eckmann), Lecture Notes in Math., **468**, Springer-Verlag, 1975, pp. 185–190.
- [20] B. Martelli and C. Petronio, Dehn filling of the “magic” 3-manifold, *Comm. Anal. Geom.*, **14** (2006), 969–1026.
- [21] S. Matsumoto, Topological entropy and Thurston’s norm of atoroidal surface bundles over the circle, *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, **34** (1987), 763–778.
- [22] H. Minakawa, Examples of pseudo-Anosov homeomorphisms with small dilatations, *J.*

- Math. Sci. Univ. Tokyo, **13** (2006), 95–111.
- [23] C. T. McMullen, Polynomial invariants for fibered 3-manifolds and Teichmüller geodesic for foliations, *Ann. Sci. École Norm. Sup. (4)*, **33** (2000), 519–560.
- [24] D. Rolfsen, *Knots and Links*, AMS Chelsea Publishing, 2003.
- [25] E. Rykken, Expanding factors for pseudo-Anosov homeomorphisms, *Michigan Math. J.*, **46** (1999), 281–296.
- [26] W. P. Thurston, A norm for the homology of 3-manifolds, *Mem. Amer. Math. Soc.*, **339** (1986), 99–130.
- [27] W. P. Thurston, Hyperbolic structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle, preprint, arXiv:math/9801045.
- [28] W. P. Thurston, *The Geometry and Topology of Three-Manifolds*, Lecture Notes, Princeton University, 1979.
- [29] R. W. Venzke, Braid forcing, hyperbolic geometry, and pseudo-Anosov sequences of low entropy, PhD thesis, California Institute of Technology, 2008, available at http://resolver.caltech.edu/Caltech_ETD:etd-05292008-085545.
- [30] P. Walters, *An Introduction to Ergodic Theory*, Grad. Texts in Math., **79**, Springer-Verlag, 1982.
- [31] A. Y. Zhironov, On the minimum dilation of pseudo-Anosov diffeomorphisms of a double torus, *Russian Math. Surveys*, **50** (1995), 223–224.

Eiko KIN

Department of Mathematics
Graduate School of Science
Osaka University
Toyonaka
Osaka 560-0043, Japan
E-mail: kin@math.sci.osaka-u.ac.jp

Mitsuhiko TAKASAWA

Department of Mathematical and Computing Sciences
Tokyo Institute of Technology
Ohokayama, Meguro
Tokyo 152-8552, Japan
E-mail: takasawa@is.titech.ac.jp