

## ON PSEUDO-PRIMALITY OF THE $n$ -TH POWER OF PRIME ENTIRE FUNCTIONS

BY GUODONG SONG AND JUE HUANG

### I. Introduction and main results.

Let  $g_0(z)$  be a transcendental entire function which is prime or pseudo-prime. We pose the following question: are the functions  $g_0(z)^n$  always pseudo-prime for  $n=2, 3, \dots$ ? The answer is affirmative if and only if  $n$  is an odd number. That is to say that  $g_0(z)^n$  is pseudo-prime if  $n$  is odd; while for even number  $n$ , there exists a prime entire function  $g_0(z)$  such that  $g_0(z)^n$  is not pseudo-prime. This assertion is contained in the following two theorems.

**THEOREM 1.** *Let  $g_0(z)$  be a pseudo-prime entire function, and  $n (\geq 3)$  be an odd number. Then  $F(z)=g_0(z)^n$  is also pseudo-prime.*

**THEOREM 2.** *The function*

$$F(z)=(\sin z)e^{\cos z} \tag{1}$$

*is prime.*

*Remark 1.* If  $F(z)$  is the function of the form (1), and  $n$  is an even number, then  $F(z)^n$  is not pseudo-prime, as is shown by the following factorization

$$F(z)^n=(\sin^n z)e^{n \cos z}=(1-w^2)^{n/2}e^{nw} \circ \cos z.$$

*Remark 2.* The function  $F(z)$  of the form (1) is also an example of prime periodic entire functions. In 1971, Gross [4] asked if there exist such functions. Later on, Ozawa [8, 9], Baker & Yang [2], Gross & Yang [6] constructed various examples of such kind of entire functions. Our example here is a much simpler one.

From Theorem 2 and Remark 1, it is easy to derive the following

**COROLLARY.** *For any polynomial  $P(z)$  of degree 2, there exists a prime entire function  $g^*(z)$  such that  $F(z)=P(g^*(z))$  is not pseudo-prime.*

The basic notions in the factorization theory of entire and meromorphic

functions, such as prime,  $E$ -prime, pseudo-prime, etc., shall not be stated here. One may find the definitions of these notions in the references.

**2. Preliminary lemmas.**

In proving our theorems we shall need several known results.

LEMMA 1 (Hayman [7]). *Let  $f(z)$  be an entire function. Then*

$$\sum_{a \neq \infty} \left(1 - \frac{1}{v(a)}\right) \leq 1,$$

where  $v(a)$  stands for the least order of almost all  $a$ -points of  $f(z)$ . Especially, there is at most one complex number  $a$  such that  $v(a) \geq 3$ .

LEMMA 2 (Edrei [3]). *Let  $g(z)$  be an entire function. If there exists an unbounded sequence  $\{a_n\}$  such that almost all the roots of  $g(z) = a_n$  ( $n=1, 2, \dots$ ) lie on one straight line, then  $g(z)$  is a polynomial of degree at most two.*

LEMMA 3 (Baker & Gross [1]). *Let  $h(z)$  be a periodic entire function of finite lower order, and  $c$  be a non-zero constant. Then*

$$H(z) = h(z) + cz$$

is prime.

LEMMA 4 (Gross [5]). *All meromorphic solutions of the functional equation*

$$f(z)^2 + g(z)^2 = 1 \tag{2}$$

are of the form

$$f(z) = \frac{2s(z)}{1+s(z)^2}, \quad g(z) = \frac{1-s(z)^2}{1+s(z)^2},$$

where  $s(z)$  is any meromorphic function. In particular, there is no non-constant  $f(z) = z^{-n}f_0(z)$  where  $n$  is a non-negative integer and  $f_0(z)$  is a polynomial satisfying equation (2).

**3. Proof of theorem 1.**

Let  $n = p_1 \cdot p_2 \cdots p_k$  with prime numbers  $p_j \geq 3$ ,  $j=1, \dots, k$ . If  $g_1(z) = g_0(z)^{p_1}$  is proved to be pseudo-prime, so is  $g_2(z) = g_1(z)^{p_2} = g_0(z)^{p_1 p_2}$ , and so on. Therefore, we may assume that  $n = p$  is a prime number. Also,  $g_0(z)$  may be assumed transcendental.

Suppose  $F(z) = g_0(z)^p = f(g(z))$  with transcendental entire functions  $f$  and  $g$ . By Lemma 1, it is easily seen that among zeros of  $f(w)$  there is at most one zero with order  $q$  such that  $(q, p) = 1$ . Hence 2 cases may occur.

(a)  $f(w)=(w-w_0)^q h(w)^p$  and  $g(z)=w_0+s(z)^p$  with transcendental entire functions  $h$  and  $s$ . Then

$$g_0(z)^p = s(z)^{qp} (h(w_0 + s(z)^p))^p$$

or

$$g_0(z) = u s(z)^q h(w_0 + s(z)^p) \quad (u^p = 1),$$

which gives a contradiction as  $g_0(z)$  is assumed to be pseudo-prime.

(b)  $f(w)=h(w)^p$  with a transcendental entire function  $h$ . Then  $g_0(z)=uh(g(z))$ . Again a contradiction.

Now let  $F=f(g)$  with  $f$  being meromorphic (not entire) and  $g$  entire, both transcendental. Then  $f$  must have exactly one pole,  $w_0$  say, which  $g$  doesn't take. And we may write

$$f(w)=(w-w_0)^{-k} f^*(w), \quad g(z)=w_0+e^{M(z)},$$

where  $k$  is a positive integer,  $f^*$  is transcendental entire with  $f^*(w_0) \neq 0$ , and  $M(z)$  is non-constant entire.

If  $f^*(w)$  has no zeros, or each zero of  $f^*$  is of order  $mp$  with a positive integer  $m$ , then  $f^*(w)=h(w)^p$  with a transcendental entire function  $h$ , which implies

$$g_0(z) = u e^{-kM(z)/p} h(w_0 + e^{M(z)}) \quad (u^p = 1)$$

or

$$g_0(z) = (u w^{-k} h(w_0 + w^p)) \circ e^{M(z)/p}.$$

But this violates the pseudo-primality of  $g_0(z)$ .

If  $f^*(w)$  has a zero,  $w_1$  say ( $w_1 \neq w_0$ ), of order  $q$  with  $(q, p)=1$  (By the same reasoning as in case(a),  $f^*$  has at most one such zero). Then  $g(z)$  must be of the form

$$g(z) = w_1 + s(z)^p$$

with an entire function  $s$ . But this is impossible, since the entire function  $g(z)$ , which has a Picard exceptional value  $w_0$ , can not have any completely ramified values.

The proof of theorem 1 is completed.

#### 4. Proof of theorem 2.

Let  $F(z)=f(g(z))$  with non-linear entire functions  $f$  and  $g$ . We discuss two cases.

Case (a).  $f$  has infinitely many zeros. Then by lemma 2,  $g(z)$  must be a polynomial of degree 2. Hence,  $F(z)$  may be expressed by

$$F(z) = f_1((z-c)^2)$$

with an entire function  $f_1$  and a constant  $c$ . This yields

$$\sin(z+c)e^{\cos(z+c)} = \sin(-z+c)e^{\cos(-z+c)}.$$

But the above equality can not hold, as is shown by substituting some special values of  $z$ .

Case (b).  $f$  has only finitely many zeros. Then we may write

$$f(w) = Q(w)e^{L(w)}$$

with a non-constant polynomial  $Q$  and entire function  $L$ . Thus

$$Q(g(z))e^{L(g(z))} = (\sin z)e^{\cos z}. \tag{3}$$

Since  $F(z) = (\sin z)e^{\cos z}$  is of infinite order and its convergent exponent of zeros, denoted by  $\rho^*(F)$ , is one, if  $L$  is a constant, then  $Q$  must have exactly one (simple) zero, *i. e.*  $Q$  is linear, which is out of our consideration. Therefore,  $L$  must be non-constant.

By considering the growth of functions in both sides of (3), we see that the order of  $g(z)$

$$\rho(g) \leq \rho(L(g)) = \rho(\cos z) = 1$$

and

$$\rho(g) = \rho(Q(g)) \geq \rho^*(Q(g)) = \rho^*(\sin z) = 1.$$

So that  $\rho(g) = 1$ .

Putting

$$s(z) = \frac{Q(g(z))}{\sin z} \exp(\cos z - L(g(z))),$$

we have  $\rho(s) \leq 1$ , which implies

$$\cos z - L(g(z)) = az + b, \quad a \text{ and } b \text{ are constants.}$$

If  $a \neq 0$ , then by lemma 3,  $L(g(z)) = \cos z - az - b$  is prime, so that  $L$  is linear. And we may write

$$g(z) = c_1 \cos z + c_2 z + b_1,$$

where  $c_1, c_2$  and  $b_1$  are constants with  $c_1 c_2 \neq 0$ . On the other hand

$$Q(g(z)) = (\sin z)e^{az+b}. \tag{4}$$

Therefore, substituting  $z = 2n\pi$  into both sides of (4), we see that the right side of (4) is 0, while the left side tends to  $\infty$ , which is a contradiction.

If  $a = 0$ , then  $L(g(z)) = \cos z - b$  and  $Q(g(z)) = b_2 \sin z$ , and we obtain an identity

$$Q^*(g(z))^2 + L^*(g(z))^2 \equiv 1 \quad \text{or} \quad Q^*(w)^2 + L^*(w)^2 \equiv 1$$

with a polynomial  $Q^*$  and an entire function  $L^*$  for every  $w \in \mathcal{C}$ , which violates lemma 4.

Up to now we have proved that  $F(z)$  is  $E$ -prime.

Now, let  $F=f(g)$  with meromorphic functions  $f$  and  $g$  ( $f$  is not entire), we discuss three cases.

Case (i).  $f$  is transcendental. Then  $g$  must be entire, and we have

$$f(w)=(w-w_0)^{-n}f_1(w), \quad g(z)=w_0+e^{M(z)}, \quad (5)$$

where  $f_1(w)$  is a transcendental entire function with  $f_1(w_0) \neq 0$ ,  $n$  is a positive integer, and  $M(z)$  is a non-constant entire function. We derive

$$\begin{aligned} F(z) &= (\sin z)e^{\cos z} = e^{-nM(z)}f_1(w_0+e^{M(z)}) \\ &= [e^{-nM(z)}f_1(w_0+e^{M(z)})] \circ M(z) \end{aligned}$$

Since  $F(z)$  is  $E$ -prime,  $M(z)$  must be linear, and we may write

$$F(z) = (\sin z)e^{\cos z} = e^{-naz}f^*(e^{az}), \quad (6)$$

where  $f^*$  is transcendental entire.

By the same argument as in case (a), we conclude that  $f^*$  has only finitely many zeros. Then we may write

$$f^*(w) = P(w)e^{N(w)}$$

with a non-constant polynomial  $P$  and entire function  $N$ . We obtain

$$(\sin z)e^{\cos z} = e^{-anz}P(e^{az})\exp(N(e^{az}))$$

Putting

$$T(z) = \frac{P(e^{az})}{\sin z} = \exp(\cos z + naz - N(e^{az})).$$

Obviously,  $\rho(T) \leq 1$ . Hence

$$\cos z + naz - N(e^{az}) = Az + B, \quad A \text{ and } B \text{ are constants.}$$

If  $A \neq na$ , then  $N$  is linear (by using lemma 3), and we would get

$$e^{az} = A_1 \cos z + A_2 z + B_1,$$

which is apparently impossible.

If  $A = na$ , then  $N(e^{az}) = \cos z - B$  and  $P(e^{az}) = B_2(\sin z)e^{naz}$ , and we would derive an identity

$$P^*(e^{az})^2 e^{-2n az} + N^*(e^{az})^2 \equiv 1 \quad \text{or} \quad \frac{P^*(w)^2}{w^{2n}} + N^*(w)^2 \equiv 1$$

with a polynomial  $P^*$  and  $a$  positive integer  $n$ . This again violates lemma 4.

Case (ii).  $f$  is rational and  $g$  entire. Then we obtain (5) and (6) with  $f_1$  (and  $f^*$ ) being a polynomial. And we may deduce that  $M(z)$  is linear. But in this case the function in the right side of (6) would be of finite order, which is also a contradiction.

Case (iii).  $f$  is rational and  $g$  meromorphic (not entire). Let  $x_0$  be a pole of  $f$ , then  $g(z)$  doesn't assume  $x_0$ , so that

$$g_1(z) = \frac{1}{g(z) - x_0}$$

is entire. Denoting

$$R(w) = f\left(\frac{1}{w} + x_0\right),$$

we get a factorization  $F = R \circ g_1$  which is equivalent to  $F = f \circ g$ . Then this case reduces to case (ii).

The proof is thus completed.

## 5. Final remark.

We propose the following questions:

(1) Does there exist an entire function  $g_0(z)$  which is prime and of *finite order* such that  $g_0(z)^2$  is not pseudo-prime?

(2) Let  $P(z)$  be a polynomial of degree  $\geq 3$  which has no right factor of the form  $(z-a)^2 + b$  with constants  $a, b$  and let  $g_0(z)$  be a pseudo-prime entire function. Can we conclude that the function  $F(z) = P(g_0(z))$  is also pseudo-prime?

## REFERENCES

- [ 1 ] BAKER, I.N. AND GROSS, F., Further results on factorization of entire functions, Proc. Symp. Pure Math. (Amer. Math. Soc., Providence, R.I.) **II** (1968), 30-35.
- [ 2 ] BAKER, I.N. AND YANG, C.C., An infinite-order periodic entire function which is prime, In: Proceedings of a Conference on Complex Analysis (Lexington 1976) pp. 7-10, Lecture Notes in Mathematics, 599, Berlin-Heidelberg-New York: Springer 1977.
- [ 3 ] EDREI, A., A meromorphic function with three radially distributed values, Trans. Amer. Math. Soc., **78** (1955), 276-293.
- [ 4 ] GROSS, F., Factorization of entire function which are periodic mod  $g$ , Indian J. Pure Appl. Math., **2** (1971), 561-571.
- [ 5 ] GROSS, F., On the equation  $f^n + g^n = 1$ , Bull. Amer. Math. Soc., **72** (1966), 86-88.
- [ 6 ] GROSS, F. AND YANG, C.C., On prime periodic entire functions, Math. Z., 174

(1980), 43-48.

- [ 7 ] HAYMAN, W. K., Meromorphic functions, Clarendon Press, Oxford, 1964.
- [ 8 ] OZAWA, M., Factorization of entire functions, Tohoku Math. J., 27 (1975), 321-336.
- [ 9 ] OZAWA, M., On the existence of prime periodic entire functions, Kodai Math. Sem. Rep., 29 (1978), 308-321.

DEPARTMENT OF MATHEMATICS  
EAST CHINA NORMAL UNIVERSITY  
SHANGHAI, CHINA

DEPARTMENT OF MATHEMATICS,  
HUAINAN MINING COLLEGE  
ANHUI, CHINA