

SUFFICIENT CONDITIONS FOR A REAL POLYNOMIAL TO BE A SUM OF SQUARES OF POLYNOMIALS

VAN DOAT DANG* AND THI THAO NGUYEN†

Abstract

In this paper, we establish new sufficient conditions for the polynomial f to be SOS in terms of the Newton polyhedron of f (Theorems 2.6 and 2.12). These new sufficient conditions include results which were proved earlier by Lasserre [13, Theorem 3], Fidalgo and Kovacec [6, Theorem 4.3], Ghasemi and Marshall [7, Theorems 2.1 and 2.3], and Ghasemi and Marshall [8, Theorem 2.3].

1. Introduction

The problem of representation of positive semidefinite polynomials (PSD) as sums of squares of polynomials (SOS) has been treated by many authors. Note that if f is SOS then the degree of f is even. In [10], Hilbert proved that $P_{2d,n} = \Sigma_{2d,n}$ if and only if $n \leq 2$ or $d = 1$ or $(n, d) = (3, 2)$, where $P_{2d,n}$ and $\Sigma_{2d,n}$ are the cone of PSD forms of degree $2d$ in n variables and the cone of SOS forms of degree $2d$ in n variables, respectively; see [2, Theorem 6.3.7]. The first explicit example for this phenomenon was given by Motzkin [18]; see [16, Proposition 1.2.2].

We denote by \mathbf{N} the set of nonnegative integers $\{0, 1, 2, \dots\}$. For $x = (x_1, \dots, x_n)$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbf{N}^n$, define $x^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ and $|\alpha| = \alpha_1 + \cdots + \alpha_n$. Using these notations, every polynomial $f \in \mathbf{R}[x]$ can be written as $f(x) = \sum_{\alpha \in \mathbf{N}^n} f_\alpha x^\alpha$, where $f_\alpha \in \mathbf{R}$ and $f_\alpha = 0$, except for finitely many α . Assume now that f is nonconstant and has even degree $2d$. Let $\Omega := \{\alpha \in \mathbf{N}^n \mid f_\alpha \neq 0\} \setminus$

2010 *Mathematics Subject Classification.* Primary 12D15; Secondary 14P99, 90C25.

Key words and phrases. Positive polynomial, sum of squares, Newton polyhedron, framework, mediated set, barycentric coordinates, optimization, geometric programming.

*This author was partially supported by Vietnam National Foundation for Science and Technology Development (NAFOSTED) grant number 101.04-2013.07.

†This author was partially supported by Vietnam National Foundation for Science and Technology Development (NAFOSTED) grant number 101.04-2014.23.

Received January 23, 2015; revised April 30, 2015.

$\{0, 2de_1, \dots, 2de_n\}$, where $e_1 = (1, 0, \dots, 0), \dots, e_n = (0, \dots, 0, 1)$. Thus f has the form

$$f(x) = f_0 + \sum_{\alpha \in \Omega} f_\alpha x^\alpha + \sum_{i=1}^n f_{2de_i} x_i^{2d}.$$

Setting

$$\begin{aligned} \Delta &:= \{\alpha \in \Omega \mid f_\alpha x^\alpha \text{ is not a sum of squares in } \mathbf{R}[x]\} \\ &= \{\alpha \in \Omega \mid f_\alpha < 0 \text{ or } \alpha_i \text{ is odd for some } i \in \{1, \dots, n\}\}. \end{aligned}$$

In [13, Theorem 3], Lasserre proved that if

$$f_0 \geq \sum_{\alpha \in \Delta} |f_\alpha| \quad \text{and} \quad f_{2de_i} \geq \sum_{\alpha \in \Delta} |f_\alpha| \frac{|\alpha|}{2d}, \quad i = 1, \dots, n,$$

then f is SOS.

Later on, some various versions on sufficient conditions of this type have been given by Fidalgo and Kovacec [6, Theorem 4.3], and Ghasemi and Marshall [7, Theorems 2.1 and 2.3] where the left-hand sides of the inequalities are kept unchanged and the right-hand sides are replaced by quantities which also depend on the coefficients f_α , $\alpha \in \Delta$. Indeed, in [7, Theorems 2.1], f is SOS if

$$f_0 \geq \sum_{\alpha \in \Delta} |f_\alpha| \frac{2d - |\alpha|}{2d} \quad \text{and} \quad f_{2de_i} \geq \sum_{\alpha \in \Delta} |f_\alpha| \frac{\alpha_i}{2d}, \quad i = 1, \dots, n;$$

and this result is an new version of Lasserre’s one.

Recently, Ghasemi and Marshall [8, Theorem 2.3] have given another sufficient condition on additional variables satisfying constrains depending on the coefficients f_α , $\alpha \in \Delta$, for the polynomial f to be SOS. Precisely, a sufficient condition for a form f of degree $2d$ to be SOS is that there exist nonnegative real numbers $a_{\alpha,i}$ for $\alpha \in \Delta$, $i = 1, \dots, n$, such that

$$(2d)^{2d} a_\alpha^\alpha = |f_\alpha|^{2d} \alpha^\alpha, \quad \alpha \in \Delta, \quad \text{and} \quad f_{2de_i} \geq \sum_{\alpha \in \Delta} a_{\alpha,i}, \quad i = 1, \dots, n,$$

where $a_\alpha^\alpha = a_{\alpha,1}^{\alpha_1} \cdots a_{\alpha,n}^{\alpha_n}$ and $\alpha^\alpha = \alpha_1^{\alpha_1} \cdots \alpha_n^{\alpha_n}$ with the convention $0^0 = 1$. This result include improvements of the above results. However, if $f_0 = f_{2de_1} = \cdots = f_{2de_n} = 0$, the conditions of [8, Theorem 2.3] imply that $\Delta = \emptyset$ and so f is obviously SOS. Thus, this criterion is overly restrictive.

We recall here some notations about Newton polyhedra. Let $\text{supp}(f) := \{\alpha \in \mathbf{N}^n \mid f_\alpha \neq 0\}$. The *Newton polyhedron* $\Gamma(f)$ of f is the convex hull of $\text{supp}(f)$ in \mathbf{R}^n . Clearly, $\Gamma(f)$ is a compact convex polyhedron of dimension at most n . A *supporting hyperplane* of $\Gamma(f)$ is a hyperplane minimizing the value of some linear function on $\Gamma(f)$. The *faces* of the boundary of the Newton polyhedron $\Gamma(f)$ are the intersection of $\Gamma(f)$ with its supporting hyperplanes.

They are compact convex polyhedra of dimension at most $n - 1$. The *vertices* are faces of dimension 0 (i.e., points).

In this paper, we establish new sufficient conditions for the polynomial f to be SOS in terms of the Newton polyhedron $\Gamma(f)$ of f (Theorems 2.6 and 2.12). Precisely, we replace the values $f_0, f_{2de_1}, \dots, f_{2de_n}$ in the left sides of the inequalities above by some values $f_u, u \in \mathcal{U}$, where \mathcal{U} is some set satisfying $V(f) \subset \mathcal{U} \subset \Gamma(f) \cap (2\mathbf{Z})^n$. Note that our new sufficient conditions include all the results mentioned above.

The paper is organized as follows. Section 2 presents the new sufficient conditions for a polynomial to be a sum of squares of polynomials (Theorem 2.6 and 2.12). In Section 3, we prove these results. As a consequence of Theorem 2.12, a lower bound of f is established in Section 4.

2. Statement of the results

Suppose $f(x_1, \dots, x_n)$ is a polynomial of degree $2d$. Let

$$\bar{f}(x_0, x_1, \dots, x_n) := x_0^{2d} f\left(\frac{x_1}{x_0}, \dots, \frac{x_n}{x_0}\right).$$

Then \bar{f} is a form of degree $2d$ called the *homogenization* of f . We have the following well-known result.

PROPOSITION 2.1 ([16, Proposition 1.2.4]). *Let f be a polynomial of degree $2d$. Then f is PSD iff \bar{f} is PSD; and f is SOS iff \bar{f} is SOS.*

We now assume f is a form, and $\Gamma(f)$ is the Newton polyhedron of f . Let

- $\mathcal{C} := \Gamma(f) \cap \mathbf{Z}^n$ denote the set of all integer points in $\Gamma(f)$,
- $\mathcal{A} := \{\frac{1}{2}(s+t) \mid s, t \in \Gamma(f) \cap (2\mathbf{Z})^n\}$ denote the set of integer points from $\Gamma(f)$ which are averages of even integer points from $\Gamma(f)$, and
- $\mathcal{V} := \mathcal{A} \setminus \{\frac{1}{2}(s+t) \mid s \neq t, s, t \in \Gamma(f) \cap (2\mathbf{Z})^n\}$ denote the set of even integer points from $\Gamma(f)$ which are not averages of distinct even integer points from $\Gamma(f)$.

From now on, $V := V(f)$ denotes the set of vertices of the Newton polyhedron of f . It follows from [9, Theorem 3.1] that if f is PSD, the set V is contained in $(2\mathbf{Z})^n$, and so $V \subset \mathcal{V} \subset \mathcal{C} \cap (2\mathbf{Z})^n$. We recall some results, which give some necessary conditions for a polynomial to be SOS.

PROPOSITION 2.2 ([4, Proposition 3.7]). *If f is SOS, then*

- (i) *for every $\alpha \in \text{supp}(f)$, we have $\alpha \in \mathcal{A}$, and*
- (ii) *for every $u \in \mathcal{V}$, $f_u \geq 0$.*

We recall here some notations, due to Reznick [23]. We use the term *framework* to denote a set $\mathcal{U} = \{u^1, \dots, u^m\}$ in which $u^i = (u^i_1, \dots, u^i_n) \in (2\mathbf{Z})^n$ with $u^i_j \geq 0$ and $\sum_{j=1}^n u^i_j = 2d$ for all i and some d . A *trellis* is a framework in

which u^1, \dots, u^m comprise the vertices of a simplex. Note that if \mathcal{U} is a trellis, then \mathcal{U} is a linearly independent set.

DEFINITION 2.3 ([23, Definition 2.1]). Let \mathcal{U} be a framework. A finite set $\mathcal{L} \subset \mathbf{Z}^n$ is called \mathcal{U} -mediated if \mathcal{L} contains \mathcal{U} , and every $v \in \mathcal{L} \setminus \mathcal{U}$ is an average of two distinct even points in \mathcal{L} .

We now assume that $V \subset \mathcal{U} \subset \mathcal{V}$. Then, \mathcal{U} is a framework. By [23, Theorem 2.2], we get the following.

THEOREM 2.4 ([23]). There is a \mathcal{U} -mediated set \mathcal{U}^* satisfying $\mathcal{A}(\mathcal{U}) := \{\frac{1}{2}(s+t) \mid s, t \in \mathcal{U}\} \subset \mathcal{U}^* \subset \mathcal{C}$ and \mathcal{U}^* contains every \mathcal{U} -mediated set.

Suppose that $V = \{u^1, \dots, u^m\}$ is a framework. If $\alpha \in \mathbf{Z}^n$ and $\alpha = \lambda_1 u^1 + \dots + \lambda_m u^m$ with $\lambda_i \geq 0$ and $\lambda_1 + \dots + \lambda_m = 1$, then $\alpha \in \mathcal{C}$; conversely, if $\alpha \in \mathcal{C}$ then at least one such $\lambda = (\lambda_1, \dots, \lambda_m)$ exists. Let

$$\Lambda(\alpha) = \{\lambda = (\lambda_1, \dots, \lambda_m) \mid \lambda_i \geq 0, \lambda_1 + \dots + \lambda_m = 1, \text{ and } \alpha = \lambda_1 u^1 + \dots + \lambda_m u^m\}.$$

If V is a trellis, then only one such $\lambda = (\lambda_1, \dots, \lambda_m)$ exists, and $(\lambda_1, \dots, \lambda_m)$ are called the *barycentric coordinates* of α with respect to V .

THEOREM 2.5 ([23, Theorem 3.3]). Suppose $V = \{u^1, \dots, u^m\}$ is a framework, $\alpha \in \mathcal{C}$ and $\lambda \in \Lambda(\alpha)$. Then, if the form $f(V, \lambda, \alpha)(x) := \lambda_1 x^{u^1} + \dots + \lambda_m x^{u^m} - x^\alpha$ is SOS then $\alpha \in V^*$.

The following is our new result, which gives a sufficient condition on the coefficients for a polynomial to be sum of squares.

THEOREM 2.6. Let f be a form in n real variables with $V \subset (2\mathbf{Z})^n$. Suppose that $V \subset \mathcal{U} \subset \mathcal{V}$ such that

- (i) $\alpha \in \mathcal{U}^*$ for all $\alpha \in \Delta$,
- (ii) $\min_{u \in \mathcal{U}} f_u \geq \sum_{\alpha \in \Delta} |f_\alpha|$.

Then f is SOS. Here, by convention, we set $\sum_{\alpha \in \Delta} |f_\alpha| := 0$ if $\Delta = \emptyset$.

COROLLARY 2.7. Let f be a form in n real variables with $V \subset (2\mathbf{Z})^n$. Suppose

- (i) $\alpha \in \mathcal{A}$ for all $\alpha \in \Delta$,
- (ii) $\min_{u \in \mathcal{V}} f_u \geq \sum_{\alpha \in \Delta} |f_\alpha|$.

Then f is SOS.

COROLLARY 2.8. Let f be a form in 3 real variables with $V \subset (2\mathbf{Z})^3$. Then, f is SOS if

$$\min_{u \in \mathcal{V}} f_u \geq \sum_{\alpha \in \Delta} |f_\alpha|.$$

Remark 2.9. Condition (ii) of Theorem 2.6 together with $f_u = 0$ for some $u \in \mathcal{U}$ implies that $\Delta = \emptyset$ and $f_u \geq 0$ for all $u \in \mathcal{U}$; in this case, f is obviously SOS. Thus, an alternative sufficient condition is given by Theorem 2.6 for $V \subset \mathcal{U} \subset \text{supp}(f) \cap \mathcal{V}$.

Next, we shall give another sufficient condition for a polynomial to be SOS. We recall the following result, due to Reznick.

THEOREM 2.10 ([23, Corollary 4.9]). *Suppose $V = \{u^1, \dots, u^m\}$ is a trellis, and $\alpha \in \mathcal{C}$. Then the form $f(V, \alpha)(x) := \lambda_1 x^{u^1} + \dots + \lambda_m x^{u^m} - x^\alpha$ is SOS if and only if $\alpha \in V^*$. Here $(\lambda_1, \dots, \lambda_m)$ are the barycentric coordinates of α with respect to V .*

We will make use Theorem 2.10 to prove the following.

COROLLARY 2.11. *Suppose that $V = \{u^1, \dots, u^m\}$ is a trellis and $\alpha \in V^*$. Then, for a form $E(x) = b_1 x^{u^1} + \dots + b_m x^{u^m} - cx^\alpha$ such that $b_i \geq 0$ for $i = 1, \dots, m$, and $c \geq 0$ if all α_i are even, the following are equivalent:*

- (i) $E \geq 0$.
- (ii) $|c| \lambda_1^{\lambda_1} \dots \lambda_m^{\lambda_m} \leq b_1^{\lambda_1} \dots b_m^{\lambda_m}$ (with the convention $0^0 = 1$).
- (iii) E is SOS.

Here $(\lambda_1, \dots, \lambda_m)$ are the barycentric coordinates of α with respect to V .

Now, we present our other result, which is an improved version of [8, Theorem 2.3]. The proof of this result is by using Corollary 2.11.

THEOREM 2.12. *Let f be a form, and $V = \{u^1, \dots, u^m\}$ be the set of vertices of $\Gamma(f)$. Suppose that V is a trellis, and $\alpha \in V^*$ for all $\alpha \in \Delta$. A sufficient condition for f to be SOS is that there exist nonnegative real numbers $a_{\alpha,i}$ for $\alpha \in \Delta$, $i = 1, \dots, m$, such that*

- (i) $a_\alpha^{\lambda_\alpha} = |f_\alpha| \lambda_\alpha^{\lambda_\alpha}$, $\alpha \in \Delta$, and
- (ii) $f_{u^i} \geq \sum_{\alpha \in \Delta} a_{\alpha,i}$, $i = 1, \dots, m$.

Here $a_\alpha = (a_{\alpha,1}, \dots, a_{\alpha,m})$, $\lambda_\alpha = (\lambda_{\alpha,1}, \dots, \lambda_{\alpha,m})$ are the barycentric coordinates of α with respect to V , $a_\alpha^{\lambda_\alpha} = a_{\alpha,1}^{\lambda_{\alpha,1}} \dots a_{\alpha,m}^{\lambda_{\alpha,m}}$, and $\lambda_\alpha^{\lambda_\alpha} = \lambda_{\alpha,1}^{\lambda_{\alpha,1}} \dots \lambda_{\alpha,m}^{\lambda_{\alpha,m}}$.

We mention some corollaries of Theorem 2.12. Corollary 2.13 generalizes known earlier results, due to Lasserre [13, Theorem 3] and Ghasemi and Marshall [7, Theorem 2.1]. Corollary 2.14 includes results which were proved earlier by Fidalgo and Kovacec [6, Theorem 4.3] and Ghasemi and Marshall [7, Theorem 2.3]. Corollary 2.15 is a generalization of the result of Ghasemi and Marshall [8, Corollary 2.7]. Each corollary will be proved by applying Theorem 2.12 for a particular choice of the lifted variables $a_{\alpha,i}$.

COROLLARY 2.13. *Suppose f is a form such that $V = \{u^1, \dots, u^m\}$ is a trellis, and $\alpha \in V^*$ for all $\alpha \in \Delta$. Then if*

$$f_{u^i} \geq \sum_{\alpha \in \Delta} |f_\alpha| \lambda_{\alpha, i}, \quad i = 1, \dots, m,$$

then f is SOS.

COROLLARY 2.14. *Suppose f is a form such that $V = \{u^1, \dots, u^m\}$ is a trellis, and $\alpha \in V^*$ for all $\alpha \in \Delta$. Then f is SOS if*

$$f_{u^i} \geq \sum_{\alpha \in \Delta} |f_\alpha| \lambda_\alpha^{\lambda_{\alpha, i}}, \quad i = 1, \dots, m.$$

COROLLARY 2.15. *Suppose f is a form such that $V = \{u^1, \dots, u^m\}$ is a trellis, and $\alpha \in V^*$ for all $\alpha \in \Delta$. Then f is SOS if*

$$\sum_{\alpha \in \Delta} \frac{|f_\alpha| \lambda_\alpha^{\lambda_{\alpha, i}}}{\prod_{i=1}^m f_{u^i}^{\lambda_{\alpha, i}}} \leq 1.$$

Notice that, by a similar argument as in [8, Remark 2.8], Corollary 2.15 is an improved version of Corollary 2.14.

COROLLARY 2.16. *Suppose f is a form such that $V = \{u^1, \dots, u^m\}$ is a trellis, and $\alpha \in V^*$ for all $\alpha \in \Delta$. Then f is SOS if*

$$f_{u^i} \geq \sum_{\alpha \in \Delta, \lambda_{\alpha, i} \neq 0} k \lambda_{\alpha, i} \left(\frac{|f_\alpha|}{k} \right)^{1/\lambda_{\alpha, i} n_\alpha}, \quad i = 1, \dots, m,$$

where $n_\alpha := |\{i \mid \lambda_{\alpha, i} \neq 0\}|$, and k is a positive real number.

3. Proof of the results

Let $\mathbf{R}[x]$ be the ring of real polynomials in the variables $x = (x_1, \dots, x_n)$, and $\mathbf{R}[x]_{2d}$ be the vector space of real polynomials of degree at most $2d$, with canonical basis $(x^\alpha) = \{x^\alpha \mid \alpha \in \mathbf{N}^n, |\alpha| \leq 2d\}$. Given a sequence $y = (y_\alpha) \subset \mathbf{R}$ indexed by the canonical basis (x^α) , let $L_y : \mathbf{R}[x]_{2d} \rightarrow \mathbf{R}$ be the linear mapping

$$f = \sum_{\alpha} f_\alpha x^\alpha \mapsto L_y(f) = \sum_{\alpha} f_\alpha y_\alpha, \quad f \in \mathbf{R}[x]_{2d},$$

and let $M_d(y)$ be the *moment* matrix with rows and columns indexed by (x^α) , and defined by

$$M_d(y)(\alpha, \beta) := L_y(x^{\alpha+\beta}) = y_{\alpha+\beta}, \quad \alpha, \beta \in \mathbf{N}^n : |\alpha|, |\beta| \leq d.$$

Let the notation $M_d(y) \succeq 0$ stand for $M_d(y)$ is positive semidefinite.

Remark 3.1. It is clear that $M_d(y) \succeq 0$ iff $L_y(f^2) \geq 0$ for all $f \in \mathbf{R}[x]_d$. Moreover, f is SOS iff $L_y(f) \geq 0$ for all y such that $M_d(y) \succeq 0$ (this is a corollary of [14, Theorem 3.1]).

LEMMA 3.2. *Let \mathcal{U} be a framework and \mathcal{L} be a \mathcal{U} -mediated set. Suppose $y = (y_\alpha) \subset \mathbf{R}$ such that $M_d(y) \succeq 0$. Then*

$$|L_y(x^\alpha)| \leq \max_{u \in \mathcal{U}} L_y(x^u) \quad \text{for all } \alpha \in \mathcal{L}.$$

Proof. First of all, we will show that if $\alpha \in \mathcal{L} \setminus \mathcal{U}$, there exists $k \geq 1$ and a sequence

$$\alpha_{i-1} = \frac{1}{2}(\alpha_i + \beta_i), \quad \alpha_i \neq \beta_i, \quad \alpha_i, \beta_i \in \mathcal{L} \cap (2\mathbf{Z})^n, \quad i = 1, \dots, k,$$

such that $\alpha_0 = \alpha$ and $\alpha_k \in \mathcal{U}$. In fact, let

$$X = \left\{ \alpha' \mid \text{there exists } k \geq 1 \text{ and a sequence } \alpha_{i-1} = \frac{1}{2}(\alpha_i + \beta_i), \alpha_i \neq \beta_i, \right. \\ \left. \alpha_i, \beta_i \in \mathcal{L} \cap (2\mathbf{Z})^n, i = 1, \dots, k, \text{ such that } \alpha_0 = \alpha \text{ and } \alpha_k = \alpha' \right\}.$$

Since X is contained in \mathcal{L} , the set X is finite, and so the convex hull of X has vertices which belong to \mathcal{U} .

Now, setting $\tau := \max_{\alpha \in \mathcal{L}} |L_y(x^\alpha)|$. Then there exists some $\gamma \in \mathcal{L}$ such that $\tau = |L_y(x^\gamma)|$. If $\gamma \in \mathcal{U}$, then

$$|L_y(x^\alpha)| \leq |L_y(x^\gamma)| = \max_{u \in \mathcal{U}} |L_y(x^u)| \quad \text{for all } \alpha \in \mathcal{L}.$$

Otherwise, by the above claim, there exists some $k \geq 1$ such that

$$\begin{aligned} \gamma &= \frac{1}{2}(\gamma_1 + \beta_1), \quad \gamma_1 \neq \beta_1 \in \mathcal{L} \cap (2\mathbf{Z})^n, \\ \gamma_1 &= \frac{1}{2}(\gamma_2 + \beta_2), \quad \gamma_2 \neq \beta_2 \in \mathcal{L} \cap (2\mathbf{Z})^n, \\ &\dots\dots\dots \\ \gamma_{k-1} &= \frac{1}{2}(\gamma_k + \beta_k), \quad \gamma_k \neq \beta_k \in \mathcal{L} \cap (2\mathbf{Z})^n, \end{aligned}$$

where $\gamma_k \in \mathcal{U}$. Since $M_d(y) \succeq 0$, we have

$$\tau = |L_y(x^\gamma)| = |L_y(x^{(1/2)(\gamma_1 + \beta_1)})| \leq \sqrt{L_y(x^{\gamma_1})L_y(x^{\beta_1})} \leq \sqrt{L_y(x^{\gamma_1})\tau}.$$

Hence $\tau \leq L_y(x^{\gamma_1})$. By repeating the above arguments, after finitely many steps, we deduce

$$\tau \leq L_y(x^{\gamma_k}) \leq \max_{u \in \mathcal{U}} L_y(x^u).$$

This complete the proof of the lemma. □

Proof of Theorem 2.6. By Remark 3.1, it is sufficient to show that $L_y(f) \geq 0$ for any y such that $M_d(y) \geq 0$.

So let y be such that $M_d(y) \geq 0$. Let $\tau := \max\{L_y(x^u) \mid u \in \mathcal{U}\}$. Then, by Lemma 3.2, we have

$$|L_y(x^\alpha)| \leq \tau \quad \text{for all } \alpha \in \mathcal{U}^*.$$

This together with Conditions (i)–(ii) implies that

$$\begin{aligned} L_y(f) &= \sum_{u \in \mathcal{U}} f_u L_y(x^u) + \sum_{\alpha \in \Delta} f_\alpha L_y(x^\alpha) + \sum_{\alpha \notin (\mathcal{U} \cup \Delta)} f_\alpha L_y(x^\alpha) \\ &\geq \left(\min_{u \in \mathcal{U}} f_u - \sum_{\alpha \in \Delta} |f_\alpha| \right) \tau \geq 0. \end{aligned} \quad \square$$

Proof of Corollary 2.7. We first prove the following.

CLAIM 3.3. *We have $\mathcal{V}^* = \mathcal{A}$.*

Proof. We first prove that $\mathcal{A} \subset \mathcal{V}^*$. Indeed, for every $\alpha \in \mathcal{A} \setminus \mathcal{V}$, we have $\alpha = \frac{1}{2}(s+t)$ for some $s \neq t, s, t \in \Gamma(f) \cap (2\mathbf{Z})^n \subset \mathcal{A} \cap (2\mathbf{Z})^n$, and hence α is an average of two distinct even points in \mathcal{A} . Thus \mathcal{A} is a \mathcal{V} -mediated set, and so $\mathcal{A} \subset \mathcal{V}^*$.

On the other hand, let $\mathcal{C}' := \mathcal{C} \setminus \mathcal{A}$, we will show that $\mathcal{C}' \cap \mathcal{V}^* = \emptyset$. Indeed, by contradiction, assume that there exists some $\alpha \in \mathcal{C}' \cap \mathcal{V}^*$. Then, we have

- $\alpha \in \mathcal{C}'$, and so $\alpha \notin \mathcal{A}$;
- $\alpha \in \mathcal{V}^*$; and hence $\alpha = \frac{1}{2}(s+t)$ for some $s, t \in \mathcal{V}^* \cap (2\mathbf{Z})^n \subset \mathcal{C} \cap (2\mathbf{Z})^n$, by Theorem 2.4; and so $\alpha \in \mathcal{A}$.

The contradiction shows that $\mathcal{C}' \cap \mathcal{V}^* = \emptyset$, and hence $\mathcal{V}^* \subset \mathcal{A}$.

Therefore, we have $\mathcal{V}^* = \mathcal{A}$. □

Now, the corollary follows immediately from Theorem 2.6 and Claim 3.3. □

Proof of Corollary 2.8. Note that \mathcal{V} is also framework in \mathbf{Z}^3 . Then it is shown in [22] that $\mathcal{A} = \mathcal{C}$. This, together with Claim 3.3, shows that $\mathcal{V}^* = \mathcal{C}$. Thus, Condition (i) of Theorem 2.6 is obviously true, and f is SOS. □

Proof of Corollary 2.11. (iii) \Rightarrow (i) is trivial, so it suffices to show that (i) \Rightarrow (ii) and (ii) \Rightarrow (iii).

First of all, notice that:

- If $c = 0$, the result is trivially true.
- If $c < 0$, by the assumption, there exists some $\alpha_i \notin 2\mathbf{Z}$. Then making a change of the variables $x_i = -y_i$, and $x_j = y_j$ for $j \neq i$, c is replaced by $-c$. In this way, we can assume that $c > 0$.

• If $\lambda_m = 0$, let

$$\tilde{E}(x) = b_1x^{u^1} + \dots + b_{m-1}x^{u^{m-1}} - cx^\alpha.$$

CLAIM 3.4. *The following are equivalent.*

- (a) $E \geq 0$.
- (b) $\tilde{E} \geq 0$.

Proof. (a) \Rightarrow (b). Suppose $E(x) \geq 0$ for all $x \in \mathbf{R}^n$. Since $\mathcal{U} = \{u^1, \dots, u^m\}$ is a trellis, $\{u^1, \dots, u^m\}$ is linearly independent. Without the loss of generality, we can assume that

$$\det \begin{bmatrix} u_1^1 & \dots & u_m^1 \\ \dots & \dots & \dots \\ u_1^m & \dots & u_m^m \end{bmatrix} \neq 0,$$

where $u^i = (u_1^i, \dots, u_n^i)$, $i = 1, \dots, m$. We will show that there exist $c_1, \dots, c_m \in \mathbf{R}_{>0}$ such that

$$\begin{cases} c_1^{u_1^1} \dots c_m^{u_m^1} = 1 \\ \dots \dots \dots \\ c_1^{u_1^{m-1}} \dots c_m^{u_m^{m-1}} = 1 \\ c_1^{u_1^m} \dots c_m^{u_m^m} = \varepsilon \end{cases}$$

for every $\varepsilon \in \mathbf{R}_{>0}$. In fact, it is clear that the system of linear equations

$$\begin{cases} u_1^1\theta_1 + \dots + u_m^1\theta_m = 0 \\ \dots \dots \dots \\ u_1^{m-1}\theta_1 + \dots + u_m^{m-1}\theta_m = 0 \\ u_1^m\theta_1 + \dots + u_m^m\theta_m = \ln \varepsilon \end{cases}$$

has one only root $(\theta_1, \dots, \theta_m) \in \mathbf{R}^m$. Let $c_i := e^{\theta_i}$, $i = 1, \dots, m$, so the claim is proved. Make a change of variables

$$\begin{cases} x_i = c_i y_i, & i = 1, \dots, m \\ x_j = y_j, & j = m + 1, \dots, n. \end{cases}$$

Then

$$\begin{aligned} E_\varepsilon(y) &:= E(x) = b_1x^{u^1} + \dots + b_mx^{u^m} - c(x^{u^1})^{\lambda_1} \dots (x^{u^{m-1}})^{\lambda_{m-1}} \\ &= b_1(c_1^{u_1^1} \dots c_m^{u_m^1})y^{u^1} + \dots + b_m(c_1^{u_1^m} \dots c_m^{u_m^m})y^{u^m} \\ &\quad - c[(c_1^{u_1^1} \dots c_m^{u_m^1})y^{u^1}]^{\lambda_1} \dots [(c_1^{u_1^{m-1}} \dots c_m^{u_m^{m-1}})y^{u^{m-1}}]^{\lambda_{m-1}} \\ &= b_1y^{u^1} + \dots + b_{m-1}y^{u^{m-1}} + \varepsilon b_my^{u^m} - cy^\alpha. \end{aligned}$$

So $E(x) < 0$ for k sufficiently large. This implies that (i) also fails. Thus the claimed implication are trivially true in this case.

Now, we can assume that $b_1, \dots, b_m, c, \lambda_1, \dots, \lambda_m \in \mathbf{R}_{>0}$. By the same arguments as above, there exist positive real numbers c_1, \dots, c_m such that

$$\begin{cases} c_1^{u_1} \cdots c_m^{u_m} = \frac{\lambda_1}{b_1} \\ \dots\dots\dots \\ c_1^{u_m} \cdots c_m^{u_m} = \frac{\lambda_m}{b_m}. \end{cases}$$

(i) \Rightarrow (ii). Assume $E(x) \geq 0$ for all $x \in \mathbf{R}^n$. Taking

$$x := (c_1, \dots, c_m, 1, \dots, 1).$$

Then, we have

$$\begin{aligned} E(x) &= b_1 x^{u_1} + \dots + b_m x^{u_m} - c(x^{u_1})^{\lambda_1} \dots (x^{u_m})^{\lambda_m} \\ &= b_1(c_1^{u_1} \cdots c_m^{u_1}) + \dots + b_m(c_1^{u_m} \cdots c_m^{u_m}) - c[c_1^{u_1} \cdots c_m^{u_1}]^{\lambda_1} \dots [c_1^{u_m} \cdots c_m^{u_m}]^{\lambda_m} \\ &= \lambda_1 + \dots + \lambda_m - c\left(\frac{\lambda_1}{b_1}\right)^{\lambda_1} \cdots \left(\frac{\lambda_m}{b_m}\right)^{\lambda_m} \\ &= 1 - c\left(\frac{\lambda_1}{b_1}\right)^{\lambda_1} \cdots \left(\frac{\lambda_m}{b_m}\right)^{\lambda_m} \\ &= 1 - c \frac{\lambda^\lambda}{b^\lambda} \geq 0. \end{aligned}$$

This proves (ii).

(ii) \Rightarrow (iii). Make a change of variables

$$\begin{cases} x_i = c_i y_i, & i = 1, \dots, m \\ x_j = y_j, & j = m + 1, \dots, n. \end{cases}$$

Let $\tilde{c} := c\left(\frac{\lambda_1}{b_1}\right)^{\lambda_1} \cdots \left(\frac{\lambda_m}{b_m}\right)^{\lambda_m}$. By (ii), we have $\tilde{c} \leq 1$. Then

$$\begin{aligned} E(x) &= b_1(c_1^{u_1} \cdots c_m^{u_1})y^{u_1} + \dots + b_m(c_1^{u_m} \cdots c_m^{u_m})y^{u_m} \\ &\quad - c[(c_1^{u_1} \cdots c_m^{u_1})y^{u_1}]^{\lambda_1} \cdots [(c_1^{u_m} \cdots c_m^{u_m})y^{u_m}]^{\lambda_m} \\ &= \lambda_1 y^{u_1} + \dots + \lambda_m y^{u_m} - c\left(\frac{\lambda_1}{b_1}\right)^{\lambda_1} \cdots \left(\frac{\lambda_m}{b_m}\right)^{\lambda_m} y^\alpha \\ &= \lambda_1 y^{u_1} + \dots + \lambda_m y^{u_m} - \tilde{c}y^\alpha \\ &= \tilde{c}(\lambda_1 y^{u_1} + \dots + \lambda_m y^{u_m} - y^\alpha) + (1 - \tilde{c})(\lambda_1 y^{u_1} + \dots + \lambda_m y^{u_m}), \end{aligned}$$

which is SOS, by Theorem 2.10. This proves (iii). □

Proof of Theorem 2.12. Suppose that such real numbers exist. Condition (i) and Corollary 2.11 imply that $\sum_{i=1}^m a_{\alpha,i}x^{u^i} + f_{\alpha}x^{\alpha}$ is SOS for every $\alpha \in \Delta$. So

$$\sum_{\alpha \in \Delta} \left(\sum_{i=1}^m a_{\alpha,i}x^{u^i} + f_{\alpha}x^{\alpha} \right)$$

is SOS. By Condition (ii), we have

$$\sum_{i=1}^m \left(f_{u^i} - \sum_{\alpha \in \Delta} a_{\alpha,i} \right) x^{u^i}$$

is SOS. Now, it is clear that $f_{\alpha}x^{\alpha}$ is SOS for every $\alpha \notin \Delta$. Therefore,

$$f(x) = \sum_{i=1}^m \left(f_{u^i} - \sum_{\alpha \in \Delta} a_{\alpha,i} \right) x^{u^i} + \sum_{\alpha \in \Delta} \left(\sum_{i=1}^m a_{\alpha,i}x^{u^i} + f_{\alpha}x^{\alpha} \right) + \sum_{\alpha \notin \Delta \cup V} f_{\alpha}x^{\alpha}$$

is SOS. □

Proof of Corollary 2.13. Apply Theorem 2.12 with

$$a_{\alpha,i} = |f_{\alpha}| \lambda_{\alpha,i} \quad \text{for all } \alpha \in \Delta, i = 1, \dots, m. \quad \square$$

Proof of Corollary 2.14. Apply Theorem 2.12 with

$$a_{\alpha,i} = |f_{\alpha}| \lambda_{\alpha}^{\lambda_{\alpha,i}} \quad \text{for all } \alpha \in \Delta, i = 1, \dots, m. \quad \square$$

Proof of Corollary 2.15. Apply Theorem 2.12 with

$$a_{\alpha,i} = \frac{|f_{\alpha}| \lambda_{\alpha}^{\lambda_{\alpha,i}} f_{u^i}}{\prod_{i=1}^m f_{u^i}^{\lambda_{\alpha,i}(x)}} \quad \text{for all } \alpha \in \Delta, i = 1, \dots, m. \quad \square$$

Proof of Corollary 2.16. Apply Theorem 2.12 with

$$a_{\alpha,i} = \begin{cases} k \lambda_{\alpha,i} \left(\frac{|f_{\alpha}|}{k} \right)^{1/\lambda_{\alpha,i} n_{\alpha}} & \text{if } \lambda_{\alpha,i} \neq 0, \\ 0 & \text{if } \lambda_{\alpha,i} = 0 \end{cases} \quad \text{for all } \alpha \in \Delta, i = 1, \dots, m. \quad \square$$

4. Application to global optimization

Let $f \in \mathbf{R}[x_1, \dots, x_n]$ be a real polynomial, and let

$$f_* := \inf \{ f(x) \mid x \in \mathbf{R}^n \}.$$

Since $f_* = \sup \{ r \in \mathbf{R} \mid f - r \text{ is PSD} \}$, finding f_* reduces to determining when $f - r$ is PSD. It is known that deciding when a polynomial is PSD is NP-hard

[1, Theorem 1.1]. Suppose $\deg f = m$ and $f = f_0 + \dots + f_m$, where f_i is a form of degree i . Then a necessary condition for $f_* \neq -\infty$ is that f_m is PSD, and a sufficient condition for $f_* \neq -\infty$ is that f_m is PD, i.e., $f_m(x) > 0$ for all $x \in \mathbf{R}^n$ (see [15, Theorem 5.1 and 5.3]). In [9, Theorem 3.1], H. V. Ha and T. S. Pham have given another necessary condition and another sufficient condition for $f_* \neq -\infty$ in terms of the Newton polyhedron of f , which are improved versions of the results above.

Let

$$f_{SOS} := \sup\{r \in \mathbf{R} \mid f - r \text{ is SOS}\}.$$

Note that $f_* \geq f_{SOS}$. It is known that if $f_{SOS} \neq -\infty$, then f_{SOS} can be computed by semidefinite programming (SDP); see [12, Section 3] and [20]. However, in practice the computation of f_{SOS} can only be carried out if the number of variables and degree are relatively small. The situation is better if f has structured sparsity; see [24, Section 3]. Note that a necessary condition for $f_{SOS} \neq -\infty$ is that $f_{2d} \in \Sigma_{2d,n}$ and a sufficient condition for $f_{SOS} \neq -\infty$ is that $f_{2d} \in \Sigma_{2d,n}^0$, where $\Sigma_{2d,n}^0$ is the interior of $\Sigma_{2d,n}$ in the vector space of forms of degree $2d$, equipped with the euclidean topology (see [17, Proposition 5.1]).

Recently, Ghasemi and Marshall have given another lower bounded f_{gp} for the polynomial f of degree $2d$ with $f_{2de_i} > 0, i = 1, \dots, n$; see [8, Corollary 3.6]. Notice that $f_{gp} \leq f_{SOS}$, and f_{gp} can be computed by geometric programming. As explained in [8, Section 3], although the lower bound found using this method is typically not as good as the lower bound found using SDP, a practical comparison confirms that the computation is faster and larger problems can be handed.

Now, by using Theorem 2.12, we will give a lower bound, which can also be computed by geometric programming, for a larger set of polynomials.

In this section, we also use the term *framework* to denote a set $\mathcal{U} = \{u^0, \dots, u^m\}$ in which $u^i = (u^i_1, \dots, u^i_n) \in (2\mathbf{Z})^n$ with $u^i_j \geq 0$, and use the term *trellis* to denote a framework in which u^0, \dots, u^m comprise the vertices of a simplex.

Let \mathcal{U} be a framework. We also use the term *\mathcal{U} -mediated* to denote a set $\mathcal{L} \subset \mathbf{Z}^n$ which contains \mathcal{U} , and every $v \in \mathcal{L} \setminus \mathcal{U}$ is an average of two distinct even points in \mathcal{L} . Also, by [23, Theorem 2.2], it is easily seen that there is a \mathcal{U} -mediated set \mathcal{U}^* satisfying $\{\frac{1}{2}(s+t) \mid s, t \in \mathcal{U}\} \subset \mathcal{U}^* \subset \text{conv}(\mathcal{U}) \cap \mathbf{Z}^n$ which contains every \mathcal{U} -mediated set.

Let V_∞ denote the set of vertices of $\Gamma_\infty(f) := \text{conv}\{\text{supp}(f) \cup 0\}$. We now assume that $V_\infty = \{u^0, \dots, u^m\}$ is a trellis with the origin point u^0 . Then for every $\alpha \in \Gamma_\infty(f)$, there is only one $\lambda(\alpha) = (\lambda_{\alpha,1}, \dots, \lambda_{\alpha,m})$ with $\lambda_{\alpha,i} \geq 0$ and $|\lambda(\alpha)| := \lambda_1(\alpha) + \dots + \lambda_m(\alpha) \leq 1$ such that

$$\alpha = \lambda_{\alpha,1}u^1 + \dots + \lambda_{\alpha,m}u^m.$$

Recall that

$$\Delta = \{\alpha \in \text{supp}(f) \mid \text{either } f_\alpha < 0 \text{ or } \alpha_i \text{ is odd for some } 1 \leq i \leq n\}.$$

THEOREM 4.1. *Let f be a nonconstant polynomial of degree $2d$, and $r \in \mathbf{R}$. Suppose that $V_\infty = \{u^0, \dots, u^m\}$ is a trellis, and $\alpha \in V_\infty^*$ for all $\alpha \in \Delta$. Then if there exist nonnegative real numbers $a_{\alpha,i}$ for $\alpha \in \Delta$, $i = 1, \dots, m$, $a_{\alpha,i} = 0$ iff $\lambda_{\alpha,i} = 0$, such that*

(i) $a_\alpha^{\lambda(\alpha)} = |f_\alpha| \lambda(\alpha)^{\lambda(\alpha)}$ for every $\alpha \in \Delta$ such that $|\lambda(\alpha)| = 1$,

(ii) $f_{u^i} \geq \sum_{\alpha \in \Delta} a_{\alpha,i}$, $i = 1, \dots, m$,

(iii) $f_{u^0} - r \geq \sum_{\alpha \in \Delta^{<1}} (1 - |\lambda(\alpha)|) \left[\frac{|f_\alpha| \lambda(\alpha)^{\lambda(\alpha)}}{a_\alpha^{\lambda(\alpha)}} \right]^{1/(1-|\lambda(\alpha)|)}$,

then $f - r$ is SOS. Here $\Delta^{<1} = \{\alpha \in \Delta \mid |\lambda(\alpha)| < 1\}$.

Proof. For every $r \in \mathbf{R}$, consider the polynomial $f - r$. By setting $\Omega := \text{supp}(f) \setminus V_\infty$, we can write

$$f(x) = f_{u^0} + \sum_{i=1}^m f_{u^i} x^{u^i} + \sum_{\alpha \in \Omega} f_\alpha x^\alpha.$$

There are two cases to consider.

CASE 1: If $r \neq f_0$, then the homogenization of $g := f - r$ is given by

$$g(x_0, x) = (f_{u^0} - r)x_0^{2d} + \sum_{i=1}^m f_{u^i} x_0^{2d-|u^i|} x^{u^i} + \sum_{\alpha \in \Omega} f_\alpha x_0^{2d-|\alpha|} x^\alpha,$$

It is not hard to show that the set of vertices of the Newton polyhedron $\Gamma(g)$ of g is the following

$$\bar{V}_\infty = \{\bar{u}^0 = (2d, 0), \dots, \bar{u}^m = (2d - |u^m|, u^m)\};$$

and moreover,

- \bar{V}_∞ is a trellis, since V_∞ is a trellis;
- for every $\alpha \in \Delta$, we have $\bar{\alpha} := (2d - |\alpha|, \alpha) \in \bar{V}_\infty^*$, and $(1 - |\lambda(\alpha)|, \lambda_{\alpha,1}, \dots, \lambda_{\alpha,m})$ are the barycentric coordinates of $\bar{\alpha}$ with respect to \bar{V}_∞ ;
- by Conditions (i), (ii), and (iii) of Theorem 4.1, there exist nonnegative real numbers $a_{\alpha,i}$ for $\alpha \in \Delta$, $i = 0, \dots, m$, such that

(i) $a_{\alpha,0}^{1-|\lambda(\alpha)|} a_\alpha^{\lambda(\alpha)} = |f_\alpha| (1 - |\lambda(\alpha)|)^{1-|\lambda(\alpha)|} \lambda(\alpha)^{\lambda(\alpha)}$ for $\alpha \in \Delta$,

(ii) $f_{u^i} \geq \sum_{\alpha \in \Delta} a_{\alpha,i}$, $i = 1, \dots, m$, and $f_{u^0} - r \geq \sum_{\alpha \in \Delta^{<1}} a_{\alpha,0}$.

Thus g satisfies the conditions of Theorem 2.12, and so g is SOS. By Proposition 2.1, $f - r$ is SOS.

CASE 2: If $r = f_0$, then $\Delta^{<1} = \emptyset$, by Condition (iii) of Theorem 4.1. Then, by Theorem 2.12, Conditions (i) and (ii) of Theorem 4.1 show that $\sum_{i=1}^m f_{u^i} x^{u^i} + \sum_{\alpha \in \Delta} f_\alpha x^\alpha$ is SOS. Therefore,

$$f - r = \left(\sum_{i=1}^m f_{u^i} x^{u^i} + \sum_{\alpha \in \Delta} f_\alpha x^\alpha \right) + \sum_{\alpha \notin \Delta} f_\alpha x^\alpha$$

is SOS. □

DEFINITION 4.2. Let f be a nonconstant polynomial of degree $2d$. Suppose $V_\infty = \{u^0, \dots, u^m\}$ is a trellis, and $\alpha \in V_\infty^*$ for all $\alpha \in \Delta$. We define

$$f_{gp} := \sup\{r \in \mathbf{R} \mid \exists a_{\alpha,i} \in \mathbf{R}^{\geq 0}, \alpha \in \Delta, i = 1, \dots, m, a_{\alpha,i} = 0 \text{ iff } \lambda_{\alpha,i} = 0$$

satisfying Conditions (i), (ii), and (iii) of Theorem 4.1\}

REMARK 4.3. 1. As a consequence of Theorem 4.1, we have $f_{gp} \leq f_{SOS}$.
 2. If $f_{u^i} < 0$ for some $i \in \{1, \dots, m\}$, then $f_{gp} = -\infty$.

COROLLARY 4.4. Let f be a nonconstant polynomial of degree $2d$ such that $V_\infty = \{u^0, \dots, u^m\}$ is a trellis. Suppose $\Omega := \text{supp}(f) \setminus V_\infty = \{\alpha\}$, and $\alpha \in V_\infty^*$ if $\alpha \in \Delta$. Then $f_* = f_{SOS} = f_{gp}$.

Proof. We write

$$f(x) = f_{u^0} + \sum_{i=1}^m f_{u^i} x^{u^i} + f_\alpha x^\alpha.$$

Note that $f_{gp} \leq f_{SOS} \leq f_*$. So it is sufficient to show that for every real number r , $f_* \geq r \Rightarrow f_{gp} \geq r$. Fix $r \in \mathbf{R}$ and assume that $f_* \geq r$. We will show $f_{gp} \geq r$; i.e., that r satisfies the constraints of Theorem 4.1. In fact, let g denote the homogenization of $f - r$; i.e.,

$$g(x_0, x) = (f_{u^0} - r)x_0^{2d} + \sum_{i=1}^m f_{u^i} x_0^{2d-|u^i|} x^{u^i} + f_\alpha x_0^{2d-|\alpha|} x^\alpha.$$

Since $f_* \geq r$, we get $g \geq 0$.

CLAIM 4.5. We have $f_{u^i} \geq 0$, $i = 1, \dots, m$, and $f_{u^0} \geq r$.

Proof. Since $V_\infty = \{u^0, \dots, u^m\}$ is a trellis, $\bar{V}_\infty = \{\bar{u}^0, \dots, \bar{u}^m\}$ is linearly independent in \mathbf{R}^{n+1} , where $\bar{u}^i = (u_0^i := 2d - |u^i|, u_1^i, \dots, u_n^i)$, $i = 0, \dots, m$. Without the loss of generality, we can assume

$$\det \begin{bmatrix} u_0^0 & \cdots & u_m^0 \\ \cdots & \cdots & \cdots \\ u_0^m & \cdots & u_m^m \end{bmatrix} \neq 0.$$

By the same arguments as in the proof of Corollary 2.11, for every $\varepsilon \in \mathbf{R}_{>0}$, there exist $c_0, \dots, c_m \in \mathbf{R}_{>0}$ such that

$$\begin{cases} c_0^{u_0^0} \cdots c_m^{u_m^0} = \varepsilon \\ \cdots \cdots \cdots \\ c_0^{u_0^{m-1}} \cdots c_m^{u_m^{m-1}} = \varepsilon \\ c_0^{u_0^m} \cdots c_m^{u_m^m} = 1. \end{cases}$$

Let $\lambda = (\lambda_1, \dots, \lambda_m)$ be such that $\alpha = \lambda_1 u^1 + \dots + \lambda_m u^m$, with $\lambda_i \geq 0$ and $\lambda_1 + \dots + \lambda_m \leq 1$. Taking $\bar{x} := (c_0, \dots, c_m, 1, \dots, 1)$, we have

$$\begin{aligned} g(\bar{x}) &= (f_{u^0} - r)\bar{x}^{\bar{\alpha}^0} + \dots + f_{u^m}\bar{x}^{\bar{\alpha}^m} + f_x(\bar{x}^{\bar{\alpha}^0})^{\lambda_0} \dots (\bar{x}^{\bar{\alpha}^m})^{\lambda_m} \quad (\lambda_0 := 1 - |\lambda|) \\ &= (f_{u^0} - r)(c_0^{\lambda_0} \dots c_m^{\lambda_m}) + \dots + f_{u^m}(c_0^{\lambda_0} \dots c_m^{\lambda_m}) \\ &\quad + f_x[c_0^{\lambda_0} \dots c_m^{\lambda_m}]^{\lambda_0} \dots [c_0^{\lambda_0} \dots c_m^{\lambda_m}]^{\lambda_m} \\ &= (f_{u^0} - r + \dots + f_{u^{m-1}})\varepsilon + f_{u^m} + f_x \varepsilon^{\lambda_0 + \dots + \lambda_{m-1}}. \end{aligned}$$

Since $\alpha \in \Omega$, $\lambda_0 + \dots + \lambda_{m-1} > 0$. Then, since $g \geq 0$, we have $f_{u^m} \geq 0$, by letting $\varepsilon \rightarrow 0$. By the same arguments as the above, we get $f_{u^i} \geq 0$ for $i = 1, \dots, m$, and $f_{u^0} \geq r$. □

To complete the proof of Corollary 4.4, let us consider two cases.

CASE 1: Assume $f_x > 0$ and all α_i are even. Then $\alpha \notin \Delta$, and so $\Delta = \emptyset$. In this case, r satisfies trivially the constraints of Theorem 4.1. So $f_{gp} \geq r$.

CASE 2: Assume either $f_x < 0$ or not all α_i are even. Then $\alpha \in \Delta$; i.e., $\Delta = \Omega = \{\alpha\}$. In this case, applying Corollary 2.11 to g , we deduce that

$$|f_x| \lambda^\lambda (1 - |\lambda|)^{1-|\lambda|} \leq \prod_{i=1}^m f_{u^i}^{\lambda_i} (f_{u^0} - r)^{1-|\lambda|}.$$

There are two subcases to consider.

- If $|\lambda| < 1$, then r satisfies the constraints of Theorem 4.1 with

$$a_{\alpha, i} = \begin{cases} f_{u^i} & \text{if } \lambda_i \neq 0, \\ 0 & \text{if } \lambda_i = 0. \end{cases}$$

- If $|\lambda(\alpha)| = 1$, then $|f_x| \lambda^\lambda \leq \prod_{i=1}^m f_{u^i}^{\lambda_i}$, and so r satisfies the constraints of Theorem 4.1 with

$$a_{\alpha, i} = \begin{cases} s f_{u^i} & \text{if } \lambda_i \neq 0, \\ 0 & \text{if } \lambda_i = 0, \end{cases}$$

where

$$s = \frac{|f_x| \lambda^\lambda}{\prod_{i=1}^m f_{u^i}^{\lambda_i}}.$$

Therefore, $f_{gp} \geq r$. □

DEFINITION 4.6 (geometric program). (1) A function $\phi : \mathbf{R}_{>0}^n \rightarrow \mathbf{R}$ of the form

$$\phi(x) = c x_1^{a_1} \dots x_n^{a_n},$$

where $c > 0$, $a_i \in \mathbf{R}$, and $x = (x_1, \dots, x_n)$, is called a *monomial function*. A sum of monomial functions, i.e., a function of the form

$$\phi(x) = \sum_{i=1}^k c_i x_1^{a_{i1}} \cdots x_n^{a_{in}},$$

where $c_i > 0$ for $i = 1, \dots, k$, is called a *posynomial function*.

(2) An optimization problem of form

$$\begin{cases} \text{Minimize} & \phi_0(x) \\ \text{Subject to} & \phi_i(x) \leq 1, \quad i = 1, \dots, m, \quad \text{and} \quad \psi_j(x) = 1, \quad j = 1, \dots, p, \end{cases}$$

where ϕ_0, \dots, ϕ_m are posynomials and ψ_0, \dots, ψ_p are monomial functions, is called a *geometric program (GP)*. The subset of $\mathbf{R}_{>0}^n$ defined by the constraints

$$\phi_i(x) \leq 1, \quad i = 1, \dots, m, \quad \text{and} \quad \psi_j(x) = 1, \quad j = 1, \dots, p,$$

is called the *feasible set* of the GP. $\phi_0(x)$ is called the *objective function*. The output of the GP is the minimum, more precisely, the infimum, of $\phi_0(x)$, taken as x runs through the feasible set. In case feasible set is empty, the output is understood to be $+\infty$.

See [3, Section 4.5] or [21, Section 3.5] for more about GPs.

THEOREM 4.7. *Let f be a nonconstant polynomial of degree $2d$. Suppose that*

- (i) $V_\infty = \{u^0, \dots, u^m\}$ is a trellis,
- (ii) $\alpha \in V_\infty^*$ for all $\alpha \in \Delta$.

Then $f_{gp} = f_0 - r^*$, where r^* is the output of the GP

$$\begin{cases} \text{Minimize} & \sum_{\alpha \in \Delta^{<1}} (1 - |\lambda(\alpha)|) \left[\frac{|f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)}}{a_\alpha^{\lambda(\alpha)}} \right]^{1/(1-|\lambda(\alpha)|)} \\ \text{Subject to} & \sum_{\alpha \in \Delta} \frac{a_{\alpha,i}}{f_{u^i}} \leq 1, \quad i = 1, \dots, m, \quad \text{and} \\ & \frac{a_\alpha^{\lambda(\alpha)}}{|f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)}} = 1, \quad \alpha \in \Delta, \quad |\lambda(\alpha)| = 1. \end{cases}$$

The variables in the program are the $a_{\alpha,i}$, $\alpha \in \Delta$, $i = 1, \dots, m$, $\lambda_i(\alpha) \neq 0$, the understanding being that $a_{\alpha,i} = 0$ iff $\lambda_i(\alpha) = 0$.

Proof. Observe that

$$\phi_0(a) := \sum_{\alpha \in \Delta^{<1}} (1 - |\lambda(\alpha)|) \left[\frac{|f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)}}{a_\alpha^{\lambda(\alpha)}} \right]^{1/(1-|\lambda(\alpha)|)}$$

and

$$\phi_i(a) := \sum_{\alpha \in \Delta} \frac{a_{\alpha,i}}{f_{u^i}}, \quad i = 1, \dots, m,$$

are posynomials in the variables $a_{\alpha,i}$, and

$$\psi_\alpha(a) := \frac{a_\alpha^{\lambda(\alpha)}}{|f_\alpha|^{\lambda(\alpha)}}, \quad \alpha \in \Delta, |\lambda(\alpha)| = 1,$$

are monomial functions in the variables $a_{\alpha,i}$. By the definition of f_{gp} , it follows immediately that $f_{gp} = f_0 - r^*$. \square

COROLLARY 4.8. *Let f be a nonconstant polynomial of degree $2d$. Suppose that*

- (i) $V_\infty = \{u^0, \dots, u^m\}$ is a trellis,
- (ii) $\alpha \in V_\infty^*$ for all $\alpha \in \Delta$.

Then if $|\lambda(\alpha)| < 1$ for each $\alpha \in \Delta$, then $f_{gp} \neq -\infty$ and $f_{gp} = f_0 - r^$, where r^* is the output of the GP*

$$\begin{cases} \text{Minimize} & \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[\frac{|f_\alpha|^{\lambda(\alpha)}}{a_\alpha^{\lambda(\alpha)}} \right]^{1/1-|\lambda(\alpha)|} \\ \text{Subject to} & \sum_{\alpha \in \Delta} \frac{a_{\alpha,i}}{f_{u^i}} \leq 1, \quad i = 1, \dots, m. \end{cases}$$

Proof. In this case, the equality constraints in the computation of r^* are vacuous and the feasible set is always nonempty, so $f_{gp} \neq -\infty$. The rest is immediate from Theorem 4.7. \square

We establish some lower bounds in terms of its coefficients, which can be obtained by evaluating the objective function of the GP in Corollary 4.8 at suitably chosen feasible points. Corollaries 4.9, 4.10, and 4.11 are improved versions of [8, Corollaries 4.1, 4.2, and 4.3] (and so [7, Theorems 3.1, 3.2, and 3.3]) respectively.

Recall that for a (univariate) polynomial of the form $p(t) = t^n - \sum_{i=0}^{n-1} a_i t^i$, where each a_i is nonnegative and at least one a_i is nonzero, there is a unique positive root of p (see [19, Theorem 1.1.3]). Let $C(p)$ denote the unique positive root of p . See [5], [11, Example 4.6.2], and [8, Proposition 1.2] for more details and upper bounds for $C(p)$.

COROLLARY 4.9. *Let f be a nonconstant polynomial of degree $2d$. Suppose that*

- (i) $V = \{u^0, \dots, u^m\}$ is a trellis,
- (ii) $\alpha \in V^*$ for all $\alpha \in \Delta$, and
- (iii) $|\lambda(\alpha)| < 1$ for each $\alpha \in \Delta$.

Then $f_{gp} \geq r_L$, where

$$r_L := f_0 - \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) |f_\alpha| k^{2D|\lambda(\alpha)|} f_V^{-\lambda(\alpha)},$$

$$k \geq \max_{i=1, \dots, m} C \left(t^{2D} - \sum_{\alpha \in \Delta} \lambda_{\alpha, i} |f_\alpha| f_{u^i}^{-|\lambda(\alpha)|} t^{2D|\lambda(\alpha)|} \right).$$

Here $f_V := (f_{u^1}, \dots, f_{u^m})$, and D is a nonnegative integer number such that $2D\lambda_{\alpha, i} \in \mathbf{Z}$ for all $\alpha \in \Delta$, $i = 1, \dots, m$.

Proof. For each $\alpha \in \Delta$ and $i = 1, \dots, m$, let

$$a_{\alpha, i} = \frac{\lambda_{\alpha, i} |f_\alpha| f_{u^i}^{1-|\lambda(\alpha)|}}{k^{2D(1-\lambda(\alpha))}}.$$

By the definition of k , for each $i = 1, \dots, m$, we have $\sum_{\alpha \in \Delta} \lambda_{\alpha, i} |f_\alpha| f_{u^i}^{-|\lambda(\alpha)|} k^{2D|\lambda(\alpha)|} \leq k^{2D}$. Hence

$$\sum_{\alpha \in \Delta} a_{\alpha, i} = \sum_{\alpha \in \Delta} \frac{\lambda_{\alpha, i} |f_\alpha| f_{u^i}^{1-|\lambda(\alpha)|}}{k^{2D(1-\lambda(\alpha))}} \leq f_{u^i}.$$

This shows that the array $(a_{\alpha, i} \mid \alpha \in \Delta, i = 1, \dots, m)$ is a feasible point for the geometric program in the statement of Corrolary 4.8. Plugging this into the objective function of the program yields

$$\begin{aligned} & \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[\frac{|f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)}}{a_\alpha^{\lambda(\alpha)}} \right]^{1/(1-|\lambda(\alpha)|)} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[|f_\alpha| \prod_{\lambda_{\alpha, i} \neq 0} \left(\frac{\lambda_{\alpha, i}}{a_{\alpha, i}} \right)^{\lambda_{\alpha, i}} \right]^{1/(1-|\lambda(\alpha)|)} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[|f_\alpha| \prod_{\lambda_{\alpha, i} \neq 0} \left(\frac{\lambda_{\alpha, i} k^{2D(1-\lambda(\alpha))}}{\lambda_{\alpha, i} |f_\alpha| f_{u^i}^{1-|\lambda(\alpha)|}} \right)^{\lambda_{\alpha, i}} \right]^{1/(1-|\lambda(\alpha)|)} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) |f_\alpha| k^{2D|\lambda(\alpha)|} \prod_{\lambda_{\alpha, i} \neq 0} f_{u^i}^{-\lambda_{\alpha, i}} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) |f_\alpha| k^{2D|\lambda(\alpha)|} f_V^{-\lambda(\alpha)}. \end{aligned}$$

So, $r_L = f_0 - \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) |f_\alpha| k^{2D|\lambda(\alpha)|} f_V^{-\lambda(\alpha)} \leq f_{gp}$. □

COROLLARY 4.10. *Let f be a nonconstant polynomial of degree $2d$. Suppose that*

- (i) $V = \{u^0, \dots, u^m\}$ is a trellis,
- (ii) $\alpha \in V^*$ for all $\alpha \in \Delta$, and
- (iii) $|\lambda(\alpha)| < 1$ for each $\alpha \in \Delta$.

Then $f_{gp} \geq r_{FK}$, where $r_{FK} := f_0 - k^{2D}$, with

$$k \geq C \left(t^{2D} - \sum_{j=1}^{2D-1} b_j t^j \right), \quad \text{and}$$

$$b_j := \left(\frac{2D-j}{2D} \right)^{(2D-j)/2D} \sum_{\alpha \in \Delta, 2D|\lambda(\alpha)|=j} |f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)} f_V^{-\lambda(\alpha)}, \quad j = 1, \dots, 2D-1.$$

Here D is a nonnegative integer number such that $2D\lambda_i(\alpha) \in \mathbf{Z}$ for all $\alpha \in \Delta$, $i = 1, \dots, m$.

Proof. For each $\alpha \in \Delta$ and $i = 1, \dots, m$, let

$$a_{\alpha,i} = (1 - |\lambda(\alpha)|)^{1-|\lambda(\alpha)|} |f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)} f_V^{-\lambda(\alpha)} f_{u^i} k^{j-2D}.$$

By the definition of k , we have $\sum_{j=1}^{2D-1} b_j k^j \leq k^{2D}$. Hence

$$\begin{aligned} \sum_{\alpha \in \Delta} a_{\alpha,i} &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|)^{1-|\lambda(\alpha)|} |f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)} f_V^{-\lambda(\alpha)} f_{u^i} k^{j-2D} \\ &= \sum_{j=1}^{2D-1} \sum_{\alpha \in \Delta, 2D|\lambda(\alpha)|=j} \left(\frac{2D-j}{2D} \right)^{(2D-j)/2D} |f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)} f_V^{-\lambda(\alpha)} f_{u^i} k^{j-2D} \\ &= f_{u^i} \sum_{j=1}^{2D-1} k^{j-2D} \left(\frac{2D-j}{2D} \right)^{(2D-j)/2D} \sum_{\alpha \in \Delta, 2D|\lambda(\alpha)|=j} |f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)} f_V^{-\lambda(\alpha)} \\ &= f_{u^i} k^{-2D} \sum_{j=1}^{2D-1} k^j \left(\frac{2D-j}{2D} \right)^{(2D-j)/2D} \sum_{\alpha \in \Delta, 2D|\lambda(\alpha)|=j} |f_\alpha| |\lambda(\alpha)|^{\lambda(\alpha)} f_V^{-\lambda(\alpha)} \\ &= f_{u^i} k^{-2D} \sum_{j=1}^{2D-1} b_j k^j \leq f_{u^i}. \end{aligned}$$

This shows that the array $(a_{\alpha,i} | \alpha \in \Delta, i = 1, \dots, m)$ is a feasible point for the geometric program in the statement of Corollary 4.8. Plugging this into the objective function of the program yields

$$\begin{aligned}
& \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[\frac{|f_\alpha| \lambda(\alpha)^{\lambda(\alpha)}}{a_\alpha^{\lambda(\alpha)}} \right]^{1/(1-|\lambda(\alpha)|)} \\
&= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[|f_\alpha| \prod_{\lambda_{\alpha,i} \neq 0} \left(\frac{\lambda_{\alpha,i}}{a_{\alpha,i}} \right)^{\lambda_{\alpha,i}} \right]^{1/(1-|\lambda(\alpha)|)} \\
&= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[|f_\alpha| \prod_{\lambda_{\alpha,i} \neq 0} \left(\frac{f_V^{\lambda(\alpha)} k^{2D-j} \lambda_{\alpha,i}}{(1 - |\lambda(\alpha)|)^{1-|\lambda(\alpha)|} |f_\alpha| \lambda(\alpha)^{\lambda(\alpha)} f_{u^i}} \right)^{\lambda_{\alpha,i}} \right]^{1/(1-|\lambda(\alpha)|)} \\
&= \sum_{j=1}^{2D-1} \sum_{\alpha \in \Delta, 2D|\lambda(\alpha)|=j} \left(\frac{2D-j}{2D} \right) \\
&\quad \times \left[|f_\alpha| \prod_{\lambda_{\alpha,i} \neq 0} \left(\frac{f_V^{\lambda(\alpha)} k^{2D-j} \lambda_{\alpha,i}}{\left(\frac{2D-j}{2D} \right)^{(2D-j)/2D} |f_\alpha| \lambda(\alpha)^{\lambda(\alpha)} f_{u^i}} \right)^{\lambda_{\alpha,i}} \right]^{2D/(2D-j)} \\
&= \sum_{j=1}^{2D-1} k^j \left(\frac{2D-j}{2D} \right)^{(2D-j)/2D} \sum_{\alpha \in \Delta, 2D|\lambda(\alpha)|=j} |f_\alpha| \lambda(\alpha)^{\lambda(\alpha)} f_V^{-\lambda(\alpha)} \\
&= \sum_{j=1}^{2D-1} b_j k^j \leq k^{2D}.
\end{aligned}$$

So, $r_{FK} = f_0 - k^{2D} \leq f_{gp}$. □

COROLLARY 4.11. *Let f be a nonconstant polynomial of degree $2d$. Suppose that*

- (i) $V = \{u^0, \dots, u^m\}$ is a trellis,
- (ii) $\alpha \in V^*$ for all $\alpha \in \Delta$, and
- (iii) $|\lambda(\alpha)| < 1$ for each $\alpha \in \Delta$.

Then

$$f_{gp} \geq r_{mt} := f_0 - \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) [|f_\alpha| t^{|\lambda(\alpha)|} \lambda(\alpha)^{\lambda(\alpha)} f_V^{-\lambda(\alpha)}]^{1/(1-|\lambda(\alpha)|)},$$

where $t := |\Delta|$.

Proof. For each $\alpha \in \Delta$ and $i = 1, \dots, m$, let $a_{\alpha,i} = \frac{f_{u^i}}{t}$. Then

$$\sum_{\alpha \in \Delta} a_{\alpha,i} = \sum_{\alpha \in \Delta} \frac{f_{u^i}}{t} = f_{u^i}.$$

This shows that the array $(a_{\alpha,i} \mid \alpha \in \Delta, i = 1, \dots, m)$ is a feasible point for the geometric program in the statement of Corollary 4.8. Plugging this into the objective function of the program yields

$$\begin{aligned} & \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[\frac{|f_{\alpha}| \lambda(\alpha)^{\lambda(\alpha)}}{a_{\alpha}^{\lambda(\alpha)}} \right]^{1/(1-|\lambda(\alpha)|)} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[|f_{\alpha}| \prod_{\lambda_{\alpha,i} \neq 0} \left(\frac{\lambda_{\alpha,i}}{a_{\alpha,i}} \right)^{\lambda_{\alpha,i}} \right]^{1/(1-|\lambda(\alpha)|)} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) \left[|f_{\alpha}| \prod_{\lambda_{\alpha,i} \neq 0} \left(\frac{t \lambda_{\alpha,i}}{f_{u^i}} \right)^{\lambda_{\alpha,i}} \right]^{1/(1-|\lambda(\alpha)|)} \\ &= \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) [|f_{\alpha}| t^{|\lambda(\alpha)|} \lambda(\alpha)^{\lambda(\alpha)} f_V^{-\lambda(\alpha)}]^{1/(1-|\lambda(\alpha)|)}. \end{aligned}$$

So, $r_{dnt} = f_0 - \sum_{\alpha \in \Delta} (1 - |\lambda(\alpha)|) [|f_{\alpha}| t^{|\lambda(\alpha)|} \lambda(\alpha)^{\lambda(\alpha)} f_V^{-\lambda(\alpha)}]^{1/(1-|\lambda(\alpha)|)} \leq f_{gp}$. \square

Acknowledgments. The authors are greatly indebted to Ha Huy Vui for suggesting the problem and for many stimulating conversations.

REFERENCES

- [1] M. BELLARE AND P. ROGAWAY, The complexity of approximating a nonlinear program, *Math. Program.* **69** (1993), 429–441.
- [2] J. BOCHNAK, M. COSTE AND M. F. ROY, *Real algebraic geometry*, *Ergeb. Math. Grenzgeb.* **36**, Springer, New York, 1998.
- [3] S. BOYD AND L. VANDENBERGHE, *Convex optimization*, Cambridge University Press, Cambridge, 2004.
- [4] M. D. CHOI, T. Y. LAM AND B. REZNICK, Sum of squares of real polynomials, *Proc. Sympos. Pure Math.* **58** (1995), 103–126.
- [5] E. DEUTSCH, Bounds for the zeros of polynomial, *Amer. Math. Monthly* **88** (1981), 205–206.
- [6] C. FIDALGO AND A. KOVACEC, Positive semidefinite diagonal minus tail forms are sum of squares, *Math. Z.* **269** (2011), 629–645.
- [7] M. GHASEMI AND M. MARSHALL, Lower bounds for polynomials in terms of its coefficients, *Arch. Math.* **95** (2010), 343–354.
- [8] M. GHASEMI AND M. MARSHALL, Lower bounds for polynomials using geometric programming, *SIAM J. Optim.* **22** (2012), 460–473.
- [9] H. V. HA AND T. S. PHAM, Minimizing polynomial functions, *Acta Math. Vietnamica* **32** (2007), 71–82.
- [10] D. HILBERT, *Über die Darstellung definiter Formen als Summe von Formenquadraten*, *Math. Ann.* **32** (1888), 342–350 (in German).
- [11] D. KNUTH, *The art of computer programming 2*, Addison-Wesley, New York, 1969.
- [12] J. B. LASSERRE, Global optimization with polynomials and the problem of moments, *SIAM J. Optim.* **11** (2001), 796–817.

- [13] J. B. LASSERRE, Sufficient conditions for a real polynomial to be a sum of squares, *Arch. Math. (Basel)* **89** (2007), 390–398.
- [14] J. B. LASSERRE AND T. NETZER, SOS approximation of nonnegative polynomial via simple high degree perturbations, *Math. Z.* **256** (2006), 99–112.
- [15] M. MARSHALL, Optimization of polynomials functions, *Canad. Math. Bull.* **46** (2008), 537–587.
- [16] M. MARSHALL, Positive polynomials and sum of squares, *Math. Surveys Monogr.* **146**, American Mathematical Society, Providence, RI, 2008.
- [17] M. MARSHALL, Representation of non-negative polynomials, degree bounds and application to optimization, *Canad. J. Math.* **61** (2009), 205–221.
- [18] T. MOTZKIN, The arithmetic-geometric inequality, *Inequalities* (O. Shisha, ed.), Academic Press, New York, 1970, 205–224.
- [19] V. V. PRASOLOV, Polynomials, *Algorithms comput. math.* **11**, Springer, New York, 2004.
- [20] P. PARRILO AND B. STURMEELS, Minimizing polynomial functions, *DIMACS Serie in Discrete Mathematics and Theoretical Computer Science* **60**, AMS, 2003, 83–99.
- [21] A. L. PERESSINI, F. E. SULLIVAN AND J. J. UHL, JR, The mathematics of nonlinear programming, *Undergrad. texts math.*, Springer, New York, 1987.
- [22] B. REZNICK, Midpoint polytopes and the map $x_i \rightarrow x_i^k$, in preparation.
- [23] B. REZNICK, Forms derived from the arithmetic-geometric inequality, *Math. Ann.* **383** (1989), 431–464.
- [24] H. WAKI, S. KIM, M. KOJIMA AND M. MURAMATSU, Sum of squarer and semidefinite program relaxations for polynomials optimization problems with structured sparsity, *SIAM J. Optim.* **17** (2006), 218–242.

Van Doat Dang
THANG LONG HIGH SCHOOL FOR GIFTED
10 TRAN PHU, DA LAT
VIETNAM
E-mail: dangvandoat@yahoo.com.vn

Thi Thao Nguyen
DEPARTMENT OF MATHEMATICS
HANOI NATIONAL UNIVERSITY OF EDUCATION
136 XUAN THUY ROAD
CAU GIAY DISTRICT, HANOI
VIETNAM
E-mail: math_thao@yahoo.com.vn