

36. On Regular Duo Rings

By Sándor LAJOS

K. Marx University of Economics, Budapest, Hungary

(Comm. by Kinjirô KUNUGI, M. J. A., March 12, 1969)

Recently the author proved some characterizations of semigroups, which are semilattices of groups (see [3]-[5]). One of these characterizations reads as follows:

Theorem 1. *A semigroup S is a semilattice of groups if and only if the relation*

$$(1) \quad R \cap L = LR$$

holds for any left ideal L and right ideal R of S .

In this paper we formulate an analogous assertion in the theory of associative rings. A ring A is called a *duo* (or *two-sided*) ring if every one-sided (left or right) ideal of A is a two-sided ideal (see Hille [1]). An associative ring A is said to be *regular* if to every element a in A there exists an element $x \in A$ such that $axa = a$ (see von Neumann [7]). Interesting characterizations of regular rings were obtained by Kovács [2] and Luh [6].

Theorem 2. *An associative ring A is a regular duo ring if and only if the relation (1) holds for any left ideal L and right ideal R of A .*

Proof. Necessity. Let A be a regular duo ring. Then A satisfies the relation

$$(2) \quad R \cap L = RL$$

for every left ideal L and right ideal R of A (see Kovács [2]). Now (2) implies (1) because each one-sided ideal of a duo ring is a two-sided ideal.

Sufficiency. Let A be an associative ring having the property (1) for any left ideal L and right ideal R of A . In case of $R = A$ the relation (1) implies

$$(3) \quad A \cap L = LA,$$

whence every left ideal L of A is also a right ideal of A . Similarly, in case of $L = A$ (1) implies that

$$(4) \quad R \cap A = AR,$$

whence the right ideal R of A is a two-sided ideal of A . Therefore A is a duo ring. Finally (1) implies (2), which is equivalent with the regularity of A .

Theorem 2 is completely proved.

References

- [1] E. Hille: Functional analysis and semi-groups. Amer. Math. Soc., Coll. Publ. New York (1948).
- [2] L. Kovács: A note on regular rings. Publ. Math. Debrecen, **4**, 465–468 (1955–56).
- [3] S. Lajos: Characterizations of semigroups, which are semilattices of groups (in Hungarian). Magyar Tud. Akad. Mat. Fiz. Oszt. Közl. (to appear).
- [4] —: Note on semigroups, which are semilattices of groups. Proc. Japan Acad., **44**, 805–806 (1968).
- [5] —: A characterization of semigroups, which are semilattices of groups (to appear).
- [6] J. Luh: A characterization of regular rings. Proc. Japan Acad., **39**, 741–742 (1963).
- [7] J. von Neumann: On regular rings. Proc. Nat. Acad. Sci. U. S. A., **22**, 707–713 (1936).