

## A MORSE FUNCTION ON GRASSMANN MANIFOLDS

THEODOR HANGAN

Studying the critical sections of a convex body Wen Tsun Wu has obtained in [2] a Morse function on a Grassmann manifold. In the sequel it will be shown that another function may be obtained by composing the embedding of this manifold into a projective space with the well known Morse function of the projective space; our work is valid only for the real and complex fields.

1. The homology of the Grassmann manifold  $G_{p,q}$  of all the  $p$ -planes of codimension  $q$  which pass through a fixed point  $0$  in an affine space  $A^n$  of dimension  $n = p + q$  was determined in 1934 by Ch. Ehresmann who gave a cell subdivision of  $G_{p,q}$ . The number of cells in his subdivision is the number  $N = \binom{n}{p}$  of combinations of  $p$  elements of the set  $\{1, \dots, n\}$ ; such a combination  $\sigma = (\sigma_1, \dots, \sigma_p)$  where  $1 \leq \sigma_1 < \dots < \sigma_p \leq n$  is called a Schubert symbol. In the cell-subdivision of  $G_{p,q}$ , with each symbol  $\sigma$  one associates a cell of dimension

$$d(\sigma) = (\sigma_1 - 1) + \dots + (\sigma_p - p).$$

Let us consider the lexicographical order in the set  $S(p, q)$  of all the Schubert symbols which correspond to the integers  $p$  and  $q$ ; this means that  $\sigma = (\sigma_1, \dots, \sigma_p) < \sigma' = (\sigma'_1, \dots, \sigma'_p)$  if and only if for the least integer  $i \leq p$  for which  $\sigma_i \neq \sigma'_i$  the inequality  $\sigma_i < \sigma'_i$  holds. We say that two symbols  $\sigma = (\sigma_1, \dots, \sigma_p)$  and  $\sigma' = (\sigma'_1, \dots, \sigma'_p)$  are *neighboring* if the sets  $\{\sigma_1, \dots, \sigma_p\}$  and  $\{\sigma'_1, \dots, \sigma'_p\}$  have exactly  $p - 1$  elements in common, or equivalently, if they differ only in what a single element is concerned. With these conventions we observe that the number  $d(\sigma)$  equals the number of those Schubert-symbols which are less than and neighboring to  $\sigma$ . Indeed, in order to obtain a new symbol less than and neighboring to  $\sigma$ , the change of  $\sigma_i$  in  $\sigma$  may be made in  $\sigma_i - i$  ways by replacing  $\sigma_i$  with a positive integer less than  $\sigma_i$  and different from  $\sigma_1, \dots, \sigma_{i-1}$ .

2. In the projective space  $P^{N-1}$  of dimension  $N - 1$  we consider homogeneous coordinates  $y_\alpha$  having as indices Schubert symbols  $\sigma \in S(p, q)$  instead of positive integers running from 1 to  $N$ .

It is known, for example from [1], that the function

$$(1) \quad f = \sum_{\sigma \in S(p, q)} c_{\sigma} |y_{\sigma}|^2,$$

where  $c_{\sigma}$  are constants, and satisfy the inequalities  $c_{\sigma} < c_{\sigma'}$ , when  $\sigma < \sigma'$  defines a Morse function on  $P^{N-1}$  when the coordinates  $y_{\sigma}$  satisfy the equation

$$(2) \quad \sum_{\sigma \in S(p, q)} |y_{\sigma}|^2 = 1.$$

The critical points of this function  $f$  correspond to the coordinate axes in the numerical  $N$ -dimensional space of the variables  $y_{\sigma}$ , and therefore may be denoted by  $A_{\sigma}$ ,  $\sigma \in S(p, q)$ . The index of the point  $A_{\sigma}$  corresponding to the  $y_{\sigma}$ -axis is equal to and twice the number of constants  $c_{\sigma'}$ , which are less than  $c_{\sigma}$ , in the real and complex cases respectively. In other words, this index equals  $n_{\sigma} - 1$  in the real case and  $2(n_{\sigma} - 1)$  in the complex case, where  $n_{\sigma}$  is the number associated with  $\sigma$  in the ordering of  $S(p, q)$ .

3. In the affine space  $A^n$  of dimension  $n$  denote by  $e_a$ ,  $a = 1, \dots, n$ , the basis vectors of the system of cartesian coordinates having the origin at 0. Consider  $p$  linearly independent vectors  $v_{\alpha}$ ,  $\alpha = 1, \dots, p$ , with components with respect to the basis  $\{e_a\}$  denoted by  $v_{\alpha}^a$  ( $v_{\alpha} = \sum_{a=1}^n v_{\alpha}^a e_a$ ) and form the determinants

$$(3) \quad v^{\sigma} = \det \|v_{\alpha}^{\sigma}\|, \quad \alpha = 1, \dots, p, \quad \sigma \in S(p, q),$$

which realize a system of Plücker coordinates for the  $p$ -plane spanned by the  $p$  vectors  $v_{\alpha}$ . The Plückerian embedding  $\pi$  of  $G_{p, q}$  in  $P^{N-1}$  is given by the equations

$$(4) \quad y_{\sigma} = v^{\sigma}.$$

Observe that when  $v_{\alpha} = e_{\sigma_{\alpha}}$  the corresponding  $p$ -plane has the only non-zero component  $y_{\sigma} = 1$  and thus the points  $A_{\sigma}$ , which are critical points for the function  $f$ , belong to the image  $\pi(G_{p, q})$ .

**Theorem.** *The function  $f \circ \pi: G_{p, q} \rightarrow R$  is a Morse function having  $N = \binom{n}{p}$  nondegenerate critical points, which are  $\pi^{-1}(A_{\sigma})$ , and the index of each such point is  $d(\sigma)$  in the real case and  $2d(\sigma)$  in the complex case.*

4. The points  $\pi^{-1}(A_{\sigma})$  are critical for the function  $f \circ \pi$  since their images  $A_{\sigma}$  are so for the function  $f$ . In order to show that the critical points  $\pi^{-1}(A_{\sigma})$  are nondegenerate and their index is  $d(\sigma)$ , we introduce a system of local coordinates on  $G_{p, q}$  in the neighborhood  $U_{\sigma}$  of the  $p$ -plane  $\pi^{-1}(A_{\sigma})$  whose points are the  $p$ -planes having a nondegenerate projection on  $\pi^{-1}(A_{\sigma})$ . Clearly, if  $e_{\bar{\sigma}_i}$ ,  $i = p + 1, \dots, p + q$ ,  $1 \leq \bar{\sigma}_i \leq n$ ,  $\bar{\sigma}_i \neq \sigma_{\alpha}$ , are the vectors of the already chosen basis in  $A^n$ , which are not in  $\pi^{-1}(A_{\sigma})$ , then the  $pq$  local coordinates  $x_{\alpha}^i$  of a point  $x$  belonging to  $U_{\sigma}$  are determined by the formulas

$$v_\alpha = e_{\sigma_\alpha} + \sum_{i=p+1}^n x_\alpha^i e_{\bar{\sigma}_i}, \quad \alpha = 1, \dots, p,$$

$v_\alpha$  being the generating vectors of  $x$ . Observe now that  $v^\sigma = 1$  and that the only determinants  $v^\rho$ ,  $\rho \in S(p, q)$ , which are linear functions of the coordinates  $x_\alpha^i$ , are those corresponding to the symbols  $\rho$  which are neighboring to  $\sigma$ . The other determinants  $v^\rho$  are homogeneous polynomials in  $x_\alpha^i$  of degree greater than one. In order for the embedding  $\pi: G_{p,q} \rightarrow P^{N-1}$  to satisfy the condition (2) we use the following formulas:

$$(5) \quad y_\rho = \frac{v^\rho}{\left( \sum_{\tau \in S(p,q)} |v^\tau|^2 \right)^{1/2}}.$$

Thus the function  $F = f \circ \pi$  becomes

$$(6) \quad F = \frac{\sum_{\rho \in S(p,q)} c_\rho |v^\rho|^2}{\sum_{\tau \in S(p,q)} |v^\tau|^2},$$

and at the origin of the system of coordinates  $x_\alpha^i$  the value of this function  $F$  is  $c_\sigma$ . This point is a critical one and the quadratic form  $F_\sigma$ , which approximates the function  $F - c_\sigma$  in the neighborhood of the origin, is

$$F_\sigma = \sum (c_{\sigma'} - c_\sigma) |v^{\sigma'}|^2,$$

where  $\sigma'$  is neighboring to  $\sigma$ .  $|v^{\sigma'}|^2$  is the square of one of the coordinates  $x_\alpha^i$  in the real case, and is its modulus  $(\text{Re } x_\alpha^i)^2 + (\text{Im } x_\alpha^i)^2$  in the complex case. Hence the last part of the theorem follows from the choice of the constants  $c_\rho$ .

5. It remains to be proved that the function  $F$  has no other critical points different from  $\pi^{-1}(A_\sigma)$ . In order to do this suppose that  $v$  is a critical point for  $F$ , and that  $\sigma$  is the least Schubert symbol having the property that the  $p$ -plane  $v$  belongs to  $U_\sigma$ . Thus the matrix of the components of a system of  $p$  vectors  $v_\alpha$  which span the  $p$ -plane  $v$  in  $U_\sigma$  is of the form

$$(7) \quad \begin{pmatrix} 0 \dots 0 & 1 & v_1^{\sigma_1+1} \dots 0 & v_1^{\sigma_2+1} \dots 0 & \dots & v_1^n \\ 0 \dots & & & 0 & 1 & v_2^{\sigma_2+1} \dots 0 & \dots & v_2^n \\ \dots & & & & & & & \\ 0 & & & & & & 0 & 1 & v_p^{\sigma_p+1} \dots & v_p^n \end{pmatrix}.$$

Clearly  $v_\alpha^a = 0$ ,  $a < \sigma_\alpha$ , and  $v_\alpha^{\sigma_\alpha} = \delta_\alpha^b$  where  $\delta_\alpha^b$  is the Kronecker symbol. Now we consider the curve  $w: (-\epsilon, \epsilon) \rightarrow G_{p,q}$  obtained by keeping the vectors  $v_2, \dots, v_p$  constant and varying only  $v_1$  in accord with the formulas

$$(8) \quad w_1^i = v_1^i + t v_1^{\sigma_1}, \quad i \neq \sigma_1; \quad w_1^{\sigma_1} = v_1^{\sigma_1} = 1, \quad w_\alpha = v_\alpha, \quad \alpha = 2, \dots, p.$$

Thus, when  $v^i, i \neq \sigma_1$ , are not all zero,  $\left(\frac{dF(w(t))}{dt}\right)_{t=0} \neq 0$  which contradicts the hypothesis that  $v$  is a critical point. Indeed, from (6) we obtain

$$(9) \quad \left(\frac{dF(w(t))}{dt}\right)_{t=0} = 2 \operatorname{Re} \frac{\left(\sum_p c_p v^p \bar{w}_0^{p'}\right) \left(\sum_\tau |v^\tau|^2\right) - \left(\sum_\tau v^\tau \bar{w}_0^{\tau'}\right) \left(\sum_p c_p |v^p|^2\right)}{\left(\sum_\tau |v^\tau|^2\right)^2},$$

where  $w_0^{p'} = \left(\frac{dw^p}{dt}\right)_{t=0}$ . From (7) and (8) we observe that if  $w_0^{p'} \neq 0$ , then the symbol  $\rho = (\rho_1, \dots, \rho_p)$  must have  $\rho_1 > \sigma_1$  and in this case  $w_0^{p'} = v^p$ . We write such a symbol in the form  $\rho = \rho_1 \bar{\rho}$  where  $\bar{\rho} \in S(p-1, q+1)$  is the Schubert symbol  $\rho = (\rho_2, \dots, \rho_p)$ . With this convention the numerator on the right-hand side of (9) then becomes

$$\begin{aligned} \mathfrak{N}_1 &= \left(\sum_{\rho_1 > \sigma_1, \bar{\rho}} c_{\rho_1 \bar{\rho}} |v^{\rho_1 \bar{\rho}}|^2\right) \left(\sum_{\tau_1 > \sigma_1, \bar{\tau}} |v^{\tau_1 \bar{\tau}}|^2 + \sum_{\bar{\tau}} |v^{\sigma_1 \bar{\tau}}|^2\right) \\ &\quad - \left(\sum_{\tau_1 > \sigma_1, \bar{\tau}} |v^{\tau_1 \bar{\tau}}|^2\right) \left(\sum_{\rho_1 > \sigma_1, \bar{\rho}} c_{\rho_1 \bar{\rho}} |v^{\rho_1 \bar{\rho}}|^2 + \sum_{\bar{\rho}} c_{\sigma_1 \bar{\rho}} |v^{\sigma_1 \bar{\rho}}|^2\right) \\ &= \sum_{\rho_1 > \sigma_1, \bar{\rho}, \bar{\tau}} (c_{\rho_1 \bar{\rho}} - c_{\sigma_1 \bar{\tau}}) |v^{\rho_1 \bar{\rho}}|^2 |v^{\sigma_1 \bar{\tau}}|^2. \end{aligned}$$

But  $c_{\rho_1 \bar{\rho}} > c_{\sigma_1 \bar{\tau}}$  since  $\rho_1 > \sigma_1$ , and as among the components  $v^{\sigma_1 \bar{\tau}}$  there is at least one different from zero (the component  $v^\sigma = 1$ ) we infer that  $\mathfrak{N}_1 = 0$  only if all the determinants  $v^{\rho_1 \bar{\rho}}$  vanish. Among these determinants  $v^\rho$  we find those, for which  $p-1$  indices in the symbol  $\rho$  coincide with  $\sigma_2, \dots, \sigma_p$  and are equal to  $\pm v^i, i > \sigma_1$ . Thus, if  $v$  is a critical point for  $F$ , then its coordinates  $v^i, i > \sigma_1$ , must vanish. The same method may be used to show that all the components  $v^\alpha, i > \sigma_\alpha$ , vanish for a critical point  $v$ . Indeed suppose that for a critical point  $v$  we have

$$(10) \quad v^\alpha = 0, i > \sigma_\alpha, \alpha = 1, \dots, k-1 < p,$$

and consider the curve  $w: (-\varepsilon, \varepsilon) \rightarrow G_{p,q}$  defined by

$$(11) \quad w_k^i = v_k^i + tv_k^i, i \neq \sigma_k, \quad w_k^{\sigma_k} = v_k^{\sigma_k}, \quad w_\beta = v_\beta, \beta \neq k.$$

From (10), (11) and (7) we infer that the components  $v^\rho$  where  $\rho$  is not of the form  $\rho = (\sigma_1, \dots, \sigma_{k-1}, \rho_k, \dots, \rho_p)$  are zero, and that the derivatives  $w_0^{p'} = \left(\frac{dw^p}{dt}\right)_{t=0}$  where  $\rho_k = \sigma_k$  are also zero. Thus  $\mathfrak{N}_1$ , now denoted by  $\mathfrak{N}_k$ , becomes

$$\begin{aligned}
\mathfrak{N}_k &= \left( \sum_{\rho_k > \sigma_k, \bar{\rho}} c_{\sigma_1 \dots \sigma_k - 1 \rho_k \bar{\rho}} |v^{\sigma_1 \dots \sigma_k - 1 \rho_k \bar{\rho}}|^2 \right) \left( \sum_{\tau_k > \sigma_k, \bar{\tau}} |v^{\sigma_1 \dots \sigma_k - 1 \tau_k \bar{\tau}}|^2 + \sum_{\bar{\tau}} |v^{\sigma_1 \dots \sigma_k \bar{\tau}}|^2 \right) \\
&\quad - \left( \sum_{\tau_k > \sigma_k, \bar{\tau}} |v^{\sigma_1 \dots \sigma_k - 1 \tau_k \bar{\tau}}|^2 \right) \left( \sum_{\rho_k > \sigma_k, \bar{\rho}} c_{\sigma_1 \dots \sigma_k - 1 \rho_k \bar{\rho}} |v^{\sigma_1 \dots \sigma_k - 1 \rho_k \bar{\rho}}|^2 \right. \\
&\quad \quad \quad \left. + \sum_{\bar{\tau}} c_{\sigma_1 \dots \sigma_k \bar{\tau}} |v^{\sigma_1 \dots \sigma_k \bar{\tau}}|^2 \right) \\
&= \sum_{\rho_k > \sigma_k, \bar{\rho}, \bar{\tau}} (c_{\sigma_1 \dots \sigma_k - 1 \rho_k \bar{\rho}} - c_{\sigma_1 \dots \sigma_k \bar{\tau}}) |v^{\sigma_1 \dots \sigma_k - 1 \rho_k \bar{\rho}}|^2 |v^{\sigma_1 \dots \sigma_k \bar{\tau}}|^2,
\end{aligned}$$

where  $\bar{\rho}$  denotes  $\rho_{k+1} \dots \rho_p$  for abbreviation. As above  $\mathfrak{N}_k = 0$  implies  $v_k^i = 0$ ,  $i > \sigma_k$ . Hence the only critical points  $F$  are the points  $\pi^{-1}(A_i)$ .

### Bibliography

- [1] J. Milnor, *Morse theory*, Ann. of Math. Studies, No. 51, Princeton University Press, Princeton, 1963.  
[2] W. T. Wu, *On critical sections of convex bodies*, Sci. Sinica 14 (1965) 1721-1728.

INSTITUTE OF MATHEMATICS  
BUCAREST, RUMANIA

