

MANIFOLDS OF NEGATIVE CURVATURE

M. GROMOV

1. Statement of results

1.1. For a Riemannian manifold V we denote by $c^+(V)$ and $c^-(V)$ respectively the upper and the lower bounds of the sectional curvature, by $\text{vol}(V)$ the volume, and by $d(V)$ the diameter.

1.2. Let V be an n -dimensional closed Riemannian manifold of negative curvature and $c^-(V) \geq -1$. If $n \geq 8$, then $\text{vol}(V) \geq C(1 + d(V))$, where the constant $C > 0$ depends only on n .

Remark. This inequality is exact: For each n there exists an infinite sequence V_i with $d(V_i) \rightarrow \infty$, $i \rightarrow \infty$, and with uniformly bounded ratio $\text{vol}(V_i)/d(V_i)$.

Proof. Take a manifold V of constant negative curvature with infinite group $H_1(V)$ (see [8]) and a sequence of its finite cyclic coverings.

For $n = 4, 5, 6, 7$ we shall prove here the following weaker result: $\text{vol}(V) \geq C(1 + d^{1/3}(V))$. Notice that arguments from § 4 show that for $n \geq 4$ an n -dimensional manifold V with $-\varepsilon \geq c^+(V) \geq c^-(V) \geq -1$, $\varepsilon > 0$, satisfies: $\text{vol}(V) \geq C(1 + d(V))$ where C depends on n and ε .

1.3. Theorem 1.2 sharpens the Margulis-Heintze theorem (see [6], [4]) stating the inequality $\text{vol}(V) \geq C = C_n$. In this paper we prove the following generalization.

1.3A. Let X be a complete simply connected manifold of negative curvature with $c^-(X) > -1$. Let Γ be a discrete group (possibly with torsion) of isometries of V . Then $\text{vol}(X/\Gamma) \geq C$, where $C > 0$ depends only on $\dim(X)$.

This fact is still true for manifolds of nonpositive curvature with $c^-(X) \geq -1$ and negative Ricci curvature (see [5]). In the homogeneous case this is the Kazhdan-Margulis theorem (see [9]).

The finiteness theorems

1.4. Combining § 1.2 with Cheeger's results (see [1], [4]) we immediately conclude:

For given $n \neq 3$ and $C > 0$ there exist only finitely many pairwise non-diffeomorphic closed n -dimensional manifolds V with $0 > c^+(V) \geq c^-(V) > -1$ and $\text{vol}(V) \leq C$.

1.5. Counter-example for $n = 3$. There exists an infinite sequence of 3-di-

mensional manifolds with uniformly bounded negative curvature and uniformly bounded volume but pairwise not isomorphic one-dimensional homology groups (although with uniformly bounded Betti numbers).

1.6. The homotopy theoretic version of Theorem 1.4 was announced in [7] by Margulis for all n . Although Margulis's statement is incorrect, his geometrical ideas are extremely fruitful and widely used in this paper.

E. Heintze proved in [6] the homotopy type finiteness theorem with diameter instead of volume. In fact the stronger result is true: For given $n = 1, 2, \dots$ and $C > 0$ there exist only finitely many pairwise non-diffeomorphic closed n -dimensional manifolds V with nonpositive curvature and with $c^-(V) > -1$ and $d(V) \leq C$. (For the proof see [5]).

Without assumption $c^+(V) \leq 0$ only the Betti numbers of V can be estimated by curvature and diameter (see [3]).

Pinching

1.7. Another standard application of § 1.2 is the following:

For given n and $C > 0$, there exists an $\varepsilon > 0$ such that under one of the following two conditions a closed n -dimensional manifold V admits a metric of constant negative curvature:

1. $\text{vol}(V) \leq C$, $n \neq 3$, $-1 \geq c^+(V) \geq c^-(V) \geq -1 - \varepsilon$, (for $n = 3$ it is unknown).
2. $|\chi(V)| \leq C$, n is even, $-1 \geq c^+(V) \geq c^-(V) \geq -1 - \varepsilon$, where χ is the Euler characteristic. (Compare with [4]).

Noncompact manifolds

1.8. Let V be a complete Riemannian manifold of negative curvature. If $c^-(V) > -\infty$ and $\text{vol}(V) < \infty$, then V has finite topological type, i.e., V is diffeomorphic to the interior of a compact manifold with boundary.

1.9. D. Kazhdan informed me recently that Margulis proved this fact for manifolds of strictly negative curvature, i.e., with $c^+(V) < 0$. In fact, Theorem 1.8 is still true for real analytic Riemannian manifolds of nonpositive curvature (see [5]), but is not so for C^∞ -manifolds (see § 5.1). For the homogeneous case see [9].

2. Groups of isometries

2.1 For an isometry $\gamma: X \rightarrow X$ we denote by $\delta_\gamma = \delta_\gamma(x)$, $x \in X$ the displacement $\text{dist}(x, \gamma x)$, and for a group Γ of isometries of X we set $\delta_\Gamma = \delta_\Gamma(x) = \min_{\substack{\gamma \in \Gamma \\ \gamma \neq e}} \delta_\gamma$, where $e \in \Gamma$ is the identity element.

An isometry γ is said to be semisimple if the function δ_γ assumes its minimum on X . If $\min_{x \in X} \delta_\gamma(x) = 0$, then a semisimple isometry is said to be elliptic and hyperbolic otherwise.

2.2. Let X be a complete simply connected manifold of negative curvature. Then for an isometry γ the functions δ_γ and δ_γ^2 are geodesically convex, and δ_γ^2 is strictly convex outside of the set where δ_γ^2 assumes its minimum. (See [2]).

2.3. If X is as above and γ is hyperbolic, then there exists a unique geodesic λ invariant under γ , and δ_γ assumes its minimum on λ . This is obvious and well known.

2.4. We say that a group Γ is almost nilpotent if it possesses a nilpotent subgroup of finite index.

2.5. Let X be as in § 2.2, and Γ an almost nilpotent group of isometries without elliptic elements. Let $\gamma \in \Gamma$ be an hyperbolic isometry, and λ a geodesic invariant under γ . Then λ is invariant under Γ , and Γ is an infinite cyclic group.

Proof. This follows immediately from § 2.3.

2.6. Let X be a complete simply connected manifold of negative curvature, and Γ an almost nilpotent group of isometries. Then there exists a smooth non-negative convex function $\varphi: X \rightarrow \mathbf{R}$ which is strictly convex at any point $x \in X$ where there is no (non-identical) element from Γ whose displacement assumes its minimum.

Proof. Take a nilpotent subgroup $N \subset \Gamma$ of finite index and any (non-identity) element γ from its center. There are only finitely many isometries $\gamma_1, \dots, \gamma_k$ conjugate to γ (in Γ). Take $\varphi = \sum_{i=1}^k \delta_{\gamma_i}^2$.

2.7. Corollary. *If X and Γ are as above, then the function δ_Γ does not assume its maximum; if Γ has no semisimple elements, then δ_Γ has no critical points (in the sense to be explained below).*

2.8. Generally our function $f(x) = \delta_\Gamma(x)$ is not smooth, but near each point $x \in X$ it can be represented as the minimum of smooth functions f_1, \dots, f_k . A point x is said to be noncritical if there exist a tangent vector t at x such that $\langle t, df_i \rangle > 0, i = 1, \dots, k$, and x is said to be critical otherwise.

3. The groups generated by small isometries

3.1. For a group Γ , isometrically acting on X , we denote by $\Gamma_\epsilon(v), \epsilon > 0, v \in V$, the subgroup generated by all $\gamma \in \Gamma$ with $\delta_\gamma(v) \leq \epsilon$.

3.2. The Margulis lemma. Let V be a complete Riemannian manifold without closed geodesics of length less than 1 and with $1 \geq c^+(X) \geq c^-(X) \geq -1$. Let Γ be a discrete group of isometries of X . Then there exists a number $\epsilon = \epsilon_n > 0$ depending only on $n = \dim X$ such that for any point $x \in X$ the group $\Gamma_\epsilon(v)$ is almost nilpotent.

For the proof and discussion see [4]. Notice that in [4] this lemma is presented in a different form, but the proof given there serves our present needs as well.

3.3. Proof of Theorem 1.3A. If $\text{vol}(X/\Gamma) < \infty$, then the function δ_Γ assumes its maximum, say, at a point $x \in X$. If $\delta_\Gamma(x) \geq \epsilon = \epsilon_n$, where ϵ_n is as above, the proof is finished. If $\delta_\Gamma(x) < \epsilon$, then the group $\Gamma_\epsilon(x)$ is almost nilpotent.

tent, and the functions δ_r and δ_{r_ε} are equal in a neighborhood of x ; but this contradicts § 2.7.

3.4. Let X be a complete simply connected n -dimensional manifold of negative curvature with $c^-(X) \geq -1$, and Γ a discrete group of isometries without elliptic elements. Let, $\gamma_1, \dots, \gamma_i, \dots \in \Gamma$ be hyperbolic isometries, and let $\Gamma_1, \dots, \Gamma_i, \dots \subset \Gamma$ be (uniquely defined) maximal cyclic subgroups containing, $\gamma_1, \dots, \gamma_i, \dots$ correspondingly. Denote the sets $(\delta_{\Gamma_i})^{-1}[0, \varepsilon]$ by $A_i \subset X$.

If the groups Γ_i are pairwise not conjugate in Γ , and the number ε is chosen equal to $\varepsilon = \varepsilon_n$ from § 3.2, then for $i \neq j$ and any $\gamma \in \Gamma$, the intersection $A_i \cap \gamma A_j$ is empty; if the intersection $A_i \cap \gamma A_i$ is not empty, then $\gamma \in \Gamma_i$.

Proof. Take $x \in A_i$. If $\gamma(x) \in A_j, j \neq i$, then the group $\Gamma_i(x)$ can not be cyclic because there are $\gamma' \in \Gamma_i$ with $\delta_{\gamma'}(v) \leq \varepsilon$ and $\gamma'' \in \Gamma_j$ with $\delta_{\gamma''}(\gamma(v)) \leq \varepsilon$ and so $\gamma', \gamma^{-1}\gamma''\gamma \in \Gamma_i(x)$. On the other hand, it follows from § 3.2 and § 2.5 that $\Gamma_i(x)$ is infinite cyclic; so the contradiction proves the first statement and an analogous argument proves the second.

3.5. Corollary. Let X, Γ and Γ_i be as above. If $\text{vol}(X/\Gamma) < \infty$ and $\min_{x \in X} \delta_{\Gamma_i} \leq \frac{1}{2}\varepsilon = \frac{1}{2}\varepsilon_n, i = 1, 2, \dots$ (ε_n is again from § 3.2), then the number of the subgroups Γ_i is finite.

Proof. The volumes of the sets $B_i = A_i/\Gamma$ are bounded away from zero, the projections $B_i \rightarrow X/\Gamma$ are, according to § 3.4, injective and their images do not intersect; therefore the number of B_i and Γ_i is finite.

3.6. Proof of Theorem 1.8. Consider the universal covering $p: X \rightarrow V$ with the group $\Gamma = \pi_1(V)$ acting on X . From § 3.5 it follows that there exists a positive number ε' such that for any $x \in X$ the group $\Gamma_i(x)$ has no hyperbolic elements, and applying § 3.2 and § 2.7 we conclude that outside of the set $X_0 = (\delta_r)^{-1}[0, \varepsilon'] \subset X, \varepsilon' = \min(\varepsilon', \varepsilon)$ and $\varepsilon = \varepsilon_n$ from § 3.2 it follows that the function δ_r has no critical points. This function is Γ -invariant and so defines a positive function f on V without critical points outside of the set $f^{-1}[0, \varepsilon'] \subset V, \varepsilon' > 0$. Since $\text{vol}(V) < \infty$ we have $f(v) \rightarrow 0$ as $v \rightarrow \infty$, and the application of the Morse theory finishes the proof. (The function $f(v)$ is not smooth, but the Morse theory is obviously applicable for the functions described in § 2.8.)

4. The volume of the tube

4.1. Let λ be a geodesic segment of length l in a manifold X , and let $\theta \in [0, l]$ be the natural parameter in λ . Let $J = J(\theta)$ be a Jacobi field normal to λ with $\langle J(0), J'(0) \rangle = 0$. Set $f(\theta) = \|J(\theta)\|$ and $g(\theta) = \|J'(\theta)\|$. Notice that $f'(\theta) \leq g(\theta)$.

If $0 \geq c^+(X) \geq c^-(X) \geq -1, f(0) \leq 1, l \geq 1$, then $f(l) \geq f(0) + Cl(\min(g(0), g^3(0)))$, where $C > 0$ depends only on $\dim X$.

Proof. The curvature is nonpositive, so $f' \geq 0, f'' \geq 0$ and $(f^2)'' \geq 2g^2$.

Curvature is bounded and so $g' \leq Kf$, where K is the norm of the curvature operator. Using the last inequality we have

$$\begin{aligned}
 |g(\theta) - g(0)| &\leq K \int_0^\theta f(\tau) d\tau \leq K\theta f(\theta) \\
 &= K\theta \left(f(0) + \int_0^\theta f'(\tau) d\tau \right) \leq K\theta(f(0) + \theta f'(\theta)) \\
 &\leq K\theta(f(0) + \theta g(\theta)),
 \end{aligned}$$

and for $\theta \leq 1$ we have $g(\theta) \geq \frac{g(0)}{1 + K} - \theta$. Integrating the inequality $(f^2)'' \geq 2g^2$ we obtain for $\theta \leq \min\left(1, l_0 = \frac{g(0)}{2 + 2K}\right)$: $f^2(\theta) \geq f^2(0) + \frac{\theta^2 g^2(0)}{8(1 + K)^2}$ and using the convexity of f we have $f(l) \geq f(0) + (l/\theta)(f(\theta) - f(0))$. Combining the last two inequalities and substituting θ by $\min(1, l_0)$ we get the needed estimate.

4.2. Let X be a complete simply connected manifold of negative curvature with $c^-(X) \geq -1$. Consider two points $x_1, x_2 \in X$ with $\text{dist}(x_1, x_2) = m$ and the geodesic μ joining x_1 and x_2 . Let t_1 and t_2 be unit tangent vectors at x_1 and x_2 normal to μ , and let α be the angle between t_1 and the vector t_2' at x_1 , which is parallel to t_2 along μ . Consider two geodesic segments λ_1, λ_2 of lengths l_1, l_2 starting from x_1, x_2 in the directions t_1, t_2 . Denote by y_1, y_2 the second ends of these segments.

If $l_1 \geq 1$, then $\text{dist}(y_1, y_2) \geq m + Cl_1 \frac{\alpha^3}{1 + m^2}$, where $C \geq 0$ depends only on $\dim X$.

This follows from the previous lemma by arguments of the standard comparison type (see [2]).

4.3. Let g be an isometry of the standard unit sphere S^{n-2} . Then it is obvious that for every $N = 1, 2, \dots$ there exist points $s_1, \dots, s_N \in S^{n-2}$ with the following property: for any $k = \dots, -1, 0, 1, \dots$, $\text{dist}(s_i, g^k s_j) \geq CN^{-1/d}$, where $i \neq j, i, j = 1, \dots, N, d = n - 2 - \text{rank}(SO(n - 1)) = n - 2 - \text{ent}\left(\frac{n - 1}{2}\right)$, and $C > 0$ depends only on n .

4.4. Let X be a manifold as in § 4.2, $\mu \subset X$ a geodesic, and let $\gamma: X \rightarrow X$ be a hyperbolic isometry keeping μ invariant. Denote by Γ the group generated by γ , and denote by $A_\epsilon \subset X, 0 \leq \epsilon \leq 1$ the set $\delta_{\Gamma^{-1}}[0, \epsilon]$.

Let $n = \dim X$, and let ∂A_ϵ be the boundary of A_ϵ . If $\text{dist}(\mu, \partial A_\epsilon) \geq 2\epsilon$, and there is a point $y \in A_\epsilon$ with $l = \text{dist}(y, \mu) \geq 3$, then $\text{vol}(A_{2\epsilon}/\Gamma) \geq Cl^{P_n} \epsilon^n$, where $C > 0$ depends on $n, P_n = 1$ for $n \geq 8, P_n = \frac{2}{3}$ for $n = 6, 7$, and $P_n = \frac{1}{3}$ for $n = 4, 5$.

Proof. Take the projection $x \in \mu$ from y to μ , and denote by S^{n-2} the sphere of all unit tangent vectors at x normal to μ . The holonomy along μ together with γ defines the isometry g of S^{n-2} . Take points $s_1, \dots, s_N \in S^{n-2}$ as in § 4.3 with $N = \text{ent}(l^{P_n})$ and the geodesic rays $\lambda_1, \dots, \lambda_N$ starting at x in directions

s_1, \dots, s_N . Take points $y_i \in \lambda_i \cap \partial A_\varepsilon$, and suppose without loss of generality that $y = y_1$. If for all i $\text{dist}(y_i, \mu) \geq \frac{1}{2}l$, then applying § 4.2 and § 4.3 we have $\text{dist}(y_i, \gamma^k y_j) \geq \beta$, $i \neq j, k = \dots, -1, 0, 1, \dots$, and $\beta > 0$ depends only on n . Thus the lemma is proved.

If there is y_i with $\text{dist}(y_i, \mu) \leq \frac{1}{2}l$ and $n \geq 3$, then obviously $\text{vol}(A_{2i}/\Gamma) \geq Cl\varepsilon^n$ which suits us as well.

4.5. Proof of Theorem 1.2. Consider the universal covering $p: X \rightarrow V$, and take isometries $\gamma_1, \dots, \gamma_i, \dots$ representing the conjugacy classes of isometries corresponding to all simple closed geodesics in X of length $\leq \frac{1}{4}\varepsilon_n$, where ε_n is from § 3.2. Take the sets $A_i = \delta_{\Gamma_i}^{-1}[0, \varepsilon_n]$, where Γ_i is the group generated by γ_i . According to § 3.4 the projections $A_i/\Gamma_i \rightarrow V$ are injective, and their images $T_i \subset V$ do not intersect (compare with § 3.5).

Take now two points $v_1, v_2 \in V$ with $\text{dist}(v_1, v_2) = d(V)$, and join them by the shortest geodesic segment ν . Consider the union T of all T_i intersecting ν and the ε_n -neighborhood U of ν . It follows from § 4.4 that the set $T \cup U$ provides enough volume to finish the proof.

5. Examples

C^∞ -manifolds of nonpositive curvature

5.1. Start with a compact C^∞ -surface $V_i, i = 1, 2, \dots$ with the following properties:

- a. V_i is diffeomorphic to the torus with two holes.
- b. $\text{vol}(V_i) \leq 100$.
- c. $0 \geq c^+(V_i) \geq c^-(V_i) \geq -1$.
- d. Boundary of V_i consists of two geodesics S'_i and S''_{i+2} of lengths $1/2^i$ and $1/2^{i+2}$.
- e. Near the boundary each manifold V_i is flat (its curvature is zero).

Denote the product $S_{i+1} \times V_i$ by W_i , where S_{i+1} denotes the circle of length $1/2^{i+1}$. Boundary of W_i consists of two tori $B'_i = S'_i \times S_{i+1}$ and $B''_i = S_{i+1} \times S''_{i+2}$. Each manifold B''_i is canonically isometric to B'_{i+1} , and by identifying all pairs of isometrical tori we obtain the manifold W' with boundary B'_i . The double W of W' is complete C^∞ -manifold with finite volume, bounded nonpositive curvature but infinitely generated group $H_1(W)$.

5.2. The previous construction provides many other examples of C^∞ -manifolds of nonpositive curvature but without real analytic metrics of nonpositive curvature. The simplest one is the boundary of $V \times V$, where V is a compact surface of positive genus with one hole and with the same geometry at the boundary as manifolds V_i from § 5.1.

Three-dimensional manifolds

5.3. Horns. We denote by \bigwedge^3 the 3-dimensional hyperbolic space with cur-

vature -1 . Consider a horosphere $S \subset \wedge^3$ and the (convex) horoball B bounded by S .

A horn is, by definition, the quotient $H = B/\Gamma$ where Γ is a discrete group of isometries isomorphic to $\mathbb{Z} \oplus \mathbb{Z}$. Boundary of H is the flat torus S/Γ .

5.4. Tubes. Consider a geodesic $\mu \subset \wedge^3$ and the set $A(l) \subset \wedge^3$ consisting of all points $a \in \wedge^3$ with $\text{dist}(a, \mu) \leq l$. A tube is, by definition, the quotient $B(l) = A(l)/\Gamma$ where Γ is an infinite cyclic group generated by a hyperbolic isometry γ keeping μ invariant. The boundary $\partial B(l)$ of the tube $B(l)$ is isometric to a flat torus, and $\text{vol}(B(l)) \leq 100 \text{vol}(\partial B(l))$.

5.5. Consider a tube $B(l)$ and a horn H , and let $I: \partial H \rightarrow \partial B(l)$ be an isometry. Using the normal geodesic coordinates we can canonically extend I to a map $J: U_\varepsilon \rightarrow B(l)$ where $U_\varepsilon \subset H$, $\varepsilon < l$, is the ε -neighborhood of ∂H . Denote by $g(J)$ the metric in U_ε induced by J , and denote by g_0 the original metric in U_ε .

It is obvious that if ε is kept fixed and $l \rightarrow \infty$, then the metric $g(J)$ C^∞ -converges to g_0 .

5.6. Let T be a flat torus, and let $h \in H_1(T)$ be an indivisible element. Then there exist a tube $B(l)$ and an isometrical imbedding $I: T \rightarrow B(l)$ which maps T isometrically onto the boundary of $B(l)$, and the kernel of the induced homomorphism $I_*: H_1(T) \rightarrow H_1(B(l))$ is generated by h . If $h_j \in H_1(T)$ is the sequence of indivisible elements and $h_j \rightarrow \infty$, then for the corresponding tubes $B(l_j)$ we also have $l_j \rightarrow \infty$.

Proof. Every tube is determined by three parameters: l and two parameters of the isometry γ (shift and rotation), and choosing these parameters in an obvious fashion we construct the needed tubes.

5.7. Take now a complete noncompact orientable 3-dimensional manifold V of curvature -1 . One can find in V a compact 3-dimensional submanifold V_0 with the boundary consisting of k flat tori T_1, \dots, T_k and with the complement $V \setminus \text{Int } V_0$ consisting of k horns bounded by these tori. According to § 5.6 we can find for every T_i a tube with the boundary isometric to T_i , and attaching these tubes to V_0 we obtain a closed manifold with corners at T_i . Moreover Lemma 5.6 shows that by this construction we can obtain infinitely many manifolds with different one-dimensional homologies. On the other hand, using § 5.5 we can smooth the corners providing our closed manifolds with metrics of uniformly bounded negative curvature. That gives the sequence V_i of the manifolds promised in § 1.5. In fact, the above construction gives the manifolds V_i with $-1 \geq c^+(V_i) \geq c^-(V_i) \geq -1 - \varepsilon_i$, where $\varepsilon_i \rightarrow 0$, as $i \rightarrow \infty$.

Bibliography

- [1] J. Cheeger, *Finiteness theorems for Riemannian manifolds*, Amer. J. Math. **92** (1970) 61–74.
- [2] J. Cheeger & D. G. Ebin, *Comparison theorems in Riemannian geometry*, North-Holland, Amsterdam, 1975.
- [3] M. Gromov, *Topology of Riemannian manifolds with small curvature and diame-*

- ter, Notices, Amer. Math. Soc. **22** (1975) A-592.
- [4] ———, *Almost flat manifolds*, J. Differential Geometry **13** (1978).
- [5] ———, *Manifolds of nonpositive curvature*, to appear.
- [6] E. Heintze, *Mannigfaltigkeiten negativer Krümmung*, Preprint, Bonn, 1976.
- [7] R. A. Margulis, *On connections between metric and topological properties of manifolds of nonpositive curvature*, Proc. the Sixth Topological Conf., Tbilisi, USSR, 1972, p. 83 (in Russian).
- [8] J. Millson, *On the first Betti number of a compact constant negatively curved manifolds*, Preprint.
- [9] M. S. Raghunathan, *Discrete subgroups of Lie groups*, Ergebnisse der Math. Vol. 68, Springer, Berlin, 1972.

STATE UNIVERSITY OF NEW YORK, STONY BROOK