

## ERRATUM TO “COMPARING RIEMANNIAN FOLIATIONS WITH TRANSVERSALLY SYMMETRIC FOLIATIONS”

FRANZ W. KAMBER, ERNST A. RUH & PHILIPPE TONDEUR

As has been pointed out by Hebda [1], the main theorem of our paper [2] is not correct as stated. In the theorem on p. 461 the assumption “ $\pi_1(K)$  finite, i.e.,  $K$  semisimple” should be added. The averaging process for the developing map  $\Phi: \tilde{P} \rightarrow \tilde{G}$  referred to in the second paragraph on p. 462 works under this assumption. All the arguments are valid for the foliation  $\mathcal{F}$  lifted to the ( $K$ -reduction of the) normal frame bundle  $P$  of  $\mathcal{F}$ , for which we produce a Cartan connection with vanishing curvature. This implies that this lifted foliation is a Lie foliation. For this we refer to [3].

### References

- [1] J. Hebda, *An example relevant to curvature pinching theorems for Riemannian foliations* (to appear).
- [2] F. Kamber, E. Ruh & P. Tondeur, *Comparing Riemannian foliations with transversally symmetric foliations*, *J. Differential Geometry* 27 (1988) 461–475.
- [3] E. Ruh & P. Tondeur, *Almost Lie foliations and the heat equation method*, Proc. VI Internat. Colloq. on Differential Geometry, Santiago de Compostela (Spain) 1988, Cursos y Congresos 61, Univ. Santiago de Compostela, 1989, 239–246.

UNIVERSITY OF ILLINOIS  
OHIO STATE UNIVERSITY  
UNIVERSITY OF ILLINOIS

