

MINIMAL SURFACES FOR HITCHIN REPRESENTATIONS

SONG DAI* & QIONGLING LI†

Abstract

Given a reductive representation $\rho : \pi_1(S) \rightarrow G$, there exists a ρ -equivariant harmonic map f from the universal cover of a fixed Riemann surface Σ to the symmetric space G/K associated to G . If the Hopf differential of f vanishes, the harmonic map is then minimal. In this paper, we investigate the properties of immersed minimal surfaces inside symmetric space associated to a subloci of Hitchin component: the q_n and q_{n-1} cases. First, we show that the pullback metric of the minimal surface dominates a constant multiple of the hyperbolic metric in the same conformal class and has a strong rigidity property. Secondly, we show that the immersed minimal surface is never tangential to any flat inside the symmetric space. As a direct corollary, the pullback metric of the minimal surface is always strictly negatively curved. In the end, we find a fully decoupled system to approximate the coupled Hitchin system.

1. Introduction

For a closed, connected, oriented surface S of genus $g \geq 2$ and a reductive Lie group G , consider the representation variety $Rep(\pi_1(S), G) = Hom(\pi_1(S), G)/G$. For $G = PSL(2, \mathbb{R})$, there are two connected components of $Rep(\pi_1(S), PSL(2, \mathbb{R}))$ are identified with Teichmüller space; and the representations in these components are called Fuchsian. For a general real split Lie group, using the unique irreducible representation $PSL(2, \mathbb{R}) \rightarrow G$, we can single out a component of $Rep(\pi_1(S), G)$, the connected component containing representations factors through Fuchsian representations, called Hitchin component for G . In particular, we denote the Hitchin component for $PSL(n, \mathbb{R})$ as Hit_n .

Fix a Riemann surface structure Σ on S . By the work of Donaldson [5] and Corlette [4], given a reductive representation $\rho : \pi_1(S) \rightarrow G$,

*The author is supported by NSFC grant No. 11601369.

†Corresponding author, supported in part by the center of excellence grant ‘Center for Quantum Geometry of Moduli Spaces’ from the Danish National Research Foundation (DNRF95).

Received June 3, 2016.

there exists a ρ -equivariant harmonic map f from the universal cover of Σ to the symmetric space G/K associated to G , where K is the maximal compact subgroup of G . If the representation ρ is a Hitchin representation, Sanders [20] showed that the corresponding equivariant harmonic map is an immersion. In particular, if the Hopf differential of the harmonic map f vanishes, then the harmonic map is conformal and, hence, is a minimal immersion.

Understanding such equivariant minimal immersions $f : \tilde{\Sigma} \rightarrow G/K$ for Hitchin representations is the main goal of this paper.

To achieve this goal, we need to use the tool of Higgs bundles. By the results of Hitchin [9] and Simpson [21], given a polystable G -Higgs bundle (E, ϕ) , there exists a Hermitian metric h (compatible with G -structure) satisfying Hitchin equation, giving rise to a flat connection $D = \nabla^h + \phi + \phi^{*h}$, the corresponding holonomy $\rho : \pi_1(S) \rightarrow G$, and a ρ -equivariant harmonic map $f : \tilde{\Sigma} \rightarrow G/K$.

Hitchin [10] gives an explicit description for the Hitchin component for $SL(n, \mathbb{R})$ in terms of Higgs bundles. Explicitly, the bundle of the Higgs bundles is $E = K^{\frac{n-1}{2}} \oplus K^{\frac{n-3}{2}} \oplus \cdots \oplus K^{\frac{1-n}{2}}$ and the Higgs field is explicitly parametrized by the holomorphic differentials $(q_2, q_3, \cdots, q_n) \in \bigoplus_{i=2}^n H^0(K^i)$. In particular, q_2 is the Hopf differential of the corresponding harmonic map.

Replacing the quadratic differential by varying the Riemann surface choice, Labourie [13] considered the Hitchin map from the total space of vector bundle over Teichmüller space with fiber at Σ as $(0, q_3, \cdots, q_n)$ to Hit_n and showed that this map is surjective. Therefore, understanding all minimal surfaces arising from the Hitchin representations will eventually give properties of the representations, not depending on the choice of Riemann surface. In the same paper, Labourie conjectured that the Hitchin map is also injective. When $n = 3$, Labourie [12] and Loftin [16] independently proved the conjecture using affine geometry. Using different methods, Labourie [14] then proved the conjecture for Hitchin representations into all rank 2 real split Lie groups: $PSL(3, \mathbb{R})$, $PSp(4, \mathbb{R})$, G_2 , and $PSL(2, \mathbb{R}) \times PSL(2, \mathbb{R})$. Equivalently, given any Hitchin representation ρ into a rank 2 real split Lie group G , there is a unique minimal surface inside G/K that is ρ -equivariant.

We restrict to consider the Higgs fields parametrized by $(0, \cdots, 0, q_n)$ and $(0, \cdots, q_{n-1}, 0)$:

$$\left(\begin{array}{cccc} 0 & & & q_n \\ 1 & 0 & & \\ & 1 & 0 & \\ & & \ddots & \\ & & & 0 \\ & & & 1 & 0 \end{array} \right), \quad \left(\begin{array}{cccc} 0 & & q_{n-1} & 0 \\ 1 & 0 & & q_{n-1} \\ & 1 & 0 & \\ & & \ddots & \\ & & & 0 \\ & & & 1 & 0 \end{array} \right).$$

In particular, $q_n = 0$ or $q_{n-1} = 0$ gives the base Fuchsian point. From the surjectivity of the Hitchin map, for any Hitchin representation, we may find a complex structure on S , such that $q_2 = 0$ in the corresponding Higgs bundle. Hence, if we vary the choice of Riemann surface, these two families of Higgs bundles give the whole Hitchin component for $PSL(3, \mathbb{R})$, $PSp(4, \mathbb{R})$, and G_2 .

If $n = 2$, the Higgs bundles of the q_2 case parametrize the whole Teichmüller component. If $n = 3$, the q_{n-1} case gives the embedding of Teichmüller component inside Hitchin component for $PSL(3, \mathbb{R})$. But in these two cases, the Hopf differential of the harmonic map does not vanish and, hence, the harmonic map is not conformal. However, we remark that in these two cases, almost all of our results can be applied. For simplicity of language, from now on, whenever referring to the q_n, q_{n-1} cases, we don't include the case that q_n (or q_{n-1}) is q_2 .

Such Higgs bundles of the q_n case were initially studied by Baraglia [1], which he called cyclic Higgs bundle. Later Collier [3] considered other Higgs bundles under finite order automorphisms, in particular, q_{n-1} case (also see [2]). These families of Higgs bundles possess particular nice properties. In both cases, the Hermitian metric h solving Hitchin equations is diagonal, i.e., $h = (h_1, h_2, \dots, h_2^{-1}, h_1^{-1})$.

We are ready to explain our main results for the q_n, q_{n-1} cases. In general, these are some families in Hitchin component. But from the above argument, we remark that our results, in fact, hold for the Hitchin component for $PSL(3, \mathbb{R})$, $PSp(4, \mathbb{R})$, and G_2 .

• **Metric Domination.** We show that the pullback metric g_f of the minimal immersion f dominates the base Fuchsian metric g_{Fuchsian} .

Theorem 1.1. *For the q_n, q_{n-1} cases*

$$g_f = 2n \operatorname{tr}(\phi\phi^*) \geq g_{\text{Fuchsian}} = \frac{1}{6}(n^4 - n^2)g_0.$$

Moreover, if equality holds at one point, then $q_n = 0$, or $q_{n-1} = 0$, respectively, which implies it is base Fuchsian.

For the precise definitions above involving the q_n, q_{n-1} cases, see Section 3.

If we integrate the pullback metric, this is closely related to the Morse function considered in Hitchin [10], which plays an important role to determine the topology of the representation variety for $PSL(n, \mathbb{R})$. The Morse function on moduli space of polystable Higgs bundles is defined as:

$$f(E, \phi) = \int_{\Sigma} \operatorname{tr}(\phi\phi^*)\sqrt{-1}dz \wedge d\bar{z}.$$

Hitchin [10] showed that in the Hitchin component, the only minimum is the base Fuchsian point. That is, consider any Higgs bundle (E, ϕ)

in Hitchin component parametrized by (q_2, q_3, \dots, q_n) , then

$$f(E, \phi) \geq \text{base Fuchsian case} = \text{topological quantity.}$$

Equality holds if and only if it is base Fuchsian.

When $q_2 = 0$, the harmonic map is a minimal immersion and the Morse function is, in fact, the area of the minimal surface up to a constant. Then

$$\text{MinArea}(\rho) \geq \text{Area}(\rho_{\text{Fuchsian}}) = \text{topological quantity,}$$

where $\text{MinArea}(\rho)$ is the minimum of the area of the pullback metric going through all the ρ -equivariant immersion. Equality holds if and only if it is the base Fuchsian representation ρ_{Fuchsian} . Labourie [14] pointed out that this is also a corollary of the entropy result for Hitchin representations by Potrie–Sambarino [19]. For the Hitchin component for rank 2 Lie group, this is reproved by Labourie [14] which he called area rigidity formula.

In fact, this area rigidity formula by Labourie inspired us to be interested in this metric domination question. We conjecture that this domination property also holds for all minimal surfaces arising from Hitchin representations.

Conjecture 1.2. (*Metric Domination Conjecture*) *Consider Higgs bundle (E, ϕ) in Hitchin component parametrized by $(0, q_3, \dots, q_n)$. On the surface Σ , the pullback metric g_f of the minimal immersion f satisfies*

$$g_f \geq g_{\text{Fuchsian}} = \frac{1}{6}(n^4 - n^2)g_0.$$

If the equality holds at one point, then it holds at every point.

Remark 1.3. For $q_2 \neq 0$, in other words, the harmonic map f is not conformal. Instead of the pullback metric g_f , we consider the $(1, 1)$ part of g_f , $g_f^{(1,1)} = 2\text{tr}(\phi\phi^{*h})$ which is conformal to the base Fuchsian metric. From our proof of Theorem 1.1, the above domination theorem is also true in non-minimal surface case for the lower rank case. More precisely, for Hitchin representations in $PSL(2, \mathbb{R})$ parametrized by q_2 , we have

$$g_f^{(1,1)} \geq g_{\text{Fuchsian}} = 2g_0.$$

Moreover, if equality holds at one point, then $q_2 = 0$, that is, the base Fuchsian case.

The domination question above is comparing Hitchin representations with the base Fuchsian points. If instead, we consider the other non-Hitchin components, it won't make sense to compare them with Fuchsian points. At first, the harmonic map is not necessarily an immersion, hence, the pullback metric can never dominate a hyperbolic metric

pointwise. A more desired question is to ask whether the representations dominate some particular special representations which are similar to the status of Fuchsian points in Hitchin components.

It is interesting to compare the domination question here with another type of domination question in terms of length spectrum. By work of Deroin and Tholozan [6] and Guéritaud, Kassel and Wolff [8], for representations of $\pi_1(S)$ into $PSL(2, \mathbb{R})$, any Fuchsian representation dominates some non-Fuchsian representation and any non-Fuchsian representation can be dominated by some Fuchsian one. In a similar spirit, Lee and Zhang [15] conjectured for any Hitchin representation $\rho : \pi_1(S) \rightarrow PSL(n, R)$, there is a Fuchsian representation whose length spectrum strictly dominated by ρ . Tholozan [23] proved this conjecture for $n = 3$ case and mentioned that Labourie pointed out the conjecture of Lee and Zhang cannot hold anymore for $n \geq 4$. The contradiction comes from Hitchin representations in $PSp(2k, R)$ and $PSO(k, k + 1)$. Therefore, in the same paper, Tholozan made some modification of the conjecture by changing Fuchsian representation by representations into $SO(n, n + 1)$ or $Sp(2n, \mathbb{R})$. Our Conjecture 1.2 here is, in fact, weaker than the conjecture by Lee and Zhang since the length spectrum of the pullback back metric is larger than the length spectrum of the representation which is Lipschitz to the distance inside the symmetric space. So even though our conjecture is already true for Hitchin representations into rank 2 Lie groups: $PSL(3, \mathbb{R})$, $PSp(4, \mathbb{R})$, and G_2 , it does not imply the conjecture of Lee and Zhang.

• **Negative Curvature.** We describe how the immersed minimal surface sits inside the symmetric space by showing that the minimal immersion is never tangential to any flat inside the symmetric space G/K .

Theorem 1.4. *For the q_n, q_{n-1} cases, we have the following results.*
 (1) *The Hitchin equation never decouples: for every point on Σ ,*

$$F^{\nabla^h} \neq 0, \quad [\phi, \phi^{*h}] \neq 0.$$

(2) *The sectional curvature $K_{G/K}(\sigma)$ is strictly negative, where σ is the tangent space of the image of f . Geometrically, the minimal immersion is never tangential to any flat inside the symmetric space.*

(3) *On each line bundle $K^{\frac{n+1-2k}{2}}$, $1 \leq k \leq n$, the Chern form of h_k :*

$$\sqrt{-1}\Theta_{h_k} = \sqrt{-1}\partial\bar{\partial}\log h_k$$

is strictly positive if $n + 1 - 2k > 0$; zero if $n + 1 - 2k = 0$; strictly negative if $n + 1 - 2k < 0$.

Remark 1.5. The phenomenon in Part (1) is in contrast to the asymptotic behavior of Hitchin equation proved in Collier and Li [2]:

along the ray tq_n (or tq_{n-1}), the Hitchin equation decouples as $t \rightarrow \infty$:

$$F^{\nabla^h} \rightarrow 0, \quad [\phi, \phi^{*h}] \rightarrow 0.$$

This asymptotic behavior is generalized by Mochizuki [17] to a much more general family of Higgs bundles.

From Gauss equation, it is easy to see that the curvature of the pullback metric g_f is always non-positive. Then, moreover, as a corollary of Theorem 1.4, we obtain the curvature is strictly negative.

Corollary 1.6. *For the q_n, q_{n-1} cases, the sectional curvature of the immersed minimal surface is strictly negative.*

We conjecture that this phenomenon is true for all minimal immersions arising from Hitchin representations.

Conjecture 1.7. (*Negative Curvature Conjecture*) *For the Hitchin representation parametrized by $(0, q_3, \dots, q_n)$, the minimal immersion is never tangential to any flat inside the symmetric space. And as a corollary, the sectional curvature of immersed minimal surface is strictly negatively curved.*

We apply our result to estimate the entropy of Hitchin representations. Given ρ a Hitchin representation into G and select a point in the symmetric space $p \in G/K$. The volume entropy of ρ is defined as

$$h(\rho) := \limsup_{R \rightarrow \infty} \frac{\log(\#\{\gamma \in \pi_1(S) \mid d(p, \rho(\gamma)(p)) \leq R\})}{R},$$

where d is the distance in G/K . Lots of progress have been made on the volume entropy of Hitchin representations. Potrie and Sambarino [19] showed that for any Hitchin representation ρ , one has $h(\rho) \leq 1$ and the equality holds only if ρ is Fuchsian. Zhang [25, 26] constructed certain sequences of Hitchin representations along which $h(\rho) \rightarrow 0$. For $n = 3$, Nie [18] showed that the entropy of Hitchin representations parametrized by tq_3 goes to zero as $t \rightarrow \infty$. Sanders [20] showed that for a Hitchin representation, the curvature of the pullback metric g_f satisfies

$$\frac{1}{\text{Area}(g_f)} \int_{\Sigma} \sqrt{-K_{g_f}} dV_{g_f} \leq h(\rho).$$

And he used this inequality to show $h(\rho) > 0$. The entropy of Hitchin representation then satisfies:

$$h(\rho) \geq \min_{\Sigma} \{\sqrt{-K_{g_f}}\}.$$

By Corollary 1.6, the sectional curvature of immersed minimal surface $K_{g_f} < 0$ for the q_n, q_{n-1} cases, we obtain

Corollary 1.8. $\min_{\Sigma} \{\sqrt{-K_{g_f}}\}$ provides a positive lower bound for the volume entropy $h(\rho)$ of Hitchin representations for the q_n, q_{n-1} cases.

In fact, since $K_{g_f} \leq K_{G/K}$, we can also use $\min_{\Sigma} \{\sqrt{-K_{G/K}}\}$ as a weaker lower bound. The term $\min_{\Sigma} \{\sqrt{-K_{G/K}}\}$ is very interesting and involves more analytical terms. We hope to show a more quantitative estimate on this term in future work.

• **Coupled Hitchin System vs Decoupled System.** We investigate the coupled Hitchin system in the q_n, q_{n-1} cases. Coupled equations are generally hard to study. If there is some way to decouple the system, it will be substantially easier to solve the system. In this paper, we find a fully decoupled system such that the solutions of the decoupled system approximate the solutions of coupled Hitchin system of equations. The decoupled system are formed of single scalar equations as follows. For n even, set $n = 2m$; for n odd, set $n = 2m + 1$.

Theorem 1.9. *There exists a unique Hermitian metric u_k, v_k , respectively, on the holomorphic line bundle $K^{\frac{n+1-2k}{2}}$ satisfying, locally*

$$\Delta \log u_k + u_k^{-\frac{2}{n+1-2k}} - (u_k^2 |q_n|^2)^{\frac{1}{2k-1}} = 0, \quad 1 \leq k \leq m,$$

$$\Delta \log v_k + v_k^{-\frac{2}{n+1-2k}} - (v_k^2 |2q_{n-1}|^2)^{\frac{1}{2k-2}} = 0, \quad 2 \leq k \leq m.$$

And the following estimates hold,

$$\max\{|q_n|^{\frac{2\alpha_k}{n}}, (\alpha_k g_0)^{\alpha_k}\} \leq u_k^{-1} < ((\max_{\Sigma} |q_n|_{g_0}^{\frac{2}{n}} + \alpha_k) g_0)^{\alpha_k},$$

$$\max\{|2q_{n-1}|^{\frac{2\alpha_k}{n-1}}, (\alpha_k g_0)^{\alpha_k}\} \leq v_k^{-1} < ((\max_{\Sigma} |2q_{n-1}|_{g_0}^{\frac{2}{n-1}} + \alpha_k) g_0)^{\alpha_k},$$

where $\alpha_k = \frac{n+1-2k}{2}$ and g_0 is the Hermitian hyperbolic metric on K^{-1} .

Remark 1.10. Such equations are natural generalization of familiar vortex-like equations. In particular, in the case $n = 2, k = 1$, this is the harmonic equation from surface to surface with given Hopf differential q_2 . In the case $n = 3, k = 1$, this gives Wang's equation [24] and the solution is the Blaschke metric for the hyperbolic affine sphere. Dumas and Wolf [7] solved such equations for the case when n is general and $k = 1$ on the complex plane.

Remark 1.11. Surprisingly, there is a geometric interpretation for the above system of equations. Let $\sigma_k = u_k^{-\frac{2}{n+1-2k}}$ be a Hermitian metric on the Riemann surface Σ . The equation for σ_k is

$$\frac{n+1-2k}{2} K_{\sigma_k} = -1 + |q_n|_{\sigma_k}^{\frac{2}{2k-1}},$$

where K_{σ_k} is the curvature of σ_k . Roughly speaking, the curvature of σ_k differs from the curvature of hyperbolic metric by the norm of q_n .

We show that u_k (or v_k) is an upper approximate of h_k . Let u_k^t (or v_k^t) be the solution for tq_n (or tq_{n-1}) in Theorem 1.9.

Theorem 1.12. *For $m \geq 2$, suppose q_n (or q_{n-1}) is not zero. Then for each compact set $K \subset \Sigma$ away from zeros of q_n (or q_{n-1}), there is a positive constant $C = C(K)$ independent of t such that*

$$\begin{aligned} q_n \text{ case:} \quad & u_k^t(1 - C||tq_n||^{-1}) \leq h_k^t < u_k^t, 1 \leq k \leq m, \\ q_{n-1} \text{ case:} \quad & v_k^t(1 - C||tq_{n-1}||^{-1}) \leq h_k^t < v_k^t, 2 \leq k \leq m. \end{aligned}$$

Remark 1.13. To prove the left direction of the above theorem, we make use the asymptotic behavior of h_k established by Collier and Li [2]: for example, for the q_n case,

$$h_k = |q_n|^{-\frac{n+1-2k}{n}}(1 + O(||q_n||^{-1}))$$

away from the zeros of q_n . And our results, in fact, improve the estimate of Collier and Li from the other direction by the estimate of u_k, v_k in Theorem 1.9.

Structure of the article. The article is organized as follows. In Section 2, we recall some fundamental results about Higgs bundles. In particular, we recall the Donaldson–Uhlenbeck–Yau correspondence and the explicit relation between Higgs bundles and related harmonic maps for further calculation. We fix some notations at the end of Section 2. In Section 3, we set up the main object and describe the q_n case and q_{n-1} case. In Section 4, we show the metric domination Theorem 1.1. In Section 5, we describe the shape of minimal surface inside the symmetric space by proving Theorem 1.4. In Section 6, we find a decoupled system to bound the Hitchin coupled system and prove Theorem 1.9 and Theorem 1.12.

Acknowledgment. The authors wish to thank Mike Wolf, Andy Sanders, Ian McIntosh, Daniele Alessandrini and Brian Collier for helpful discussions and comments. The second author acknowledges support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network).

2. Preliminaries and notations

In this section, we recall some results in (principal) Higgs bundle and harmonic map. A good reference is the thesis of Baraglia [1], Section 2.1.

Let Σ be a closed Riemann surface with genus $g \geq 2$. Let $\pi_1 = \pi_1(\Sigma, p)$ be the fundamental group. Let G be a complex semisimple Lie group, K be a compact real form of G . Let $\mathfrak{g}, \mathfrak{k}$ be the corresponding Lie algebra. Then $\mathfrak{g} = \mathfrak{k} \otimes \mathbb{C}$ as a real Lie algebra. Let θ be the anti-linear involution of \mathfrak{g} induced from the conjugation of \mathbb{C} . Let $B_{\mathfrak{g}}$ be the

complex Killing form of \mathfrak{g} and $B_{\mathfrak{k}}$ be the real Killing form of \mathfrak{k} . Then $B_{\mathfrak{g}}$ is the complex linear extension of $B_{\mathfrak{k}}$. So without confusing, we just use B to denote Killing form of \mathfrak{g} or \mathfrak{k} . Then $H(X, Y) := -B(X, \theta Y)$ gives an Hermitian metric on \mathfrak{g} . Then we have an orthogonal decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{k}^{\perp}$ with respect to H , where $\mathfrak{k}^{\perp} = \sqrt{-1}\mathfrak{k}$. This decomposition also gives a direct sum as Ad_K module. And notice that $[\mathfrak{k}^{\perp}, \mathfrak{k}^{\perp}] \subseteq \mathfrak{k}$. For example, $G = SL(n, \mathbb{C})$, $K = SU(n)$, then $B(X, Y) = 2n \operatorname{tr}_{\mathbb{C}}(XY)$, where $X, Y \in \mathfrak{sl}(n, \mathbb{C})$.

We can establish a correspondence between the following two moduli spaces.

Betti Moduli Space: Equivalent classes $[\rho]$ of reductive representations, $\rho : \pi_1 \rightarrow G$. Here reductive means that the induced representation on \mathfrak{g} is a direct sum of irreducible representations. For two representations ρ_1, ρ_2 , they are equivalent if and only if there exists $g \in G$, such that $\rho_1 = g\rho_2g^{-1}$.

Higgs Moduli Space: Gauge equivalent classes of harmonic G -Higgs. More precisely, let P be a principal G bundle over Σ . Suppose we have a K principal bundle reduction $i : P_K \hookrightarrow P$, or equivalently, a section of $P \times_l G/K$, where l is the left multiplication. Denote $AdP^c = P \times_{Ad_G} \mathfrak{g}$. This reduction also gives an identification

$$AdP^c = P \times_{Ad_G} (\mathfrak{k} \otimes \mathbb{C}) = P_K \times_{Ad_K} (\mathfrak{k} \otimes \mathbb{C}) = (P_K \times_{Ad_K} \mathfrak{k}) \otimes \mathbb{C}.$$

Notice that θ and B are both Ad_K invariant on $\mathfrak{k} \otimes \mathbb{C}$. So as a complex vector bundle AdP^c , we can define θ and B on AdP^c , and then Hermitian metric H . Let $X \in AdP^c$, denote $X^* = -\theta(X)$ for the sake that the adjoint of ad_X is $ad_{-\theta(X)}$ with respect to H . Let ϕ be a section of $AdP^c \otimes K$, where K is the canonical line bundle (don't confuse with the compact subgroup K). Locally, suppose $\phi = Xdz$, then we define $\phi^* = X^*d\bar{z} \in AdP^c \otimes \bar{K}$ and $[\phi, \phi^*] = [X, X^*]dz \wedge d\bar{z}$. Let A be a principal connection on P_K , ∇_A be the corresponding connection on associated bundles of P_K . Denote $\bar{\partial}_A = \nabla_A^{0,1} \otimes 1 + 1 \otimes \bar{\partial}$, which gives a holomorphic structure on $AdP^c \otimes K$. Denote F_A be the curvature of A . Then the Hitchin equations are given by

$$\begin{aligned} (1) \quad & F_A + [\phi, \phi^*] = 0, \\ (2) \quad & \bar{\partial}_A \phi = 0. \end{aligned}$$

For a harmonic G -Higgs bundle, we mean the data $(\Sigma, G, K, P, P_K, \phi, A)$, or briefly (ϕ, A) , satisfying equations (1) and (2). Notice that $A + \phi + \phi^*$ gives a principal connection on P , and equation (1) is equivalent to this connection being flat. Given Σ, G, K , for two harmonic G -Higgs bundles, (ϕ_1, A_1) and (ϕ_2, A_2) , they are equivalent if and only if there exists a K -gauge transformation α over Σ (automatically G -gauge transformation), such that $\alpha^*\phi_2 = \phi_1, \alpha^*A_2 = A_1$.

Before we discuss the relation between Betti Moduli Space and Higgs Moduli Space, we first consider the relation between (vector) Higgs bundle and harmonic G -Higgs bundle.

Definition 2.1. Let Σ be a Riemann surface, E be a holomorphic vector bundle over Σ . Let ϕ be a holomorphic section of $End(E) \otimes K$. We call (E, ϕ) a Higgs bundle over Σ .

Given a harmonic G -Higgs bundle and suppose G acts on \mathbb{C}^n . Then the associated bundle gives a complex vector bundle E , and $\nabla_A^{0,1}$ gives a holomorphic structure of E . By definition, ϕ is a section of $End(E) \otimes K$. And by equation (2), ϕ is holomorphic.

Conversely, under some assumptions, one can obtain a harmonic G -Higgs bundle from Higgs bundle. Denote $\mu_E = \frac{\deg(E)}{\text{rank}(E)}$ be the slope of E , where $\deg(E)$ is the degree of E , and $\text{rank}(E)$ is the complex rank of E . We call (E, ϕ) is stable if for any proper ϕ -invariant holomorphic subbundle F , $\mu_F < \mu_E$. We call (E, Φ) is polystable if (E, Φ) is a direct sum of stable Higgs bundles of the same slope. Let $G = SL(n, \mathbb{C})$, $K = SU(n)$. We have the following result.

Theorem 2.2. (Hitchin [9] and Simpson [21]) *Let (E, ϕ) be a polystable Higgs bundle with structure group $SL(n, \mathbb{C})$. Then there exists a Hermitian metric h on E compatible with the $SL(n, \mathbb{C})$ structure, such that*

$$(3) \quad F^{\nabla^h} + [\phi, \phi^{*h}] = 0,$$

where F^{∇^h} is the Chern connection of h , in local holomorphic trivialization,

$$F^{\nabla^h} = \bar{\partial}(h^{-1}\partial h),$$

and ϕ^{*h} is the adjoint of ϕ with respect to h , in the sense that

$$h(\phi(u), v) = h(u, \phi^{*h}(v)), \quad u, v \in \Gamma(E)$$

in local frame, $\phi^{*h} = \bar{h}^{-1}\bar{\phi}^\top\bar{h}$, and the bracket is the commutator of $End(E)$. We regard both F^{∇^h} and $[\phi, \phi^{*h}]$ as sections of $(K \wedge \bar{K}) \otimes End(E)$. This gives rise to a flat connection $\nabla^h + \phi + \phi^{*h}$.

Notice that the Hermitian metric h gives a reduction to $SU(n)$ bundle, and ∇^h gives a principal $SU(n)$ connection. Then clearly we obtain a harmonic G -Higgs bundle.

Now we establish the correspondence between Betti Moduli Space and Higgs Moduli Space. Given a harmonic G -Higgs bundle, since $A + \phi + \phi^*$ is a flat principal connection on P , the monodromy gives a representation $\rho: \pi_1 \rightarrow G$. One can show that ρ is reductive. And the monodromy descends to the equivalent classes.

Given a reductive representation ρ , we have an associated principal G bundle $\tilde{\Sigma} \times_\rho G$, denoted as P , where $\tilde{\Sigma}$ is the universal cover of Σ ,

regarded as the π_1 principal bundle over Σ . Now we want to find a reduction from P to a principal K bundle P_K . The reduction should satisfy some condition which will be clarified below. Let $i : P_K \hookrightarrow P$ be the reduction, which is equivalent to a ρ -equivariant map $f : \tilde{\Sigma} \rightarrow G/K$. Notice that the Maurer–Cartan form ω of G gives a flat connection on P , we still use ω to denote the connection. Consider $i^*\omega$, which is a \mathfrak{g} valued one form on P_K . Decomposing $i^*\omega = A + \Phi$ from $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{k}^\perp$, where A is \mathfrak{k} valued and Φ is \mathfrak{k}^\perp valued. Then A is a principal connection on P_K and Φ is a section of $T^*M \otimes (P_K \times_{Ad_K} \mathfrak{k}^\perp)$. Consider

$$\begin{aligned} T^*M \otimes (P_K \times_{Ad_K} \mathfrak{k}^\perp) \otimes \mathbb{C} &= (T^*M \otimes \mathbb{C}) \otimes (P_K \times_{Ad_K} \mathfrak{k}^\perp \otimes \mathbb{C}) \\ &= (K \oplus \bar{K}) \otimes AdP^c. \end{aligned}$$

Regard Φ as a section of $T^*M \otimes (P_K \times_{Ad_K} \mathfrak{k}^\perp) \otimes \mathbb{C}$, and consider the decomposition $(K \oplus \bar{K}) \otimes AdP^c$. Let $\Phi = \phi + \phi^*$, where ϕ is a section of $K \otimes AdP^c$ and ϕ^* is a section of $\bar{K} \otimes AdP^c$. Notice that $\theta(\Phi) = -\Phi$, so $\phi^* = -\theta(\phi)$. Now we obtain all the desired data. Equation (1) follows from the flatness of ω . By direct calculation (see [1]), equation (2) is equivalent to f being harmonic, where the conformal metric on $\tilde{\Sigma}$ is induced from the complex structure and the metric on G/K is induced from the Killing form of \mathfrak{g} . So our requirement on f is just that f is ρ -equivariant and harmonic. From Donaldson [5] and Corlette [4], the existence of such f is equivalent to ρ being reductive. And f is unique up to the composition of the centralizer of $\rho(\pi_1)$. We see the above construction descends to the equivalence classes. So we have the bijection between Betti Moduli Space and Higgs Moduli Space.

So far, given a reductive representation $\rho : \pi_1 \rightarrow G$, we have a ρ -equivariant harmonic map $f : \tilde{\Sigma} \rightarrow G/K$. Let \hat{g} be the metric on G/K induced from Killing form of \mathfrak{g} . More precisely, let B be the Killing form of \mathfrak{g} . Consider the tangent bundle $T(G/K) = G \times_{Ad_K} \mathfrak{k}^\perp$ over G/K . Since B is Ad_K -invariant and positive definite on \mathfrak{k}^\perp , we have a well defined Riemannian metric on G/K . Locally, suppose U is a neighborhood of $p \in G/K$, $X, Y \in T_pU$. Let $\tilde{U}, \tilde{p}, \tilde{X}, \tilde{Y}$ be a lift to G . Using the Maurer–Cartan form, we have a decomposition of $T_{\tilde{p}}G$ according to $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{k}^\perp$. Then $\hat{g}(X, Y) = B(\tilde{X}^\perp, \tilde{Y}^\perp)$, where \tilde{X}^\perp is the \mathfrak{k}^\perp component of \tilde{X} .

The Higgs bundle and the corresponding harmonic map are related as follows.

Locally, choose a lift from Σ to $\tilde{\Sigma}$ and a lift from G/K to G . And lift f to \tilde{f} as a map from $\tilde{\Sigma}$ to G . For $x \in \Sigma$, we see that,

$$(\tilde{f}_* \tilde{X})^\perp = \Phi(X),$$

where $X \in T_x\Sigma$. Hence,

$$\left(\tilde{f}_* \frac{\partial}{\partial z}\right)^\perp = \phi\left(\frac{\partial}{\partial z}\right), \quad \left(\tilde{f}_* \frac{\partial}{\partial \bar{z}}\right)^\perp = \phi^*\left(\frac{\partial}{\partial \bar{z}}\right).$$

We consider the pullback metric g_f on Σ , $g_f = \pi_* f^* \hat{g}$, where π is the covering map $\pi : \tilde{\Sigma} \rightarrow \Sigma$. Since f is ρ -equivariant and \hat{g} is G -invariant, g_f is well defined. Then $\forall X, Y \in T\Sigma$,

$$g_f(X, Y) = B(\Phi(X), \Phi(Y)).$$

In particular, for $G = SL(n, \mathbb{C})$, $K = SU(n)$, we have

$$\text{Hopf}(f) = g_f^{2,0} = 2n \text{tr}_{\mathbb{C}}(\phi\phi).$$

If $\text{Hopf}(f) = 0$, then as a section of $K \otimes \bar{K}$, the Hermitian metric is

$$g_f = g_f^{1,1} = 2n \text{tr}_{\mathbb{C}}(\phi\phi^{*h}).$$

(In fact, $\phi\phi^{*h}$ is real.) And the corresponding Riemannian metric is $g_f + \bar{g}_f$.

Now we fix some notations used throughout this paper.

Let g be an Hermitian metric on K^{-1} , where K is the canonical line bundle. We can also regard g as a section of $K \otimes \bar{K}$. In local coordinate, $g = g dz \otimes d\bar{z}$. We abuse the same notation g to denote both the metric and the local function. Similarly, let h be an Hermitian metric on K^{-l} , denoted by $h = h dz^{\otimes l} \otimes d\bar{z}^{\otimes l}$. Notice that as a local function, h is not globally well defined, but if we set $h = ag^l$, then a is a globally defined function.

Let q_n be an n -differential. Locally, denote $q_n = q_n dz^n$. Define a local function $|q_n|^2 = q_n \bar{q}_n$, which is corresponding to the Hermitian metric on K^{-n} . Here are some notations used later.

- (1) $|q_n|_g^2$ as the square of the norm of q_n with respect to the metric g , which is globally defined, in local coordinate, $|q_n|_g^2 = |q_n|^2 g^{-n}$;
- (2) $\|q_n\| = \int_{\Sigma} |q_n|^{\frac{2}{n}}$, which will be used in Section 6;
- (3) g_0 as the unique Hermitian hyperbolic metric compatible with the complex structure;
- (4) g_{Fuchsian} as the Hermitian metric corresponding to the base Fuchsian point, a multiple of g_0 ;
- (5) $\Delta = \partial_z \partial_{\bar{z}} = \frac{1}{4}(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2})$, which is locally defined;
- (6) $\Delta_g = g^{-1} \Delta$, noting that Δ_g is globally defined.

3. Rewriting Hitchin equations in two subclasses for Hitchin representations

In Section 2, we relate the moduli space of reductive representations $\rho \in \text{Hom}(\pi_1, SL(n, \mathbb{C}))$ to the moduli space of harmonic $SL(n, \mathbb{C})$ -Higgs bundles. Under this bijection, we can also describe the moduli space of reductive representations $\rho \in \text{Hom}(\pi_1, SL(n, \mathbb{R}))$ in the setting of Higgs bundles. In fact, in [10], Hitchin gives a parametrization of the Hitchin component of the $SL(n, \mathbb{R})$ -Higgs bundle moduli space using

The pullback metric g_f of the minimal immersion on the surface is

$$g_f = 2n \operatorname{tr}_{\mathbb{C}}(\phi \phi^{*h}) = 2n(h_1^2 |q_n|^2 + 2 \sum_{k=1}^{m-1} h_k^{-1} h_{k+1} + h_m^{-2}).$$

For $n = 2m + 1$ odd, the situation is similar. Notice that there is a trivial bundle K^0 in the middle of E , we see that the metric on K^0 is 1, i.e., the standard Hermitian metric on \mathbb{C} . Then the last equation of the system of equations becomes

$$\Delta \log h_m + h_m^{-1} - h_{m-1}^{-1} h_m = 0.$$

If $q_n = 0$, then

$$h_k^{-1} h_{k+1} = \frac{1}{2} k(n-k) g_0, \quad 1 \leq k \leq m-1, \quad h_m^{-1} = \frac{1}{8} (n^2 - 1) g_0.$$

The last term in the pullback metric g_f of the minimal immersion is $2h_m^{-1}$ instead of h_m^{-2} .

q_{n-1} **Case:** The Higgs field ϕ is a holomorphic section of $\operatorname{End}(E) \otimes K$ of the form

$$\begin{pmatrix} 0 & & q_{n-1} & 0 \\ 1 & 0 & & q_{n-1} \\ & 1 & 0 & \\ & & \ddots & \\ & & & 0 \\ & & & 1 & 0 \end{pmatrix},$$

where q_{n-1} is a holomorphic section of K^{n-1} . From [2], the Hermitian metric h is given by

$$h = \operatorname{diag}(h_1, h_2, \dots, h_2^{-1}, h_1^{-1}).$$

By direct calculation, the Hopf differential $\operatorname{Hopf}(f) = 2n \operatorname{tr}_{\mathbb{C}}(\phi^2)$ vanishes and, hence, f is a minimal immersion. Equation (3) gives the following system of equations.

For $n = 2m$ even,

$$\begin{aligned} \Delta \log h_1 + h_1^{-1} h_2 - h_1 h_2 |q_{n-1}|^2 &= 0, \\ \Delta \log h_2 + h_2^{-1} h_3 - h_1^{-1} h_2 - h_1 h_2 |q_{n-1}|^2 &= 0, \\ \Delta \log h_k + h_k^{-1} h_{k+1} - h_{k-1}^{-1} h_k &= 0, \quad 3 \leq k \leq m-1, \\ \Delta \log h_m + h_m^{-2} - h_{m-1}^{-1} h_m &= 0. \end{aligned}$$

When $q_{n-1} = 0$, h_k is the same as in the q_n case, $n = 2m$.

The pullback metric of minimal immersion is

$$g_f = 2n \operatorname{tr}_{\mathbb{C}}(\phi \phi^{*h}) = 2n(2h_1 h_2 |q_{n-1}|^2 + 2 \sum_{k=1}^{m-1} h_k^{-1} h_{k+1} + h_m^{-2}).$$

For $n = 2m + 1$ is odd, the last equation of the system of equations is

$$\Delta \log h_m + h_m^{-1} - h_{m-1}^{-1} h_m = 0.$$

When $q_{n-1} = 0$, h_k is the same as in the q_n case, $n = 2m + 1$.

The last term in the pullback metric g_f is $2h_m^{-1}$ instead of h_m^{-2} .

Remark 3.1. For the situation $q_n = 0$ in q_n case (or $q_{n-1} = 0$ in q_{n-1} case), it is the base Fuchsian point. We denote the pullback metric g_f in this situation as g_{Fuchsian} . It is computed in Section 4 that $g_{\text{Fuchsian}} = \frac{1}{6}(n^4 - n^2)g_0$, where g_0 is the unique Hermitian hyperbolic metric.

4. Domination of pullback metric

Let g_0 be the unique hyperbolic metric on Riemann surface Σ with constant curvature -1 and compatible with the complex structure. Regarding g_0 as a section of $K \otimes \bar{K}$, we set $g_0 = g_0 dz \otimes d\bar{z}$. (If we regard g_0 as an Hermitian metric on $T\Sigma$, then $g_0 = g_0(dz \otimes d\bar{z} + d\bar{z} \otimes dz) = 2g_0(dx^2 + dy^2)$.)

Locally, g_0 satisfies

$$\Delta \log g_0 - g_0 = 0,$$

where $\Delta = \partial_z \partial_{\bar{z}} = \frac{1}{4}(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2})$ is the coordinate Laplacian.

In this section, we show that the pullback metric of minimal immersion for the q_n, q_{n-1} cases dominates the base Fuchsian metric g_{Fuchsian} . More precisely,

Theorem 4.1. *For the q_n, q_{n-1} cases*

$$g_f \geq g_{\text{Fuchsian}} = \frac{1}{6}(n^4 - n^2)g_0.$$

Moreover, if equality holds at one point, then $q_n = 0$, or $q_{n-1} = 0$, respectively, which implies h_k is a suitable power of g_0 , as described in Section 3.

Remark 4.2. In fact, we prove the rigidity result for every term in g_f . For example, for the q_n case, $n = 2m$, if one of the following equalities hold at one point,

$$h_k^{-1} h_{k+1} \geq (km - \frac{1}{2}k^2)g_0, \quad 1 \leq k \leq m-1, \quad h_m^{-2} \geq \frac{1}{2}m^2 g_0,$$

the rigidity result then holds.

Proof of Theorem 4.1. We only prove the theorem for the q_n case, $n = 2m$. For other cases, the proof is similar. Write $h_k = a_k g_0^{k-\frac{1}{2}-\frac{n}{2}}$, $1 \leq k \leq m$ and $|q_n|^2 = a_0 g_0^n$. Then $a_k, 0 \leq k \leq m$ are globally well-defined functions. They satisfy

$$\begin{aligned} \Delta \log a_1 + \left(\frac{1}{2} - \frac{n}{2}\right) \Delta \log g_0 + a_1^{-1} a_2 g_0 - a_1^2 a_0 g_0 &= 0, \\ \Delta \log a_k + \left(k - \frac{n+1}{2}\right) \Delta \log g_0 + (a_k^{-1} a_{k+1} - a_{k-1}^{-1} a_k) g_0 &= 0, \\ &\text{for } 2 \leq k \leq m-1, \\ \Delta \log a_m - \frac{1}{2} \Delta \log g_0 + (a_m^{-2} - a_{m-1}^{-1} a_m) g_0 &= 0. \end{aligned}$$

Then g_f becomes $2n(a_1^2 a_0 + 2 \sum_{k=1}^{m-1} a_k^{-1} a_{k+1} + a_m^{-2}) g_0$.

Let $b_k = a_k^{-1} a_{k+1}$, $1 \leq k \leq m-1$ and $b_m = a_m^{-2}$. Then

$$a_k = b_m^{-\frac{1}{2}} b_{m-1}^{-1} \cdots b_k^{-1}, \quad 1 \leq k \leq m-1, \quad a_m = b_m^{-\frac{1}{2}}.$$

So

$$\begin{aligned} \Delta \log a_k &= -\frac{1}{2} \Delta \log b_m - \Delta \log b_{m-1} - \cdots - \Delta \log b_k, \quad 1 \leq k \leq m-1, \\ \Delta \log a_m &= -\frac{1}{2} \Delta \log b_m. \end{aligned}$$

Plug these terms into the system of equations above, using $\Delta \log g_0 = g_0$, we have

$$\begin{aligned} \frac{1}{2} \Delta \log b_m + \sum_{j=1}^{m-1} \Delta \log b_j - \left(\frac{1}{2} - m + b_1\right) g_0 &\leq 0, \\ \frac{1}{2} \Delta \log b_m + \sum_{j=k}^{m-1} \Delta \log b_j - \left(k - \frac{1}{2} - m + b_k - b_{k-1}\right) g_0 &= 0, \\ &\text{for } 2 \leq k \leq m-1, \\ \frac{1}{2} \Delta \log b_m - \left(-\frac{1}{2} + b_m - b_{m-1}\right) g_0 &= 0. \end{aligned}$$

Then

$$\begin{aligned} \Delta \log b_1 + (1 + b_2 - 2b_1) g_0 &\leq 0, \\ (4) \quad \Delta \log b_k + (1 + b_{k+1} + b_{k-1} - 2b_k) g_0 &= 0, \quad 2 \leq k \leq m-1, \\ \Delta \log b_m + (1 + 2b_{m-1} - 2b_m) g_0 &= 0. \end{aligned}$$

Let x_k be a minimizer of b_k . Then

$$\begin{aligned} 2b_1 &\geq 1 + b_2(x_1), \\ 2b_k &\geq 1 + b_{k+1}(x_k) + b_{k-1}(x_k), \quad 2 \leq k \leq m-1, \\ 2b_m &\geq 1 + 2b_{m-1}(x_m). \end{aligned}$$

First consider

$$\begin{aligned} 2b_{m-1} &\geq 1 + b_m(x_{m-1}) + b_{m-2}(x_{m-1}), \\ 2b_m &\geq 1 + 2b_{m-1}(x_m). \end{aligned}$$

We see $b_m(x_{m-1}) \geq 2 + b_{m-2}(x_{m-1})$ and then $2b_{m-1} \geq 3 + 2b_{m-2}(x_{m-1})$. Then, similarly, consider $k = m-1, m-2$. We obtain

$$\begin{aligned} b_{m-1}(x_{m-2}) &\geq 4 + b_{m-3}(x_{m-2}), \\ 2b_{m-2} &\geq 5 + 2b_{m-3}(x_{m-2}). \end{aligned}$$

Follow this procedure until we obtain

$$\begin{aligned} b_3(x_2) &\geq 2(m-2) + b_1(x_2), \\ 2b_2 &\geq 2m-3 + 2b_1(x_2). \end{aligned}$$

From $2b_1 \geq 1 + b_2(x_1)$, we have $b_2(x_1) \geq 2m-2$ and then $2b_1 \geq 2m-1$. Similarly,

$$2b_k \geq (2m-1) + \cdots + (2m - (2k-1)), \quad 1 \leq k \leq m.$$

So we obtain

$$2b_k \geq 2km - k^2, \quad 1 \leq k \leq m.$$

Recall that $\sum_{k=1}^{m-1} k^2 = \frac{1}{6}(m-1)m(2m-1)$. Hence, finally,

$$g_f = 2n(h_1^2 |q_{2m}|^2) + 2 \sum_{k=1}^{m-1} h_k^{-1} h_{k+1} + h_m^{-2} \geq \frac{n^4 - n^2}{6} g_0.$$

For rigidity, suppose $b_k = km - \frac{1}{2}k^2$ holds at one point p for some k , let $\log b_k = \Omega_k$, then from (4),

$$\Delta \Omega_k + 2(km - \frac{1}{2}k^2 - e^{\Omega_k})g_0 \leq 0.$$

Then one may apply the strong maximum principle to finish the proof. More precisely, in a local coordinate chart,

$$\begin{aligned} &\Delta(\Omega_k - \log(km - \frac{1}{2}k^2)) \\ &- 2(\Omega_k - \log(km - \frac{1}{2}k^2)) \left(\int_0^1 e^{t\Omega_k + (1-t)\log(km - \frac{1}{2}k^2)} dt \right) g_0 \leq 0. \end{aligned}$$

This is from the fact

$$e^{\Omega_k} - e^{\log(km - \frac{1}{2}k^2)} = \frac{d}{dt} \Big|_0^1 e^{t\Omega_k + (1-t)\log(km - \frac{1}{2}k^2)}.$$

Notice that $\int_0^1 e^{t\Omega_k + (1-t)\log(km - \frac{1}{2}k^2)} dt \geq 0$ and $\Omega_k - \log(km - \frac{1}{2}k^2) \geq 0$, then by the strong maximum principle [11], if $\Omega_k - \log(km - \frac{1}{2}k^2) = 0$ at one point p , then $\Omega_k - \log(km - \frac{1}{2}k^2) = 0$ everywhere. Then b_k equals $km - \frac{1}{2}k^2$ identically for all $1 \leq k \leq m$. From the origin system of equations, we see that q_n must be zero. q.e.d.

5. Shape of minimal surface inside symmetric space

In this section, we investigate the shape of minimal surface Σ inside the symmetric space G/K . In particular, we show that the tangent space of Σ is never tangential to any flat inside G/K . Recall $G = SL(n, \mathbb{C})$, $K = SU(n)$.

Theorem 5.1. *For the q_n, q_{n-1} cases, we have the following results.*

(1) *The Hitchin equation never decouples: for every point on Σ ,*

$$F^{\nabla^h} \neq 0, \quad [\phi, \phi^{*h}] \neq 0.$$

(2) *The sectional curvature $K_{G/K}(\sigma)$ is strictly negative, where σ is the tangent space of the image of g_f . Geometrically, the minimal immersion is never tangential to any flat inside the symmetric space.*

(3) *On each line bundle $K^{\frac{n+1-2k}{2}}$, $1 \leq k \leq n$, the Chern form of h_k :*

$$\sqrt{-1}\Theta_{h_k} = \sqrt{-1}\bar{\partial}\partial \log h_k$$

is strictly positive if $n+1-2k > 0$; zero if $n+1-2k = 0$; strictly negative if $n+1-2k < 0$.

Before we prove Theorem 5.1, an immediate corollary is as follows.

Corollary 5.2. *For the q_n, q_{n-1} cases, the sectional curvature K_{g_f} of Σ equipped with the induced metric g_f is strictly negative.*

Proof of Corollary 5.2. Given the ρ -equivariant harmonic map $f : \tilde{\Sigma} \rightarrow G/K$, we want to investigate that, as an immersed submanifold inside the symmetric space G/K , how $f(\tilde{\Sigma})$ interacts with the symmetric space. Let ∇ denote the Levi-Civita connection of G/K and ∇^T denote the component of ∇ tangential to the image of f . Then the second fundamental form is the symmetric 2-tensor with values in the normal bundle given by

$$II(X, Y) = \nabla_X Y - \nabla_X^T Y,$$

where $X, Y \in \Gamma(f^*T(G/K))$ are tangent to the image of f . Let $\{e_1, e_2\}$ be an orthonormal basis such that

$$II(e_1, e_2) = 0.$$

The immersion is minimal if the trace of II with respect to g_f is vanishing. Therefore,

$$II(e_1, e_1) + II(e_2, e_2) = 0.$$

Recall the Gauss equation, let X, Y be orthonormal vector fields on Σ , then

$$K_{g_f}(X, Y) = K_{G/K}(X, Y) + \langle II(X, X), II(Y, Y) \rangle - |II(X, Y)|^2.$$

Therefore, the sectional curvature K_{g_f} and $K_{G/K}$ measured in Σ and in G/K , respectively, are related by Gauss equation:

$$K_{g_f} = K_{G/K} - \frac{1}{2} \|II\|_{g_f}^2.$$

By Theorem 5.1, $K_{G/K} < 0$. Therefore, $K_{g_f} < 0$.

q.e.d.

Proof of Theorem 5.1. We first discover the following inequalities for the 2-tensors appearing in the Hitchin system of equations. We will prove Lemma 5.3, 5.4 later.

Lemma 5.3. *For the q_n case, on the whole surface Σ , if $n = 2m$*

$$h_1^2|q_n|^2 < h_1^{-1}h_2 < h_2^{-1}h_3 < \cdots < h_{k-1}^{-1}h_k < \cdots < h_{m-1}^{-1}h_m < h_m^{-2}.$$

If $n = 2m + 1$, the last term is replaced by h_m^{-1} .

Lemma 5.4. *For the q_{n-1} case, on the whole surface Σ , if $n = 2m$*

$$h_1h_2|q_{n-1}|^2 < h_1^{-1}h_2,$$

$$h_1h_2|q_{n-1}|^2 + h_1^{-1}h_2 < h_2^{-1}h_3 < \cdots < h_{m-1}^{-1}h_m < h_m^{-2}.$$

If n is odd, the last term is replaced by h_m^{-1} .

For the q_n case, the $(1,1)$ -entry of $[\phi, \phi^{*h}]$ is $h_1^2|q_n|^2 - h_1^{-1}h_2$. It is strictly negative everywhere by Lemma 5.3. For the q_{n-1} case, the $(1,1)$ -entry of $[\phi, \phi^{*h}]$ is $h_1h_2|q_{n-1}|^2 - h_1^{-1}h_2$. It is strictly negative everywhere by Lemma 5.4. Therefore, Part (1) follows.

For part (2), the sectional curvature in the symmetric space G/K for the 2-dimensional subspace spanned by Y, Z is

$$K_{G/K}(Y, Z) := -\frac{B([Y_0, Z_0], [Y_0, Z_0])}{B(Y_0, Y_0)B(Z_0, Z_0)} = -\frac{|[Y_0, Z_0]|^2}{|Y_0|^2|Z_0|^2},$$

where $Y_0 = \Phi(Y) \in \mathfrak{k}^\perp$ and $Z_0 = \Phi(Z) \in \mathfrak{k}^\perp$ from Section 2.

The tangent space of the minimal immersion at $f(p)$ inside G/K is spanned by $Y = f_*(\frac{\partial}{\partial x})$ and $Z = f_*(\frac{\partial}{\partial y})$ with $Y_0 = \Phi(\frac{\partial}{\partial x}) = (\phi + \phi^*)(\frac{\partial}{\partial x}) = \phi(\frac{\partial}{\partial z}) + \phi^*(\frac{\partial}{\partial \bar{z}})$, and $Z_0 = \Phi(\frac{\partial}{\partial y}) = (\phi + \phi^*)(\frac{\partial}{\partial y}) = \sqrt{-1}\phi(\frac{\partial}{\partial z}) - \sqrt{-1}\phi^*(\frac{\partial}{\partial \bar{z}})$. Hence,

$$[Y_0, Z_0] = -2\sqrt{-1}[\phi(\frac{\partial}{\partial z}), \phi^*(\frac{\partial}{\partial \bar{z}})] = -2\sqrt{-1}[\phi, \phi^*](\frac{\partial}{\partial z}, \frac{\partial}{\partial \bar{z}}).$$

By part (1), $[\phi, \phi^*] \neq 0$, hence, $K_{G/K} < 0$. Therefore, Part (2) follows.

Part (3) directly follows from Lemma 5.3, 5.4. q.e.d.

Proof of Lemma 5.3. We focus on the case $n = 2m$. Along the proof, we single out each time the only differences for the case $n = 2m + 1$ and show that the proof works as well in this case. Let $h_k = a_k g_0^{\frac{k-n+1}{2}}$, $1 \leq k \leq m$. Let $|q_n|^2 = a_0 g_0^n$. Then $a_k, 0 \leq k \leq m$ are globally defined functions.

The situation $q_n = 0$ is clear. We then assume q_n is not identically zero. Since q_n is holomorphic, q_n only has discrete zeros. Denote $U = \{p \in \Sigma | q_n(p) \neq 0\}$. Then a_k satisfies, locally

$$\Delta \log a_1 + (1 - \frac{n+1}{2} + a_1^{-1}a_2 - a_1^2 a_0)g_0 = 0,$$

$$\begin{aligned}\Delta \log a_k + \left(k - \frac{n+1}{2} + a_k^{-1} a_{k+1} - a_{k-1}^{-1} a_k\right) g_0 &= 0, \quad 2 \leq k \leq m-1, \\ \Delta \log a_m + \left(m - \frac{n+1}{2} + a_m^{-2} - a_{m-1}^{-1} a_m\right) g_0 &= 0.\end{aligned}$$

For the case $n = 2m + 1$, in the last equation, a_m^{-2} is replaced by a_m^{-1} .

Let $\Omega_k = \log a_k$, $k = 0, 1, \dots, m$. Notice that at the zero point of q_n , Ω_0 goes to $-\infty$ continuously. Then the system of equations becomes,

$$\begin{aligned}\Delta \Omega_1 + \left(1 - \frac{n+1}{2} + e^{-\Omega_1 + \Omega_2} - e^{2\Omega_1 + \Omega_0}\right) g_0 &= 0, \quad \text{in } U, \\ \Delta \Omega_k + \left(k - \frac{n+1}{2} + e^{-\Omega_k + \Omega_{k+1}} - e^{-\Omega_{k-1} + \Omega_k}\right) g_0 &= 0, \quad 2 \leq k \leq m-1, \\ \Delta \Omega_m + \left(m - \frac{n+1}{2} + e^{-2\Omega_m} - e^{-\Omega_{m-1} + \Omega_m}\right) g_0 &= 0.\end{aligned}$$

For the case $n = 2m + 1$, in the last equation, $e^{-2\Omega_m}$ is replaced by $e^{-\Omega_m}$.

Let $f_1 = 2\Omega_1 + \Omega_0$, $f_k = -\Omega_{k-1} + \Omega_k$, $2 \leq k \leq m$, $f_{m+1} = -2\Omega_m$. For the case $n = 2m + 1$, let $f_{m+1} = -\Omega_m$. Notice that, since q_n is holomorphic, away from zeros of q_n ,

$$\begin{aligned}\Delta \Omega_0 &= \Delta \log a_0 = \Delta \log(|q_n|^2 g_0^{-n}) \\ &= \Delta \log |q_n|^2 - n \Delta \log g_0 = -n g_0.\end{aligned}$$

Then we obtain

$$\begin{aligned}\Delta f_1 + (1 + 2e^{f_2} - 2e^{f_1}) g_0 &= 0, \quad \text{in } U, \\ \Delta f_k + (1 + e^{f_{k+1}} - 2e^{f_k} + e^{f_{k-1}}) g_0 &= 0, \quad 2 \leq k \leq m, \\ \Delta f_{m+1} + (1 - 2e^{f_{m+1}} + 2e^{f_m}) g_0 &= 0.\end{aligned}$$

For the case $n = 2m + 1$, the last equation is replaced by

$$\Delta f_{m+1} + (1 - e^{f_{m+1}} + e^{f_m}) g_0 = 0.$$

Then

$$\begin{aligned}(5) \quad \Delta(f_1 - f_2) + (-3(e^{f_1} - e^{f_2}) + (e^{f_2} - e^{f_3})) g_0 &= 0, \quad \text{in } U, \\ \Delta(f_k - f_{k+1}) + ((e^{f_{k-1}} - e^{f_k}) + (e^{f_{k+1}} - e^{f_{k+2}}) - 2(e^{f_k} - e^{f_{k+1}})) g_0 &= 0, \\ &\text{for } 2 \leq k \leq m-1, \\ \Delta(f_m - f_{m+1}) + (-3(e^{f_m} - e^{f_{m+1}}) + (e^{f_{m-1}} - e^{f_m})) g_0 &= 0.\end{aligned}$$

For the case $n = 2m + 1$, the last equation is replaced by

$$\Delta(f_m - f_{m+1}) + (-2(e^{f_m} - e^{f_{m+1}}) + (e^{f_{m-1}} - e^{f_m})) g_0 = 0.$$

Let A_k be the maximum of $e^{f_k - f_{k+1}}$, $1 \leq k \leq m$.

Step 1: We first show $A_m \leq 1$. From the compactness of Σ , A_k , $2 \leq k \leq m$ must be achieved at some p_k . For A_1 , since $f_1 - f_2$ goes

to $-\infty$ around the zeros of q_n , A_1 must be achieved at some $p_1 \in U$. Notice that $A_k \geq 0$, $1 \leq k \leq m$. Then by the maximum principle, at p_1

$$\begin{aligned} & -3(e^{f_1(p_1)} - e^{f_2(p_1)}) + (e^{f_2(p_1)} - e^{f_3(p_1)}) \geq 0 \\ \Rightarrow & -3(e^{(f_1-f_2)(p_1)} - 1) + (1 - e^{-(f_2-f_3)(p_1)}) \geq 0 \\ \Rightarrow & -3(A_1 - 1) + (1 - A_2^{-1}) \geq 0 \\ \Rightarrow & 2(A_1 - 1) \leq (1 - A_2^{-1}) - (A_1 - 1). \end{aligned}$$

At p_k , $2 \leq k \leq m-1$

$$\begin{aligned} & (-2(e^{f_k(p_k)} - e^{f_{k+1}}) + (e^{f_{k-1}} - e^{f_k}) + (e^{f_{k+1}} - e^{f_{k+2}}))(p_k) \geq 0 \\ \Rightarrow & -2(e^{(f_k-f_{k+1})(p_k)} - 1) + (e^{(f_{k-1}-f_k)(p_k)} - 1)e^{(f_k-f_{k+1})(p_k)} \\ & + (1 - e^{-(f_{k+1}-f_{k+2})(p_k)}) \geq 0 \\ \Rightarrow & -2(A_k - 1) + (A_{k-1} - 1)A_k + (1 - A_{k+1}^{-1}) \geq 0 \\ \Rightarrow & (1 - A_k^{-1}) - (A_{k-1} - 1) \leq A_k^{-1}((1 - A_{k+1}^{-1}) - (A_k - 1)). \end{aligned}$$

Similarly, at p_m , we obtain

$$(1 - A_m^{-1}) - (A_{m-1} - 1) \leq -2(1 - A_m^{-1}).$$

For $n = 2m + 1$ case, at p_m , instead we obtain

$$(1 - A_m^{-1}) - (A_{m-1} - 1) \leq -(1 - A_m^{-1}).$$

Let $B_1 = 2(A_1 - 1)$, $B_k = (1 - A_k^{-1}) - (A_{k-1} - 1)$, $2 \leq k \leq m$. Then

$$\begin{aligned} B_1 & \leq B_2, \\ B_k & \leq A_k^{-1}B_{k+1}, \quad 2 \leq k \leq m-1, \\ B_m & \leq -2(1 - A_m^{-1}). \end{aligned}$$

For $n = 2m + 1$ case, the last inequality is replaced by

$$B_m \leq -(1 - A_m^{-1}).$$

On the other hand,

$$\begin{aligned} 1 - A_m^{-1} & = B_m + (A_{m-1} - 1) \\ & = B_m + A_{m-1}(1 - A_{m-1}^{-1}) \\ & = B_m + A_{m-1}(B_{m-1} + (A_{m-2} - 1)) \\ & \quad \vdots \\ & = B_m + A_{m-1}B_{m-1} + A_{m-1}A_{m-2}B_{m-2} + \cdots \\ & \quad + A_{m-1}A_{m-2} \cdots A_2B_2 + A_{m-1}A_{m-2} \cdots A_2(A_1 - 1) \\ & \leq \underbrace{B_m + B_m + \cdots + B_m}_{m-1 \text{ terms}} + \frac{1}{2}B_m \\ & \leq -2(m - \frac{1}{2})(1 - A_m^{-1}). \end{aligned}$$

For $n = 2m + 1$ case, the inequality is replaced by

$$1 - A_m^{-1} \leq -(m - \frac{1}{2})(1 - A_m^{-1}).$$

Thus, we obtain $A_m \leq 1$.

Step 2: Now we claim $A_1 < 1$ and $A_m < 1$. To prove $A_1 < 1$ by contradiction, assume $A_1 \geq 1$. Then $B_1 \geq 0$. And then $B_k \geq 0$ for $2 \leq k \leq m$. In other words, $1 - A_k^{-1} \geq A_{k-1} - 1$ for $2 \leq k \leq m$. From $A_1 \geq 1$, we see $A_k \geq 1$ for $2 \leq k \leq m$. In particular, $A_m \geq 1$. So A_m has to be equal to 1. It turns out $A_k = 1$ for every k . We use the strong maximum principle to get contradiction. Consider equation (5)

$$\Delta(f_1 - f_2) + (-3(e^{f_1 - f_2} - 1) + (1 - e^{-(f_2 - f_3)}))e^{f_2}g_0 = 0, \quad \text{in } U.$$

Since $e^{f_2 - f_3} \leq 1$, we have

$$\Delta(f_1 - f_2) - 3(e^{f_1 - f_2} - 1)e^{f_2}g_0 \geq 0 \quad \text{in } U.$$

Let $u = f_1 - f_2$. Then

$$\Delta u - (3e^{f_2}g_0 \int_0^1 e^{tu} dt)u \geq 0 \quad \text{in } U.$$

By the assumption, u achieves its maximum 0 in U . Then by the strong maximum principle, u has to be 0 identically in U . But q_n must have zeros. So u has to take value $-\infty$, which is a contradiction to $ueqnarrray * uiv0$. So we have proved $A_m < 1$ by contradiction. But since $A_1 \geq 1$ implies $A_m \geq 1$, we also see $A_1 < 1$ by contradiction as well, and $e^{f_1} < e^{f_2}$ everywhere on Σ .

Step 3: Now we begin to show $A_k < 1$, for all $1 \leq k \leq m$. Assume there is some $1 \leq k_0 \leq m - 1$ such that $B_{k_0} \leq 0, B_{k_0+1} > 0$. The proof below works in the other two cases $B_1 > 0$ and $B_m \leq 0$ as well. Since $B_k \leq A_k^{-1}B_{k+1}$, for all $1 \leq k \leq m - 1$, then

$$\begin{aligned} B_k &\leq 0, & 1 \leq k \leq k_0, \\ B_k &> 0, & k_0 + 1 \leq k \leq m. \end{aligned}$$

Then $B_k = (1 - A_k^{-1}) - (A_{k-1} - 1) \leq 0$ for $1 \leq k \leq k_0$ gives

$$1 - A_k^{-1} \leq A_{k-1} - 1, \quad 1 \leq k \leq k_0.$$

Since $A_1 < 1$, then $A_1 - 1 < 0$, hence,

$$A_k - 1 < 0, \quad 1 \leq k \leq k_0.$$

And $B_k = (1 - A_k^{-1}) - (A_{k-1} - 1) > 0$ for $k_0 + 1 \leq k \leq m$ gives

$$1 - A_{k+1}^{-1} > A_k - 1, \quad k_0 + 1 \leq k \leq m.$$

Since $A_m - 1 < 0$, we have

$$A_k - 1 < 0, \quad k_0 + 1 \leq k \leq m.$$

Therefore, we obtain $A_k < 1$, for all $1 \leq k \leq m$.

q.e.d.

Proof of Lemma 5.4. We only prove for the case $n = 2m$. The case for $n = 2m + 1$ is proved in a similar way as in Lemma 5.3. As above, let $h_k = a_k g_0^{k-\frac{1}{2}-m}$, $1 \leq k \leq m$. Let $|q_{n-1}|^2 = a_0 g_0^{n-1}$. Then a_k satisfies, locally

$$\begin{aligned} \Delta \log a_1 + \left(\frac{1}{2} - m + a_1^{-1} a_2 - a_1 a_2 a_0\right) g_0 &= 0, \\ \Delta \log a_2 + \left(\frac{3}{2} - m + a_2^{-1} a_3 - a_1^{-1} a_2 - a_1 a_2 a_0\right) g_0 &= 0, \\ \Delta \log a_k + \left(k - \frac{1}{2} - m + a_k^{-1} a_{k+1} - a_{k-1}^{-1} a_k\right) g_0 &= 0, \quad 3 \leq k \leq m-1, \\ \Delta \log a_m + \left(-\frac{1}{2} + a_m^{-2} - a_{m-1}^{-1} a_m\right) g_0 &= 0. \end{aligned}$$

Denote $U = \{p \in \Sigma | q_{n-1}(p) \neq 0\}$. Let $\Omega_k = \log a_k$, $k = 0, 1, \dots, m$. Then

$$\begin{aligned} \Delta \Omega_1 + \left(\frac{1}{2} - m + e^{-\Omega_1 + \Omega_2} - e^{\Omega_1 + \Omega_2 + \Omega_0}\right) g_0 &= 0, \quad \text{in } U, \\ \Delta \Omega_2 + \left(\frac{3}{2} - m + e^{-\Omega_2 + \Omega_3} - e^{-\Omega_1 + \Omega_2} - e^{\Omega_1 + \Omega_2 + \Omega_0}\right) g_0 &= 0, \quad \text{in } U, \\ \Delta \Omega_k + \left(k - \frac{1}{2} - m + e^{-\Omega_k + \Omega_{k+1}} - e^{-\Omega_{k-1} + \Omega_k}\right) g_0 &= 0, \\ &\text{for } 3 \leq k \leq m-1, \\ \Delta \Omega_m + \left(-\frac{1}{2} + e^{-2\Omega_m} - e^{-\Omega_{m-1} + \Omega_m}\right) g_0 &= 0. \end{aligned}$$

Let $f_1 = \Omega_1 + \Omega_2 + \Omega_0$, $f_k = -\Omega_{k-1} + \Omega_k$, $2 \leq k \leq m$, $f_{m+1} = -2\Omega_m$. Notice that $\Delta \Omega_0 = (1-n)g_0$. Then we obtain

$$\begin{aligned} \Delta f_1 + (1 + e^{f_3} - 2e^{f_1}) g_0 &= 0, \quad \text{in } U, \\ \Delta f_2 + (1 + e^{f_3} - 2e^{f_2}) g_0 &= 0, \\ \Delta f_3 + (1 + e^{f_4} - 2e^{f_3} + e^{f_2} + e^{f_1}) g_0 &= 0, \\ \Delta f_k + (1 + e^{f_{k+1}} - 2e^{f_k} + e^{f_{k-1}}) g_0 &= 0, \quad 4 \leq k \leq m, \\ \Delta f_{m+1} + (1 - 2e^{f_{m+1}} + 2e^{f_m}) g_0 &= 0. \end{aligned}$$

Then we have the following system

$$\begin{aligned} \Delta(f_3 - f_4) + (-2(e^{f_3} - e^{f_4}) + (e^{f_4} - e^{f_5}) + (e^{f_1} + e^{f_2} - e^{f_3})) g_0 &= 0, \\ \Delta(f_k - f_{k+1}) + ((e^{f_{k-1}} - e^{f_k}) + (e^{f_{k+1}} - e^{f_{k+2}}) - 2(e^{f_k} - e^{f_{k+1}})) g_0 &= 0, \\ &\text{for } 3 \leq k \leq m-1, \\ \Delta(f_m - f_{m+1}) + (-3(e^{f_m} - e^{f_{m+1}}) + (e^{f_{m-1}} - e^{f_m})) g_0 &= 0. \end{aligned}$$

Let A_k be the maximum of $e^{f_k - f_{k+1}}$, $k = 1$ and $3 \leq k \leq m$, and A_2 be the maximum of $e^{f_1 - f_3} + e^{f_2 - f_3}$.

Step 1: We show $A_1 < 1$. We only look at the first equation of the Hitchin equation

$$\Delta\Omega_1 + \left(\frac{1}{2} - m + e^{-\Omega_1+\Omega_2} - e^{\Omega_1+\Omega_2+\Omega_0}\right)g_0 = 0, \quad \text{in } U.$$

Consider the function $-\frac{1}{2}\Omega_0 = \log|q_{n-1}|^{-1}g_0^{\frac{n-1}{2}}$, it satisfies the above equation. Since it is ∞ at zeros of q_{n-1} , it gives an upper bound for Ω_1 . By the strong maximum principle, we have $\Omega_1 < \log|q_{n-1}|^{-1}g_0^{\frac{n-1}{2}}$. Hence, $e^{2\Omega_1}|q_{n-1}|^2 < 1$, then $A_1 < 1$.

Step 2: We claim that $A_m < 1$. For other A_k , we apply the same argument in Lemma 5.3 to the above system. Let $B_k = (1 - A_k^{-1}) - (A_{k-1} - 1)$, $3 \leq k \leq m$. Then

$$(6) \quad \begin{aligned} B_k &\leq A_k^{-1}B_{k+1}, \quad 3 \leq k \leq m-1, \\ B_m &\leq -2(1 - A_m^{-1}). \end{aligned}$$

Let $B_2 = A_2 - 1$. Now we start to obtain inequality for B_2 . Applying the inequality

$$\Delta \log(e^f + 1) \geq \frac{e^f}{e^f + 1} \Delta f,$$

we have

$$\begin{aligned} &\frac{1}{g_0} \Delta \log(e^{f_1-f_3} + e^{f_2-f_3}) \\ &= \Delta \log(e^{f_1-f_2} + 1) + \Delta \log e^{f_2-f_3} \\ &\geq \frac{e^{f_1-f_2}}{e^{f_1-f_2} + 1} \Delta(f_1 - f_2) + \Delta(f_2 - f_3) \\ &= \frac{e^{f_1-f_2}}{e^{f_1-f_2} + 1} 2(e^{f_1} - e^{f_2}) + (e^{f_1} + 3e^{f_2} - 3e^{f_3} + e^{f_4}) \\ &= e^{f_2} \frac{(e^{f_1-f_2} - 1)^2}{e^{f_1-f_2} + 1} + (e^{f_1} - e^{f_2}) + (e^{f_1} + 3e^{f_2} - 3e^{f_3} + e^{f_4}) \\ &> 2(e^{f_1} + e^{f_2} - e^{f_3}) - (e^{f_3} - e^{f_4}), \quad \text{By Step 1} \\ &= 2e^{f_3}(e^{f_1-f_3} + e^{f_2-f_3} - 1) - e^{f_3}(1 - e^{-f_3+f_4}). \end{aligned}$$

Hence, at maximum of $e^{f_1-f_3} + e^{f_2-f_3}$, we have

$$2(A_2 - 1) < 1 - A_3^{-1}.$$

Hence,

$$(7) \quad B_2 = A_2 - 1 < (1 - A_3^{-1}) - (A_2 - 1) = B_3.$$

We apply the same argument:

$$\begin{aligned} &1 - A_m^{-1} \\ &= B_m + (A_{m-1} - 1) \\ &= B_m + A_{m-1}(B_{m-1} + (A_{m-2} - 1)) \end{aligned}$$

$$\begin{aligned}
 & \dots \\
 & = B_m + A_{m-1}B_{m-1} + A_{m-1}A_{m-2}B_{m-2} + \dots \\
 & \quad + A_{m-1}A_{m-2} \dots A_3B_3 + A_{m-1}A_{m-2} \dots A_3(A_2 - 1) \\
 & \quad \text{Applying Inequality (6),(7)} \\
 & < \underbrace{B_m + B_m + \dots + B_m}_{m-1 \text{ terms}} \\
 & < -2(m-1)(1 - A_m^{-1}).
 \end{aligned}$$

Therefore, $A_m < 1$.

Step 3: By applying the same process as in Lemma 5.3, we show $A_k < 1$ and finish the proof. q.e.d.

6. Comparison to decoupled equations

In this section, we compare the solution to coupled Hitchin equations by the solutions to decoupled vortex-like equations. For n even, set $n = 2m$; for n odd, set $n = 2m + 1$. Throughout this section, we assume $m \geq 2$, since for $m = 1$ the Hitchin system of equations is a single equation which is already decoupled.

Theorem 6.1. *There exists a unique Hermitian metric u_k, v_k , respectively, on the holomorphic line bundle $K^{\frac{n+1-2k}{2}}$ satisfying, locally*

$$(8) \quad \Delta \log u_k + u_k^{-\frac{2}{n+1-2k}} - (u_k^2 |q_n|^2)^{\frac{1}{2k-1}} = 0, \quad 1 \leq k \leq m,$$

$$(9) \quad \Delta \log v_k + v_k^{-\frac{2}{n+1-2k}} - (v_k^2 |2q_{n-1}|^2)^{\frac{1}{2k-2}} = 0, \quad 2 \leq k \leq m.$$

And the following estimates hold,

$$\max\{|q_n|^{\frac{2\alpha_k}{n}}, (\alpha_k g_0)^{\alpha_k}\} \leq u_k^{-1} < ((\max_{\Sigma} |q_n|_{g_0}^{\frac{2}{n}} + \alpha_k) g_0)^{\alpha_k},$$

$$\max\{|2q_{n-1}|^{\frac{2\alpha_k}{n-1}}, (\alpha_k g_0)^{\alpha_k}\} \leq v_k^{-1} < ((\max_{\Sigma} |2q_{n-1}|_{g_0}^{\frac{2}{n-1}} + \alpha_k) g_0)^{\alpha_k},$$

where $\alpha_k = \frac{n+1-2k}{2}$ and g_0 is the unique Hermitian hyperbolic metric on K^{-1} .

Remark 6.2. Such equations are natural generalization of familiar vortex-like equations. In particular, in the case $n = 2, k = 1$, this is the harmonic equation from surface to surface with given Hopf differential q_2 . In the case $n = 3, k = 1$, this gives Wang's equation [24] and the solution is the Blaschke metric for the hyperbolic affine sphere. Dumas and Wolf [7] solved such equations for the case when n is general and $k = 1$ on the complex plane.

Remark 6.3. Let $u_k = b_k g^{\frac{2k-n-1}{2}}$, then equation (8) becomes

$$\Delta_g \log b_k - \frac{2k-n-1}{2} K_g + b_k^{-\frac{2}{2(m-k)+1}} - (b_k^2 |q_n|^2)^{\frac{1}{2k-1}} = 0,$$

where $K_g = -\Delta_g \log g$ is the sectional curvature of g . In particular, there is a geometric interpretation for the equation (8). Let $\sigma_k = u_k^{-\frac{2}{n+1-2k}}$. Then σ_k is an Hermitian metric on the Riemann surface Σ . The equation for σ_k is

$$\frac{n+1-2k}{2} K_{\sigma_k} = -1 + |q_n|_{\sigma_k}^{\frac{2}{2k-1}}.$$

Proof of Theorem 6.1. Again, we only prove the case $n = 2m$ for the q_n case. The proof for other cases is similar. Take the Hermitian hyperbolic metric g_0 on Σ . Let $\sigma_k = u_k^{-\frac{2}{n+1-2k}}$ and $\sigma_k = g_0 e^{\eta_k}$ with η_k being a C^∞ function on Σ . Then equation (8) is equivalent to

$$\frac{n+1-2k}{2} \Delta_{g_0} \eta_k - e^\eta + |q_n|_{g_0}^{\frac{2}{2k-1}} e^{-\frac{n+1-2k}{2k-1} \eta_k} + \frac{n+1-2k}{2} = 0,$$

for $1 \leq k \leq m$. Then showing the first part of Theorem 6.1 reduces to show the existence and uniqueness of η_k which follows from the following lemma. The proof is similar to Proposition 4.0.2 in Loftin [16].

Lemma 6.4. *Let (M, g) be a closed Riemannian manifold, f be a nonnegative C^∞ function on M , and a, b, c be positive real numbers. Then the equation*

$$a \Delta_g \eta - e^\eta + f(x) e^{-c\eta} + b = 0, \quad 1 \leq k \leq m$$

has a unique C^∞ solution, such that

$$\log b \leq \eta < \log(G^{\frac{1}{c+1}} + b),$$

where G denotes the maximum value of $f(x)$. If f is not identically zero, then $\eta > \log b$.

Proof of Lemma 6.4. For the existence, it is sufficient to find a subsolution and a supersolution for this equation (see Schoen and Yau [22]). First, the constant function $\eta = \log b$ is a subsolution by direct calculation. Second, set m to be the smallest positive root of the equation

$$x^{c+1} - bx^c - G = 0.$$

Then $\eta = \log m$ satisfies

$$a \Delta_g \eta - e^\eta + f(x) e^{-c\eta} + b = -m + f(x) m^{-c} + b \leq 0.$$

Therefore, $s = \log m$ is a supersolution. Then there is a smooth solution η to the equation satisfying

$$\log b \leq \eta \leq \log m < \log(G^{\frac{1}{c+1}} + b).$$

If f is not zero, the strong maximum principle implies $\eta > \log b$. The uniqueness comes from the maximum principle. And by the standard elliptic theory, we obtain the smoothness of the solution. q.e.d.

Theorem 6.1 then follows from the above Lemma directly except to show $u_k \leq |q_n|^{-\frac{n+1-2k}{n}}$. In fact, we observe that $|q_n|^{-\frac{n+1-2k}{n}}$ is also a solution to equation (8) outside the zero locus of q_n . Then applying the strong maximum principle, we finish the proof. q.e.d.

We now compare the solution to the coupled Hitchin system of equations to such Hermitian metrics u_k and v_k satisfying decoupled equations.

Theorem 6.5. *For $m \geq 2$,*

$$\begin{aligned} q_n \text{ case:} \quad & h_k < u_k, \quad 1 \leq k \leq m, \\ q_{n-1} \text{ case:} \quad & h_k < v_k, \quad 2 \leq k \leq m. \end{aligned}$$

Remark 6.6. If $m = 1$, for the q_n case, then the Hitchin equation coincides with the equation for u_1 . Therefore, $h_1 = u_1$. In this sense, u_k is a natural generalization to bound the coupled Hitchin system.

Proof of Theorem 6.5. Again, we only prove the case $n = 2m$ for the q_n case. The proof for the other cases is similar. From Lemma 5.3

$$(h_k^{-1}h_{k+1})^{2(m-k)+1} < (h_k^{-1}h_{k+1})^2 \cdots (h_{m-1}^{-1}h_m)^2 \cdot h_m^{-2} = h_k^{-2},$$

hence,

$$h_k^{-1}h_{k+1} < h_k^{-\frac{2}{2(m-k)+1}}, \quad 1 \leq k \leq m-1.$$

From Lemma 5.3

$$(h_{k-1}^{-1}h_k)^{2k-1} > (h_1^2|q_{2m}|^2)(h_1^{-1}h_2)^2 \cdots (h_{k-1}^{-1}h_k)^2 = h_k^2|q_{2m}|^2,$$

hence,

$$h_{k-1}^{-1}h_k > (h_k^2|q_{2m}|^2)^{\frac{1}{2k-1}}, \quad 2 \leq k \leq m.$$

Therefore, the system of equations

$$\begin{aligned} \Delta \log h_1 + h_1^{-1}h_2 - h_1^2|q_n|^2 &= 0, \\ \Delta \log h_k + h_k^{-1}h_{k+1} - h_{k-1}^{-1}h_k &= 0, \quad 2 \leq k \leq m-1, \\ \Delta \log h_m + h_m^{-2} - h_{m-1}^{-1}h_m &= 0, \end{aligned}$$

implies that

$$\Delta \log h_k + h_k^{-\frac{2}{2(m-k)+1}} - (h_k^2|q_{2m}|^2)^{\frac{1}{2k-1}} > 0, \quad 1 \leq k \leq m.$$

Then applying the strong maximum principle, we obtain $h_k < u_k$. q.e.d.

Remark 6.7. For h_1 in the q_{n-1} case, from Lemma 5.4, one can show that $h_1 < |q_{n-1}|^{-1}$.

Now fixing q_n (or q_{n-1}), we study the asymptotic behavior of h_k, u_k, v_k along the ray tq_n (or tq_{n-1}) when t approaches to infinity. For h_k , Collier and Li [2] proved the following result.

Theorem 6.8. (Collier-Li [2]) Suppose q_n (or q_{n-1}) is not zero. Then for each compact set $K \subset \Sigma$ away from zeros of q_n (or q_{n-1}), there is a positive constant $C = C(K)$ independent of t such that at each point $p \in K$,

$$q_n \text{ case: } h_k = |q_n|^{-\frac{n+1-2k}{n}}(1 + O(\|q_n\|^{-1})), \quad 1 \leq k \leq m,$$

$$q_{n-1} \text{ case: } h_k = (|2q_{n-1}|)^{-\frac{n+1-2k}{n-1}}(1 + O(\|q_{n-1}\|^{-1})), \quad 2 \leq k \leq m,$$

where $\|q_n\| = \int_{\Sigma} |q_n|^{\frac{2}{n}}$ and $f \in O(\|q_n\|^{-1})$ means $|f|/\|q_n\|^{-1} \leq C$.

Remark 6.9. By Theorem 6.1 and Theorem 6.5, $h_k < |q_n|^{-\frac{n+1-2k}{n}}$, so we improve the above result in one direction.

Let u_k^t (or v_k^t) be the solution for tq_n (or tq_{n-1}) in Theorem 6.1 (or tq_{n-1}). Combining Theorem 6.5 and Theorem 6.8, we obtain

Theorem 6.10. Suppose q_n (or q_{n-1}) is not zero. Then for each compact set $K \subset \Sigma$ away from zeros of q_n (or q_{n-1}), there is a positive constant $C = C(K)$ independent of t such that

$$q_n \text{ case: } u_k^t(1 - C\|tq_n\|^{-1}) \leq h_k^t < u_k^t, \quad 1 \leq k \leq m,$$

$$q_{n-1} \text{ case: } v_k^t(1 - C\|tq_{n-1}\|^{-1}) \leq h_k^t < v_k^t, \quad 2 \leq k \leq m.$$

Remark 6.11. The theorem above shows that u_k (or v_k) is an upper approximation of h_k .

Finally, we prove the following asymptotic estimates of u_k, v_k . It can be derived from Theorem 6.1, Theorem 6.5 and Theorem 6.8. Here we give a direct proof not relying on Theorem 6.8.

Theorem 6.12. Suppose q_n (or q_{n-1}) is not zero. Then for each compact set $K \subset \Sigma$ away from zeros of q_n (or q_{n-1}), there is a positive constant $C = C(K)$ independent of t such that

$$|tq_n|^{-\frac{n+1-2k}{n}}(1 - C\|tq_n\|^{-1}) \leq u_k^t < |tq_n|^{-\frac{n+1-2k}{n}}, \quad 1 \leq k \leq m,$$

$$|2tq_{n-1}|^{-\frac{n+1-2k}{n-1}}(1 - C\|tq_{n-1}\|^{-1}) \leq v_k^t < |2tq_{n-1}|^{-\frac{n+1-2k}{n-1}}, \quad 2 \leq k \leq m.$$

Proof of Theorem 6.12. For simplicity, we drop the superscript t of u_k^t, v_k^t . We only prove $u_k \geq |tq_n|^{-\frac{n+1-2k}{n}}(1 - C\|tq_n\|^{-1})$. Let K be a compact subset of Σ which does not contain any zeros of q_n . Choose a background metric g on the surface Σ defined as follows: $g = \frac{|q_n|^{\frac{2}{n}}}{\|q_n\|}$ on K and $g \geq \frac{|q_n|^{\frac{2}{n}}}{\|q_n\|}$ outside K . Let $u_k = b_k g^{\frac{2k-n-1}{2}}$, where b_k is a positive function on Σ satisfying

$$\Delta_g \log b_k - \frac{2k-n-1}{2} K_g + b_k^{-\frac{2}{n+1-2k}} - (b_k^2 |tq_n|_g^2)^{\frac{1}{2k-1}} = 0.$$

At the minimum of b_k ,

$$\frac{n+1-2k}{2}K_g + b_k^{-\frac{2}{n+1-2k}} - (b_k^2|tq_n|_g^2)^{\frac{1}{2k-1}} \leq 0.$$

Let $M = \max_{\Sigma} \frac{n+1-2k}{2}|K_g|$ and, hence,

$$-M + b_k^{-\frac{2}{n+1-2k}} \leq (b_k^2|tq_n|_g^2)^{\frac{1}{2k-1}} \leq b_k^{\frac{2}{2k-1}} \|tq_n\|^{\frac{n}{2k-1}}.$$

Let $x = \|tq_n\| b_k^{\frac{2}{n+1-2k}}$, then x satisfies

$$x^{\frac{n}{2k-1}} + \frac{M}{\|tq_n\|}x - 1 \geq 0.$$

Therefore, $\|tq_n\| b_k^{\frac{2}{n+1-2k}} = x \geq 1 - C\|tq_n\|^{-1}$ at the minimum of b_k , also the minimum of x .

Hence, globally on the surface, $\|tq_n\| b_k^{\frac{2}{n+1-2k}} = x \geq 1 - C\|tq_n\|^{-1}$. Then

$$\begin{aligned} u_k &= (x\|tq_n\|^{-1})^{\frac{n+1-2k}{2}} g^{-\frac{n+1-2k}{2}} \\ &\geq x^{\frac{n+1-2k}{2}} |tq_n|^{-\frac{n+1-2k}{n}} \\ &\geq |tq_n|^{-\frac{n+1-2k}{n}} (1 - C\|tq_n\|^{-1}). \end{aligned} \quad \text{q.e.d.}$$

Remark 6.13. It is an interesting question that how C varies when K approaches to $\Sigma - Z$, where Z is the set of zeros of q_n .

References

- [1] D. Baraglia, *G₂ Geometry and integrable system*, thesis, arXiv:1002.1767v2, 2010.
- [2] B. Collier and Q. Li, *Asymptotics of Higgs bundles in the Hitchin component*, Adv. Math. 307 (2017), 488–558, MR3590524, Zbl 06670884.
- [3] B. Collier, *Finite order automorphism of Higgs bundles: theory and application*, thesis, 2016.
- [4] K. Corlette, *Flat G-bundles with canonical metrics*, J. Diff. Geom. 28 (1988), no. 3, 361–382, MR965220, Zbl 0676.58007.
- [5] S. K. Donaldson, *Twisted harmonic maps and the self-duality equations*, Proc. London Math. Soc. (3) 55 (1987), no. 1, 127–131, MR0887285, Zbl 0634.53046.
- [6] B. Deroin and N. Tholozan, *Dominating surface group representations by Fuchsian ones*, Int. Math. Res. Not. IMRN 2016, no. 13, 4145–4166, MR3544632.
- [7] D. Dumas and M. Wolf, *Polynomial cubic differentials and convex polygons in the projective plane*, Geom. Funct. Anal. 25 (2015), no. 6, 1734–1798, MR3432157, Zbl 1335.30013.
- [8] F. Guéritaud, F. Kassel and M. Wolff, *Compact anti-de Sitter 3-manifolds and folded hyperbolic structures on surfaces*, Pacific J. Math. 275 (2015), no. 2, 325–359, MR3347373, Zbl 1346.57005.
- [9] N. J. Hitchin, *The self-duality equations on a Riemann surface*, Proc. London Math. Soc. (3) 55 (1987), no. 1, 59–126, MR0887284, Zbl 0634.53045.

- [10] N. J. Hitchin, *Lie groups and Teichmüller space*, Topology 31 (1992), no. 3, 449–473, MR1174252, Zbl 0769.32008.
- [11] J. Jost, *Partial differential equations*. Second edition. Graduate Texts in Mathematics, 214. Springer, New York, 2007.
- [12] F. Labourie, *Flat projective structures on surfaces and cubic holomorphic differentials*, Pure Appl. Math. Q. 3 (2007), no. 4, Special Issue: In honor of Grigory Margulis. Part 1, 1057–1099. MR2402597, Zbl 1158.32006.
- [13] F. Labourie, *Cross ratios, Anosov representations and the energy functional on Teichmüller space*, Ann. Sci. Éc. Norm. Supér. (4) 41 (2008), no. 3, 437–469. MR2482204, Zbl 1160.37021.
- [14] F. Labourie, *Cyclic surfaces and Hitchin components in rank 2*, Ann. of Math. (2) 185 (2017), no. 1, 1–58, MR3583351, Zbl 06686583.
- [15] G. Lee and T. Zhang, *Collar lemma for Hitchin representations*, Geom. Topol. 21 (2017), no. 4, 2243–2280. MR3654108, Zbl 1367.57010.
- [16] J. Loftin, *Affine spheres and convex RP^n -manifolds*, Amer. J. Math. 123 (2001), no. 2, 255–274. MR1828223, Zbl 0997.53010.
- [17] T. Mochizuki, *Asymptotic behavior of certain families of harmonic bundles on Riemann surfaces*, J. Topol. 9 (2016), no. 4, 1021–1073. MR3620459, Zbl 06761416.
- [18] X. Nie, *Entropy degeneration of convex projective surfaces*, Conform. Geom. Dyn. 19 (2015), 318–322, MR3432325, Zbl 1335.52004.
- [19] R. Potrie and A. Sambarino, *Eigenvalues and entropy of a Hitchin representation*, Invent. Math. 209 (2017), no. 3, 885–925. MR3681396, Zbl 06786973.
- [20] A. Sanders, *Hitchin harmonic maps are immersions*, arXiv:1407.4513, 2014.
- [21] C. Simpson, *Constructing variations of Hodge structure using Yang–Mills theory and applications to uniformization*, J. Amer. Math. Soc. 1 (1988), no. 4, 867–918. MR0944577, Zbl 0669.58008.
- [22] R. Schoen and S. T. Yau, *Lectures on differential geometry*, International Press, 1994.
- [23] N. Tholozan, *Entropy of Hilbert metrics and length spectrum of Hitchin representations in $PSL(3, \mathbb{R})$* , Duke Math. J. 166 (2017), no. 7, 1377–1403. MR3649358, Zbl 1375.53022.
- [24] C. P. Wang, *Some examples of complete hyperbolic affine 2-Spheres in \mathbb{R}^3* , Global differential geometry and global analysis (Berlin, 1990), 271–280, Lecture Notes in Math., 1481, Springer, Berlin, 1991, MR1178538, Zbl 0743.53004.
- [25] T. Zhang, *The degeneration of convex $\mathbb{R}P^2$ structures on surfaces*, Proc. Lond. Math. Soc. (3) 111 (2015), no. 5, 967–1012, MR3477227, Zbl 1351.30029.
- [26] T. Zhang, *Degeneration of Hitchin representations along internal sequences*, Geom. Func. Anal. 25 (2015), no. 5, 1588–1645, MR3426063, Zbl 1327.30054.

CENTER FOR APPLIED MATHEMATICS AT TIANJIN UNIVERSITY
 TIANJIN UNIVERSITY
 NO.92 WEIJINLU NANKAI DISTRICT
 TIANJIN
 P.R. CHINA 300072
E-mail address: song.dai@tju.edu.cn

CENTRE FOR QUANTUM GEOMETRY OF MODULI SPACES (QGM)

DEPARTMENT OF MATHEMATICS, AARHUS UNIVERSITY
NY MUNKEGADE 118, BLDG. 1530
8000 AARHUS C
DENMARK

DEPARTMENT OF MATHEMATICS
CALIFORNIA INSTITUTE OF TECHNOLOGY
1200 EAST CALIFORNIA BOULEVARD
PASADENA, CA 91125
USA

E-mail address: qionglng.li@gmail.com