



# On Generalized Type 1 Logistic Distribution

Mohammad Ahsanullah

Department of Management Sciences, Rider University Lawrenceville, New Jersey, 08648, USA

Received: October 02, 2013; Accepted: 15 November 15, 2013

Copyright © 2013, Afrika Statistika. All rights reserved

**Abstract.** Some distributional properties of the generalized type 1 logistic distribution are given. Based on these distributional property a characterization of this distribution is presented.

**Résumé.** Quelques propriétés relatives à la loi de distribution de variables aléatoires suivant la loi logistique généralisée de Type 1 sont données. Nous en déduisons une caractérisation de ce type de distribution.

**Key words:** Conditional Expectation; Reversed Hazard Rate; Characterization.

**AMS 2010 Mathematics Subject Classification :** 62E10, 6215, 62E20.

---

## 1. Introduction

We say that an absolutely continuous random variable (rv) has a generalized type I logistic distribution ( $GLD_1(\alpha)$ ) if its cumulative distribution function (*cdf*)  $F(x)$  is as follows:

$$F(x) = \frac{1}{(1 + e^{-x})^\alpha} \quad -\infty < x < \infty, \alpha > 0. \quad (1)$$

The corresponding probability density function (*pdf*) is given by

$$f(x) = \frac{\alpha e^{-x}}{(1 + e^{-x})^{\alpha+1}}. \quad (2)$$


The generalized logistic distribution is also called exponentiation logistic distribution. If  $\alpha = 1$ , then generalized logistic distribution is the standard logistic distribution (SLD). The standard logistic distribution is similar in shape to normal distribution. It is symmetric around zero and bell shaped. The pdf of  $GLD_1(\alpha)$  is an increasing function for  $x < \ln \alpha$  and it is a decreasing function for  $x > \ln \alpha$ . It is unimodal with mode at  $\ln \alpha$ . It is log-concave

---

for all values of  $\alpha$ . and  $X - \ln \alpha$  behaves like a type I extreme value( maximum) distribution. The Generalized logistic distribution can be either left or right skewed (when parameter  $\alpha$  is less than 1 or greater than 1 respectively) or symmetric ( $\alpha = 1$ ).  $GLD_1(\alpha)$  has been used in growth models and modelling of data. In this paper we will present some distributional property of  $GLD_1(\alpha)$  and a characterization of it based on conditional expectation of  $X$ .

## 2. Main Results

The hazard rate of  $GLD_1(\alpha)$  can be bathtub type or an increasing function depending on  $\alpha$ . The reversed hazard rate (RHR)  $\tau(x) (= f(x)/F(x))$  is  $\tau(x) = \alpha/(1 + e^x)$ .  $\tau(x)$  is monotonically decreasing from  $\infty$  to 0 as  $x$  increases from  $-\infty$  to  $\infty$ . The following is the graph of  $(\tau(x)/\alpha)$ .



**Fig. 1.**  $\tau(x)/\alpha$ .

Suppose  $Y$  is distributed as a power function distribution with  $f(y) = \alpha y^{\alpha-1}$ ,  $0 < y < 1$ ,  $\alpha > 0$ , then  $X = \ln \frac{Y}{1-Y}$  is distributed as type I generalized logistic distribution. Let  $\varphi(t)$  be the characteristic function of  $GLD_1(\alpha)$ , then

$$\varphi(t) = \mathbb{E}(e^{itX}) = \int_{-\infty}^{\infty} e^{-(1-it)x} (1 + e^{-x})^{-(\alpha+1)} dx = \frac{\Gamma(1-it)\Gamma(\alpha+it)}{\Gamma(\alpha)}. \quad (3)$$

We can easily calculate the mean, variance and other moments from  $\varphi(t)$

$$\mathbb{E}(X) = \Psi(\alpha) - \Psi(1) = \sum_{k=1}^{\alpha} \frac{1}{k}, \quad (4)$$

if  $\alpha$  is an integer.


$$Var(x) = \Psi'(x) + \Psi'(1)$$

where  $\Psi(x)$  and  $\Psi'(x)$  are the digamma and the polygamma function respectively. The mean is an increasing function of  $\alpha$ . The mean increases to  $\infty$  and the variance decreases to  $\Psi'(1)(= \frac{\pi^2}{6})$  as  $\alpha \rightarrow \infty$ . The percentile points,  $x_p = -\ln(p^{-1/\alpha} - 1)$  and the median,  $M = -\ln(2^{1/\alpha} - 1)$ . Table 1 gives the mean, median and quartile points of  $GLD_1(1)$ .

$\alpha$	Mean	first quartile	Mode	Median	Third quartile
0.5	-1.3863	-2.7081	-0.69315	-1.0986	0.25131
1	0	-1.9086	0	0	1.0986
5	2.0833	1.1410	1.6094	1.9058	2.8264
10	2.8290	1.9058	2.3026	2.6342	3.5341
15	3.2516	2.3349	2.7081	3.0514	3.9443
25	3.7760	2.8644	3.2189	3.5715	4.4590
50	4.4792	3.5715	3.9120	4.2716	5.1550
100	5.1774	4.2718	4.6052	4.9683	5.8496

**Table 1.**

From the table we see that mean is less (greater) than median for  $\alpha < 1(\alpha > 1)$ . For  $\alpha = 1$ , mean and median coincide. The mean is always less than the mode for all  $\alpha \neq 1$ . The graph of  $y = Var(X)$  is given in Figure 2.



**Fig. 2.**  $Var(X)$ .

From Figure 2, it is evident that  $Var(X)$  decrease to  $\frac{\pi^2}{6}$  as  $\alpha \rightarrow \infty$ .

Let  $X_1, X_2, \dots, X_n$  be  $n$  independent and identically distributed random variables. Suppose  $X_{1,n} \leq \dots \leq X_{n,n}$  be the corresponding order statistics. [Ahsanullah et al. \(2013\)](#) gave some distributional properties and characterizations of logistic distribution ( $GD$ ). They gave the following the characteristic function,  $\phi_{k,n}(t)$  (Result 1 of  $X_{r,n}$ ,  $1 \leq k \leq n$  when  $X$ 's are SLD). Here we give in Theorem 1 a generalization of the above result for  $GLD_1$ .

$$\phi_{k,n}(t) = \frac{B(k + it, n - k - it + 1)}{B(k, n - k + 1)},$$

where

$$B(p, q) = \frac{\Gamma(p)\Gamma(q)}{\Gamma(p + q)}, \quad p, q > 0.$$

**Theorem 1.** *If  $X$ 's are from  $GDL_1(\alpha)$  with cdf as given in (3) and  $\Psi_{k,n}(t)$  be the characteristic function of  $X_{k,n}$ , then.*

$$\Psi_{k,n}(t) = \frac{\alpha}{B(k, n - k + 1)} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^j B(1 - it, \alpha(n + j) + it)$$

*Proof.* The pdf  $f_{k,n}(x)$  of a  $k$ th order statistics from cdf  $F(x)$  and pdf  $f(x)$  is (see [Ahsanullah et al., 2013](#))

$$f_{k,n}(x) = C_{k,n}(F(x))^{k-1}(1 - F(x))^{n-k} f(x),$$

where  $C_{r,n} = n!/(k - 1)!(n - k)!$ .

Substituting the pdf of  $GLD_1(\alpha)$  from (3), we obtain

$$\begin{aligned} \Psi_{k,n}(t) &= C_{k,n} \int_{-\infty}^{\infty} e^{itx} \left( \frac{1}{(1 + e^{-x})^\alpha} \right)^{k-1} \left( 1 - \frac{1}{(1 + e^{-x})^\alpha} \right)^{n-k} \frac{\alpha e^{-x}}{(1 + e^{-x})^{\alpha+1}} dx \\ &= C_{k,n} \int_{-\infty}^{\infty} \sum_{j=0}^{n-k} (-1)^{n-k-j} \binom{n-k}{j} e^{itx} \left( \frac{1}{(1 + e^{-x})^\alpha} \right)^{n-j} \frac{\alpha e^{-x}}{1 + e^{-x}} dx \\ &= C_{k,n} \alpha \int_0^1 \sum_{j=0}^{n-k} (-1)^{n-k-j} \binom{n-k}{j} (u)^{\alpha(n-j)+it-1} (1 - u)^{-it} du \\ &= \frac{\alpha}{B(k, n - k + 1)} \sum_{j=0}^{n-k} \binom{n-k}{j} (-1)^{n-k-j} B(1 - it, \alpha(n - j) + it). \end{aligned}$$

□

**Remark 1.** For  $k = n$ ,  $\Psi_{n,n}(t) = \frac{\alpha B(n\alpha + it, 1 - it)}{B(n, 1)}$ . It is known (see [Ahsanullah and Kirmani, 2000](#), page 10) that if for a continuous random variable with cdf  $F(x)$ , the inverse function  $F^{-1}$  satisfies the condition

$$\lim_{c \rightarrow 0} \frac{F^{-1}(1 - c) - F^{-1}(1 - 2c)}{F^{-1}(1 - 2c) - F^{-1}(1 - 4c)} = 1,$$

then  $F$  belongs to the domain of attraction of the type 1 (extreme) value distribution. For  $GLD_1(\alpha)$ ,

$$F^{-1}(x) = -\ln(x^{-1/\alpha} - 1)$$

and

$$\lim_{c \rightarrow 0} \frac{F^{-1}(1-c) - F^{-1}(1-2c)}{F^{-1}(1-2c) - F^{-1}(1-4c)} = 1$$

Thus  $GLD_1(\alpha)$  belongs to the domain of attraction of the type  $I$  extreme(maximum) distribution;

$$P(X_{n,n} \leq x + \ln(n\alpha)) = \left(\frac{1}{1 + e^{-(x+n\alpha)}}\right)^{n\alpha} = \left(1 + \frac{e^{-x}}{n\alpha}\right)^{-n\alpha} \rightarrow e^{-e^{-x}}$$

as  $n \rightarrow \infty$ .

The following theorem give a characterization of  $GLD_1(\alpha)$ .

**Theorem 2.** *Suppose  $X$  is an absolutely continuous (with respect to Lebesgue measure) random variable with cdf  $F(x)$  and pdf  $f(x)$ . We assume that  $E(X)$  exists. Then  $X$  has the  $GLD_1(\alpha)$  distribution if and only if*

$$\mathbb{E}(X|X \leq x) = g(x)\tau(x),$$

where

$$\tau(x) = \frac{f(x)}{F(x)}, \quad g(x) = \frac{x(1 + e^x)}{\alpha} - I_\alpha(x)$$

and

$$I_\alpha(x) = \frac{e^x(1 + e^{-x})^{\alpha+1}}{\alpha} \int_{-\infty}^x (1 + e^{-u})^{-\alpha} du, \alpha > 0$$

for all  $x$ ,  $-\infty < x < \infty$ .

To proof the Theorem, we need the following lemma.

**Lemma 1.** *Suppose  $X$  is an absolutely continuous (with respect to Lebesgue measure) random variable with cdf  $F(x)$  and pdf  $f(x)$ . We assume that  $E(X)$  exists. Then if  $\mathbb{E}(X|X \leq x) = g(x)\tau(x)$  where  $\tau(x) = f(x)/F(x)$ , then*

$$f(x) = f(x) = ce^{\int \frac{x-g'(x)}{g(x)} dx}.$$

*Proof.* We have

$$g(x) = \frac{\mathbb{E}(X|X \leq x)}{\tau(x)} = \frac{\int_{-\infty}^x uf(u)du}{f(x)} = \frac{x F(x)}{f(x)} + \frac{\int_{-\infty}^x F(u)du}{f(x)}. \quad (5)$$

On simplification, we obtain from (5).

$$xf(x) + \int_{-\infty}^x F(u)du = f(x)g(x). \quad (6)$$

On differentiating both sides of (6) with respect to  $x$ , we have  $f(x) + xf'(x) - f(x) = f(x)g'(x) + f'(x)g(x)$  i.e.

$$\frac{f'(x)}{f(x)} = \frac{x - g'(x)}{g(x)}. \tag{7}$$

On integrating, we obtain from (7),

$$f(x) = ce^{\int \frac{x-g'(x)}{g(x)} dx} \tag{8}$$

where the constant  $C$  is determined by using the boundary condition  $F(-\infty) = 0$ , and  $F(\infty) = 1$ .  $\square$

**Proof of Theorem 2.**

Suppose that

$$f(x) = \frac{\alpha e^{-x}}{(1 + e^{-x})^{\alpha+1}}$$

then

$$\begin{aligned} g(x) &= \frac{\mathbb{E}(X|X \leq x)}{f(x)} \\ &= \frac{\int_{-\infty}^x \frac{\alpha e^{-u}}{(1+e^{-u})^{\alpha+1}} du}{\frac{\alpha e^{-x}}{(1+e^{-x})^{\alpha+1}}} = \frac{x e^x (1 + e^{-x})}{\alpha} - \frac{\int_{-\infty}^x \frac{1}{(1+e^{-u})^\alpha} du}{\frac{\alpha e^{-x}}{(1+e^{-x})^{\alpha+1}}} \\ &= \frac{x(1 + e^x)}{\alpha} - I_\alpha(x) \end{aligned}$$

where

$$I(x) = \frac{e^x(1 + e^{-x})^{\alpha+1}}{\alpha} \int_{-\infty}^x (1 + e^{-u})^{-\alpha} du'.$$

Suppose

$$g(x) = \frac{x(1 + e^x)}{\alpha} - I_\alpha(x)$$

then

$$\begin{aligned} g'(x) &= \frac{1 + e^{-x}}{\alpha} + \frac{x e^x}{\alpha} - \frac{e^x(1 + e^{-x})^{\alpha+1}}{\alpha} (1 + e^{-x})^{-\alpha} - I_\alpha(x) \left\{ 1 - (\alpha + 1) \frac{-e^{-x}}{1 + e^{-x}} \right\} \\ &= \frac{x e^x}{\alpha} + \frac{e^x(1 + e^{-x})^\alpha}{\alpha} I_\alpha(x) (1 - \alpha e^{-x}). \end{aligned}$$

We have now

$$\begin{aligned} x - g'(x) &= \frac{\alpha - e^x}{\alpha} x - I_\alpha(x) \frac{\alpha - e^x}{1 + e^{-x}} \\ &= \frac{\alpha - e^x}{1 + e^x} \left[ \frac{x(1 + e^x)}{\alpha} - I_\alpha(x) \right] \\ &= \frac{\alpha - e^x}{1 + e^x} g(x). \end{aligned}$$

Thus

$$\frac{f'(x)}{f(x)} = \frac{x - g'(x)}{g(x)} = \frac{\alpha - e^x}{1 + e^x}. \quad (9)$$

On integrating, we obtain from (9)

$$f(x) = ce^{\int \frac{x-e^x}{1+e^x} dx} = c \frac{e^{-x}}{(1 + e^{-x})^{\alpha+1}}. \quad (10)$$

Using the condition  $\int_{-\infty}^{\infty} f(x)dx = 1$ , we obtain

$$f(x) = \frac{\alpha e^{-x}}{(1 + e^{-x})^{\alpha+1}}. \quad (11)$$

□

### References

- Ahsanullah, M. and Kirmani, S.N.U.A., 2008. *Topics in Extreme Value*. Nova Publishers, New-York, USA.
- Ahsanullah, M., Nevzorov, V.B. and Shakil, M., 2013. *An Introduction to Order Statistics*. Atlantis Press, Paris, France.
- Ahsanullah, M., Yanev, G.P and Onica, C., 2012. Characterizations of Logistic Distribution through Order Statistics with Independent Exponential Shift. *Economic Quality Control.*, **27**. 85-96.
- Lin, G.D. and Hu, C.Y., 2008. On characterizations of the logistic distribution. *J. Statistical Planning. Inference*, **138**. 1147-1156.