

THE LIGHTCONE GAUSS MAP OF A SPACELIKE SURFACE IN MINKOWSKI 4-SPACE *

SHYUICHI IZUMIYA[†], DONGHE PEI[‡], AND MARÍA DEL CARMEN ROMERO FUSTER[§]

Abstract. We study the geometry of the spacelike surfaces in Minkowski 4-space through their generic contact with lightlike hyperplanes.

1. Introduction. Inspired in the results obtained by Kossowski [10, 11] for submanifolds of Minkowski space, we introduced in [7] the notion of lightcone Gauss map of a spacelike curve in Minkowski 3-space. We studied the singularities of lightcone Gauss maps of spacelike curves and established the relationships between such singularities and the geometric invariants of these curves under the action of Lorentz group. Our aim in this paper is to develop the analogous theory for spacelike surfaces in Minkowski 4-space. To do this we need to work out local differential geometry tools for spacelike surfaces in Minkowski 4-space similar to those of surfaces in Euclidean 4-space [12]. As it was to be expected, the situation presents certain peculiarities when compared with the Euclidean case. For instance, it is always possible to choose, in our case, two lightlike normal directions along the surface as a frame of its normal bundle. By using this, we define a Lorentzian invariant, $\mathcal{K}_l(1, \pm 1)$, called *lightlike Gaussian curvature* of the spacelike surface. We introduce in §3 the notion of lightcone height function which is useful to show that the lightcone Gauss map has a singular point if and only if the lightlike Gaussian curvature vanishes at such point. Moreover, we show that the lightcone Gauss map is a constant map if and only if the surface is contained in a lightlike hyperplane, so we can view the singularities of the lightcone Gauss map as an estimate of the contacts of the surface with lightlike hyperplanes. In §4 we define the notion of lightcone pedal surface of a spacelike surface and show that its singularities are in correspondence with that of the lightcone Gauss map of the surface. We apply Montaldi's methods [16] to the extended lightcone height function in order to study the generic contacts between spacelike surfaces and lightlike hyperplanes in §5. Finally, in §6, we give a generic classification of lightcone pedal surfaces and lightcone Gauss maps (cf., Theorem 6.1 and Corollary 6.3). Briefly speaking, generic singularities of lightcone pedal surfaces are cuspidaledges and swallowtails (cf., Fig. 1) and the corresponding singularities for lightcone Gauss maps are folds and cusps respectively (cf., Fig. 2). We also analyze the geometrical meaning of cuspidaledges, cross caps and swallowtails of lightcone pedal surfaces.

We observe that any surface M in Euclidean space \mathbb{R}^3 can be viewed as a spacelike surface in \mathbb{R}_1^4 . In this case the lightlike Gaussian curvature is the (Euclidean) Gaussian curvature of the surface. In this sense, our theory is a generalization of the theory of cusps of Gauss maps developed in [2]. On the other hand, the notion of hyperbolic Gauss maps for surfaces in the Poincaré disk was introduced in [5]. The singularities of the hyperbolic Gauss maps for hypersurfaces in Hyperbolic n -space

*Received April 6, 2002; accepted for publication February 12, 2004. Work partially supported by DGCYT grant no. BFM2000-1110.

[†]Department of Mathematics, Hokkaido University, Sapporo 060-0810, Japan (izumiya@math.sci.hokudai.ac.jp).

[‡]Department of Mathematics, Northeast Normal University, Changchun 130024, P.R. China (northlcd@public.cc.jl.cn, pei@math.sci.hokudai.ac.jp).

[§] Departament de Geometria i Topologia, Universitat de València, 46100 Burjassot (València), Espanya (carmen.romero@post.uv.es).




Fig. 1.




Fig. 2.

$H_+^n(-1)$ were studied in [7] under the framework of Minkowski model. The hypersurfaces in Hyperbolic n -space $H_+^n(-1)$ are a particular case of spacelike submanifolds of codimension 2 of Minkowski $(n+1)$ -space \mathbb{R}_1^{n+1} . We remark that in the case $n=3$ the hyperbolic Gaussian curvature is equal to the lightlike Gaussian curvature and hence our theory is also a generalization of the singularity theory for hyperbolic Gauss maps for surfaces. We give in §7 some examples of spacelike surfaces of 4-dimensional Minkowski space that are not included in any of these two categories.

Motivated by physical applications [6] and the previous work of Kossowski [10, 11] we are preparing a further article dealing with singularities of lightlike hypersurfaces in Minkowski 4-space [9] in which we shall push forward the topics treated in the present paper.

We shall assume throughout the whole paper that all the maps and manifolds are C^∞ unless the contrary is explicitly stated.

2. Local differential geometry of spacelike surfaces. In this section we introduce the basic geometrical tools for the study of spacelike surfaces in Minkowski 4-space in an analogous way to the theory developed in [12] for surfaces in Euclidean

4-space.

Let $\mathbb{R}^4 = \{(x_1, x_2, x_3, x_4) | x_1, x_2, x_3, x_4 \in \mathbb{R}\}$ be a 4-dimensional vector space. For any vectors $\mathbf{x} = (x_1, x_2, x_3, x_4)$ $\mathbf{y} = (y_1, y_2, y_3, y_4)$ in \mathbb{R}^4 , the *pseudo scalar product* of \mathbf{x} and \mathbf{y} is defined to be $\langle \mathbf{x}, \mathbf{y} \rangle = -x_1y_1 + x_2y_2 + x_3y_3 + x_4y_4$. We call $(\mathbb{R}^4, \langle \cdot, \cdot \rangle)$ a *Minkowski 4-space*. We write \mathbb{R}_1^4 instead of $(\mathbb{R}^4, \langle \cdot, \cdot \rangle)$.

We say that a vector \mathbf{x} in $\mathbb{R}_1^4 \setminus \{\mathbf{0}\}$ is *spacelike*, *lightlike* or *timelike* if $\langle \mathbf{x}, \mathbf{x} \rangle > 0, = 0$ or < 0 respectively. The norm of the vector $\mathbf{x} \in \mathbb{R}_1^4$ is defined by $\|\mathbf{x}\| = \sqrt{|\langle \mathbf{x}, \mathbf{x} \rangle|}$. For a lightlike vector $\mathbf{n} \in \mathbb{R}_1^4$ and a real number c , we define the *lightlike hyperplane* with pseudo normal \mathbf{n} by

$$LHP(\mathbf{n}, c) = \{\mathbf{x} \in \mathbb{R}_1^4 | \langle \mathbf{x}, \mathbf{n} \rangle = c\}.$$

Let \mathbb{R}_1^4 be an oriented and timelike oriented space (i.e., a 4-volume form dV , and future time-like vector field, have been chosen), and $\mathbf{X} : U \rightarrow \mathbb{R}_1^4$ a regular surface (i.e., an immersion), where $U \subset \mathbb{R}^2$ is an open subset. We denote by $M = \mathbf{X}(U)$ and identify M and U by the immersion \mathbf{X} .

We say that M is a *spacelike surface* if the tangent plane T_pM of M is a spacelike plane (i.e., consists of spacelike vectors) for any point $p \in M$. In this case, the normal space N_pM is a timelike plane (i.e., Lorentz plane) (cf.,[18]). Let $\{\mathbf{e}_3(x, y), \mathbf{e}_4(x, y); p = (x, y)\}$ be an orthonormal frame of T_pM and $\{\mathbf{e}_1(x, y), \mathbf{e}_2(x, y); p = (x, y)\}$ a pseudo-orthonormal frame of N_pM . Here, $\mathbf{e}_1(p)$ is a timelike vector and $\mathbf{e}_i; i = 2, 3, 4$ are spacelike vectors.

We shall now establish the fundamental formula for a spacelike surface in \mathbb{R}_1^4 by means of similar notions to those of Little [12].

We can write $d\mathbf{X} = \sum_{i=1}^4 \omega_i \mathbf{e}_i$ and $d\mathbf{e}_i = \sum_{j=1}^4 \omega_{ij} \mathbf{e}_j; i = 1, 2, 3, 4$. where ω_i and ω_{ij} are 1-forms given by $\omega_i = \delta(\mathbf{e}_i) \langle d\mathbf{X}, \mathbf{e}_i \rangle$ and $\omega_{ij} = \delta(\mathbf{e}_j) \langle d\mathbf{e}_i, \mathbf{e}_j \rangle$, with

$$\delta(\mathbf{e}_i) = \text{Sign}(\mathbf{e}_i) = \begin{cases} 1 & i = 2, 3, 4 \\ -1 & i = 1 \end{cases}.$$

We have the Codazzi type equations:

$$\begin{cases} d\omega_i = \sum_{j=1}^4 \delta(\mathbf{e}_i) \delta(\mathbf{e}_j) \omega_{ij} \wedge \omega_j \\ d\omega_{ij} = \sum_{k=1}^4 \omega_{ik} \wedge \omega_{kj}, \end{cases} \tag{1}$$

where d is exterior derivative.

In fact, we have

$$\begin{aligned} d\omega_i &= \delta(\mathbf{e}_i) \langle d^2 \mathbf{X}, \mathbf{e}_i \rangle - \delta(\mathbf{e}_i) d\mathbf{X} \wedge d\mathbf{e}_i = -\delta(\mathbf{e}_i) \left(\sum_{j=1}^4 \omega_j \mathbf{e}_i \wedge d\mathbf{e}_i \right) \\ &= \sum_{j=1}^4 -\delta(\mathbf{e}_i) (\omega_j \wedge \langle d\mathbf{e}_i, \mathbf{e}_j \rangle) = \sum_{j=1}^4 -\delta(\mathbf{e}_i) (\omega_j \wedge \omega_{ij}) \delta(\mathbf{e}_j) \\ &= \sum_{j=1}^4 \delta(\mathbf{e}_i) \delta(\mathbf{e}_j) \omega_{ij} \wedge \omega_j \end{aligned}$$

and

$$\begin{aligned}
d\omega_{ij} &= \delta(\mathbf{e}_j)\langle d^2\mathbf{e}_i, \mathbf{e}_j \rangle - \delta(\mathbf{e}_j)d\mathbf{e}_i \wedge d\mathbf{e}_j = -\delta(\mathbf{e}_j)d\mathbf{e}_i \wedge d\mathbf{e}_j \\
&= -\delta(\mathbf{e}_j)\left(\sum_{k=1}^4 \omega_{ik}\mathbf{e}_k \wedge \sum_{k=1}^4 \omega_{jk}\mathbf{e}_k\right) \\
&= -\delta(\mathbf{e}_j)\delta(\mathbf{e}_k)\sum_{k=1}^4 -\delta(\mathbf{e}_j)\delta(\mathbf{e}_k)\omega_{ik} \wedge \omega_{kj} \\
&= \sum_{k=1}^4 \omega_{ik} \wedge \omega_{kj}.
\end{aligned}$$

Since $\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \delta_{ij}\delta(\mathbf{e}_j)$ (where δ_{ij} is Kronecker's delta), we get

$$\langle d\mathbf{e}_i, \mathbf{e}_j \rangle + \langle \mathbf{e}_i, d\mathbf{e}_j \rangle = 0,$$

that is $\delta(\mathbf{e}_j)\omega_{ij} + \delta(\mathbf{e}_i)\omega_{ji} = 0$. And thus

$$\omega_{ij} = -\delta(\mathbf{e}_i)\delta(\mathbf{e}_j)\omega_{ji}. \quad (2)$$

In particular, $\omega_{ii} = 0$; $i = 1, 2, 3, 4$.

It follows from the fact $\langle d\mathbf{X}, \mathbf{e}_1 \rangle = \langle d\mathbf{X}, \mathbf{e}_2 \rangle = 0$ that

$$\omega_1 = \omega_2 = 0. \quad (3)$$

Therefore we have

$$\begin{cases} 0 = d\omega_1 = \sum_{j=1}^4 \delta(\mathbf{e}_1)\delta(\mathbf{e}_j)\omega_{1j} \wedge \omega_j = \sum_{j=3}^4 \delta(\mathbf{e}_j)\omega_{1j} \wedge \omega_j = -\omega_{13} \wedge \omega_3 - \omega_{14} \wedge \omega_4 \\ 0 = d\omega_2 = \sum_{j=1}^4 \delta(\mathbf{e}_2)\delta(\mathbf{e}_j)\omega_{2j} \wedge \omega_j = \sum_{j=3}^4 \delta(\mathbf{e}_j)\omega_{2j} \wedge \omega_j = \omega_{23} \wedge \omega_3 + \omega_{24} \wedge \omega_4. \end{cases}$$

By Cartan's lemma, we can write

$$\begin{cases} \omega_{13} = a\omega_3 + b\omega_4, \quad \omega_{14} = b\omega_3 + c\omega_4 \\ \omega_{23} = e\omega_3 + f\omega_4, \quad \omega_{24} = f\omega_3 + g\omega_4. \end{cases} \quad (4)$$

for appropriate functions a, b, c, e, f and g .

Since $\langle d\mathbf{X}, \mathbf{e}_1 \rangle = \langle d\mathbf{X}, \mathbf{e}_2 \rangle = 0$,

$$\begin{aligned}
\langle d^2\mathbf{X}, \mathbf{e}_1 \rangle &= -\langle d\mathbf{X}, d\mathbf{e}_1 \rangle \\
&= -\left\langle \sum_{i=1}^4 \omega_i\mathbf{e}_i, \sum_{j=1}^4 \omega_{1j}\mathbf{e}_j \right\rangle = -\left\langle \sum_{i=3}^4 \omega_i\mathbf{e}_i, \sum_{j=2}^4 \omega_{1j}\mathbf{e}_j \right\rangle \\
&= -(\omega_3\omega_{13} + \omega_4\omega_{14}) = -\omega_3(a\omega_3 + b\omega_4) - \omega_4(b\omega_3 + c\omega_4) \\
&= -(a\omega_3^2 + 2b\omega_3\omega_4 + c\omega_4^2),
\end{aligned}$$

and

$$\begin{aligned} \langle d^2 \mathbf{X}, \mathbf{e}_2 \rangle &= -\langle d\mathbf{X}, d\mathbf{e}_2 \rangle \\ &= -\left\langle \sum_{i=3}^4 \omega_i \mathbf{e}_i, \omega_{21} \mathbf{e}_1 + \omega_{23} \mathbf{e}_3 + \omega_{24} \mathbf{e}_4 \right\rangle = -(\omega_3 \omega_{23} + \omega_4 \omega_{24}) \\ &= -(e\omega_3^2 + 2f\omega_3\omega_4 + g\omega_4^2). \end{aligned}$$

This means that

$$-\langle d^2 \mathbf{X}, \mathbf{e}_1 \rangle \mathbf{e}_1 + \langle d^2 \mathbf{X}, \mathbf{e}_2 \rangle \mathbf{e}_2 = (a\omega_3^2 + 2b\omega_3\omega_4 + c\omega_4^2) \mathbf{e}_1 - (e\omega_3^2 + 2f\omega_3\omega_4 + g\omega_4^2) \mathbf{e}_2 \tag{5}$$

By using equations (2), we may write

$$d \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \\ \mathbf{e}_4 \end{pmatrix} = \begin{pmatrix} 0 & \omega_{12} & \omega_{13} & \omega_{14} \\ \omega_{12} & 0 & \omega_{23} & \omega_{24} \\ \omega_{13} & -\omega_{23} & 0 & \omega_{34} \\ \omega_{14} & -\omega_{24} & -\omega_{34} & 0 \end{pmatrix} \begin{pmatrix} \mathbf{e}_1 \\ \mathbf{e}_2 \\ \mathbf{e}_3 \\ \mathbf{e}_4 \end{pmatrix}. \tag{6}$$

And a straight forward calculation leads us to the following equations:

$$d \begin{pmatrix} \mathbf{e}_1 - \mathbf{e}_2 \\ \mathbf{e}_1 + \mathbf{e}_2 \\ \mathbf{e}_3 \\ \mathbf{e}_4 \end{pmatrix} = \begin{pmatrix} 0 & -\omega_{12} & \omega_{13} - \omega_{23} & \omega_{14} - \omega_{24} \\ \omega_{12} & 0 & \omega_{13} + \omega_{23} & \omega_{14} + \omega_{24} \\ \frac{\omega_{13} - \omega_{23}}{2} & \frac{\omega_{13} + \omega_{23}}{2} & 0 & \omega_{34} \\ \frac{\omega_{14} - \omega_{24}}{2} & \frac{\omega_{14} + \omega_{24}}{2} & -\omega_{34} & 0 \end{pmatrix} \begin{pmatrix} \mathbf{e}_1 + \mathbf{e}_2 \\ \mathbf{e}_1 - \mathbf{e}_2 \\ \mathbf{e}_3 \\ \mathbf{e}_4 \end{pmatrix} \tag{7}$$

On the other hand, we define

$$LC_p = \{ \mathbf{x} \in \mathbb{R}_1^4 \mid \mathbf{x} = -(x_1 - p_1)^2 + \sum_{i=2}^4 (x_i - p_i)^2 = 0 \}$$

and

$$S_+^2 = \{ \mathbf{x} = (x_1, x_2, x_3, x_4) \in LC =: LC_0 \mid x_1 = 1 \},$$

where $p = (p_1, p_2, p_3, p_4) \in \mathbb{R}_1^4$. We call S_+^2 a *lightlike unit sphere* and $LC_p^* = LC_p \setminus \{p\}$ a lightcone at the vertex p . Given any lightlike vector $\mathbf{x} = (x_1, x_2, x_3, x_4)$, we have $\tilde{\mathbf{x}} = (1, \frac{x_2}{x_1}, \frac{x_3}{x_1}, \frac{x_4}{x_1}) \in S_+^2$.

Let $\mathbf{e}_1 = (a_1, a_2, a_3, a_4)$ and $\mathbf{e}_2 = (b_1, b_2, b_3, b_4)$ and consider

$\mathbf{e}_1 \pm \mathbf{e}_2 = (a_1 \pm b_1) \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}$, and thus

$$d(\mathbf{e}_1 \pm \mathbf{e}_2) = d(a_1 \pm b_1) \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2} + (a_1 \pm b_1) d\widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}.$$

Hence

$$\begin{aligned} (a_1 + b_1) d\widetilde{\mathbf{e}_1 + \mathbf{e}_2} &= d(\mathbf{e}_1 + \mathbf{e}_2) - d(a_1 + b_1) \widetilde{\mathbf{e}_1 + \mathbf{e}_2} \\ &= \omega_{12}(\mathbf{e}_1 + \mathbf{e}_2) + (\omega_{13} + \omega_{23})\mathbf{e}_3 + (\omega_{14} + \omega_{24})\mathbf{e}_4 - d(a_1 + b_1) \widetilde{\mathbf{e}_1 + \mathbf{e}_2} \\ &= (a_1 + b_1)\omega_{12} \widetilde{\mathbf{e}_1 + \mathbf{e}_2} + (\omega_{13} + \omega_{23})\mathbf{e}_3 + (\omega_{14} + \omega_{24})\mathbf{e}_4 \\ &\quad - d(a_1 + b_1) \widetilde{\mathbf{e}_1 + \mathbf{e}_2} \\ &= [(a_1 + b_1)\omega_{12} - d(a_1 + b_1)] \widetilde{\mathbf{e}_1 + \mathbf{e}_2} + (\omega_{13} + \omega_{23})\mathbf{e}_3 + (\omega_{14} + \omega_{24})\mathbf{e}_4. \end{aligned}$$

which leads us to the equality

$$d\widetilde{e_1 + e_2} = [\omega_{12} - d\ln(a_1 + b_1)]\widetilde{e_1 + e_2} + \frac{(\omega_{13} + \omega_{23})e_3}{a_1 + b_1} + \frac{(\omega_{14} + \omega_{24})e_4}{a_1 + b_1}.$$

Similarly we have

$$d\widetilde{e_1 - e_2} = [-\omega_{12} - d\ln(a_1 - b_1)]\widetilde{e_1 - e_2} + \frac{(\omega_{13} - \omega_{23})e_3}{a_1 - b_1} + \frac{(\omega_{14} - \omega_{24})e_4}{a_1 - b_1}.$$

And finally we arrive to the the following fundamental formula:

$$d \begin{pmatrix} \widetilde{e_1 - e_2} \\ \widetilde{e_1 + e_2} \\ e_3 \\ e_4 \end{pmatrix} = \begin{pmatrix} 0 & -\omega_{12} - \frac{d(a_1 - b_1)}{a_1 - b_1} & \frac{\omega_{13} - \omega_{23}}{a_1 - b_1} & \frac{\omega_{14} - \omega_{24}}{a_1 - b_1} \\ \omega_{12} - \frac{d(a_1 + b_1)}{a_1 + b_1} & 0 & \frac{\omega_{13} + \omega_{23}}{a_1 + b_1} & \frac{\omega_{14} + \omega_{24}}{a_1 + b_1} \\ \frac{\omega_{13} - \omega_{23}}{2} & \frac{\omega_{13} + \omega_{23}}{2} & 0 & \omega_{34} \\ \frac{\omega_{14} - \omega_{24}}{2} & \frac{\omega_{14} + \omega_{24}}{2} & -\omega_{34} & 0 \end{pmatrix} \begin{pmatrix} \widetilde{e_1 + e_2} \\ \widetilde{e_1 - e_2} \\ e_3 \\ e_4 \end{pmatrix} \tag{8}$$

We now define the *curvature ellipse* at a given point $p \in M$ as the image of the *normal curvature vector map* $\eta : S^1 \rightarrow \mathbb{R}_1^2$ of M , where S^1 is the unit sphere in T_pM and we identify N_pM with \mathbb{R}_1^2 .

Let $\gamma : I \rightarrow \mathbb{R}_1^4$ be a parameterization by arclength of the regular spacelike curve in \mathbb{R}_1^4 obtained by intersecting M with the 3-plane determined by a unit direction $\mathbf{v} \in T_pM$ and N_pM . We clearly have that $\mathbf{v} = \gamma'(s)$ and $p \in \gamma(I)$. The normal curvature vector of γ lies in N_pM and is given by

$$\begin{aligned} \eta(\mathbf{v}) &= -\langle d^2\gamma/ds^2(p), \mathbf{e}_1 \rangle \mathbf{e}_1 + \langle d^2\gamma/ds^2(p), \mathbf{e}_2 \rangle \mathbf{e}_2 \\ &= (a \cos^2 \theta + 2b \cos \theta \sin \theta + c \sin^2 \theta) \mathbf{e}_1 - (e \cos^2 \theta + 2f \cos \theta \sin \theta + g \sin^2 \theta) \mathbf{e}_2, \end{aligned}$$

where $\mathbf{v} = \cos \theta \mathbf{e}_3 + \sin \theta \mathbf{e}_4 \in T_pM$ and $\{\mathbf{e}_3(p), \mathbf{e}_4(p); p\}$ is an orthonormal frame of T_pM .

We define the *mean curvature vector* \mathfrak{H} as

$$\mathfrak{H} = \frac{1}{2}(a + c)\mathbf{e}_1 - \frac{1}{2}(e + g)\mathbf{e}_2. \tag{9}$$

We clearly have

$$\eta(\theta) = \left(\frac{1}{2}(a - c) \cos 2\theta + b \sin 2\theta\right)\mathbf{e}_1 - \left(\frac{1}{2}(e - g) \cos 2\theta + f \sin 2\theta\right)\mathbf{e}_2 + \mathfrak{H}.$$

Or equivalently

$$\eta(\theta) - \mathfrak{H}(\theta) = \begin{pmatrix} \frac{1}{2}(a - c) & b \\ -\frac{1}{2}(e - g) & -f \end{pmatrix} \cdot \begin{pmatrix} \cos 2\theta \\ \sin 2\theta \end{pmatrix}$$

which shows that varying θ from 0 to π , the vector $\eta(\theta)$ describes an ellipse in N_pM .

Given $\mathbf{v} = x\mathbf{e}_1 + y\mathbf{e}_2 \in N_pM$, we have $d\mathbf{v} = dx\mathbf{e}_1 + x d\mathbf{e}_1 + dy\mathbf{e}_2 + y d\mathbf{e}_2$. and then

$$\begin{aligned} \langle d\mathbf{v}, \mathbf{e}_3 \rangle \wedge \langle d\mathbf{v}, \mathbf{e}_4 \rangle &= [(ax + ey)(cx + gy) - (bx + fy)^2]\omega_3 \wedge \omega_4 \\ &= [(ac - b^2)x^2 + (ec + ag - 2bf)xy + (eg - f^2)y^2]\omega_3 \wedge \omega_4. \end{aligned}$$

We define a function \mathcal{K}_l as follows

$$\mathcal{K}_l(x, y) = (ac - b^2)x^2 + (ec + ag - 2bf)xy + (eg - f^2)y^2. \quad (10)$$

On the other hand, we define two maps

$$LG_M^\pm : M \longrightarrow S_+^2$$

by $LG_M^\pm(x, y) = \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(x, y)$. Each one of these maps shall be called *lightcone Gauss map* of $\mathbf{X}(U) = M$.

3. Lightcone height functions on spacelike surfaces. In this section we introduce the notion of lightcone height functions on space like surfaces which, as we shall see, is useful for the study of singularities of the lightcone Gauss maps.

Given a spacelike surface $M(= \mathbf{X}(U))$ we define the function

$$H : M \times S_+^2 \longrightarrow \mathbb{R}$$

as $H((x, y), \boldsymbol{\lambda}) = \langle \mathbf{X}(x, y), \boldsymbol{\lambda} \rangle$, where $\boldsymbol{\lambda} = (1, \lambda_2, \lambda_3, \lambda_4) \in S_+^2$. We call H the *lightcone height function* on the spacelike surface M . We denote that $h_{\lambda_0}(x, y) = H(x, y, \boldsymbol{\lambda}_0)$, for any fixed $\boldsymbol{\lambda}_0 \in S_+^2$. The singularities of this function are characterized as follows.

PROPOSITION 3.1. *Let M be a spacelike surface and $H : M \times S_+^2 \longrightarrow \mathbb{R}$ a lightcone height function. Then we have the following assertions:*

- (1) $(\partial h_\lambda / \partial x)(p_0) = (\partial h_\lambda / \partial y)(p_0) = 0$ if and only if $\boldsymbol{\lambda} = \mu(\mathbf{e}_1 \pm \mathbf{e}_2)(p_0) = \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(p_0)$. Here $\mathbf{e}_1(p_0) = (a_1, a_2, a_3, a_4)$, $\mathbf{e}_2(p_0) = (b_1, b_2, b_3, b_4)$ and $\mu = \frac{1}{a_1 \pm b_1}$; $(p_0) = (x_0, y_0) \in M$,
- (2) $(\partial h_\lambda / \partial x)(p_0) = (\partial h_\lambda / \partial y)(p_0) = \det \mathcal{H}(h_\lambda)(p_0) = 0$ if and only if

$$\boldsymbol{\lambda} = \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(p_0) \text{ and } \mathcal{K}_l(1, \pm 1)(p_0) = 0.$$

Here, $\det \mathcal{H}(h_\lambda)(x, y)$ is the determinant of the Hessian matrix of h_λ at (x, y) .

Proof. It follows from a straight forward calculation that $(\partial h_\lambda / \partial x)(p_0) = (\partial h_\lambda / \partial y)(p_0) = 0$ if and only if

$$\langle \mathbf{X}_x, \boldsymbol{\lambda} \rangle(p_0) = \langle \mathbf{X}_y, \boldsymbol{\lambda} \rangle(p_0) = 0.$$

This is equivalent to the condition that $\boldsymbol{\lambda} \in N_{p_0}M$ and $\boldsymbol{\lambda} \in S_+^2$ which means that $\boldsymbol{\lambda} = \mu(\mathbf{e}_1 \pm \mathbf{e}_2) = \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}$.

On the other hand, if we choose local coordinates such that \mathbf{X} is given in the Monge form $\mathbf{X}(x, y) = (f_1(x, y), f_2(x, y), x, y)$ and $\mathbf{e}_1(p_0) = (1, 0, 0, 0)$ and $\mathbf{e}_2(p_0) = (0, 1, 0, 0)$, since

$$\det \mathcal{H}(h_\lambda)(x, y) = \begin{vmatrix} \langle \mathbf{X}_{xx}, \boldsymbol{\lambda} \rangle & \langle \mathbf{X}_{xy}, \boldsymbol{\lambda} \rangle \\ \langle \mathbf{X}_{xy}, \boldsymbol{\lambda} \rangle & \langle \mathbf{X}_{yy}, \boldsymbol{\lambda} \rangle \end{vmatrix} = 0$$

and $\boldsymbol{\lambda}(p_0) = (1, \pm 1, 0, 0)$, we have

$$\begin{aligned} & \begin{vmatrix} \langle (f_{1_{xx}}, f_{2_{xx}}, 0, 0), (1, \pm 1, 0, 0) \rangle & \langle (f_{1_{xy}}, f_{2_{xy}}, 0, 0), (1, \pm 1, 0, 0) \rangle \\ \langle (f_{1_{xy}}, f_{2_{xy}}, 0, 0), (1, \pm 1, 0, 0) \rangle & \langle (f_{1_{yy}}, f_{2_{yy}}, 0, 0), (1, \pm 1, 0, 0) \rangle \end{vmatrix} \\ &= \begin{vmatrix} -a \pm e & -b \pm f \\ -b \pm f & -c \pm g \end{vmatrix} = 0. \end{aligned}$$

But this is equivalent to the condition

$$(ac + eg) \pm (2bf - ag - ec) - (b^2 + f^2) = (ac - b^2) + (eg - f^2) \pm (2bf - ag - ce) = 0$$

and $\lambda(p_0) = (1, \pm 1, 0, 0)$, namely $\mathcal{K}_l(1, \pm 1)(x, y) = 0$ and $\lambda(p_0) = (1, \pm 1, 0, 0)$. \square

As a corollary of Proposition 3.1, we have the following theorem.

THEOREM 3.2. *Under the assumption of Proposition 2.1, we have that the following conditions are equivalent:*

- (1) $p \in M$ is a degenerate singular point of the lightcone height function h_λ .
- (2) (p, λ) is a singular point of the lightcone Gauss map LG_M^\pm .
- (3) $\mathcal{K}_l(1, \pm 1)(p) = 0$.

Proof. Let's consider the subset

$$\Sigma(H) = \left\{ (p, \lambda) \in M \times S_+^2 \mid \frac{\partial h_\lambda}{\partial x}(p) = \frac{\partial h_\lambda}{\partial y}(p) = 0 \right\}.$$

which from Proposition 3.1, (1) can be also written as

$$\Sigma(H) = \{ (p, \lambda) \in M \times S_+^2 \mid \lambda = \widetilde{e_1 \pm e_2}(p) \}.$$

We now observe that the restriction, $\pi[\Sigma(H)]$, of the canonical projection $\pi : M \times S_+^2 \rightarrow S_+^2$ can be identified with the lightcone Gauss map LG_M^\pm . We can easily see, under this identification, that the condition (1) is equivalent to the condition (2).

That the condition (2) is equivalent to the condition (3) follows immediately from Proposition 3.1 (2). \square

THEOREM 3.3. *Let M be a spacelike surface in Minkowski 4-space.*

- (1) *The lightcone Gauss map LG_M^+ (respectively, LG_M^-) is constant if and only if there exists a unique lightlike hyperplane $HLP(\mathbf{v}^+, c^+)$ (respectively, $HLP(\mathbf{v}^-, c^-)$) such that $M \subset HLP(\mathbf{v}^+, c^+)$ (respectively, $M \subset HLP(\mathbf{v}^-, c^-)$)*
- (2) *Both maps, LG_M^+ and LG_M^- , are constant if and only if M is a spacelike plane. In this case, the intersection of the lightlike hyperplanes*

$$HLP(\widetilde{e_1 + e_2}, c^+) \cap HLP(\widetilde{e_1 - e_2}, c^-)$$

is the spacelike plane M .

Proof. (1) We prove the assertion for the assumption $LG_M^+(x, y) = \widetilde{e_1 + e_2}(x, y)$ is constant. The other case can be shown analogously. In this case we have

$$d\langle \mathbf{X}, \widetilde{e_1 + e_2} \rangle = \langle d\mathbf{X}, \widetilde{e_1 + e_2} \rangle + \langle \mathbf{X}, d(\widetilde{e_1 + e_2}) \rangle = 0.$$

Therefore, $\langle \mathbf{X}, \widetilde{e_1 + e_2} \rangle \equiv c^+$. This means that $M = \mathbf{X}(U) \subset HLP(\mathbf{v}^+, c^+)$, where $\mathbf{v}^+ = \widetilde{e_1 + e_2}(x, y)$. Conversely, suppose that there exists a lightlike vector \mathbf{v} and a real number c such that $\mathbf{X}(U) = M \subset HLP(\mathbf{v}, c)$. Since $\langle \mathbf{X}(x, y), \mathbf{v} \rangle = c$, we have $d\langle \mathbf{X}(x, y), \mathbf{v} \rangle = 0$. This means that \mathbf{v} is a lightlike normal vector of M . Thus we have $\widetilde{\mathbf{v}} = \widetilde{e_1 \pm e_2}(x, y)$ which completes the proof of (1).

Since $\mathbf{v}^+ \notin HLP(\mathbf{v}^-, c^-)$ and $\mathbf{v}^- \notin HLP(\mathbf{v}^+, c^+)$, $HLP(\mathbf{v}^-, c^-)$ and $HLP(\mathbf{v}^+, c^+)$ intersect transversally. By the assertion (1), both of the lightcone Gauss maps LG_M^+ and LG_M^- are constant if and only if $M \subset HLP(\mathbf{v}^+, c^+) \cap HLP(\mathbf{v}^-, c^-)$. Since the intersection must be a spacelike plane we have proven assertion (2). \square

We say that a point $p = \mathbf{X}(x_0, y_0)$ is a *lightlike parabolic point* of M if $\mathcal{K}_l(1, 1)(p) = 0$ or $\mathcal{K}_l(1, -1)(p) = 0$.

4. The lightcone pedal surface of a spacelike surface. In this section we associate to M a singular hypersurface contained in the positive lightcone

$$LC_+^* = \{ \mathbf{x} = (x_1, x_2, x_3, x_4) \in LC_0 \mid x_1 > 0 \}$$

whose singularities correspond to those of the lightcone Gauss map of M . To do this we consider a family of functions

$$\tilde{H} : M \times LC_+^* \longrightarrow \mathbb{R}$$

defined by

$$\tilde{H}((x, y), \mathbf{v}) = \langle \mathbf{X}(x, y), \tilde{\mathbf{v}} \rangle - v_1,$$

where $\mathbf{v} = (v_1, v_2, v_3, v_4)$. We call \tilde{H} the *extended lightcone height function* of $M = \mathbf{X}(U)$.

An immediate consequence of Proposition 3.1 is the following.

PROPOSITION 4.1. *Let M be a spacelike surface and $\tilde{H} : M \times LC_+^* \longrightarrow \mathbb{R}$ the extended lightcone height function of M . For $p_0 = \mathbf{X}(x_0, y_0)$ and $\mathbf{v}_0 \in LC_+^*$, we have the following:*

(1) $\tilde{H}(p_0, \mathbf{v}_0) = (\partial\tilde{H}/\partial x)(p_0, \mathbf{v}_0) = (\partial\tilde{H}/\partial y)(p_0, \mathbf{v}_0) = 0$ if and only if

$$\tilde{\mathbf{v}}_0 = \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(p_0) \text{ and } v_1 = \langle \mathbf{X}(x_0, y_0), \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(p_0) \rangle.$$

(2)

$$\tilde{H}(p_0, \mathbf{v}_0) = \frac{\partial\tilde{H}}{\partial x}(p_0, \mathbf{v}_0) = \frac{\partial\tilde{H}}{\partial y}(p_0, \mathbf{v}_0) = \det\mathcal{H}(\tilde{h}_v)(p_0) = 0$$

if and only if

$$\tilde{\mathbf{v}}_0 = \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(p_0), \quad v_1 = \langle \mathbf{X}(x_0, y_0), \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(p_0) \rangle \text{ and } \mathcal{K}_l(1, \pm 1)(p_0) = 0.$$

Here, $\tilde{h}_v(x, y) = \tilde{H}((x, y), \mathbf{v})$.

The above result implies that the discriminant set of the extended lightcone height function \tilde{H} is

$$\mathcal{D}_{\tilde{H}} = \left\{ \mathbf{v} \in LC_+^* \mid \mathbf{v} = \langle \mathbf{X}(x, y), \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(x, y) \rangle \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(x, y) \text{ for some } (x, y) \in U \right\}.$$

In view of this, we associate to M a couple of singular surfaces, $LP_M^\pm \subset LC_+^*$, called *lightcone pedal surfaces* of $\mathbf{X}(U) = M$. These are defined by

$$LP_M^\pm(p) = LP_M^\pm(x, y) = \langle \mathbf{X}(x, y), \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(x, y) \rangle \widetilde{\mathbf{e}_1 \pm \mathbf{e}_2}(x, y).$$

We observe that each singularity of each one of the lightcone pedal surfaces corresponds to a singularity of the lightcone Gauss map. This correspondence has an interesting explanation in terms of the Symplectic and Contact Geometry methods that we analyze in the following paragraphs.

Given a point $\mathbf{v} = (v_1, v_2, v_3, v_4) \in LC_+^*$, we have that $v_1 = \sqrt{v_2^2 + v_3^2 + v_4^2}$. We take coordinates (v_2, v_3, v_4) on the manifold LC_+^* and consider the projective

cotangent bundle $\pi : PT^*(LC_+^*) \rightarrow LC_+^*$ with its canonical contact structure. Consider also the tangent bundle $\tau : TPPT^*(LC_+^*) \rightarrow PT^*(LC_+^*)$ and the differential, $d\pi : TPPT^*(LC_+^*) \rightarrow TLC_+^*$, of π . For any $X \in TPPT^*(LC_+^*)$, there exists an element $\alpha \in T^*(LC_+^*)$ such that $\tau(X) = [\alpha]$. For an element $V \in T_x(LC_+^*)$, the property $\alpha(V) = 0$ does not depend on the choice of representative of the class $[\alpha]$. Thus we can define the canonical contact structure on $PT^*(LC_+^*)$ as

$$K = \{X \in TPPT^*(LC_+^*) \mid \tau(X)(d\pi(X)) = 0\}.$$

In the coordinates (v_2, v_3, v_4) we have the trivialization $PT^*(LC_+^*) \cong LC_+^* \times P(\mathbb{R}^2)^*$. If $[\xi_2 : \xi_3 : \xi_4]$ represent the homogeneous coordinate of the dual projective space $P(\mathbb{R}^2)^*$, we shall call

$$((v_2, v_3, v_4), [\xi_2 : \xi_3 : \xi_4])$$

homogeneous coordinates for $PT^*(LC_+^*)$.

It is easy to show that $X \in K_{(x, [\xi])}$ if and only if $\sum_{i=2}^4 \mu_i \xi_i = 0$, where $d\pi(X) = \sum_{i=2}^4 \mu_i \frac{\partial}{\partial v_i}$. An immersion $i : L \rightarrow PT^*(LC_+^*)$ is said to be a Legendrian immersion if $\dim L = 2$ and $di_q(T_q L) \subset K_{i(q)}$ for any $q \in L$. We also call the map $\pi \circ i$ the Legendrian map and the set $W(i) = \text{image } \pi \circ i$ the wave front of i . Moreover, i (or, the image of i) is called the Legendrian lift of $W(i)$.

In order to lighten the study of the lightcone pedal surface, we include here a quick survey on the Legendrian singularity theory developed by Arnol'd and Zakalyukin [1, 22]. It is enough to consider, for our purposes, the 3-dimensional case instead of the whole general theory.

Let $F : (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \rightarrow (\mathbb{R}, \mathbf{0})$ be a function germ. We say that F is a Morse family if the map

$$\Delta^* F = \left(F, \frac{\partial F}{\partial q_1}, \dots, \frac{\partial F}{\partial q_k} \right) : (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \rightarrow (\mathbb{R}^k \times \mathbb{R}, \mathbf{0})$$

is non-singular, where $(q, x) = (q_1, \dots, q_k, x_1, x_2, x_3) \in (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0})$. In this case we have a smooth 2-dimensional submanifold

$$\Sigma_*(F) = \left\{ (q, x) \in (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \mid F(q, x) = \frac{\partial F}{\partial q_1}(q, x) = \dots = \frac{\partial F}{\partial q_k}(q, x) = 0 \right\}$$

and the map germ $\Phi_F : (\Sigma_*(F), \mathbf{0}) \rightarrow PT^*\mathbb{R}^3$ defined by

$$\Phi_F(q, x) = \left(x, \left[\frac{\partial F}{\partial x_1}(q, x) : \frac{\partial F}{\partial x_2}(q, x) : \frac{\partial F}{\partial x_3}(q, x) \right] \right)$$

is a Legendrian immersion. Then we have the following fundamental theorem ([1, 22]).

PROPOSITION 4.2. All Legendrian submanifold germs in $PT^*\mathbb{R}^3$ are constructed by the above method.

We call F a generating family of Φ_F . The corresponding wave front is

$$W(\Phi_F) = \left\{ x \in \mathbb{R}^3 \mid \text{there exists } q \in \mathbb{R}^k \text{ such that } F(q, x) = \frac{\partial F}{\partial q_1}(q, x) = \dots = \frac{\partial F}{\partial q_k}(q, x) = 0 \right\}.$$

We have, by definition, that $\mathcal{D}_F = W(\Phi_F)$. It then follows from the previous arguments that the lightcone pedal surface LP_M^\pm is the discriminant set of the extended lightcone height function \tilde{H} .

PROPOSITION 4.3. *The extended lightcone height function \tilde{H} is a Morse family.*

Proof. Consider the family of functions

$$\bar{H} : U \times S_+^2 \times \mathbb{R} \longrightarrow \mathbb{R}$$

given by $\bar{H}((x, y), \mathbf{w}, r) = \langle \mathbf{X}(x, y), \mathbf{w} \rangle - r$, and a C^∞ -diffeomorphism, $\Phi : U \times S_+^2 \times \mathbb{R} \longrightarrow LC_+^*$ defined by $\Phi((x, y), \mathbf{w}, r) = ((x, y), r\mathbf{w})$. Then we have that $\tilde{H} = \bar{H} \circ \Phi$ and it is enough to show that \bar{H} is a Morse family. Given $\mathbf{w} = (1, w_2, w_3, w_4) \in S_+^2$, we have $w_2 = \sqrt{1 - w_3^2 - w_4^2}$, so that

$$H((x, y), \mathbf{w}, r) = -x_1(x, y) + x_2(x, y)\sqrt{1 - w_3^2 - w_4^2} + x_3(x, y)w_3 + x_4(x, y)w_4 - r,$$

where $\mathbf{X}(x, y) = (x_1(x, y), x_2(x, y), x_3(x, y), x_4(x, y))$. We now prove that the map

$$\Delta^* \bar{H} = \left(\frac{\partial \bar{H}}{\partial x}, \frac{\partial \bar{H}}{\partial y}, \bar{H} \right)$$

is non-singular at any point. The Jacobian matrix of $\Delta^* \bar{H}$ is given as follows:

$$\begin{pmatrix} \langle \mathbf{X}_{xx}, \mathbf{w} \rangle & \langle \mathbf{X}_{xy}, \mathbf{w} \rangle & -x_{2,x} \frac{w_3}{w_2} + x_{3,x} & -x_{2,x} \frac{w_4}{w_2} + x_{4,x} & 0 \\ \langle \mathbf{X}_{xy}, \mathbf{w} \rangle & \langle \mathbf{X}_{yy}, \mathbf{w} \rangle & -x_{2,y} \frac{w_3}{w_2} + x_{3,y} & -x_{2,y} \frac{w_4}{w_2} + x_{4,y} & 0 \\ \langle \mathbf{X}_x, \mathbf{w} \rangle & \langle \mathbf{X}_y, \mathbf{w} \rangle & -x_2 \frac{w_3}{w_2} + x_3 & -x_2 \frac{w_4}{w_2} + x_4 & -1 \end{pmatrix}.$$

It follows now from a straight forward calculation that the determinant of the matrix

$$A = \begin{pmatrix} -x_{2,x} \frac{w_3}{w_2} + x_{3,x} & -x_{2,x} \frac{w_4}{w_2} + x_{4,x} \\ -x_{2,y} \frac{w_3}{w_2} + x_{3,y} & -x_{2,y} \frac{w_4}{w_2} + x_{4,y} \end{pmatrix}.$$

is equal to

$$\frac{w_4}{w_2}(x_{2,x}x_{3,y} - x_{3,x}x_{2,y}) + \frac{w_3}{w_2}(x_{x,4}x_{2,y} - x_{2,x}x_{4,y}) + (x_{3,x}x_{4,y} - x_{4,x}x_{3,x}).$$

Since $\mathbf{X}(U) = M$ is a spacelike surface, the surface parameterized by $(x_2(x, y), x_3(x, y), x_4(x, y))$ in Euclidean space is everywhere regular and we can interpret that the above determinant vanishes if and only if the vector (w_2, w_3, w_4) is tangent to this surface. But it is impossible because $\mathbf{X}(U) = M$ is a spacelike surface.

It follows from Proposition 4.3 that the lightcone pedal surfaces LP_M^\pm are wave fronts and the extended lightcone height function \tilde{H} provides a generating family for the Legendrian lifts of LP_M^\pm .

5. Contact with lightlike hyperplanes. In this section we give geometrical interpretations to the singularities of the lightcone Gauss map and the lightcone pedal surface of $\mathbf{X}(U) = M$. This is done, following the classical differential geometry methods, through the analysis of the contacts of a spacelike surface with the lightlike hyperplanes. We first include a brief review of the theory of contact due to Montaldi ([16]).

Let X_i, Y_i ($i = 1, 2$) be submanifolds of \mathbb{R}^n with $\dim X_1 = \dim X_2$ and $\dim Y_1 = \dim Y_2$. We say that *the contact of X_1 and Y_1 at y_1 is of the same type as the contact of X_2 and Y_2 at y_2* if there is a diffeomorphism germ $\Phi : (\mathbb{R}^n, y_1) \rightarrow (\mathbb{R}^n, y_2)$ such that $\Phi(X_1) = X_2$ and $\Phi(Y_1) = Y_2$. In this case we write $K(X_1, Y_1; y_1) = K(X_2, Y_2; y_2)$. We can, clearly, replace \mathbb{R}^n by any manifold in this definition. Montaldi gives a characterization of the notion of contact by using the terminology of singularity theory.

THEOREM 5.1. ([16]) *Let X_i, Y_i ($i = 1, 2$) be submanifolds of \mathbb{R}^n with $\dim X_1 = \dim X_2$ and $\dim Y_1 = \dim Y_2$. Let $g_i : (X_i, x_i) \rightarrow (\mathbb{R}^n, y_i)$ be immersion germs and $f_i : (\mathbb{R}^n, y_i) \rightarrow (\mathbb{R}^p, 0)$ be submersion germs with $(Y_i, y_i) = (f_i^{-1}(0), y_i)$. Then*

$$K(X_1, Y_1; y_1) = K(X_2, Y_2; y_2)$$

if and only if $f_1 \circ g_1$ and $f_2 \circ g_2$ are \mathcal{K} -equivalent.

We now consider a function $\mathcal{H} : \mathbb{R}_1^4 \times LC_+^* \rightarrow \mathbb{R}$ defined by $\mathcal{H}(\mathbf{x}, \mathbf{v}) = \langle \mathbf{x}, \tilde{\mathbf{v}} \rangle - v_1$. Given $\mathbf{v}_0 \in LC_+^*$, we denote $\mathfrak{h}_{\mathbf{v}_0}(\mathbf{x}) = \mathcal{H}(\mathbf{x}, \mathbf{v}_0)$ and observe that $\mathfrak{h}_{\mathbf{v}_0}^{-1}(0) = LHP(\tilde{\mathbf{v}}_0, v_{0,1})$ defines a lightlike hyperplane. For any $p_0 = \mathbf{X}(x_0, y_0) \in U$, we consider the lightlike vector $\mathbf{v}_0^\pm = \mathbf{e}_1 \pm \mathbf{e}_2(p_0)$ and $c^\pm = \langle \mathbf{X}(x_0, y_0), \mathbf{v}^\pm \rangle$, and we have

$$\mathfrak{h}_{\mathbf{v}_0^\pm} \circ \mathbf{X}(p_0) = \mathcal{H} \circ (\mathbf{X} \times id_{LC_+^*})(p_0, \mathbf{v}_0^\pm) = H((x_0, y_0), \tilde{\mathbf{v}}_0^\pm) - c^\pm = 0.$$

We also have the relations

$$\frac{\partial \mathfrak{h}_{\mathbf{v}_0^\pm} \circ \mathbf{X}}{\partial x}(p_0) = \frac{\partial H}{\partial x}((p_0), \mathbf{v}_0^\pm) = 0$$

and

$$\frac{\partial \mathfrak{h}_{\mathbf{v}_0^\pm} \circ \mathbf{X}}{\partial y}(p_0) = \frac{\partial H}{\partial y}((p_0), \mathbf{v}_0^\pm) = 0.$$

which imply that the lightlike hyperplane $\mathfrak{h}_{\mathbf{v}_0^\pm}^{-1}(0) = LHP(\mathbf{v}_0^\pm, c^\pm)$ is tangent to $M = \mathbf{X}(U)$ at $p_0 = \mathbf{X}(x_0, y_0)$. In this case, we call each $LHP(\mathbf{v}_0^\pm, c^\pm)$ the *tangent lightlike hyperplane* of $M = \mathbf{X}(U)$ at $p_0 = \mathbf{X}(x_0, y_0)$. Moreover, the intersection

$$LHP(\mathbf{v}_0^\pm, c^+) \cap LHP(\mathbf{v}_0^\pm, c^-)$$

is the tangent plane of M at p_0 . Let $\mathbf{v}_1, \mathbf{v}_2$ be lightlike vectors. Clearly, if $\mathbf{v}_1, \mathbf{v}_2$ are linearly dependent the lightlike hyperplanes $LHP(\mathbf{v}_1, c_1)$ and $LHP(\mathbf{v}_2, c_2)$ are parallel. We have the following simple lemma.

LEMMA 5.2. *Let $\mathbf{X} : U \rightarrow \mathbb{R}_1^4$ be a spacelike surface and $\sigma = \pm$. Given two points $p_1 = \mathbf{X}(x_1, y_1), p_2 = \mathbf{X}(x_2, y_2)$ in $M = \mathbf{X}(U)$, the following assertions hold:*
 (1) $LG_M^\sigma(p_1) = LG_M^\sigma(p_2)$ if and only if $LHP(\mathbf{v}_1^\sigma, c_1^\sigma)$ and $LHP(\mathbf{v}_2^\sigma, c_2^\sigma)$ are parallel.

(2) $LP_M^\sigma(p_1) = LP_M^\sigma(p_2)$ if and only if $LHP(\mathbf{v}_1^\sigma, c_1^\sigma) = LHP(\mathbf{v}_2^\sigma, c_2^\sigma)$.
 Here, $v_i^\pm = \widetilde{e_1 \pm e_2}(p_i)$ and $c_i^\pm = \langle \mathbf{X}(x_i, y_i), \mathbf{v}_i^\pm \rangle$ for $i = 1, 2$.

On the other hand, given any map $f : N \rightarrow P$, we denote its set of singular points by $\Sigma(f)$ and put $D(f) = f(\Sigma(f))$. We call $f|_{\Sigma(f)} : \Sigma(f) \rightarrow D(f)$ the critical part of the map f . For any Morse family $F : (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \rightarrow (\mathbb{R}, \mathbf{0})$, $(F^{-1}(0), \mathbf{0})$ is a smooth hypersurface, so we can define a smooth map germ $\pi_F : (F^{-1}(0), \mathbf{0}) \rightarrow (\mathbb{R}, 0)$ by $\pi_F(q, x) = x$. We can easily show that $\Sigma_*(F) = \Sigma(\pi_F)$. Therefore, the corresponding Legendrian map $\pi \circ \Phi_F$ is the critical part of π_F .

We introduce the following equivalence relation among Legendrian immersion germs: Let $i : (L, p) \subset (PT^*\mathbb{R}^3, p)$ and $i' : (L', p') \subset (PT^*\mathbb{R}^3, p')$ be Legendrian immersion germs. We say that i and i' are Legendrian equivalent if there exists a contact diffeomorphism germ $H : (PT^*\mathbb{R}^3, p) \rightarrow (PT^*\mathbb{R}^3, p')$ such that H preserves fibers of π and $H(L) = L'$. A Legendrian immersion germ into $PT^*\mathbb{R}^3$ at a point is said to be Legendrian stable if for every map with the given germ there is a neighbourhood in the space of Legendrian immersions (in the Whitney C^∞ topology) and a neighbourhood of the original point such that each Legendrian immersion belonging to the first neighbourhood has in the second neighbourhood a point at which its germ is Legendrian equivalent to the original germ.

Since the Legendrian lift $i : (L, p) \subset (PT^*\mathbb{R}^3, p)$ is uniquely determined on the regular part of the wave front $W(i)$, we have the following simple but significant property of Legendrian immersion germs:

PROPOSITION 5.3. *Let $i : (L, p) \subset (PT^*\mathbb{R}^3, p)$ and $i' : (L', p') \subset (PT^*\mathbb{R}^3, p')$ be Legendrian immersion germs such that the regular sets of $\pi \circ i, \pi \circ i'$ are dense respectively. Then i, i' are Legendrian equivalent if and only if wave front sets $W(i), W(i')$ are diffeomorphic as set germs.*

This result has been firstly pointed out by Zakalyukin [23]. The assumption in the above proposition is a generic condition for i, i' . In particular, if i, i' are Legendrian stable, then they satisfy this assumption.

We can interpret the Legendrian equivalence by using the notion of generating families. We denote by \mathcal{E}_n the local ring of function germs $(\mathbb{R}^n, \mathbf{0}) \rightarrow \mathbb{R}$ with the unique maximal ideal $\mathfrak{M}_n = \{h \in \mathcal{E}_n \mid h(0) = 0\}$. Let $F, G : (\mathbb{R}^k \times \mathbb{R}^n, \mathbf{0}) \rightarrow (\mathbb{R}, \mathbf{0})$ be function germs. We say that F and G are P - \mathcal{K} -equivalent if there exists a diffeomorphism germ $\Psi : (\mathbb{R}^k \times \mathbb{R}^n, \mathbf{0}) \rightarrow (\mathbb{R}^k \times \mathbb{R}^n, \mathbf{0})$ of the form $\Psi(x, u) = (\psi_1(q, x), \psi_2(x))$ for $(q, x) \in (\mathbb{R}^k \times \mathbb{R}^n, \mathbf{0})$ such that $\Psi^*(\langle F \rangle_{\mathcal{E}_{k+n}}) = \langle G \rangle_{\mathcal{E}_{k+n}}$. Here $\Psi^* : \mathcal{E}_{k+n} \rightarrow \mathcal{E}_{k+n}$ is the pull back \mathbb{R} -algebra isomorphism defined by $\Psi^*(h) = h \circ \Psi$.

Let $F : (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \rightarrow (\mathbb{R}, \mathbf{0})$ a function germ. We say that F is a \mathcal{K} -versal deformation of $f = F|_{\mathbb{R}^k \times \{\mathbf{0}\}}$ if

$$\mathcal{E}_k = T_e(\mathcal{K})(f) + \left\langle \frac{\partial F}{\partial x_1} |_{\mathbb{R}^k \times \{\mathbf{0}\}}, \frac{\partial F}{\partial x_2} |_{\mathbb{R}^k \times \{\mathbf{0}\}}, \frac{\partial F}{\partial x_3} |_{\mathbb{R}^k \times \{\mathbf{0}\}} \right\rangle_{\mathbb{R}},$$

where

$$T_e(\mathcal{K})(f) = \left\langle \frac{\partial f}{\partial q_1}, \dots, \frac{\partial f}{\partial q_k}, f \right\rangle_{\mathcal{E}_k}.$$

(See [14].)

The main result in Arnol'd-Zakalyukin's theory[1, 22] is the following:

THEOREM 5.4. *Let $F, G : (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \rightarrow (\mathbb{R}, 0)$ be Morse families. Then*

- (1) Φ_F and Φ_G are Legendrian equivalent if and only if F, G are $P\mathcal{K}$ -equivalent.
- (2) Φ_F is Legendrian stable if and only if F is a \mathcal{K} -versal deformation of $F|_{\mathbb{R}^k \times \{\mathbf{0}\}}$.

Since F, G are function germs on the common space germ $(\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0})$, we do not need the notion of stable $P\mathcal{K}$ -equivalences (cf., [1]). By the uniqueness result of the \mathcal{K} -versal deformation of a function germ, Proposition 5.2 and Theorem 5.3, we have the following classification result of Legendrian stable germs (cf., [7]). For any map germ $f : (\mathbb{R}^n, \mathbf{0}) \rightarrow (\mathbb{R}^p, \mathbf{0})$, we define the local ring of f by $Q(f) = \mathcal{E}_n/f^*(\mathfrak{M}_p)\mathcal{E}_n$.

PROPOSITION 5.5. *Let $F, G : (\mathbb{R}^k \times \mathbb{R}^3, \mathbf{0}) \rightarrow (\mathbb{R}, 0)$ be Morse families. Suppose that Φ_F, Φ_G are Legendrian stable. The the following conditions are equivalent.*

- (1) $(W(\Phi_F), \mathbf{0})$ and $(W(\Phi_G), \mathbf{0})$ are diffeomorphic as germs.
- (2) Φ_F and Φ_G are Legendrian equivalent.
- (3) $Q(f)$ and $Q(g)$ are isomorphic as \mathbb{R} -algebras, where $f = F|_{\mathbb{R}^k \times \{\mathbf{0}\}}$, $g = G|_{\mathbb{R}^k \times \{\mathbf{0}\}}$.

Proof. Since Φ_F, Φ_G are Legendrian stable, they satisfy the generic condition of Proposition 5.3, so that the conditions (1) and (2) are equivalent. The condition (3) implies that f, g are \mathcal{K} -equivalent[14, 15]. By the uniqueness of the \mathcal{K} -versal deformation of a function germ, F, G are $P\mathcal{K}$ -equivalent. This means that the condition (2) holds. By Theorem 5.4, the condition (2) implies the condition (3). \square

We apply now these tools to the study of the contact between spacelike surfaces and lightlike hyperplanes.

Let $LP_{M,i}^\sigma : (U, (x_i, y_i)) \rightarrow (LC_+^*, \mathbf{v}_i^\sigma)$ ($i = 1, 2$) be two lightcone pedal surface germs respectively associated to spacelike surface germs $\mathbf{X}_i : (U, (x_i, y_i)) \rightarrow (\mathbb{R}_1^4, p_i)$, where $\sigma = \pm$. We say that $LP_{M,1}^\sigma$ and $LP_{M,2}^\sigma$ are \mathcal{A} -equivalent if there exist diffeomorphism germs $\phi : (U, (x_1, y_1)) \rightarrow (U, (x_2, y_2))$ and $\Phi : (LC_+^*, \mathbf{v}_1^\sigma) \rightarrow (LC_+^*, \mathbf{v}_2^\sigma)$ such that $\Phi \circ LP_{M,1}^\sigma = LP_{M,2}^\sigma \circ \phi$. If the both of the regular sets of $LP_{M,i}^\sigma$ are dense in $(U, (x_i, y_i))$, it follows from Proposition 5.5 that $LP_{M,1}^\sigma$ and $LP_{M,2}^\sigma$ are \mathcal{A} -equivalent if and only if the corresponding Legendrian lift germs are Legendrian equivalent. This condition is also equivalent, by Theorem 5.4, to the condition that the two generating families \tilde{H}_1 and \tilde{H}_2 are $P\mathcal{K}$ -equivalent. Here, $\tilde{H}_i : (U \times LC_+^*, ((x_i, y_i), \mathbf{v}_i^\sigma)) \rightarrow \mathbb{R}$ is the extended lightlike height function germ of \mathbf{X}_i .

On the other hand, we denote that $h_{i,v_i^\sigma}(u) = H_i(u, \mathbf{v}_i^\sigma)$ and we have $h_{i,v_i^\pm}(u) = \mathfrak{h}_{v_i^\pm} \circ \mathbf{x}_i(u)$. By Theorem 5.1, $K(\mathbf{X}_1(U), HLP(\mathbf{v}^\sigma, -1), \mathbf{v}_1^\sigma) = K(\mathbf{x}_2(U), HLP(\mathbf{v}^\sigma, -1), \mathbf{v}_2^\sigma)$ if and only if \tilde{h}_{1,v_1} and \tilde{h}_{1,v_2} are \mathcal{K} -equivalent. So we can apply the previous arguments to our situation. We denote $Q^\sigma(\mathbf{X}, (x_0, y_0))$ the local ring of the function germ $\tilde{h}_{v_0^\sigma} : (U, (x_0, y_0)) \rightarrow \mathbb{R}$, where $v_0^\sigma = LP_M^\sigma(x_0, y_0)$. We remark that we can explicitly write the local ring as follows:

$$Q^\pm(\mathbf{X}, (x_0, y_0)) = \frac{C_{(x_0, y_0)}^\infty(U)}{\langle \langle \mathbf{X}(x, y), \mathbf{e}_1 \pm \mathbf{e}_2(x_0, y_0) \rangle - 1 \rangle_{C_{(x_0, y_0)}^\infty(U)}}$$

where $C_{(x_0, y_0)}^\infty(U)$ is the local ring of function germs at (x_0, y_0) with the unique maximal ideal $\mathfrak{M}_{(x_0, y_0)}(U)$.

THEOREM 5.6. *Let $\mathbf{X}_i : (U, (x_i, y_i)) \rightarrow (\mathbb{R}_1^4, \mathbf{X}_i((x_i, y_i)))$ ($i = 1, 2$) be spacelike surface germs such that the corresponding Legendrian lift germs are Legendrian stable and $\sigma = \pm$. Then the following conditions are equivalent:*

- (1) The lightcone pedal surface germs $LP_{M,1}^\sigma$ and $LP_{M,2}^\sigma$ are \mathcal{A} -equivalent.

- (2) \tilde{H}_1 and \tilde{H}_2 are $P\mathcal{K}$ -equivalent.
- (3) \tilde{h}_{1,v_1} and \tilde{h}_{1,v_2} are \mathcal{K} -equivalent.
- (4) $K(\mathbf{X}_1(U), HLP(\mathbf{v}^\sigma, -1), \mathbf{v}_1^\sigma) = K(\mathbf{x}_2(U), HLP(\mathbf{v}^\sigma, -1), \mathbf{v}_2^\sigma)$
- (5) $Q^\sigma(\mathbf{X}_1, (x_1, y_1))$ and $Q^\sigma(\mathbf{X}_2, (x_2, y_2))$ are isomorphic as \mathbb{R} -algebras.

Proof. The previous arguments (mainly by Theorem 5.1) imply that conditions (3) and (4) are equivalent. The other assertions follow from Proposition 5.5. \square

Given a spacelike surface germ $\mathbf{X} : (U, (x_0, y_0)) \rightarrow (\mathbb{R}_1^4, \mathbf{X}(x_0, y_0))$, we call each set

$$(\mathbf{X}^{-1}(LHP(\mathbf{v}^\pm, c^\pm)), (x_0, y_0))$$

a *tangent lightlike hyperplane indicatrix germ* of \mathbf{X} , where $\mathbf{v}^\pm = \mathbf{e}_1 \pm \mathbf{e}_2(x_0, y_0)$ and $c^\pm = \langle \mathbf{X}(x_0, y_0), \mathbf{v}^\pm \rangle$. Moreover, in view of the above results, we can borrow some basic invariants, such as the \mathcal{K} -invariants for function germs, from the singularity theory on function germs. The local ring of a function germ is a complete \mathcal{K} -invariant for generic function germs. It isn't, however, a numerical invariant. On the other hand, the \mathcal{K} -codimension (or, Tyurina number) of a function germ is a numerical \mathcal{K} -invariant of function germs [14]. We denote it by

$$\text{L-ord}^\pm(\mathbf{X}, (x_0, y_0)) = \dim_{\mathbb{R}} \frac{C_{(x_0, y_0)}^\infty(U)}{\langle \tilde{h}_{\mathbf{v}_0^\pm}(x, y), \tilde{h}_{\mathbf{v}_0^\pm, x}(x, y), \tilde{h}_{\mathbf{v}_0^\pm, y}(x, y) \rangle}.$$

$\text{L-ord}^\sigma(\mathbf{x}, u_0)$ is usually called *the \mathcal{K} -codimension of $\tilde{h}_{\mathbf{v}_0^\sigma}$* , where $\sigma = \pm$. However, we shall refer here to it as the *order of contact with the tangent lightlike hyperplane* at $\mathbf{X}(x_0, y_0)$. We also have the notion of corank of function germs,

$$\text{L-corank}^\sigma(\mathbf{X}, (x_0, y_0)) = 2 - \text{rank Hess}(\tilde{h}_{\mathbf{v}_0^\sigma}(x_0, y_0)),$$

where $\mathbf{v}_0^\pm = \mathbf{e}_1 \pm \mathbf{e}_2(x_0, y_0)$.

Then Proposition 4.1 tells us that $\mathbf{X}(x_0, y_0)$ is a L^σ -parabolic point if and only if

$$\text{L-corank}^\sigma(\mathbf{X}, (x_0, y_0)) \geq 1.$$

Moreover, $\mathbf{X}(x_0, y_0)$ is a lightlike umbilic point if and only if

$$\text{L-corank}^\sigma(\mathbf{X}, (x_0, y_0)) = 2.$$

On the other hand, a function germ $f : (\mathbb{R}^{n-1}, \mathbf{a}) \rightarrow \mathbb{R}$ has the A_k -type singularity if f is \mathcal{K} -equivalent to the germ $\pm u_1^2 \pm \dots \pm u_{n-2}^2 + u_{n-1}^{k+1}$. If $\text{L-corank}^\sigma(\mathbf{X}, (x_0, y_0)) = 1$, the extended lightcone height function $\tilde{h}_{\mathbf{v}_0^\sigma}$ has, generically, a singularity of type A_k at (x_0, y_0) . In this case, we have that $\text{H-ord}^\sigma(\mathbf{x}, u_0) = k$. The number k is equal to the order of contact in the classical sense (cf., [4]). This justifies why we call $\text{L-ord}^\sigma(\mathbf{X}, (x_0, y_0))$ the order of contact with the tangent lightlike hyperplane at $\mathbf{X}(x_0, y_0)$.

6. Classification of singularities of lightcone Gauss maps and lightcone pedal surfaces. This section is devoted to the study of the generic singularities of lightcone Gauss maps and lightcone pedal surfaces. We denote by $\text{Emb}_s(U, \mathbb{R}_1^4)$ the space of spacelike embeddings with the Whitney C^∞ -topology, where $U \subset \mathbb{R}^2$ is an open subset. The set of lightlike parabolic points (i.e., $\{p \in M \mid \mathcal{K}_\ell(1 \pm 1)(p) = 0\}$) is called a *lightlike parabolic set*.

THEOREM 6.1. *There exists an open dense subset $\mathcal{O} \subset \text{Emb}_s(U, \mathbb{R}_1^4)$ such that for any $\mathbf{X} \in \mathcal{O}$, the following conditions hold:*

(1) *Each component of the lightlike parabolic set is a regular curve. We call such a curve the lightlike parabolic curve.*

(2) *The lightcone pedal surface LP_M^σ is a cuspidal edge at each point of the lightlike parabolic curve except at isolated points. At these points LP_M^σ is a swallowtail.*

Here, a map germ $f : (\mathbb{R}^2, \mathbf{a}) \rightarrow (\mathbb{R}^3, \mathbf{b})$ is called a cuspidal edge if it is \mathcal{A} -equivalent to the germ (u_1, u_2^2, u_2^3) (cf., Fig. 1) and a swallowtail if it is \mathcal{A} -equivalent to the germ $(3u_1^4 + u_1^2u_2, 4u_1^3 + 2u_1u_2, u_2)$ (cf., Fig.1).

In order to prove Theorem 6.1, we consider the function $\mathcal{H} : \mathbb{R}_1^4 \times LC_+^* \rightarrow \mathbb{R}$ given in §5. We claim that \mathcal{H}_v is a submersion for any $\mathbf{v} \in LC_+^*$, where $\mathcal{H}_v(\mathbf{x}) = \mathcal{H}(\mathbf{x}, \mathbf{v})$. For any $\mathbf{X} \in \text{Emb}_s(U, \mathbb{R}_1^4)$, we have $\tilde{H} = \mathcal{H} \circ (\mathbf{X} \times id_{LC_+^*})$. We also have the ℓ -jet extension

$$j_1^\ell \tilde{H} : U \times LC_+^* \rightarrow J^\ell(U, \mathbb{R})$$

defined by $j_1^\ell \tilde{H}((x, y), \mathbf{v}) = j^\ell \tilde{h}_v(x, y)$. Consider the trivialization $J^\ell(U, \mathbb{R}) \equiv U \times \mathbb{R} \times J^\ell(2, 1)$. Given any submanifold $Q \subset J^\ell(2, 1)$, we denote $\tilde{Q} = U \times \{0\} \times Q$. Then we have the following proposition as a corollary of Lemma 6 in Wassermann [20]. (See also Montaldi [17] and Looijenga [13]).

PROPOSITION 6.2. *Let Q be a submanifold of $J^\ell(2, 1)$. Then the set*

$$T_Q = \{ \mathbf{x} \in \text{Emb}_s(U, \mathbb{R}_1^4) \mid j_1^\ell H \text{ is transversal to } \tilde{Q} \}$$

is a residual subset of $\text{Emb}_s(U, \mathbb{R}_1^4)$. If Q is a closed subset, then T_Q is open.

The proof of Theorem 6.1 is easily obtained from this by considering the \mathcal{K} -orbits in $J^\ell(2, 1)$, so we omit the detailed discussion. The assertion of Theorem 6.1 can be interpreted in the sense that the Legendrian lift of the lightcone pedal surface LP_M^\pm of $\mathbf{X} \in \mathcal{O}$ is Legendrian stable at each point. Since the Legendrian lift is the Legendrian covering of the Lagrangian lift of LG_M^\pm , it has been known that the corresponding singularities of LG_M^\pm are folds or cusps [1]. Hence, we have the following corollary.

COROLLARY 6.3. *Let $\mathcal{O} \subset \text{Emb}_s(U, \mathbb{R}_1^4)$ be the open dense subset provided by Theorem 6.1. Given any $\mathbf{X} \in \mathcal{O}$, the following assertions hold:*

(1) *A lightlike parabolic point $(x_0, y_0) \in U$ is a fold of the lightcone Gauss map LG_M^σ if and only if it is a cuspidal edge of the lightcone pedal surface LP_M^σ .*

(2) *A lightlike parabolic point $(x_0, y_0) \in U$ is a cusp of the lightcone Gauss map LG_M^σ if and only if it is a swallowtail of the lightcone pedal surface LP_M^σ .*

Here, a map germ $f : (\mathbb{R}^2, \mathbf{a}) \rightarrow (\mathbb{R}^2, \mathbf{b})$ is called a fold if it is \mathcal{A} -equivalent to the germ (u_1, u_2^2) and a cusp if it is \mathcal{A} -equivalent to the germ $(u_1, u_2^3 + u_1u_2)$.

Following the terminology of Whitney [21], we say that a surface $\mathbf{X} : U \rightarrow \mathbb{R}_1^4$ has an excellent lightcone pedal surface LP_M^σ if the Legendrian lift of LP_M^σ is a stable Legendrian immersion at each point. In this case, the lightcone pedal surface LP_M^σ has only cuspidal edges and swallowtails as singularities. Theorem 6.1 implies that a spacelike surface with an excellent lightcone pedal surface is generic in the space of all spacelike surfaces in \mathbb{R}_1^4 . We now analyze the geometric meanings of cuspidal edges and swallowtails of the lightcone pedal surface. We have the following results analogous to those of Banchoff et al [2].

THEOREM 6.4. Let $LP_M^\sigma : (U, (x_0, y_0)) \longrightarrow (\mathbb{R}_1^4, p_0)$ be the excellent lightcone pedal surface of a spacelike surface \mathbf{X} and $\tilde{h}_{v_0^\sigma} : (U, (x_0, y_0)) \longrightarrow \mathbb{R}$ be the extended lightcone height function germ at $v_0^\pm = e_1 \pm e_2(p_0)$, where $\sigma = \pm$. Then we have the following:

- (1) (x_0, y_0) is a lightlike parabolic point of \mathbf{X} if and only if $L\text{-corank}^\sigma(\mathbf{X}, (x_0, y_0)) = 1$.
- (2) If (x_0, y_0) is a lightlike parabolic point of \mathbf{X} , then $\tilde{h}_{v_0^\sigma}$ has the A_k -type singularity for $k = 2, 3$.
- (3) Suppose that (x_0, y_0) is a lightlike parabolic point of \mathbf{X} . Then the following conditions are equivalent:
 - (a) LP_M^σ has a cuspidal edge at (x_0, y_0)
 - (b) $\tilde{h}_{v_0^\sigma}$ has an A_2 -type singularity.
 - (c) $L\text{-ord}^\sigma(\mathbf{X}, (x_0, y_0)) = 2$.
 - (d) The tangent lightlike hyperplane indicatrix is an ordinary cusp, where a curve $C \subset \mathbb{R}^2$ is said to be an ordinary cusp if it is diffeomorphic to the curve given by $\{(u_1, u_2) \mid u_1^2 - u_2^3 = 0\}$.
 - (e) For each $\varepsilon > 0$, there exist two distinct points $(x_i, y_i) \in U$ ($i = 1, 2$) such that

$$\|(x_0, y_0) - (x_i, y_i)\| < \varepsilon$$

for $i = 1, 2$, both of (x_i, y_i) are not lightlike parabolic points and the tangent lightlike hyperplanes to $M = \mathbf{x}(U)$ at (x_i, y_i) are parallel.

- (4) Suppose that (x_0, y_0) is a lightlike parabolic point of \mathbf{X} . Then the following conditions are equivalent:
 - (a) LP_M^σ has a swallowtail at (x_0, y_0)
 - (b) $\tilde{h}_{v_0^\sigma}$ has an A_3 -type singularity.
 - (c) $L\text{-ord}^\sigma(\mathbf{X}, (x_0, y_0)) = 3$.
 - (d) The tangent lightlike hyperplane indicatrix is a point or a tachnodal, where a curve $C \subset \mathbb{R}^2$ is called a tachnodal if it is diffeomorphic to the curve given by $\{(u_1, u_2) \mid u_1^2 - u_2^4 = 0\}$.
 - (e) For each $\varepsilon > 0$, there exist three distinct points $(x_i, y_i) \in U$ ($i = 1, 2, 3$) such that

$$\|(x_0, y_0) - (x_i, y_i)\| < \varepsilon$$

for $i = 1, 2, 3$ and the tangent lightlike hyperplanes to $M = \mathbf{x}(U)$ at (x_i, y_i) are parallel.

- (f) For each $\varepsilon > 0$, there exist two distinct points $(x_i, y_i) \in U$ ($i = 1, 2$) such that

$$\|(x_0, y_0) - (x_i, y_i)\| < \varepsilon$$

for $i = 1, 2$ and the tangent lightlike hyperplanes to $M = \mathbf{x}(U)$ at (x_i, y_i) are equal.

Proof. We have shown that (x_0, y_0) is a lightlike parabolic point if and only if

$$L\text{-corank}^\sigma(\mathbf{X}, (x_0, y_0)) \geq 1.$$

We have that $L\text{-corank}^\sigma(\mathbf{X}, (x_0, y_0)) \leq 2$. Since the extended lightcone height function germ, $\tilde{H} : (U \times, LC_+^*, ((x_0, y_0), v_0)) \longrightarrow \mathbb{R}$, can be considered as a generating family of the Legendrian lift of LP_M^σ , $\tilde{h}_{v_0^\sigma}$ has only A_k -type singularities ($k = 1, 2, 3$). This means that the corank of the Hessian matrix of $\tilde{h}_{v_0^\sigma}$ at a lightlike parabolic point is 1. The assertion (2) follows analogously. By the same reason, the conditions (3);(a),(b),(c) (respectively, (4); (a),(b),(c)) are equivalent. If the height function germ $h_{v_0^\sigma}$ has

the A_2 -type singularity, then it is \mathcal{K} -equivalent to the germ $\pm u_1^2 + u_2^3$. Since the \mathcal{K} -equivalence preserves the diffeomorphism type of zero level sets, the tangent lightlike hyperplane indicatrix is diffeomorphic to the curve given by $\pm u_1^2 + u_2^3 = 0$ which is an ordinary cusp. The normal form for the A_3 -type singularity is given by $\pm u_1^2 + u_2^4$, so the tangent lightlike hyperplane indicatrix is diffeomorphic to the curve $\pm u_1^2 + u_2^4 = 0$. This means that the condition (3),(d) (respectively, (4),(d)) is also equivalent to the other conditions.

Suppose that (x_0, y_0) is a lightlike parabolic point, then the lightcone Gauss map has only folds or cusps as singularities. If the point (x_0, y_0) is a fold point, there is a neighbourhood of (x_0, y_0) at which the lightcone Gauss map is 2 to 1 except at the lightlike parabolic curve (i.e, fold curve). By Lemma 5.2, the condition (3), (e) is satisfied. If the point (x_0, y_0) is a cusp, the critical value set is an ordinary cusp. By considering its normal form we see that the lightcone Gauss map is 3 to 1 inside the region of the critical value. Moreover, the point (x_0, y_0) lies in the closure of this region from which it we get condition (4),(e). We also observe that nearby of a cusp point, there are 2 to 1 points which approach to (x_0, y_0) . However, one of these points is always a lightlike parabolic point. Since no other singularities appear in this case, we have that the condition (3), (e) (respectively, (4),(e)) characterizes a fold (respectively, a cusp).

If we consider the lightcone pedal surface instead of the lightcone Gauss map we have that its only singularities are cuspidal edges and swallowtails. we observe that we can always find a self intersection curve (cf., Fig. 1) approaching a swallowtail point (x_0, y_0) . This means that there are two distinct point (x_i, y_i) ($i = 1, 2$) such that $LP_M^\sigma(x_1, y_1) = LP_M^\sigma(x_2, y_2)$, which by Lemma 5.2, means that the tangent lightlike hyperplanes to $M = \mathbf{X}(U)$ at $(x_i$ and y_i are the same. Since no other singularities present this particularity, we can conclude that the condition (4),(f) characterizes the swallowtail points of PL_M^σ and the proof is completed. \square

More detailed properties of spacelike surfaces in Minkowski 4-space will be discussed elsewhere.

7. Examples. Surfaces contained in the Euclidean space $\mathbb{R}^3 = \{(0, x_2, x_3, x_4) \mid x_i \in \mathbb{R}\}$ or in the hyperbolic space $H_+^3(-1) = \{\mathbf{x} \in \mathbb{R}_1^4 \mid \langle \mathbf{x}, \mathbf{x} \rangle = -1\}$ provide special examples of spacelike surfaces. We remark that the singularity theory for the lightcone Gauss map of such surfaces coincides with the singularity theory for ordinary (Euclidean) Gauss maps [2, 3] or hyperbolic Gauss maps [8] respectively. In order to illustrate our results we propose next some examples of spacelike surfaces which lie neither in the Euclidean space \mathbb{R}^3 nor in the hyperbolic space $H_+^3(-1)$. Given an immersion in the Monge form

$$\mathbf{X}(x, y) = (f_1(x, y), f_2(x, y), x, y)$$

with $f_i(\underline{0}, 0) = 0$ and $\partial f_i / \partial x(0, 0) = \partial f_i / \partial y(0, 0) = 0$, ($i = 1, 2$), we can choose $\mathbf{v}_0^\pm = \mathbf{e}_1 \pm \mathbf{e}_2(0, 0) = (1, \pm 1, 0, 0)$ as in §3. Therefore, the lightcone height function is given by

$$h_{\mathbf{v}_0^\pm}(x, y) = -f_1(x, y) \pm f_2(x, y),$$

so that $h_{\mathbf{v}_0^\pm}(0, 0) = 0$. Moreover, the lightlike tangent indicatrix at the origin $\underline{0}$ is

$$\mathbf{X}^{-1}(LHP(\mathbf{v}^\pm, 0)) = \{(x, y) \in U \mid -f_1(x, y) \pm f_2(x, y) = 0\}.$$

Since $\mathbf{X}(U) = M$ is a spacelike surface, $f_i(x, y)$ ($i = 1, 2$) must satisfy the condition

$$\left(\frac{\partial f_2}{\partial x}(x, y)\right)^2 + 1 > \left(\frac{\partial f_1}{\partial x}(x, y)\right)^2, \quad \left(\frac{\partial f_2}{\partial y}(x, y)\right)^2 + 1 > \left(\frac{\partial f_1}{\partial y}(x, y)\right)^2.$$

We observe that choosing $f_1 = \pm f_2$, ensures that this condition is satisfied.

EXAMPLE 7.1. Take the functions $f_1(x, y) = -\frac{1}{3}x^3 + \frac{1}{2}y^2$ and $f_2(x, y) = -f_1(x, y)$, then

$$h_{v_0^+}(x, y) = \left(\frac{2}{3}x^3 - y^2\right).$$

So the lightlike tangent indicatrix is the ordinary cusp $y^2 = \frac{2}{3}x^3$. It follows that the lightcone pedal surface LP_M^+ is the cuspidal edge at the origin.

EXAMPLE 7.2. Take now $f_1(x, y) = \frac{1}{2}(x^2 - y^4)$ and $f_2(x, y) = -f_1(x, y)$, then

$$h_{v_0^+}(x, y) = (x^2 - y^4).$$

Clearly, the lightlike tangent indicatrix is the tacnode $x^2 = y^4$ and the lightcone pedal surface LP_M^+ has a swallowtail at the origin in this case.

REFERENCES

- [1] V. I. ARNOL'D, S. M. GUSEIN-ZADE AND A. N. VARCHENKO, *Singularities of Differentiable Maps vol. I*, Birkhäuser (1986).
- [2] T. BANCHOFF, T. GAFFNEY AND C. MCCRORY, *Cusps of Gauss Mappings*, Research Notes in Mathematics, 55 Pitman, London (1982).
- [3] D. BLEECKER AND L. WILSON, *Stability of Gauss maps*, Illinois J. Math., 22 (1978), pp. 279–289.
- [4] J. W. BRUCE AND P. J. GIBLIN, *Curves and singularities (second edition)*, Cambridge University press, (1992).
- [5] C. L. EPSTEIN, *The hyperbolic Gauss map and quasiconformal reflections*, J. Reine Angew. Math., 372 (1986), pp. 96–135.
- [6] J. EHLERS AND E. T. NEWMAN, *The theory of caustics and wave front singularities with physical applications*, Journal of Mathematical Physics, 41 (2000), pp. 3344–3378.
- [7] S. IZUMIYA, D-H. PEI AND T. SANO, *The lightcone Gauss map and the lightcone developable of a spacelike curve in Minkowski 3-space*, Glasgow. Math. J., 42 (2000), pp. 75–89.
- [8] S. IZUMIYA, D-H. PEI AND T. SANO, *Singularities of hyperbolic Gauss maps*, Proceedings of the London Math. Soc., 86 (2003), pp. 485–512.
- [9] S. IZUMIYA, M. KOSSOWSKI, D. PEI AND M. C. ROMERO FUSTER, *Singularities of lightlike hypersurfaces in Minkowski 4-space*, in preparation.
- [10] M. KOSSOWSKI, *The S^2 -valued Gauss maps and split total curvature of space-like codimension-2 surface in Minkowski space*, J. London Math. Soc.(2), 40 (1989), pp. 179–192.
- [11] M. KOSSOWSKI, *The intrinsic conformal structure and Gauss map of a light-like hypersurface in Minkowski space*, Transactions of the American mathematical society, 316:1 (1989), pp. 369–383.
- [12] J. A. LITTLE, *On singularities of submanifolds of high dimensional Euclidean space*, Annali Mat.Pura et Appl. (ser. 4A), 83 (1969), pp. 261–336.
- [13] E. J. N. LOOIJENGA, *Structural stability of smooth families of C^∞ -functions*, Thesis, Univ. Amsterdam (1974).
- [14] J. MARTINET, *Singularities of Smooth Functions and Maps*, London Math. Soc. Lecture Note Series, Cambridge Univ. Press, 58 (1982).
- [15] J. N. MATHER, *Stability of C^∞ -mappings IV: Classification of stable germs by \mathbb{R} algebras*, Publ. Math. I.H.E.S., 37 (1970), pp. 223–248.
- [16] J. A. MONTALDI, *On contact between submanifolds*, Michigan Math. J., 33 (1986), pp. 81–85.

- [17] J. A. MONTALDI, *On generic composites of maps*, Bull. London Math. Soc., 23 (1991), pp. 81–85.
- [18] B. O'NEILL, *Semi-Riemannian Geometry*, Academic Press, New York (1983).
- [19] I. PORTEOUS, *The normal singularities of submanifold*, J. Diff. Geom., 5 (1971), pp. 543–564.
- [20] G. WASSERMANN, *Stability of Caustics*, Math. Ann., 2210 (1975), pp. 443–50.
- [21] H. WHITNEY, *On singularities of mappings of Euclidean spaces I*, Ann. of Math., 62 (1955), pp. 374–410.
- [22] V. M. ZAKALYUKIN, *Lagrangian and Legendrian singularities*, Funct. Anal. Appl., 10 (1976), pp. 23–31.
- [23] V. M. ZAKALYUKIN, *Reconstructions of fronts and caustics depending one parameter and versality of mappings*, J. Sov. Math., 27 (1984), pp. 2713–2735.