

Among the orbits produced by a central force varying as the n th power of the radius vector are included the isogonal trajectories of the curves

$$(13) \quad y' = -\tan \frac{1}{2}(n+1)\theta.$$

This construction yields in the case $n = -2$ (Newtonian law) parabolas with focus at the origin; in the case $n = 1$ (elastic law) equilateral hyperbolas with center at the origin; in the case $n = -5$ the circles through the origin; and in the case $n = -3$ equiangular spirals with pole at the origin.

COLUMBIA UNIVERSITY.

ON THE EQUATIONS OF QUARTIC SURFACES IN TERMS OF QUADRATIC FORMS.

BY DR. C. H. SISAM.

(Read before the American Mathematical Society, September 5, 1907.)

THE quartic surfaces whose equations are of the form

$$\phi_2(A, B, C, D) = 0,$$

where ϕ_2 , A , B , C and D are quaternary quadratic forms, were the subject of a paper by H. Durrande in the *Nouvelles Annales*.* By counting the number of constants involved, Durrande concluded that the most general quartic surface could be represented by an equation of the above form. He recognized, however, that his reasoning was not rigorous.

It will here be shown that the coefficients of the quartic surface determined by this equation are not independent, but are subject to a single condition. It will also be shown that the equation of a general quartic surface can be written in the form

$$\theta_2(A, B, C, D, E) = 0,$$

where θ_2 is a quinary quadratic form.

Let $\phi_2 = 0$ be reduced to the form

$$(1) \quad A^2 + B^2 + C^2 + D^2 = 0,$$

where

$$A \equiv \sum a_{ij} x_i x_j \quad (i \leq j \leq 4)$$

* Durrande, *Nouvelles Annales*, ser. 2, vol. 9, p. 410.

tically. This may be seen by substituting the particular values $\alpha_{11} = b_{22} = c_{33} = d_{44} = 1$, the remaining α_{ij} , etc., being zero. Hence these quantities F_{ijkl} are independent.

Finally, consider the quartic

$$\theta_2(A, B, C, D, E) = 0.$$

Let this equation be reduced to

$$A^2 + B^2 + C^2 + D^2 + E^2 = 0.$$

Equate the coefficients of the terms of this equation to the corresponding terms of

$$\sum k_{ijkl} x_i x_j x_k x_l = 0 \quad (i \leq j \leq k \leq l \leq 4)$$

and determine the jacobian matrix as before. That the determinants of this matrix do not all vanish identically is seen by taking for A, B , etc., the particular expressions

$$A \equiv x_1^2, \quad B \equiv x_2^2, \quad C \equiv x_3^2, \quad D \equiv x_4^2, \quad E \equiv x_1 x_2.$$

Since these determinants do not vanish identically the k_{ijkl} are independent. Hence *the equation of an arbitrary quartic surface can be put into the form*

$$A^2 + B^2 + C^2 + D^2 + E^2 = 0.$$

URBANA, ILL.,
September, 1907.

SYMBOLIC LOGIC.

L'Algèbre de la Logique. Par LOUIS COUTURAT. Collection Scientia, No. 24. Gauthier-Villars, Paris, 1905. 100 pp.

Symbolic Logic and its Applications. By HUGH MACCOLL. Longmans, Green, and Co., London, 1906. xi + 141 pp.

The Development of Symbolic Logic; a Critical-Historical Study of the Logical Calculus. By A. T. SHEARMAN. Williams and Norgate, London, 1906. xi + 242 pp.

SYMBOLIC logic is in the interesting though somewhat precarious state of being little known, less used, and much scorned by the majority of mathematicians and philosophers, for whom it might supposedly offer a region of intimate contact and