

of a homeomorphism in the sense of Antoine.* From Theorem 2, p. 394, of the paper just cited, we can obtain a theorem for A-extending a homeomorphism between two subsets of spheres.

UNIVERSITY OF BUFFALO

A PROPERTY OF THE SOLUTIONS OF $t^2 - du^2 = 4$

BY GORDON PALL

Let p be any odd prime not dividing d . The integral solutions $t_i, u_i, (i=0, \pm 1, \dots)$, † of $t^2 - du^2 = 4$ have the following property.

THEOREM. *Let $m+n=r+s$. Let v stand for t or u . Then $v_m + v_n \equiv v_r + v_s \pmod{p}$ if and only if the terms are congruent in pairs; ‡ the same holds for each of*

$$v_m - v_n \equiv v_r - v_s, \quad v_m + v_n \equiv -(v_r + v_s), \quad v_m - v_n \equiv -(v_r - v_s).$$

For if $m+n$ is even and $v=u$, we can write $m=h+i, n=h-i, r=h+j, s=h-j$, whence

$$u_m + u_n = u_h t_i, \quad u_r + u_s = u_h t_j;$$

if $u_h=0$, then $u_m \equiv -u_n$; if $t_i \equiv t_j$, known conditions for two u 's or t 's to be congruent show that $u_m = u_r$ or u_s . The remaining cases are similar. If $m+n$ is odd, we transpose terms, and find with a little attention to parities ($u_i = -u_{-i}, t_i = t_{-i}$) one or other of the former cases.

MCGILL UNIVERSITY

* H. M. Gehman, *On extending a correspondence in the sense of Antoine*, American Journal of Mathematics, vol. 51 (1929), pp. 385-396.

† For notations see, for example, Pall, Transactions of this Society, vol. 35 (1933), p. 501.

‡ That is, $v_m \equiv -v_n, v_r, \text{ or } v_s$.