

# INTRINSIC METRICS ON COMPLEX MANIFOLDS

BY SHOSHICHI KOBAYASHI<sup>1</sup>

Communicated by S. Smale, January 17, 1967

**1. Definition of intrinsic pseudometric.** Let  $M$  be a (connected) complex manifold. We shall define a pseudometric  $d$  on  $M$  in a natural manner so that it depends only on the complex structure of  $M$  and nothing else.

Let  $D$  be the open unit disk in the complex plane and  $\rho$  the distance on  $D$  defined by the Poincaré-Bergman metric of  $D$ . Given two points  $p$  and  $q$  of  $M$ , choose the following objects:

- (1) points  $p = p_0, p_1, \dots, p_{k-1}, p_k = q$  of  $M$  and
- (2) points  $a_1, \dots, a_k, b_1, \dots, b_k$  of  $D$  and holomorphic mappings  $f_1, \dots, f_k$  of  $D$  into  $M$  such that  $f_i(a_i) = p_{i-1}$  and  $f_i(b_i) = p_i$  for  $i = 1, \dots, k$ .

For each choice of points and mappings satisfying (1) and (2), consider the number  $\rho(a_1, b_1) + \dots + \rho(a_k, b_k)$ . Let  $d(p, q)$  be the infimum of the numbers obtained in this manner for all possible choices. It is easy to verify that  $d$  is a pseudometric on  $M$  in the sense that

$$d(p, q) \geq 0, \quad d(p, q) = d(q, p), \quad d(p, q) + d(q, r) \geq d(p, r)$$

for  $p, q, r \in M$ . The following two propositions are immediate from the definition of  $d$ .

**PROPOSITION 1.** *Let  $M$  and  $N$  be two complex manifolds and  $d_M$  and  $d_N$  the intrinsic pseudometrics of  $M$  and  $N$ . Then every holomorphic mapping  $f: M \rightarrow N$  is distance-decreasing in the sense that*

$$d_M(p, q) \geq d_N(f(p), f(q)) \quad \text{for } p, q \in M.$$

*In particular, every holomorphic transformation of  $M$  is distance-preserving with respect to  $d_M$ .*

**PROPOSITION 2.** *For the complex Euclidean space  $C^n$ , the pseudometric  $d$  is trivial, i.e.,  $d(p, q) = 0$  for all  $p, q \in C^n$ .*

The following proposition follows from the Schwarz-Pick lemma.

**PROPOSITION 3.** *For the unit disk  $D$ , the pseudometric  $d$  coincides with the distance  $\rho$  defined by the Poincaré-Bergman metric.*

---

<sup>1</sup> Supported partially by NSF Grant GP-5798.

The following theorem follows from a generalized Schwarz-Pick lemma (see [3] and [4]).<sup>2</sup>

**THEOREM 4.** *Let  $M$  be a hermitian manifold whose holomorphic sectional curvature is negative and bounded away from zero. Then its pseudometric  $d$  is a metric, i.e.,  $d(p, q) = 0$  implies  $p = q$ .*

Theorem 4 applies to all bounded domains of  $C^n$  as well as to all Riemann surfaces of hyperbolic type.

**2. Relationship with Carathéodory metric.** Following Carathéodory [2] we define another pseudometric  $d'$  on a complex manifold  $M$ . Given two points  $p$  and  $q$  of  $M$ , let  $d'(p, q)$  be the supremum of  $\rho(f(p), f(q))$  for all holomorphic mappings  $f$  of  $M$  into the unit disk  $D$ .

It is easy to see that Propositions 1, 2 and 3 above hold also for the Carathéodory pseudometric  $d'$ . A necessary and sufficient condition for  $d'$  to be a metric on  $M$  is that there are sufficiently many bounded holomorphic functions on  $M$  so that they separate the points of  $M$ .

The following proposition is immediate from Propositions 1 and 3 and shows that whenever the Carathéodory pseudometric  $d'$  is a metric, our pseudometric  $d$  is also a metric.

**PROPOSITION 5.** *For any complex manifold  $M$ ,  $d$  is greater than or equal to  $d'$ , i.e.,*

$$d(p, q) \geq d'(p, q) \quad \text{for } p, q \in M.$$

**3. Applications.** The following theorem which follows from Propositions 1 and 2 and Theorem 4 may be considered as a generalization of Picard Theorem which states that an entire function with more than one finite lacunary value reduces to a constant function.

**THEOREM 6.** *Let  $M$  be an  $n$ -dimensional complex manifold which admits a hermitian metric whose holomorphic sectional curvature is negative and bounded away from zero. Then every holomorphic mapping  $f$  of  $C^m$  into  $M$  is a constant map.*

**COROLLARY.** *The complex Euclidean space  $C^m$  does not admit a hermitian metric whose holomorphic sectional curvature is negative and bounded away from zero.*

The condition "bounded away from zero" is essential. In fact,  $C$

---

<sup>2</sup> The results proved for Kähler manifolds in [4] hold for hermitian manifolds (with respect to the hermitian connection in the sense of Chern). Proofs there remain valid in the hermitian case.

(and hence  $C^m$ ) admits a complete Kähler metric of negative holomorphic sectional curvature, e.g.,

$$(1 + z\bar{z}) dzd\bar{z}.$$

**THEOREM 7.** *Let  $M$  be a hermitian manifold whose holomorphic sectional curvature is negative and bounded away from zero. Then the group of holomorphic transformations of  $M$  is a Lie group with compact isotropy subgroups (with respect to the compact-open topology).*

In fact, the group in question is a closed subgroup of the group of isometries of  $M$  with respect to the intrinsic metric  $d$  introduced above. By a classical theorem of Van Dantzig and van der Waerden (see Theorem 4.7 and Corollary 4.8 of Chapter I in [5]) the group of isometries of a locally compact metric space is locally compact and its isotropy subgroups are all compact with respect to the compact-open topology. Theorem 7 follows now from a well-known theorem of Bochner-Montgomery [1].

**COROLLARY 8.** *If, in Theorem 7,  $M$  is moreover compact, then the group of holomorphic transformations of  $M$  is a finite group.*

The results in this section have been obtained by Wu [6] using the notion of normal families.

#### BIBLIOGRAPHY

1. S. Bochner and D. Montgomery, *Locally compact groups of differentiable transformations*, Ann. of Math. 47 (1946), 639–653.
2. C. Carathéodory, *Über das Schwarzsche Lemma bei analytischen Funktionen von zwei komplexen Veränderlichen*, Math. Ann. 97 (1926), 76–98.
3. H. Grauert and H. Reckziegel, *Hermiteische Metriken und normale Familien holomorpher Abbildungen*, Math. Z. 89 (1965), 108–125.
4. S. Kobayashi, *Distance, holomorphic mappings and the Schwarz's lemma*, J. Math. Soc. Japan (to appear).
5. S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Interscience Tracts No. 15, Interscience, New York, 1963.
6. H. H. Wu, *Normal families of holomorphic mappings and the theorem of Bloch in several complex variables* (to appear).

UNIVERSITY OF CALIFORNIA, BERKELEY