

AUTOMORPHISMS OF A FREE ASSOCIATIVE ALGEBRA OF RANK 2

BY ANASTASIA CZERNIAKIEWICZ

Communicated by Hyman Bass, May 28, 1971

We announce here that the answer to the following conjecture [3, p. 197] is in the affirmative:

If R is a Generalized Euclidean Domain then every automorphism of the free associative algebra of rank 2 over R is tame, i.e. a product of elementary automorphisms.

We state here the necessary steps to prove the conjecture; detailed proofs will appear in [4] and [5].

Notation. R stands for a commutative domain with 1;

$R\langle x, y \rangle$ is the free associative algebra of rank 2 over R , on the *free* generators x and y ;

$R(\tilde{x}, \tilde{y})$ is the polynomial algebra over R on the *commuting* indeterminates \tilde{x} and \tilde{y} .

We write $R\langle x, y \rangle$ as a bigraded algebra

$$R\langle x, y \rangle = \bigoplus_{r, s \geq 0} \mathfrak{A}_r^s$$

where the subindex denotes the homogeneous degree and the upper index stands for the degree in x . We will write $P = \sum P_r^s$ where $P_r^s \in \mathfrak{A}_r^s$ for every $P \in R\langle x, y \rangle$.

The elementary automorphisms of $R\langle x, y \rangle$ are by definition the following:

- (i) $\sigma \in \text{Aut}_R(R\langle x, y \rangle)$; $\sigma(x) = y$; $\sigma(y) = x$.
- (ii) $\varphi_{\alpha, \beta} \in \text{Aut}_R(R\langle x, y \rangle)$, α, β units of R ;

$$\varphi_{\alpha, \beta}(x) = \alpha x; \quad \varphi_{\alpha, \beta}(y) = \beta y.$$

- (iii) $\tau_{f(y)} \in \text{Aut}_R(R\langle x, y \rangle)$, where $f(y)$ is any polynomial that does not depend on x ;

$$\tau_{f(y)}(x) = x + f(y); \quad \tau_{f(y)}(y) = y.$$

In a completely parallel way one defines the elementary automorphisms of $R(\tilde{x}, \tilde{y})$.

AMS 1970 subject classifications. Primary 16A06, 16A72; Secondary 20F55, 16A02.

Key words and phrases. Free associative algebra, endomorphisms, automorphisms, elementary automorphisms, tame automorphisms, wild automorphisms, polynomial rings, euclidean domains.

THEOREM 1. *The map*

$$\text{Aut}_R(R\langle x, y \rangle) \rightarrow \text{Aut}_R(R(\bar{x}, \bar{y}))$$

induced by the abelianization functor is a monomorphism.

The proof of Theorem 1 is an immediate corollary of the more technical result:

THEOREM 2. *Let $P, Q, E \in R\langle x, y \rangle$ satisfy the following requirements:*

- (i) $P_0^0 = Q_0^0 = 0, E_0 = E_1 = 0.$
- (ii) $P_n^0 = 0$ for all $n \geq 1; Q_m^0 = 0$ for all $m \geq 2; E_r^0 = 0$ for all $r \geq 2.$
- (iii) $E(P, Q) = xy - yx.$

Then we conclude

$$P = P_1^1 = \alpha x; \quad Q = Q_1^0 + \sum_n Q_n^n = \beta y + f(x);$$

$$E = (\alpha\beta)^{-1}(xy - yx), \quad \alpha, \beta \text{ are units of } R.$$

The proof of Theorem 2 is obtained with slight modifications from the proof of the main theorem in [4].

In fact, for every rational number $\lambda \geq 0$ we define

$$\chi_\lambda = \left\{ P_a^\alpha; a > 1, \alpha \geq 1, \frac{\alpha - 1}{a - 1} = \lambda \right\}$$

$$\cup \left\{ Q_b^\beta; b > 1, \beta \geq 0, \frac{\beta}{b - 1} = \lambda \right\}$$

$$\cup \left\{ E_m^\mu; m > 2, \mu \geq 1, \frac{\mu - 1}{m - 2} = \lambda \right\}.$$

As we have only a finite set of rational numbers λ for which $\chi_\lambda \neq \{0\}$ we use the ordering of the rational numbers to prove inductively that if $\chi_\lambda = \{0\}$ for all $\lambda < \lambda_0$ then $\chi_{\lambda_0} = \{0\}$.

To achieve this purpose we exhibit a relation of algebraic dependence between two elements of χ_{λ_0} and using a result of P. M. Cohn [2] about homogeneous elements of $R\langle x, y \rangle$ we conclude $\chi_{\lambda_0} = \{0\}$.

COROLLARY. *If R is a generalized euclidean domain then every automorphism of $R\langle x, y \rangle$ is tame.*

In fact, let ϕ be an automorphism of $R\langle x, y \rangle$. Using a theorem of Jung [6] that says that every automorphism of $R(\bar{x}, \bar{y})$ is tame, we can assume that, modulo a tame automorphism of $R\langle x, y \rangle$, the map $\text{Aut}_R(R\langle x, y \rangle) \rightarrow \text{Aut}_R(R(\bar{x}, \bar{y}))$ carries ϕ into the identity. Hence using

Theorem 1 it follows that ϕ must be the identity of $\text{Aut}_R(R\langle x, y \rangle)$.

REFERENCES

1. P. M. Cohn, *Free associative algebras*, Bull. London Math. Soc. **1** (1969), 1–39; correction, *ibid.* **1** (1969), 218. MR **39** #2800; #5614.
2. ———, *On subsemigroups of free semigroups*, Proc. Amer. Math. Soc. **13** (1962), 347–351. MR **25** #3111.
3. ———, *Lecture notes on free rings*, Yale University, New Haven, Conn., 1962.
4. A. Czerniakiewicz, *Automorphisms of a free associative algebra of rank 2* Thesis, New York University, 1970.
5. ———, *Automorphisms of a free associative algebra of rank 2*, Trans. Amer. Math. Soc. **160** (1971), 393–401.
6. H. W. E. Jung, *Über ganze birationale Transformationen der Ebene*, J. Reine Angew. Math. **184** (1942), 161–174. MR **5**, 74.

COLUMBIA UNIVERSITY, NEW YORK, NEW YORK 10027