

Reduction of Constrained Mechanical Systems and Stability of Relative Equilibria

Charles-Michel Marle

Institut de Mathématiques de Jussieu, Université Pierre et Marie Curie, 4, place Jussieu, F-75252 Paris cedex 05, France. email: marle@mathp6.jussieu.fr

Received: 1 September 1994/in revised form: 15 April 1995

Abstract: A mechanical system with perfect constraints can be described, under some mild assumptions, as a constrained Hamiltonian system (M, Ω, H, D, W) : (M, Ω) (the *phase space*) is a symplectic manifold, H (the *Hamiltonian*) a smooth function on M , D (the *constraint submanifold*) a submanifold of M , and W (the *projection bundle*) a vector sub-bundle of $T_D M$, the reduced tangent bundle along D . We prove that when these data satisfy some suitable conditions, the time evolution of the system is governed by a well defined differential equation on D . We define constrained Hamiltonian systems with symmetry, and prove a reduction theorem. Application of that theorem is illustrated on the example of a convex heavy body rolling without slipping on a horizontal plane. Two other simple examples show that constrained mechanical systems with symmetry may have an attractive (or repulsive) set of relative equilibria.

1. Introduction

Let (M, Ω) be a symplectic manifold on which a Lie group G acts by a Hamiltonian action, with an equivariant momentum map J . Let μ be a regular (or weakly regular) value of J . Under some general assumptions, the very important concept of *symplectic reduction*, due to K. Meyer [24], J. Marsden and A. Weinstein [22], allows us to obtain from these data a new symplectic manifold $(P_\mu = J^{-1}(\mu)/G_\mu, \Omega_P)$, called the *reduced symplectic manifold* at μ . Any smooth G -invariant Hamiltonian H on M induces a smooth reduced Hamiltonian H_P on P_μ , and the integral curves of the Hamiltonian vector field X_H associated with H contained in $J^{-1}(\mu)$ project onto the integral curves of the Hamiltonian vector field X_{H_P} associated with H_P . In particular, relative equilibria of X_H contained in $J^{-1}(\mu)$ project onto equilibria of X_{H_P} , i.e., onto points on P_μ where X_{H_P} vanishes. Stability properties of these relative equilibria are closely related to stability properties of the corresponding equilibria in P_μ .

Symplectic reduction plays an important part in symplectic geometry, in analytical mechanics and in mathematical physics. It has been extended to vari-

ous more general situations, in which the symplectic manifold (M, Ω) is replaced by a Poisson manifold (J. Marsden and T. Ratiu [21]), a contact or a cosymplectic manifold (C. Albert [2]), a Jacobi manifold (K. Mikami [25], J. Nunes da Costa [27]). M. Kummer has given a nice description of the reduced symplectic manifold when (M, Ω) is the cotangent bundle of a principal G -bundle [14]. A. Weinstein has used the symplectic reduction procedure [39] in the minimal coupling construction introduced by S. Sternberg [30]. In this paper, in view of applications to mechanical systems with constraints, we introduce still another extension of the reduction concept. We were led to that extension by an attempt to understand the very remarkable stability properties of some nonholonomic mechanical systems such as the celtic stones, discussed a century ago by G.T. Walker [36, 37]. The reader is referred to J. Walker [38] for a more recent and very clear (although nonmathematical) presentation of these properties. A different approach of the reduction procedure for nonholonomic mechanical systems was proposed earlier by L. Bates and J. Sniatycki [4].

The paper is organized as follows.

In Sect. 2, we present an intrinsic formulation of the dynamics of a constrained mechanical system. It is equivalent to the formulation of A.M. Vershik and L.D. Faddeev (see [33, 34, 35] and the references therein), i.e., it leads to the same equations, but it is presented under a somewhat different manner more convenient for our particular application.

Section 3 considers an action of a Lie group on a constrained mechanical system, defines the reduction procedure and proves its main properties (Theorem 3.5).

Section 4 illustrates the reduction procedure with the classical example of a heavy convex body which rolls without slipping on a horizontal plane. If the body has a symmetry axis, and if the principal directions of curvature of the surface of the body at the lowest point on that axis differ from the principal directions of inertia, the stability properties of stationary rotations around that axis, set in a vertical position, may be quite complicated, very different from those of the stationary motions of a heavy body around a fixed point. The geometric setting presented here may help to understand these properties, which have been already investigated analytically by G.T. Walker almost a century ago [37], and more recently for example by A.V. Karapetian [9, 10] and A.P. Markeev [17] in the linear approximation, M. Pascal for the asymptotic behaviour [28].

Finally Sect. 5 deals with two simple examples. The first one shows that a constrained mechanical system, even when the constraint is holonomic, may have two sets of relative equilibria, one of which is attractive and the other repulsive. An unconstrained conservative mechanical system, whose relative equilibria are neither attractive nor repulsive, cannot exhibit such a behaviour. The second example shows that a constrained Hamiltonian system, in the sense of Definition 3.1, may have an attractive equilibrium.

For the concept of Poisson manifold, the reader is referred to A. Lichnerowicz [16] and A. Weinstein [40]. The general definitions, notations and sign conventions about symplectic and Poisson manifolds are those of [15], not very different from those of Abraham and Marsden [1].

2. The Dynamics of Constrained Dynamical Systems: An Intrinsic Formulation

We present here an intrinsic formulation of the dynamics of a constrained mechanical system similar to that of A.M. Vershik and L.D. Faddeev [33, 34, 35]. This formulation is valid for very general constraints, which may be holonomic or non-holonomic, and eventually not linear in the velocities. For the sake of simplicity we present it for time-independent constraints, but its generalization to time dependent constraints, and even to active constraints [18, 19, 7] should be easy. For constraints linear in the velocities, this formulation is equivalent to the classical one, which may be found for example in the treatise of Whittaker [42]. We do not follow here the other approach to nonholonomic constraints (the so called “vakonomic” approach) proposed by V.I. Arnold, V.V. Kozlov and A.I. Neishtadt [3] which leads to equations of motion not equivalent to the classical ones. A very thorough discussion of constrained systems may be found in Chapter 1 of the book by N. Woodhouse [43]. The reader is referred to S. Benenti [5], F. Cardin and G. Zanzotto [6], P. Dazord [7], J. Koiller [11], P.S. Krishnaprasad [12] and R. Yang [13, 44], E. Massa and E. Pagani [23], for other approaches to the dynamics of constrained systems and applications.

2.1. Definition. *A constrained mechanical system is a triple (Q, L, C) , where Q is a smooth manifold (the **configuration space**), $L : TQ \rightarrow \mathbf{R}$ a smooth function (the **Lagrangian**), and C a smooth submanifold of the tangent bundle TQ (the **constraint submanifold** of the system). The tangent bundle TQ and the cotangent bundle T^*Q of the configuration space are called the **space of kinematical states** and the **phase space**, respectively.*

*We shall denote by $p : TQ \rightarrow Q$ and $q : T^*Q \rightarrow Q$ the canonical projections.*

2.2. The Lagrange differential. We recall that the Lagrangian L defines a map $\Delta(L)$, called the *Lagrange differential* of L . This map is defined on the space $J^2(\mathbf{R}, 0; Q)$ of 2-jets at the origin of smooth parametrized curves in Q . It takes its values in the cotangent bundle T^*Q , and it is fibered over Q . This map can be defined in a coordinate-free manner [8, 29, 31, 32], but for simplicity we recall here only its expression in local coordinates. Let (x^1, \dots, x^n) be the local coordinates in a chart of Q , $(x^1, \dots, x^n, v^1, \dots, v^n)$, $(x^1, \dots, x^n, v^1, \dots, v^n, a^1, \dots, a^n)$ and $(x^1, \dots, x^n, p_1, \dots, p_n)$ the local coordinates in the associated charts of TQ , $J^2(\mathbf{R}, 0; Q)$ and T^*Q , respectively. Let $c : I \rightarrow Q$ be a smooth parametrized curve in Q , with I an open interval in \mathbf{R} such that $0 \in I$. We shall write

$$x^i(t) = x^i \circ c(t), \quad v^i(t) = \frac{d}{dt}(x^i \circ c(t)), \quad a^i(t) = \frac{d^2}{dt^2}(x^i \circ c(t)).$$

The local expression of the Lagrangian is a function L of the local coordinates $(x^1, \dots, x^n, v^1, \dots, v^n)$ in TQ . Then the local coordinates $(x^1, \dots, x^n, p_1, \dots, p_n)$ of $\Delta(L)(j_0^2 c)$ are

$$x^i = x^i \circ c(0),$$

$$p_i = \left(\frac{d}{ds} \left(\frac{\partial L}{\partial v^i}(x(s), v(s)) \right) - \frac{\partial L}{\partial x^i}(x(s), v(s)) \right) \Big|_{s=0},$$

where $x(s)$ and $v(s)$ stand for $(x^1(s), \dots, x^n(s))$ and $(v^1(s), \dots, v^n(s))$, respectively. As is well known, the x^i and p_i given by these formulae depend only on the local coordinates $x^i(0), v^i(0)$ and $a^i(0)$, $i \leq i \leq n$, of the 2-jet j_0^2c of c at the origin.

2.3. *The Lagrange Equations of Motion.* In the Lagrangian formalism, the equations of motion of the constrained mechanical system (Q, L, C) can be written as

$$\Delta(L) = f, \tag{1}$$

where $\Delta(L)$ is the Lagrange differential of L discussed in Section 2.2. On the right hand side, f is the constraint force. It is in general unknown, but must satisfy some conditions, which depend on the physical properties of the constraint. These conditions will be indicated in Section 2.6 for a particular class of constraints (the so called *perfect* constraints). A *motion* of the system, i.e., a *solution* of (1), is a smooth curve $c : I \rightarrow C$, where I is an open interval in \mathbf{R} , such that, for each $t \in I$, $\frac{dc(t)}{dt}$ lies in C and the constraint force $f(t)$, defined by

$$\Delta(L)(j^2c(t)) = f(t), \tag{2}$$

satisfies these conditions.

2.4. *The Legendre Transformation.* We recall that, associated with the Lagrangian L , there is a map $\mathcal{L} : TQ \rightarrow T^*Q$, called the *Legendre transformation*. This map is such that, for each $v \in TQ$, $\mathcal{L}(v) \in T_{p(v)}^*Q$ is the differential at v of the restriction of L to the tangent space $T_{p(v)}Q$.

2.5. **Definitions.** Let (Q, L, C) be a constrained mechanical system.

1. The constraint C is said to be **regular** if there exists a submanifold Q_1 of Q such that the constraint submanifold C is contained in TQ_1 (considered as a submanifold of TQ) and that the restriction to C of the canonical projection $p : TQ \rightarrow Q$ is a submersion of C onto Q_1 .
2. The Lagrangian L is said to be **regular** if the Legendre transformation $\mathcal{L} : TQ \rightarrow T^*Q$ associated with L is a diffeomorphism.
3. The constrained system (Q, L, C) is said to be **regular** if the constraint and the Lagrangian are both regular.

2.6. *Examples and Remarks.* Let (Q, L, C) be a constrained mechanical system.

1. The constraint is said to be *purely geometric* if there exists a submanifold Q_1 of Q such that $C = TQ_1$. A purely geometric constraint is automatically regular. Such a constraint can be eliminated by choosing Q_1 as configuration space instead of Q .
2. The constraint is said to be *purely kinematic* if $p(C) = Q$, i.e., if every point of Q is a possible configuration of the system. Such a constraint is regular if the restriction to C of the canonical projection $p : TQ \rightarrow Q$ is a submersion of C onto Q .
3. Very often, the constraint submanifold C is a vector sub-bundle of TQ . Such a constraint is purely kinematic and regular. When in addition the sub-bundle C is completely integrable, we shall say that the constraint is *holonomic*. In such a case C defines a foliation of Q . If the configuration of the system is in a leaf Q_1 of that foliation at a particular time t_0 , it is in Q_1 for all times. The constraint submanifold C may then be replaced by TQ_1 , which is now a purely geometric

constraint. However, this procedure is suitable only to deal with the motions of the system for which the configuration is in the particular leaf Q_1 of the foliation of Q determined by C ; it is not suitable to deal with stability properties, since motions of the system for which the configuration is in leaves of that foliation near Q_1 , but distinct from Q_1 , must be taken into account.

4. When the constraint is regular, by taking $Q_1 = p(C)$ instead of Q as configuration space, we can replace the system by a completely equivalent one in which the constraint is purely kinematic and regular. However, in many examples, such as the one dealt with in Sect. 4, it is more convenient to keep as configuration space a manifold Q strictly larger than the set $Q_1 = p(C)$ of all the possible configurations compatible with the constraint.

2.7. Admissible Infinitesimal Virtual Displacements. We assume now that (Q, L, C) is a regular constrained mechanical system. Let $v \in C$ be any point of the constraint submanifold, and $x = p(v) \in Q$. Since $p|_C$ is a submersion of C onto the submanifold $Q_1 = p(C)$ of Q , $C_x = C \cap T_x Q$ is a submanifold of $T_x Q_1$. Therefore the space $T_v C_x$, tangent at v to a submanifold of a vector space, can be considered as a vector subspace of that vector space $T_x Q_1$, which is itself a vector subspace of $T_x Q$. The vector space $T_v C_x$ is called the space of *admissible infinitesimal virtual displacements* at v . Its annihilator $V_v = (T_v C_x)^0$ is the vector subspace of all $\eta \in T_x^* Q$ such that $\langle \eta, w \rangle = 0$ for all $w \in T_v C_x$. We are led to the following definition.

2.8. Definition. *The constraint is said to be perfect if, for each $v \in C$, $x = p(v) \in Q$, the constraint force f takes its value in the annihilator $V_v = (T_v C_x)^0$ of the space of admissible infinitesimal virtual displacements $T_v C_x$ when the kinematical state of the system is v .*

The above definition is in agreement with the principle (called the d'Alembert-Lagrange principle in Arnold, Kozlov and Neishtadt [3]) which states that when a constraint is perfect, the virtual work of the constraint force for any infinitesimal virtual displacement compatible with the constraint vanishes.

2.9. The Map λ . For any vector bundle (E, π, Q) over the base manifold Q , let $E \times_Q E$ be the fibered product of E with itself over Q , and $\lambda : E \times_Q E \rightarrow TE$ be the map such that, for any $x \in Q$, v and $w \in E_x$, $\lambda(v, w)$ is the vector equal to w , tangent at v to the fiber E_x . Since E_x is a vector space, we have identified it with its tangent space at v , $T_v E_x$. The map λ is an injective vector bundle map ($E \times_Q E$ being considered as a vector bundle over E , with canonical projection $(v, w) \mapsto w$), which maps $E \times_Q E$ onto the vertical bundle $VE = \ker(T\pi)$. In the following definition we use the map λ for the cotangent bundle (T^*Q, q, Q) .

2.10. Definitions. *Let us assume that the constrained mechanical system (Q, L, C) is regular.*

1. *The image $D = \mathcal{L}(C)$ of the constraint submanifold C by the Legendre transformation will be called the **Hamiltonian constraint submanifold**.*

2. *For any $p \in D$, the vector subspace W_p of the tangent space $T_p(T^*Q)$, defined by*

$$W_p = \{ \lambda(p, \eta) \mid \eta \in V_{\mathcal{L}^{-1}(p)} \},$$

will be called the **projection space** at p . The union

$$W = \bigcup_{p \in D} W_p$$

of all the projection spaces at points of D is a vector sub-bundle of $T_D(T^*Q)$, which will be called the **projection bundle**.

We observe that W is a subset of the vertical bundle $\ker(Tq)$.

2.11. Definition. Let $v \in TQ$ be a kinematic state, $x = p(v) \in Q$, and $L_x = L|_{T_xQ}$. The Lagrangian L is said to be **normal** at v if the matrix

$$\left(\frac{\partial^2 L_x}{\partial v^i \partial v^j}(v) \right), \quad 1 \leq i, j \leq n,$$

of second partial derivatives of L_x at v , with respect to the coordinates v^1, \dots, v^n in T_xQ associated with a basis of that vector space, is positive definite.

This definition has indeed an intrinsic meaning, since the positive definiteness of that matrix does not depend on the choice of the basis of T_xQ .

2.12. Proposition. We assume that the constrained mechanical system (Q, L, C) is regular and that the Lagrangian L is normal at every point of the constraint submanifold C . Then the sum of the vector sub-bundles TD and W of $T_D(T^*Q)$ (restriction of the tangent bundle $T(T^*Q)$ to the Hamiltonian constraint submanifold D) is a direct sum, and

$$TD \oplus W = (T_D q)^{-1}(TQ_1),$$

where $Q_1 = p(C)$ is the submanifold of Q introduced in Definition 2.5.

Proof. Let $\dim Q = n$, $\dim Q_1 = m \leq n$. Since the property to be proven is local, we can use a chart of Q adapted to the submanifold Q_1 , with local coordinates $(x) = (x^1, \dots, x^n)$. In that chart Q_1 is locally defined by the equations

$$x_i = 0 \quad \text{for } m + 1 \leq i \leq n.$$

Let $(x, v) = (x^1, \dots, x^n, v^1, \dots, v^n)$ and $(x, p) = (x^1, \dots, x^n, p_1, \dots, p_n)$ be the local coordinates in the associated charts of TQ and T^*Q respectively. The constraint submanifold C is contained in TQ_1 , therefore it can be locally defined by the $2(n - m) + k$ equations

$$x^i = 0 \quad \text{and} \quad v^i = 0 \quad \text{for } m + 1 \leq i \leq n, \quad \Phi^\alpha(x, v) = 0, \quad 1 \leq \alpha \leq k.$$

Since $q|_C$ is a submersion of C onto Q_1 , the functions Φ^α can be chosen in such a way that the matrix

$$\left(\frac{\partial \Phi^\alpha}{\partial v^j} \right), \quad 1 \leq \alpha \leq k, \quad 1 \leq j \leq m, \tag{3}$$

is of rank k at all points of C . The tangent space at (x, v) to the constraint submanifold C is the set of all vectors

$$\sum_{i=1}^m \left(A^i \frac{\partial}{\partial x^i} + B^i \frac{\partial}{\partial v^i} \right)$$

whose components A^i and B^i satisfy

$$\sum_{i=1}^m \left(A^i \frac{\partial \Phi^\alpha}{\partial x^i} + B^i \frac{\partial \Phi^\alpha}{\partial v^i} \right) = 0, \quad 1 \leq \alpha \leq k. \tag{4}$$

Since $D = \mathcal{L}(C)$, and since \mathcal{L} maps $(x, v) \in C$ onto $(x, p) \in D$, with

$$p_i = \frac{\partial L(x, v)}{\partial v^i},$$

the tangent space at (x, p) to the Hamiltonian constraint submanifold D is the set of all vectors

$$\sum_{i=1}^m \left(A^i \left(\frac{\partial}{\partial x^i} + \sum_{j=1}^n \frac{\partial^2 L}{\partial v^j \partial x^i} \frac{\partial}{\partial p_j} \right) + B^i \sum_{j=1}^n \frac{\partial^2 L}{\partial v^j \partial v^i} \frac{\partial}{\partial p_j} \right)$$

with A^i and B^i satisfying (4).

The fiber $W_{(x,p)}$ of the projection bundle at point $(x, p) \in D$ is generated by the $n - m + k$ vectors

$$\frac{\partial}{\partial p_i} \text{ for } m + 1 \leq i \leq n \quad \text{and} \quad \sum_{i=1}^m \frac{\partial \Phi^\alpha}{\partial v^i} \frac{\partial}{\partial p_i} \text{ for } 1 \leq \alpha \leq k.$$

The fiber at $(x, p) \in D$ of $(T_D q)^{-1}(TQ_1)$ is generated by the vectors

$$\frac{\partial}{\partial x^i} \text{ for } 1 \leq i \leq m \quad \text{and} \quad \frac{\partial}{\partial p_j} \text{ for } 1 \leq j \leq n.$$

That fiber is of dimension $n + m$, and it contains $T_{(x,p)}D$ and $W_{(x,p)}$, whose dimensions are $2m - k$ and $n - m + k$, respectively. Therefore we have only to prove that $W_{(x,p)} \cap T_{(x,p)}D = \{0\}$, i.e., that the only λ_α ($1 \leq \alpha \leq k$), μ_j ($m + 1 \leq j \leq n$), A^i and B^i ($1 \leq i \leq m$) which satisfy (4) and

$$\begin{aligned} & \sum_{\alpha=1}^k \left(\lambda_\alpha \sum_{j=1}^n \frac{\partial \Phi^\alpha}{\partial v^j} \frac{\partial}{\partial p_j} \right) + \sum_{j=m+1}^n \mu_j \frac{\partial}{\partial p_j} \\ &= \sum_{i=1}^m A^i \frac{\partial}{\partial x^i} + \sum_{i=1}^m \sum_{j=1}^n \left(A^i \frac{\partial^2 L}{\partial v^j \partial x^i} + B^i \frac{\partial^2 L}{\partial v^j \partial v^i} \right) \frac{\partial}{\partial p_j} \end{aligned} \tag{5}$$

vanish identically. But (5) implies

$$A^i = 0 \quad \text{for } 1 \leq i \leq m,$$

and

$$\sum_{\alpha=1}^k \lambda_\alpha \frac{\partial \Phi^\alpha}{\partial v^j} + \mu_j = \sum_{i=1}^m B^i \frac{\partial^2 L}{\partial v^j \partial v^i} \quad \text{for } 1 \leq j \leq n, \tag{6}$$

where we have set $\mu_j = 0$ for $1 \leq j \leq m$. Since the A^i vanish, (4) implies

$$\sum_{i=1}^m B^i \frac{\partial \Phi^\alpha}{\partial v^i} = 0. \tag{7}$$

Multiplying Eq. (6) by B^j , summing up over j for $1 \leq j \leq m$ and using (7), we obtain

$$\sum_{i=1}^m \sum_{j=1}^m B^i B^j \frac{\partial^2 L}{\partial v^i \partial v^j} = 0. \tag{8}$$

Since L is normal at all points of C , the above equality implies that $B^i = 0$ for all i , $1 \leq i \leq m$. Then, since the matrix (3) is of rank k at all points of C , (6) implies that $\lambda^\alpha = 0$ for all α , $1 \leq \alpha \leq k$, and that $\mu_j = 0$ for all j , $m + 1 \leq j \leq n$. \square

2.13. Remark. The property of L used in the proof is slightly less restrictive than the property stated in Definition 2.11, since the indices i and j in Eq. (8) satisfy $1 \leq i, j \leq m$, instead of $1 \leq i, j \leq n$. Therefore it should be possible to weaken a little Definition 2.11.

*2.14. The Symplectic Structure of T^*Q and the Hamiltonian Vector field X_H .* The assumptions are the same as in Proposition 2.10. We define the Hamiltonian $H : T^*Q \rightarrow \mathbf{R}$ of the system by setting

$$H = (i(Z)dL - L) \circ \mathcal{L}^{-1},$$

where Z is the Liouville vector field on TQ . In local coordinates

$$Z = \sum_{i=1}^n v^i \frac{\partial}{\partial v^i}, \quad H(x, p) = \sum_{i=1}^n v^i p_i - L(x, v),$$

where, in the right-hand side, the local coordinates (x, v) in TQ must be expressed in terms of the local coordinates (x, p) in T^*Q , by means of the diffeomorphism \mathcal{L}^{-1} .

Let α be the Liouville 1-form and $\Omega = d\alpha$ the canonical symplectic 2-form on T^*Q . In local coordinates,

$$\alpha = \sum_{i=1}^n p_i dx^i, \quad \Omega = \sum_{i=1}^n dp_i \wedge dx^i.$$

Let X_H be the Hamiltonian vector field associated with H , i.e., the vector field such that

$$i(X_H)\Omega = -dH.$$

2.15. Proposition. *Under the assumptions of Proposition 2.12, the restriction $X_H|_D$ of the Hamiltonian vector field to the Hamiltonian constraint submanifold D splits into a sum*

$$X_H|_D = X_D + X_W,$$

where X_D is a smooth vector field tangent to D and X_W a smooth section of the projection bundle W . The vector field X_D will be called the constrained Hamiltonian vector field, and the opposite $-X_W$ of X_W , the constraint force field.

Proof. In view of Proposition 2.12, it is enough to prove that $X_H|_D$ is a section of $TD \oplus W = (T_D q)^{-1}(TQ_1)$, or equivalently that, for any $z \in D$, $Tq(X_H(z))$ is tangent to the submanifold Q_1 . As in the proof of Proposition 2.12, we use a chart of Q adapted to the submanifold Q_1 , and the associated charts of TQ and T^*Q . The vector field X_H may be written as

$$X_H = \sum_{i=1}^n \left(\frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i} - \frac{\partial H}{\partial x^i} \frac{\partial}{\partial p_i} \right).$$

Let (x, v) be a point of C . Since $C \subset TQ_1$, we have $v^i = 0$ for $m + 1 \leq i \leq n$. Let $(x, p) \in D$ be the image of (x, v) by the Legendre transformation. We have, for all $i, 1 \leq i \leq n$,

$$\frac{\partial H}{\partial p_i} = v^i,$$

therefore

$$\frac{\partial H}{\partial p_i} = 0 \quad \text{for } m + 1 \leq i \leq n.$$

This shows that the projection of $X_H|_D$ on Q is tangent to Q_1 . \square

2.16. Theorem. *We assume that the constrained mechanical system (Q, L, C) is regular, that the Lagrangian L is normal at every point of the constraint submanifold C , and that the constraint is perfect. A parametrized curve $c : I \rightarrow Q$ (with I an open interval in \mathbf{R}) is a motion of the system if and only if its canonical lift \tilde{c} to the cotangent bundle, defined by*

$$\tilde{c} = \mathcal{L} \circ \frac{dc}{dt},$$

*is an integral curve of the constrained Hamiltonian vector field X_D . When such is the case, the constraint force $f(t)$ at each time $t \in I$ is the unique element in $T_{\tilde{c}(t)}^*Q$ such that*

$$\lambda(\tilde{c}(t), f(t)) = -X_W(\tilde{c}(t)).$$

Proof. In local coordinates (x^i, v^i) in TQ , (x^i, p_i) in T^*Q , related by the Legendre transformation \mathcal{L} , it is well known that the constrained Lagrange equations

$$\frac{d}{dt} \frac{\partial L}{\partial v^i} - \frac{\partial L}{\partial x^i} = f_i, \quad 1 \leq i \leq n,$$

are equivalent to the constrained Hamilton equations

$$\begin{aligned} \frac{dx^i}{dt} &= \frac{\partial H}{\partial p_i}, \\ \frac{dp_i}{dt} &= -\frac{\partial H}{\partial x^i} + f_i. \end{aligned}$$

Let us assume that $t \mapsto c(t)$ is a motion of the system. Its canonical lift $t \mapsto \tilde{c}(t)$ to the cotangent bundle lies in the Hamiltonian constraint submanifold D . Therefore, for each $t \in I$, the vector

$$\sum_{i=1}^n \left(\frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i} + \left(-\frac{\partial H}{\partial x^i} + f_i \right) \frac{\partial}{\partial p_i} \right) (\tilde{c}(t)) = X_H(\tilde{c}(t)) + \left(\sum_{i=1}^n f_i \frac{\partial}{\partial p_i} \right) (\tilde{c}(t))$$

is tangent to D . Since the constraint is perfect, the second term in the right-hand side of the above expression is in $W_{\tilde{c}(t)}$, and we have

$$\begin{aligned} X_D(\tilde{c}(t)) &= \sum_{i=1}^n \left(\frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i} + \left(-\frac{\partial H}{\partial x^i} + f_i \right) \frac{\partial}{\partial p_i} \right) (\tilde{c}(t)), \\ X_W(\tilde{c}(t)) &= - \left(\sum_{i=1}^n f_i \frac{\partial}{\partial p_i} \right) (\tilde{c}(t)). \end{aligned}$$

This proves that $t \mapsto \tilde{c}(t)$ is an integral curve of X_D and that, for each $t \in I$, $\lambda(\tilde{c}(t), f(t)) = -X_W(\tilde{c}(t))$.

Conversely, if $t \mapsto \tilde{c}(t)$ is an integral curve of X_D , the constrained Hamilton equations above are satisfied, with

$$\sum_{i=1}^n f_i \frac{\partial}{\partial p_i} = -X_W .$$

Since X_W is a section of W , the condition which states that the constraint is perfect is satisfied and the projection on Q of the curve $t \mapsto \tilde{c}(t)$ is a motion of the system. \square

2.17. Remarks.

1. Contrary to what happens for unconstrained, time independent systems, the Hamiltonian H may not be a first integral of X_D , i.e., may not remain constant during a motion of the system. This property is due to the fact that even when the constraint is time independent, it may give energy to (or withdraw energy from) the system. However, when the constraint submanifold C is tangent to the Liouville vector field of TQ (in particular, when C is a vector sub-bundle of TQ), one can prove that H is a first integral of X_D .

2. In [7], P. Dazord introduces a formalism for constrained mechanical systems different from that used here. In his formalism, the geometric properties of the constraint are described by a submersion f of the configuration manifold Q onto another manifold S , and a map $Y : TQ \rightarrow TS$, fibered over f , eventually nonlinear on the fibers. A parametrized curve $t \mapsto x(t)$ in Q is said to be *compatible with the constraint* when it satisfies

$$\frac{d}{dt} f(x(t)) = Y \left(\frac{dx(t)}{dt} \right) .$$

In other words, the subset C of TQ of kinematical states compatible with the constraint is

$$C = \{v \in TQ \mid (Tf - Y)(v) = 0\} .$$

Dazord does not assume that C is a submanifold of TQ . He says that the constraint is *holonomic* when $Y = 0$, *almost holonomic* when Y factors through $f \circ p : TQ \rightarrow S$, *semi-holonomic* when Y factors through $p : TQ \rightarrow Q$. In order to describe the mechanical properties of the constraint, he introduces a vector sub-bundle F of $p^*(TQ)$ (the vector bundle over TQ inverse image of the tangent bundle TQ by the map $p : TQ \rightarrow Q$). The sub-bundle F yields the space of infinitesimal virtual displacements for which the work of the constraint force vanishes. Dazord says that the constraint is *perfect* when

$$F = \{(v, w) \in p^*(TQ) \equiv TQ \oplus TQ \mid w \in \ker(Tf)\} .$$

Observe that Y does not play any part in the definition of perfect constraints. For semi-holonomic constraints (in Dazord's language), and *a fortiori* for almost holonomic or holonomic constraints, one can easily see that Dazord's definition of perfect constraints is equivalent to our Definition 2.8, and that Dazord's formalism and ours lead to the same equations. But for more general constraints, the two formalisms are not equivalent, even in the definition of perfect constraints. Of course, the formalism developed here deals only with *passive* constraints, while Dazord's

formalism (which extends some ideas introduced in [18 and 19] for geometric constraints) can include some *active* constraints.

3. Lie Group Actions on Constrained Systems, Reduction and Relative Equilibria

In this section we consider the action of a Lie group on a constrained mechanical system. We define *relative equilibria* (also called *stationary motions*) of the system. We introduce a reduction procedure which will allow us to obtain a reduced constrained system, and we use that reduced system for discussing the stability of relative equilibria. The reader is referred to J.E. Marsden [20] for a thorough discussion of various kinds of reduction and many applications, and to R. Montgomery, J.E. Marsden and T. Ratiu [26] for the determination of the Poisson structure on the quotient of the cotangent bundle to a principal G -bundle.

For the needs of the reduction procedure to be defined, we must first introduce a slight generalization of the concept of constrained Hamiltonian system, in which the phase space will be a general Poisson manifold, instead of a cotangent bundle equipped with its canonical symplectic structure and the associated Poisson structure.

3.1. Definition. *A constrained Hamiltonian system is a 5-tuple (M, Λ, H, D, W) , in which*

- (M, Λ) is a Poisson manifold (the **phase space**),
- $H : M \rightarrow \mathbf{R}$ is a smooth function (the **Hamiltonian**),
- D is a submanifold of M (the **Hamiltonian constraint submanifold**),
- W is a vector sub-bundle of $T_D M$ (the **projection bundle**) such that $TD \cap W = \{0\}$ and that the Hamiltonian vector field $X_H = \Lambda^\sharp(dH)$, restricted to the submanifold D , is a section of $TD \oplus W$.

We set

$$X_H|_D = X_D + X_W,$$

where X_D is a smooth vector field on D (called the *constrained Hamiltonian vector field*), and X_W a smooth section of the projection bundle W (its opposite $-X_W$ is called the *constraint force field*). Integral curves of X_D are called *motions* of the system.

We observe that the property which states that $TD \cap W = \{0\}$ and that $X_H|_D$ is a section of $TD \oplus W$, in Definition 3.1, is equivalent to the *pairing condition* of Dazord [7].

3.2. Definition. *A symmetry of the constrained Hamiltonian system (M, Λ, H, D, W) is a left action $\Phi : G \times M \rightarrow M$ of a Lie group G on the phase space M , which leaves invariant the Poisson tensor Λ , the Hamiltonian H , the Hamiltonian constraint submanifold D , and the projection bundle W .*

3.3. Remarks. The symmetry Φ is, in many examples, a Hamiltonian action, i.e., an action such that the infinitesimal generator of the restriction of Φ to every one-parameter subgroup of G is a Hamiltonian vector field. In such a case, there exists a momentum map $J : M \rightarrow \mathcal{G}^*$, where \mathcal{G}^* is the dual of the Lie algebra \mathcal{G} of G , such that, for any $X \in \mathcal{G}$, the infinitesimal generator of the restriction of Φ to $\{\exp(tX) | t \in \mathbf{R}\}$ is the Hamiltonian vector field $\Lambda^\sharp(d\langle J, X \rangle)$. For classical mechanical systems with constraints discussed in Sect. 2, the phase space M is a cotangent

bundle T^*Q , its Poisson structure is the structure associated with its canonical symplectic structure $\Omega = d\alpha$, and the action Φ is, in most cases, the canonical lift to T^*Q of an action of a Lie group G on the configuration space Q . Such an action leaves invariant the Liouville 1-form α of T^*Q , and therefore is Hamiltonian, with an Ad^* -equivariant momentum map. However, for constrained Hamiltonian systems, the momentum map does not play a part as important as for unconstrained systems, because in general it does not remain constant during a motion of the system: Noether's theorem does not apply. Moreover, a Hamiltonian action of a connected Lie group on a Poisson manifold leaves invariant every symplectic leaf of that manifold. The symmetries of constrained Hamiltonian systems obtained by reduction very often do not leave invariant the symplectic leaves of the phase space, but instead they map every symplectic leaf onto another symplectic leaf. For these reasons, in Definition 3.2, the symmetry Φ is a Poisson action which may not be Hamiltonian.

3.4. Definition. Let (M, Λ, H, D, W) be a constrained Hamiltonian system with a symmetry $\Phi : G \times M \rightarrow M$. A **relative equilibrium** (also called **stationary motion**) of the system is a motion $c : \mathbf{R} \rightarrow D$ of the system, i.e., an integral curve of the constrained Hamiltonian vector field X_D , which is also an orbit, under the action Φ , of a one-parameter subgroup of G .

3.5. Theorem. Let (M, Λ, H, D, W) be a constrained Hamiltonian system with a symmetry $\Phi : G \times M \rightarrow M$. We assume that the set \widehat{M} of orbits of the action Φ has a smooth manifold structure such that the canonical projection $\pi : M \rightarrow \widehat{M}$ is a submersion. Then:

1. $\widehat{D} = \pi(D)$ is a submanifold of \widehat{M} , and the image by π of the bundle W is a vector sub-bundle \widehat{W} of $T_{\widehat{D}}\widehat{M}$, such that $T\widehat{D} \cap \widehat{W} = \{0\}$. Moreover, there exists on \widehat{M} a unique Poisson structure $\widehat{\Lambda}$ such that π is a Poisson map, and a unique smooth function $\widehat{H} : \widehat{M} \rightarrow \mathbf{R}$, such that $H = \widehat{H} \circ \pi$.
2. $(\widehat{M}, \widehat{\Lambda}, \widehat{H}, \widehat{D}, \widehat{W})$ is a constrained Hamiltonian system, which will be called the reduced constrained system associated with (M, Λ, H, D, W) and the symmetry Φ .
3. For each motion c of the constrained Hamiltonian system (M, Λ, H, D, W) , $\widehat{c} = \pi \circ c$ is a motion of the reduced system $(\widehat{M}, \widehat{\Lambda}, \widehat{H}, \widehat{D}, \widehat{W})$.
4. A motion c of (M, Λ, H, D, W) is a relative equilibrium if and only if $\widehat{c} = \pi \circ c$ is constant, i.e., is an equilibrium point of the reduced system.

Proof.

1. For any point $\widehat{a} \in \widehat{D}$, there exists a smooth section $s : \widehat{U} \rightarrow M$ of the submersion π , defined on an open neighbourhood \widehat{U} of \widehat{a} in \widehat{M} . Then s is a diffeomorphism of \widehat{U} onto the submanifold $U = s(\widehat{U})$ of M . Moreover, U intersects D transversally, because for each point $x \in U \cap D$, $T_x D$ contains $\ker(T_x \pi)$ which is complementary to $T_x U$ in $T_x M$. Therefore $U \cap D$ is a submanifold of U . Since D is Φ -invariant, s maps $\widehat{U} \cap \widehat{D}$ onto $U \cap D$, which proves that $\widehat{U} \cap \widehat{D}$ is a submanifold of \widehat{U} . We conclude that \widehat{D} is a submanifold of \widehat{M} .

Let \widehat{x} be a point of \widehat{D} , and x be a point such that $\pi(x) = \widehat{x}$. By using the invariance of W , we see that $T_x \pi(W_x)$ does not depend on the choice of $x \in \pi^{-1}(\widehat{x})$. Therefore we can set $\widehat{W}_{\widehat{x}} = T_x \pi(W_x)$. We have $T_x D \cap W_x = \{0\}$ and $\ker(T_x \pi) \subset T_x D$; therefore the restriction of $T_x \pi$ to W_x is an isomorphism of W_x onto $\widehat{W}_{\widehat{x}}$, which

proves that $\widehat{W} = \bigcup_{\widehat{x} \in \widehat{D}} \widehat{W}_{\widehat{x}}$ is vector sub-bundle of $T_{\widehat{D}}\widehat{M}$. Moreover, $T_x\pi(T_xD) = T_{\widehat{x}}\widehat{D} \cap \widehat{W}_{\widehat{x}} = \{0\}$.

By assumption, Φ leaves A invariant. Therefore the Poisson bracket of two Φ -invariant smooth functions on M is a Φ -invariant function. This implies [15, Chapter III, Proposition 9.4] that there exists a unique Poisson structure $\widehat{\Lambda}$ on \widehat{W} such that π is a Poisson map.

Smooth functions on \widehat{M} are in one-to-one correspondence with Φ -invariant smooth functions on M . Since H is Φ -invariant, there exists a unique smooth function \widehat{H} on \widehat{M} such that $H = \widehat{H} \circ \pi$.

2. Since π is a Poisson map, the Hamiltonian vector field X_H on the Poisson manifold (M, A) projects onto the hamiltonian vector field $X_{\widehat{H}}$ on the Poisson manifold $(\widehat{M}, \widehat{\Lambda})$. Therefore $X_{\widehat{H}}|_{\widehat{D}}$ is a section of $T\widehat{D} \oplus \widehat{W}$, and $(\widehat{M}, \widehat{\Lambda}, \widehat{H}, \widehat{D}, \widehat{W})$ is a constrained Hamiltonian system.

3. The two terms X_D and X_W of the restriction of X_H to D project onto $X_{\widehat{D}}$ and $X_{\widehat{W}}$, respectively. Therefore, for each motion c of (M, A, H, D, W) , $\widehat{c} = \pi \circ c$ is a motion of the reduced system $(\widehat{M}, \widehat{\Lambda}, \widehat{H}, \widehat{D}, \widehat{W})$.

4. Let c be a relative equilibrium of (M, A, H, D, W) . Then by the very definition of a relative equilibrium, $\pi \circ c$ is constant, i.e. is an equilibrium point of the reduced system. Conversely, let $c : I \rightarrow M$ be a motion of (M, A, H, D, W) such that $\pi \circ c$ is constant. Let $t_0 \in I$, and $x_0 = x(t_0)$. For any $t \in I$, $x(t)$ is an element of the orbit $\Phi(G, x_0)$, which may be identified with an homogeneous space of G . Therefore there exists an element $X \in \mathcal{G}$ such that

$$X_D(x_0) = \left. \frac{d}{ds} \Phi(\exp(sX), x_0) \right|_{s=0} .$$

Using the Φ -invariance of the constrained Hamiltonian vector field X_D , we see that the parametrized curve $t \mapsto \Phi(\exp(t - t_0)X, x_0)$ is an integral curve of X_D which takes the value x_0 for $t = t_0$. By unicity of maximal integral curves, that parametrized curve is equal to c . \square

4. The Rolling Stone

4.1. Configuration, Kinematical States, Phase Space and Hamiltonian. Let us consider a solid heavy body, bounded by a smooth, strictly convex surface, supported by a fixed solid horizontal plane. We assume that during its motion, the body remains in contact with the plane, that the friction at the contact point prevents the body from slipping, and that the corresponding constraint is perfect.

We first define the configuration space of the system without constraint. Let E be the physical space; it is an affine Euclidean three-dimensional space. Let G_1 be the group of Euclidean displacements of E , i.e., the group of affine Euclidean orientation-preserving mappings of E onto itself. As a reference configuration, we choose a particular position P_0 of the solid body in E . Then for any other position P of the body in E , there is a unique $g \in G_1$ such that $gP_0 = P$; conversely, for any $g \in G_1$, gP_0 is a position of the body in E .

Therefore, the group G_1 can be considered as the configuration space of the system. The space of kinematical states is the tangent bundle TG_1 , and the phase space the cotangent bundle T^*G_1 .

In order to be able to consider E as a vector space, let us choose as origin an arbitrary point $O \in E$. Then the group G_1 can be identified with the semi-direct product $E \times \mathbf{SO}(E)$, the first factor being the subgroup of translations of E , and the second the stabilizer of O in G_1 , i.e., the subgroup of rotations of E around its origin O . An element of G_1 is a pair (a, g) , with $a \in E, g \in \mathbf{SO}(E)$, and the mapping of E onto itself which corresponds to that element is

$$x \mapsto a + gx .$$

The composition law of G_1 is

$$(a_1, g_1)(a_2, g_2) = (a_1 + g_1 a_2, g_1 g_2) .$$

The Lie algebra \mathcal{G}_1 of G_1 is the semi-direct product $E \times \mathfrak{so}(E)$. The first factor E is the Lie algebra of the subgroup of translations of E , and the second $\mathfrak{so}(E)$ the Lie algebra of the subgroup of rotations of E around its origin. The bracket in $\mathcal{G}_1 = E \times \mathfrak{so}(E)$ is given by

$$[(a_1, X_1), (a_2, X_2)] = (X_1 a_2 - X_2 a_1, [X_1, X_2]) ,$$

where, in the right-hand side, X_1 and $X_2 \in \mathfrak{so}(E)$ are considered as linear endomorphisms of E .

The kinetic energy of the system is

$$T(w) = \frac{1}{2}K(w, w), \quad w \in TG_1 ,$$

where K is a left-invariant Riemannian metric on G_1 . The potential energy is

$$U(a, g) = m\gamma(a + gC|e_3), \quad (a, g) \in G_1 ,$$

where $(|)$ is the scalar product in E, C the position in E of the center of mass of the body when it is in the reference position P_0 , e_3 the unit vertical vector pointing upwards, m the total mass of the body and γ the gravity acceleration. The Lagrangian of the system is therefore

$$L(w) = \frac{1}{2}K(w, w) - U(p(w)) ,$$

where $p : TG_1 \rightarrow G_1$ is the canonical projection.

By left translation, the tangent bundle TG_1 can be identified with $G_1 \times \mathcal{G}_1$. Since G_1 and \mathcal{G}_1 are themselves identified with $E \times \mathbf{SO}(E)$ and $E \times \mathfrak{so}(E)$, respectively, the space of kinematical states is $E \times \mathbf{SO}(E) \times E \times \mathfrak{so}(E)$. A kinematical state will therefore be denoted by (a, g, v, X) , with $a \in E, g \in \mathbf{SO}(E), v \in E, X \in \mathfrak{so}(E)$. We observe that v and X can be easily identified, respectively, with the velocity of the point of the body which is situated at the origin when the body is in its reference position, and with the angular velocity of the motion of the body around that point, both expressed in a frame attached to the body.

Let us now choose the reference position P_0 of the solid body in such a way that when it is in that position, its center of mass is at the origin O of E . Then by König's theorem, the kinetic energy splits into a sum

$$T(a, g, v, X) = \frac{1}{2}m(v|v) + \frac{1}{2}I(X, X) .$$

The first term of that sum, $(1/2)m(v|v)$, is the kinetic energy of the translation motion of the center of mass of the body. The second, $(1/2)I(X, X)$, is the kinetic energy of the rotation of the body around its center of mass. The bilinear form I on the Lie algebra $\mathfrak{so}(E)$ is symmetric positive definite; it is called the *inertia form* of the body at its center of mass.

The phase space T^*G_1 can be identified with the product $E \times \mathbf{SO}(E) \times E^* \times (\mathfrak{so}(E))^*$. An element of that space will be denoted by (a, g, Π, M) , where $a \in E$, $g \in \mathbf{SO}(E)$, $\Pi \in E^*$, $M \in (\mathfrak{so}(E))^*$. It can be seen easily that Π is the linear momentum of the motion of the center of mass of the body, and M the angular momentum of the motion of the body around its center of mass, both expressed in a frame attached to the body.

The Legendre transformation $\mathcal{L} : TG_1 \rightarrow T^*G_1$ is the map

$$(a, g, v, X) \mapsto (a, g, \Pi, M), \quad \text{with } \Pi = \mu(v), \quad M = I(X),$$

where $\mu : E \rightarrow E^*$ and $I : \mathfrak{so}(E) \rightarrow (\mathfrak{so}(E))^*$ are the isomorphisms defined by

$$\langle \mu(v), w \rangle = m(v|w), \quad v \text{ and } w \in E, \quad \langle I(X), Y \rangle = I(X, Y), \quad X \text{ and } Y \in \mathfrak{so}(E).$$

The Hamiltonian H of the system can be written as

$$H(a, g, \Pi, M) = \frac{1}{2} \langle \Pi, \mu^{-1}(\Pi) \rangle + \frac{1}{2} \langle M, I^{-1}(M) \rangle + \gamma \langle \mu(a), e_3 \rangle.$$

4.2. The Constraint Submanifold and the Projection Bundle . When the solid body is in its reference position P_0 , let Σ be the surface which bounds that body, considered as a 2-dimensional submanifold of E . We recall that the Gauss map, defined on Σ , with values in the unit sphere S^2 of E , associates with each point of Σ the unit vector normal to Σ at that point, oriented outwards. Since Σ is smooth and strictly convex, the Gauss map is a diffeomorphism. We will denote by $\Gamma : S^2 \rightarrow \Sigma$ the inverse of the Gauss map.

We assume now that the origin O has been chosen in the horizontal solid plane F on which the moving body is rolling. The constraint submanifold C is the set of kinematical states $(a, g, v, X) \in E \times \mathbf{SO}(E) \times \mathfrak{so}(E)$ which satisfy the following two conditions:

$$(a|e_3) = (\Gamma \circ g^{-1}(-e_3)|g^{-1}(-e_3)), \tag{9}$$

$$v + X \circ \Gamma \circ g^{-1}(-e_3) = 0. \tag{10}$$

Equation (9) states that the lowest point of the body is in the horizontal plane F . Equation (10) states that the velocity of a material point of the moving body vanishes when that point comes into contact with the horizontal plane F on which the body is rolling.

It is easy to verify that the constraint is regular in the sense of Definition 2.5. The submanifold denoted by Q_1 in that definition is the subset of elements $(a, g) \in E \times \mathbf{SO}(E)$ which satisfy (9). We observe that Q_1 and the constraint submanifold C can be identified with $F \times \mathbf{SO}(E)$ and $F \times \mathbf{SO}(E) \times \mathfrak{so}(E)$, respectively, since (9) and (10) determine $(a|e_3)$ in terms of g , and v in terms of g and X . The variable which runs over the factor F is the orthogonal projection on F of the center of mass of the moving body.

The Hamiltonian constraint submanifold D is the subset of elements $(a, g, \Pi, M) \in E \times \mathbf{SO}(E) \times E^* \times (\mathfrak{so}(E))^*$ which satisfy (9) and

$$\Pi + \mu \circ (I^{-1}M) \circ \Gamma \circ g^{-1}(-e_3) = 0. \quad (11)$$

It can be identified with $F \times \mathbf{SO}(E) \times (\mathfrak{so}(E))^*$.

For each $g \in \mathbf{SO}(E)$, we will denote by $\varphi_g : E^* \rightarrow (\mathfrak{so}(E))^*$ the linear map defined by

$$\langle \varphi_g \Pi', Y \rangle = \langle \Pi', Y \circ g^{-1}(-e_3) \rangle, \quad \Pi' \in E^*, \quad Y \in \mathfrak{so}(E).$$

Then it is easy to see that the fiber of the projection bundle over a point (a, g, Π, M) of D is the set of vectors tangent at that point to $E \times \mathbf{SO}(E) \times E^* \times (\mathfrak{so}(E))^*$, whose projections on the first two factors E and $\mathbf{SO}(E)$ vanish, and whose projections Π' on E^* and M' on $(\mathfrak{so}(E))^*$ satisfy

$$M' = \varphi_g \Pi'.$$

4.3. The Evolution Equations for the Constrained System. By using well known formulas for the Liouville 1-form on T^*G_1 and its differential (which may be found for example in [15]), we obtain the evolution equations of the constrained mechanical system:

$$\begin{cases} \frac{da}{dt} = g\mu^{-1}\Pi, \\ \frac{dg}{dt} = gI^{-1}M, \\ \frac{d\Pi}{dt} = {}^t(I^{-1}M)\Pi - \gamma\mu g^{-1}(e_3) + \Pi', \\ \frac{dM}{dt} = -\text{ad}_{I^{-1}M}^* M + \varphi_g \Pi'. \end{cases}$$

In the third equation above, ${}^t(I^{-1}M)$ is the linear endomorphism of E^* transpose of $I^{-1}M \in \mathfrak{so}(E)$, considered as a linear endomorphism of E . We recall our sign convention about the coadjoint action, used in the fourth equation above:

$$\langle \text{ad}_X^* M, Y \rangle = \langle M, -\text{ad}_X Y \rangle = \langle M, [Y, X] \rangle, \quad M \in (\mathfrak{so}(E))^*, \quad X \text{ and } Y \in \mathfrak{so}(E).$$

The element $\Pi' \in (\mathfrak{so}(E))^*$ must be chosen in such a way that for every $t \in \mathbf{R}$, $(a(t), g(t), \Pi(t), M(t))$ remains in the Hamiltonian constraint submanifold D defined by Eqs. (9) and (11). We shall say (maybe a little improperly) that Π' is the *constraint force*, since it determines that force.

4.4. The Reduced Constrained Mechanical System. The symmetry group G of the system is the subgroup of $G_1 = E \times \mathbf{SO}(E)$ generated by translations parallel to the horizontal plane F and by rotations around the vertical axis through the origin O . Elements of G are pairs $(b, h) \in E \times \mathbf{SO}(E)$ such that

$$(b|e) = 0, \quad h e_3 = e_3.$$

The symmetry group G acts on G_1 by multiplication on the left,

$$((b, h), (a, g)) \mapsto (b + ha, hg).$$

It acts on the phase space T^*G_1 by the canonical lift of that action to the cotangent bundle. If we identify T^*G_1 with $E \times \mathbf{SO}(E) \times E^* \times (\mathfrak{so}(E))^*$ by left translations,

that canonical lift is given by

$$((b, h), (a, g, \Pi, M)) \mapsto (b + ha, hg, \Pi, M).$$

The set \widehat{M} of orbits of that action can be identified with $\mathbf{R} \times S^2 \times E^* \times (\mathfrak{so}(E))^*$, and the canonical projection of $M = E \times \mathbf{SO}(E) \times E^* \times (\mathfrak{so}(E))^*$ onto \widehat{M} is

$$(a, g, \Pi, M) \mapsto (z = (a|e_3), n = g^{-1}(-e_3), \Pi, M).$$

The Poisson structure on \widehat{M} for which the canonical projection is a Poisson map is easily derived from the formulas for the symplectic 2-form on T^*G_1 . The Poisson bracket of two smooth functions F_1 and F_2 at a point (z, n, Π, M) of \widehat{M} is

$$\begin{aligned} \{F_1, F_2\} &= (d_{\Pi}F_1|e_3)\frac{\partial F_2}{\partial z} - (d_{\Pi}F_2|e_3)\frac{\partial F_1}{\partial z} \\ &\quad + \langle d_nF_1, (d_MF_2)n \rangle - \langle d_nF_2, (d_MF_1)n \rangle \\ &\quad + \langle \Pi, (d_MF_1)(d_{\Pi}F_2) - (d_MF_2)(d_{\Pi}F_1) \rangle \\ &\quad + \langle M, [d_MF_1, d_MF_2] \rangle. \end{aligned}$$

In this formula, $d_nF_i \in T_n^*S^2$, $d_{\Pi}F_i \in E$, $d_MF_i \in \mathfrak{so}(E)$ are the partial differentials of F_i ($i = 1$ or 2), at point (z, n, Π, M) , with respect to the variables $n \in S^2$, $\Pi \in E^*$, $M \in (\mathfrak{so}(E))^*$, respectively.

The reduced Hamiltonian is

$$\widehat{H}(z, n, \Pi, M) = \frac{1}{2}\langle \Pi, \mu^{-1}\Pi \rangle + \frac{1}{2}\langle M, I^{-1}M \rangle + m\gamma z.$$

The reduced Hamiltonian constraint submanifold \widehat{D} is the subset of elements (z, n, Π, M) in \widehat{M} which satisfy

$$z = (\Gamma(n)|n), \quad (12)$$

$$\Pi + \mu \circ (I^{-1}M) \circ \Gamma(n) = 0. \quad (13)$$

For each $b \in E$, we will denote by $\psi_b : E^* \rightarrow (\mathfrak{so}(E))^*$ the linear map defined by

$$\langle \psi_b \Pi', Y \rangle = \langle \Pi', Yb \rangle, \quad \Pi' \in E^*, \quad Y \in \mathfrak{so}(E). \quad (14)$$

Then the fiber of the reduced projection bundle \widehat{W} over a point (z, n, Π, M) of \widehat{D} , is the set of vectors tangent to \widehat{M} at that point whose projections on the first two factors \mathbf{R} and S^2 vanish, and whose projections Π' on the third factor E^* , and M' on the fourth factor $(\mathfrak{so}(E))^*$, satisfy

$$M' = \psi_{\Gamma(n)}\Pi'.$$

The evolution equations for the reduced constrained Hamiltonian system are

$$\begin{cases} \frac{dz}{dt} = -(\mu^{-1}\Pi|n), \\ \frac{dn}{dt} = -(I^{-1}M)n, \\ \frac{d\Pi}{dt} = (I^{-1}M)\Pi + \gamma\mu n + \Pi', \\ \frac{dM}{dt} = -\text{ad}_{I^{-1}M}^*M + \psi_{\Gamma(n)}\Pi'. \end{cases} \quad (15)$$

The constraint force Π' must be chosen in such a way that for all $t \in \mathbf{R}$, $(z(t), n(t), \Pi(t), M(t))$ lies in the reduced constraint submanifold \widehat{D} , defined by Eqs. (11) and (12).

We observe that the first Eq. (15), which expresses dz/dt , is in fact a consequence of the other three Eqs. (15) and of Eqs. (12) and (13), which define \widehat{D} . Therefore the really useful equations which govern the time evolution of the reduced system are (12), (13), and the last three Eqs. (15).

We observe also that the reduced Hamiltonian \widehat{H} is a first integral of the evolution equations.

4.5. Elimination of the Constraint Force. Let us introduce

$$P = M - \psi_{\Gamma(n)}\Pi . \quad (16)$$

This new variable, defined on the reduced Hamiltonian constraint submanifold, has a clear mechanical meaning: it is the angular momentum of the solid body at its contact point with the horizontal plane on which it is rolling, expressed in a frame attached to the body.

By using Eqs. (13) and (15), we obtain

$$\frac{dP}{dt} = -\text{ad}_{I^{-1}M}^*P - \gamma\psi_{\Gamma(n)} \circ \mu(n) - \psi_{T_n\Gamma((I^{-1}M)_n)} \circ \mu \circ (I^{-1}M) \circ \Gamma(n) . \quad (17)$$

We have denoted by $T_n\Gamma : T_nS^2 \rightarrow T_{\Gamma(n)}E \equiv E$ the tangent map to the inverse Γ of the Gauss map at $n \in S^2$. We have used the identity, which follows easily from the definition (14) of ψ ,

$$\psi_{\Gamma(n)}({}^t(I^{-1}M)\Pi) = -\text{ad}_{I^{-1}M}^*\psi_{\Gamma(n)}\Pi - \psi_{(I^{-1}M) \circ \Gamma(n)}\Pi ,$$

and the fact, which follows from Eq. (13) and the skew-symmetry of elements of $\mathfrak{so}(E)$, that

$$\psi_{(I^{-1}M) \circ \Gamma(n)}\Pi = \psi_{(I^{-1}M) \circ \Gamma(n)}(-\mu \circ (I^{-1}M) \circ \Gamma(n)) = 0 .$$

Now we use the fact that the Euclidean vector space E is of dimension 3. We know that once a particular orientation of E is chosen, we can define on that space a composition law, called the vector product and denoted by $(x, y) \mapsto x \times y$. We recall that for any positively oriented orthonormal basis (e_1, e_2, e_3) of E , we have

$$e_1 \times e_2 = e_3, \quad e_2 \times e_3 = e_1, \quad e_3 \times e_1 = e_2 ,$$

and that E , when equipped with that composition law, is a Lie algebra. We recall the well known formulae which relate the vector and scalar products on E :

$$\begin{aligned} (x \times y|z) &= (x|y \times z), & x, y \text{ and } z \in E . \\ x \times (y \times z) &= (x|z)y - (x|y)z , \end{aligned}$$

We know also that the map $\zeta : \mathfrak{so}(E) \rightarrow E$, defined by

$$(\zeta^{-1}x)y = x \times y, \quad x \text{ and } y \in E ,$$

is a Lie algebra homomorphism.

Using the above formulae and Eq. (14), we obtain for any $b \in E$ a new formula for the linear map $\psi_b : E^* \rightarrow (\mathfrak{so}(E))^*$:

$$\psi_b = \text{ad}_{\zeta^{-1}b}^* \circ {}^t\zeta.$$

Then, for any $Y \in \mathfrak{so}(E)$, we deduce from Eqs. (13) and (16):

$$\begin{aligned} \langle P, Y \rangle &= \langle M, Y \rangle + \langle \text{ad}_{\zeta^{-1}\Gamma(n)}^* \circ {}^t\zeta \circ \mu \circ (I^{-1}M) \circ \Gamma(n), Y \rangle \\ &= \langle M, Y \rangle - m \langle \zeta(I^{-1}M) \times \Gamma(n) | \Gamma(n) \times \zeta(Y) \rangle \\ &= \langle M, Y \rangle + m \langle \Gamma(n) \times (\zeta(I^{-1}M) \times \Gamma(n)) | \zeta(Y) \rangle \\ &= \langle M, Y \rangle + (\Gamma(n) | \Gamma(n)) \langle {}^t\zeta \circ \mu \circ \zeta(I^{-1}M), Y \rangle \\ &\quad - (\Gamma(n) | \zeta(I^{-1}M)) \langle {}^t\zeta \circ \mu \circ \Gamma(n), Y \rangle. \end{aligned}$$

Therefore we have

$$P = J(n)(I^{-1}M) - (\Gamma(n) | \zeta(I^{-1}M)) {}^t\zeta \circ \mu \circ \Gamma(n),$$

where $J(n) : \mathfrak{so}(E) \rightarrow (\mathfrak{so}(E))^*$ is the linear map

$$J(n) = I + (\Gamma(n) | \Gamma(n)) {}^t\zeta \circ \mu \circ \zeta. \tag{18}$$

For any Y and $Z \in \mathfrak{so}(E)$, we have

$$\langle J(n)Y, Z \rangle = \langle IY, Z \rangle + m(\Gamma(n) | \Gamma(n)) \langle \zeta Y | \zeta Z \rangle,$$

which proves that $J(n)$ is symmetric positive definite (with respect to the scalar product on $\mathfrak{so}(E)$ for which ζ is an isometry). Therefore, $J(n)$ is invertible, and we have

$$J(n)^{-1}(P) = I^{-1}M - (\Gamma(n) | \zeta(I^{-1}M)) J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n). \tag{19}$$

By applying ζ and taking the scalar product with $\Gamma(n)$, we obtain

$$(\Gamma(n) | \zeta \circ J(n)^{-1}(P)) = (\Gamma(n) | \zeta(I^{-1}M)) (1 - (\Gamma(n) | \zeta \circ J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n))).$$

Using the isomorphism ζ and the Euclidean structure of E , we can identify E , $\mathfrak{so}(E)$ and their duals. Let x_i be the components of $\Gamma(n)$ in an orthonormal basis of E in which I is diagonal, with diagonal components $I_i, i = 1, 2, 3$. In that basis, $J(n)$ is also diagonal, with diagonal components

$$J_i = I_i + m(x_1^2 + x_2^2 + x_3^2), \quad i = 1, 2, 3.$$

The components of $\zeta \circ J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n)$ are:

$$\frac{mx_i}{J_i} = \frac{mx_i}{I_i + m(x_1^2 + x_2^2 + x_3^2)},$$

and we have

$$\begin{aligned} 1 - (\Gamma(n) | \zeta \circ J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n)) &= 1 - m \sum_{i=1}^3 \frac{x_i^2}{I_i + m(x_1^2 + x_2^2 + x_3^2)} \\ &\geq 1 - \frac{m(x_1^2 + x_2^2 + x_3^2)}{\inf(I_1, I_2, I_3) + m(x_1^2 + x_2^2 + x_3^2)} > 0, \end{aligned}$$

since $\inf(I_1, I_2, I_3) > 0$. We may therefore write

$$(\Gamma(n)|_{\zeta}(I^{-1}M)) = \frac{(\Gamma(n)|_{\zeta} \circ J(n)^{-1}(P))}{1 - (\Gamma(n)|_{\zeta} \circ J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n))},$$

and, using (19),

$$I^{-1}M = J(n)^{-1}(P) - \frac{(\Gamma(n)|_{\zeta} \circ J(n)^{-1}(P))}{1 - (\Gamma(n)|_{\zeta} \circ J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n))} J(n)^{-1} \circ {}^t\zeta \circ \mu \circ \Gamma(n).$$

The above equation yields $I^{-1}M$ in terms of P and n only. Let us replace $I^{-1}M$ by that expression in the second Eq. (15) and in Eq. (17). We obtain a differential equation for $(n, P) \in S^2 \times (\mathbf{so}(E))$, which is the evolution equation for the reduced constrained mechanical system.

5. Two Simple Examples

5.1. First example: Description of the System. The configuration space of the system is the cylinder $Q = \mathbf{R} \times S^1$. We denote by (x, θ) and $(x, \dot{x}, \dot{\theta})$ the usual local coordinates on Q and TQ , respectively. Observe that $\theta \in S^1 = \mathbf{R}/2\pi\mathbf{Z}$ is an angle rather than a real.

We choose for the constraint submanifold

$$C = \{(x, \theta, \dot{x}, \dot{\theta}) \in TQ \mid \dot{x} + x\dot{\theta} = 0\}.$$

It is a rank 1, therefore integrable sub-bundle of TQ (the constraint is holonomic), and the leaves of the foliation of Q defined by C are the images of immersions of \mathbf{R} into Q of the form

$$\theta \mapsto (ce^{-\theta}, \theta \text{ modulo } 2\pi),$$

where c is a constant. The leaf $c = 0$ is a circle S^1 , and all the other leaves are (for their own manifold structure) diffeomorphic to \mathbf{R} .

For the Lagrangian, we choose

$$L(x, \theta, \dot{x}, \dot{\theta}) = \frac{1}{2}(\dot{x}^2 + \dot{\theta}^2).$$

The mechanical system (Q, L, C) has a very simple mechanical interpretation: it describes the motion of a material point of unit mass, constrained on a cylinder on which a system of grooves imposes the relation $\dot{x} + x\dot{\theta} = 0$ between the two components of the velocity. Alternatively, the system can also be physically realized by a skate moving on a cylinder, with a mechanism which imposes the angle made by the blade of the skate with the axis of the cylinder, as a function of the coordinate x of the skate along that axis.

We denote by $(x, \theta, p_x, p_\theta)$ the coordinates on the phase space T^*Q . The Legendre transformation is

$$(x, \theta, \dot{x}, \dot{\theta}) \mapsto (x, \theta, p_x = \dot{x}, p_\theta = \dot{\theta}),$$

and the Hamiltonian of the system is

$$H(x, \theta, p_x, p_\theta) = \frac{1}{2}(p_x^2 + p_\theta^2).$$

The Hamiltonian constraint submanifold is

$$D = \{(x, \theta, p_x, p_0) \in T^*Q \mid p_x + xp_0 = 0\},$$

and the projection bundle W is generated by the vector field along D

$$\frac{\partial}{\partial p_x} + x \frac{\partial}{\partial p_0}.$$

The Hamiltonian vector field X_H and the constrained Hamiltonian vector field X_D are, respectively,

$$X_H = p_x \frac{\partial}{\partial x} + p_0 \frac{\partial}{\partial \theta}, \quad X_D = p_x \frac{\partial}{\partial x} + p_0 \frac{\partial}{\partial \theta} + \lambda \left(\frac{\partial}{\partial p_x} + x \frac{\partial}{\partial p_0} \right),$$

where λ must be chosen so that X_D be tangent to D . We obtain

$$\lambda = -\frac{p_x p_0}{1 + x^2}.$$

The equations of motion are therefore

$$\begin{cases} \frac{dx}{dt} = p_x, \\ \frac{d\theta}{dt} = p_0, \end{cases} \quad \begin{cases} \frac{dp_x}{dt} = -\frac{p_x p_0}{1 + x^2}, \\ \frac{dp_0}{dt} = -\frac{x p_x p_0}{1 + x^2}. \end{cases} \quad (20)$$

We must also add to the system the constraint equation

$$p_x + xp_0 = 0, \quad (21)$$

and choose Cauchy data which satisfy that equation. The corresponding solution of (20) will then satisfy (21) for all t .

5.2. First example: The S^1 action and the Reduced System. We consider the action $\Phi : S^1 \times Q \rightarrow Q$,

$$(\alpha, (x, \theta)) \mapsto (x, \alpha + \theta).$$

Its canonical lifts to TQ and T^*Q are, respectively,

$$(\alpha, (x, \theta, \dot{x}, \dot{\theta})) \mapsto (x, \alpha + \theta, \dot{x}, \dot{\theta})$$

and

$$(\alpha, (x, \theta, p_x, p_0)) \mapsto (x, \alpha + \theta, p_x, p_0).$$

The constraint submanifold C and the hamiltonian constraint submanifold D are invariant under these actions, as well as the foliation of Q defined by C . However, the leaves of that foliation are individually not invariant in general.

The quotient Poisson manifold $\widehat{M} = T^*Q/S^1$ is diffeomorphic to \mathbf{R}^3 (coordinates x, p_x, p_0), with the Poisson tensor

$$\widehat{A} = \frac{\partial}{\partial p_x} \wedge \frac{\partial}{\partial x}.$$

The expression of the reduced Hamiltonian \widehat{H} coincides with that of H , since H does not depend on θ . The reduced hamiltonian constraint submanifold \widehat{D} is

$$\widehat{D} = \{(x, p_x, p_\theta) \in \widehat{M} \mid p_x + xp_\theta = 0\}.$$

It can be identified with \mathbf{R}^2 , with x and p_θ as coordinates. The reduced projection bundle \widehat{W} is generated by the vector field along \widehat{D} ,

$$\frac{\partial}{\partial p_x} + x \frac{\partial}{\partial p_\theta}.$$

With x and p_θ as coordinates on \widehat{D} , the reduced equations of motion are

$$\begin{cases} \frac{dx}{dt} = p_x, \\ \frac{dp_\theta}{dt} = \frac{x^2 p_\theta^2}{1 + x^2}. \end{cases}$$

Since \widehat{H} is a first integral, each integral curve satisfies

$$(1 + x^2)p_\theta^2 = \text{constant}.$$

All the points of the first coordinate axis ($x, p_\theta = 0$), including the origin ($x = 0, p_\theta = 0$) are equilibria of the reduced system, which correspond to true equilibria of the nonreduced system. All the points of the second axis ($x = 0, p_\theta \neq 0$), except the origin, are equilibria of the reduced system which correspond to relative equilibria (stationary motions) of the nonreduced system. It is easy to see that in \widehat{D} , the half axis $\{(x = 0, p_\theta > 0)\}$ is attractive (its attractive basin is the half plane $p_\theta > 0$), and that the other half axis $\{(x = 0, p_\theta < 0)\}$ is repulsive.

5.3. Another Example. We indicate here a simple example of constrained Hamiltonian system in the sense of Definition 3.1, with a stable and attractive equilibrium. We do not use here the reduction procedure: our example should be considered as an already reduced system. The reader is referred to A. Weinstein [41] for a general discussion of stability of equilibria in Poisson Hamiltonian systems.

The phase space is \mathbf{R}^3 (coordinates x, y, z), with the Lie–Poisson structure of $(\mathfrak{so}(3))^*$,

$$A = z \frac{\partial}{\partial x} \wedge \frac{\partial}{\partial y} + x \frac{\partial}{\partial y} \wedge \frac{\partial}{\partial z} + y \frac{\partial}{\partial z} \wedge \frac{\partial}{\partial x}.$$

The Hamiltonian of the system is the linear form

$$H(x, y, z) = \alpha x + \beta y + \gamma z, \quad \alpha, \beta \text{ and } \gamma \in \mathbf{R}, \text{ constants}.$$

The Hamiltonian constraint submanifold D is the plane $z = 0$. The projection bundle W is generated by the vector field along D

$$\frac{\partial}{\partial z} + a \frac{\partial}{\partial x}, \quad a \in \mathbf{R} \text{ constant}.$$

For the constrained Hamiltonian vector field X_D we obtain

$$X_D = (-a\beta x + (\gamma + \alpha x)y) \frac{\partial}{\partial x} - \gamma x \frac{\partial}{\partial y}.$$

It is easy to see that if the constants a, α and γ satisfy

$$\gamma(a\alpha + \gamma) > 0,$$

the origin is a stable and attractive equilibrium of X_D .

Acknowledgements. Several years ago, Rudolf Schmid first spoke to me about the curious behaviour of celtic stones. More recently J.J. Duistermaat suggested that it would be interesting to explain the attractiveness of their stationary motions by geometric means and indicated to me Ref. [9, 10 and 17]. I have had very stimulating discussions with Pierre Dazord about his work [7], with Richard Cushman who indicated to me Ref. [4], and with Paulette Libermann, who suggested to me several important improvements. Marco Favretti indicated to me Ref. [33]. Jean-Pierre Marco read carefully preliminary versions of this work and drew my attention to the fact that attractive relative equilibria can exist in mechanical systems with holonomic constraints. Jean-Pierre Françoise, Daniel Leborgne, Izu Vaisman and Pierre Molino encouraged me by their interest in this work and gave me the opportunity to speak about Mechanics in their seminars. I address my warmest thanks to them all.

References

1. Abraham, R., Marsden, J.E., Foundations of mechanics. Reading, MA: Benjamin, 1978
2. Albert, C.: Le théorème de réduction de Marsden–Weinstein en géométrie cosymplectique et de contact. *J. Geom. and Phys.* **6**, 627–649 (1989)
3. Arnold, V.I., Kozlov, V.V., Neishtadt, A.I.: Mathematical aspects of classical and celestial mechanics. In: Dynamical systems III, Arnold, V.I. (ed.), Berlin, Heidelberg, New York: Springer, 1988
4. Bates, L., Sniatycki, J.: Nonholonomic reduction. *Rep. Math. Phys.* **32**, 99–115 (1991)
5. Benenti, S.: Geometrical aspects of the dynamics of non-holonomic systems. *Journées relativistes, Chambéry*, 14–16 May 1987
6. Cardin, F. and Zanzotto, G.: On constrained mechanical systems: d’Alembert’s and Gauss’ principles. *J. Math. Phys.* **30**, 7, 1473–1479 (1989)
7. Dazord, P.: Mécanique hamiltonienne en présence de contraintes. *Illinois J. Math.* **38**, 1, 148–175 (1994)
8. Godbillon, C.: Géométrie différentielle et mécanique analytique. Paris: Hermann, 1969
9. Karapetian, A.V.: On the problem of steady motions stability of nonholonomic systems. *Prikl. Matem. Mekhan.* **44**, 3, 418–426 (1980) (in Russian). English translation: *PMM USSR* **44**, 295–300 (1980)
10. Karapetian, A.V.: On realizing nonholonomic constraints by viscous friction forces and celtic stones stability. *Prikl. Matem. Mekhan.* **45**, 1, 42–51 (1981) (in Russian). English translation: *PMM USSR* **45**, 30–36 (1981)
11. Koiller, J.: Reduction of some classical non-holonomic systems with symmetry. *Arch. Rational Mech. Anal.* **118**, 133–148 (1992)
12. Krishnaprasad, P.S.: Eulerian many-body problems. In: Dynamics and control of multi-body systems. Marsden, J.E., Krishnaprasad, P.S., Simo, J.C. (eds.), Contemporary Mathematics **97**, Providence, RI: American Mathematical Society, 1989, pp. 187–208
13. Krishnaprasad, P.S., Yang, R.: Geometric phase, anholonomy, and optimal movement. *Proc. of the 1991 IEEE International conference on Robotics and Automation* **3**, 2185–2189 (1991)
14. Kummer, M.: On the construction of the reduced phase space of a Hamiltonian system with symmetry. *Indiana Univ. Math. J.* **30**, 2, 281–291 (1981)
15. Libermann, P., Marle, C.-M.: Symplectic geometry and analytical mechanics. Dordrecht: D. Reidel Publishing Company, 1987
16. Lichnerowicz, A.: Les variétés de Poisson et leurs algèbres de Lie associées. *J. Differ. Geom.* **12**, 253–300 (1977)
17. Markeev, A.P.: On the dynamics of a solid on an absolutely rough plane. *Prikl. Matem. Mekhan.* **47**, 4, 575–582 (1983) (in Russian). English translation. *PMM USSR* **47**, 4, 473–478 (1983)

18. Marle, C.-M.: Sur la géométrie des systèmes mécaniques à liaisons actives. *C. R. Acad. Sci. Paris* **311**, I, 839–845 (1990)
19. Marle, C.-M.: Géométrie des systèmes mécaniques à liaisons actives. In: *Symplectic geometry and mathematical physics*, Donato, P., Duval, C., Elhadad, J., Tuynman, G.M. (eds.), Boston: Birkhäuser, 1991, pp. 260–287
20. Marsden, J.E.: *Lectures on mechanics*. London Math. Soc. Lect. Notes Series **174**, Cambridge: Cambridge University Press, 1992
21. Marsden, J.E. and Ratiu, T.S.: Reduction of Poisson manifolds. *Lett. Math. Phys.* **11**, 161–170 (1986)
22. Marsden, J., Weinstein, A.: Reduction of symplectic manifolds with symmetry. *Rep. Math. Phys.* **5**, 121–130 (1974)
23. Massa, E., Pagani, E.: Classical dynamics of non-holonomic systems: A geometric approach. *Ann. Inst. Henri Poincaré, Physique Théor.* **55**, 1, 511–544 (1991)
24. Meyer, K.R.: Symmetry and integrals in mechanics. In: *Dynamical systems*, Peixoto, M. (ed.), New York: Academic Press, 1973, pp. 259–273
25. Mikami, K.: Reduction of local Lie algebra structures. *Proc. Am. Math. Soc.* **105**, 3, 686–691 (1989)
26. Montgomery, R., Marsden, J.E., Ratiu, T.: Gauged Lie–Poisson structures. In: *Fluids and plasmas: Geometry and dynamics*. Contemporary mathematics **28**, Providence, RI: Am. Math. Soc., 1984, pp. 101–114
27. Nunes da Costa, J.: Réduction des variétés de Jacobi. *C. R. Acad. Sc. Paris* **308**, I, 101–103 (1989)
28. Pascal, M.: The asymptotic solution of the equations of motion of the celtic stone. *Prikl. Matem. Mekhan.* **47**, 2, 321–329 (1983) (in Russian). English translation: *PMM USSR* **47**, 2, 269–276 (1984)
29. Souriau, J.-M.: *Structure des systèmes dynamiques*. Paris: Dunod, 1969
30. Sternberg, S.: On minimal coupling and the symplectic mechanics of a classical particle in the presence of a Yang–Mills field. *Proc. Nat. Acad. Sci. USA* **74**, 5253–5254 (1977)
31. Tulczyjew, W.M.: Sur la différentielle de Lagrange. *C.R. Acad. Sc. Paris* **280**, A, 1295–1298 (1975)
32. Tulczyjew, W.M.: *Geometric formulations of physical theories*. Napoli, Italy: Bibliopolis, 1989
33. Vershik, A.M.: Classical and non-classical dynamics with constraints. In: *Global analysis, Studies and Applications I*, Yu. G. Borisovich and Yu. E. Gliklikh, ed., *Lecture Notes in Math.* **1108**, Berlin: Springer-Verlag, 1984, pp. 278–301
34. Vershik, A.M., Faddeev, L.D.: Lagrangian mechanics in invariant form. *Sel. Math. Sov* **1**, 339–350 (1981)
35. Vershik, A.M., Gershkovich, Y.Ya.: Nonholonomic dynamical systems, geometry of distributions and variational problems. In: *Dynamical systems VII*, Arnold, V.I. and Novikov, S.P. (eds.), Berlin, Heidelberg, New York: Springer, 1994, pp. 1–81
36. Walker, G.T.: On the curious dynamical properties of celts. *Proceedings of the Cambridge Philosophical Society*, vol. **8**, years 1892–1895, p. 305
37. Walker, G.T.: On a dynamical top. *Quart. J. Pure and Appl. Math.* **28**, 175–184 (1896)
38. Walker, J.: The mysterious “rattleback”: A stone that spins in one direction and then reverses. *Scientific American* **241**, 4, 172–184 (1979)
39. Weinstein, A.: A universal phase space for particles in Yang–Mills field. *Letters Math. Phys.* **2**, 417–420 (1978)
40. Weinstein, A.: The local structure of Poisson manifolds. *J. Differ. Geom.* **18**, 523–557 (1983) and **22**, 255 (1985)
41. Weinstein, A.: Stability of Poisson–Hamilton equilibria. In: *Fluids and plasmas: Geometry and dynamics*, Contemporary Mathematics **28**, Providence RI: Am. Math. Soc., 1984, pp. 3–13
42. Whittaker, E.T.: *A treatise on the analytical dynamics of particles and rigid bodies*. Cambridge: Cambridge University Press, 1904 (fourth edition 1937, reissued with Foreword 1988)
43. Woodhouse, N.: *Geometric quantization*. Oxford: Oxford University Press, 1980
44. Yang, R.: *Nonholonomic geometry, Mechanics and Control*. Ph.D. Dissertation, University of Maryland, 1992