

Comments on a Recent Solution to Wightman's Axioms

Karl-Henning Rehren

II. Institut für Theoretische Physik, Universität Hamburg,
Luruper Chaussee 149, D-22761 Hamburg, Germany.
E-mail: rehren@x4u2.desy.de

Received: July 25, 1995 / Accepted: September 25, 1995

Abstract: A class of exact Wightman functionals satisfying all fundamental physical requirements in an arbitrary number of space-time dimensions, which bear the appearance of describing interacting fields, was recently constructed by C. Read [1]. It is shown here, that the construction can be considerably generalized, and that even the enlarged class belongs to the Borchers class of a system of generalized free fields.

1. Introduction

Ever since Wightman's formulation of the axioms [2] to be satisfied by the collection of n -point functions of local quantum fields, there has been a discomfoting lack of models. Apart from models with polynomial interaction in two and three space-time dimensions, there are essentially only constructions based upon free fields and generalized free fields [3] available. These constructions involve Wick polynomials of derivatives of a given field, as well as so-called p - and s -products [4] (i.e., pointwise products resp. sums of independent fields in different Hilbert spaces). Although one can easily produce non-vanishing truncated Wightman functionals, such models do not describe interacting particles.

We recall the well-known list of axioms for a hermitian scalar field, referring to the standard literature [2] for the precise formulation: Positivity and Hermiticity permit to reconstruct a Hilbert space containing the cyclic vacuum vector, and an (in general unbounded) hermitian field $\phi(f)$ on this Hilbert space whose vacuum correlation functions are given by the Wightman functional. Poincaré Invariance of the Wightman functional ensures the invariance of the vacuum vector along with the Poincaré covariance of the reconstructed field. The Spectrum Condition and Cluster Property ensure the positivity of the energy spectrum

and the uniqueness of the vacuum. Finally, Locality expresses the commutativity of field operators $\phi(f)$ smeared in causally disconnected regions of space-time.

In a recent paper [1], a new class of solutions was presented which satisfy all Wightman axioms in any number $d + 1$ of space-time dimensions. The models are based on Feynman-like rules without at the same time being perturbative; instead, every n -point function is obtained as a sum over finitely many graphs. Some of the free input parameters of the models play a similar role as coupling constants in ordinary Feynman rules (although it will become clear in the course of this communication that they rather determine the structure of the field as a Wick polynomial), while the remaining parameters serve as appropriate cutoffs to ensure convergence of all sums and integrals involved. Unlike regulators in perturbative approaches, these cutoffs need not be removed in the end. They comprise a smooth space-time cutoff function, a mass distribution in a finite mass interval, and a numerical limitation of the number of vertices.

In the present contribution, we intend to shed new light onto these models. It is found that the class of models [1] can be considerably extended by a generalization which essentially makes the “interaction polynomial” obsolete. We shall then reduce the entire construction to Wick products of generalized free fields, and discuss aspects of the sharp mass limit of the extended class of models in comparison with the original class.

2. A Quick Review of the Original Construction

The definition given in ref. [1] of the n -point functions

$$\mathcal{W}_n(f_1, \dots, f_n) \equiv \langle \Omega, \phi(f_1) \cdots \phi(f_n) \Omega \rangle \tag{1}$$

is the following (in somewhat schematic notation, to be presently specified).

$$\begin{aligned} \mathcal{W}_n(f_1, \dots, f_n) := & \sum_{\text{banded graphs}} \frac{1}{G_{\text{inter}}!} \int \left(\prod_{e \in E_{\text{inter}}} dg dp \theta(p^0) |\varphi(gp)|^2 \right) \times \\ & \times \prod_{i=1}^n \left[\hat{f}_i(P_i[p]) \cdot \frac{1}{G_{i,\text{dom}}!} \int \left(\prod_{e \in E_{i,\text{dom}}} dq \theta(q^0) |\varphi(q)|^2 \right) \prod_{v \in V_i} (r! a_r \delta_v(q, gp)) \right], \end{aligned} \tag{2}$$

where the sum extends over a class of “banded graphs.” A banded graph is a (possibly disconnected) graph which contains one connected subgraph (“band graph” or simply “band”) for every field entry $\phi(f_i)$ in Eq. (1) such that the sets V_i of vertices of the n bands are disjoint and exhaust all vertices of the full graph; every band has a distinguished “external” vertex of degree 1. The internal vertices are of degree $2 \leq r \leq R$ for some finite number R . In each band the number s_i of vertices is limited by some finite number S , and the vertices are labelled $1, \dots, s_i$. Inequivalent labellings of the vertices of the same abstract graph are considered as different banded graphs to be summed over. The banded graph has no external lines and no edges connecting a vertex to itself.

Edges which connect vertices of the same band will be called “domestic;” they are oriented from the vertex with lower ordinal number to the vertex with higher ordinal number; if a domestic edge connects to the external vertex of the band, then it may carry both orientations. (This restriction on the orientations of domestic edges in [1] seems not really to be necessary. We view it as another model input parameter.) Edges which connect vertices of different bands will be called “interband;” they are oriented from the band with lower ordinal number to the band with higher ordinal number. The integration rules are the following.

(i) Associated with every domestic edge $e \in E_{\text{dom}}$ is a momentum variable $q \in \mathbb{R}^{d,1}$, to be integrated over Minkowski space $\mathbb{R}^{d,1}$ with the measure $dq \theta(q^0) |\varphi(q)|^2$. Here θ is the Heaviside step function for the energy, and $|\varphi(k)|^2 = \rho(k^2) \cdot |\hat{\Theta}(k)|^2$ consists of a smooth mass distribution ρ with support in a mass interval $m_0^2 \leq k^2 \leq m_1^2$, where $0 < m_0 < m_1$, and the square modulus of the Fourier transform of a real space-time cutoff function $\Theta(x)$ in the Schwartz space $\mathcal{S}(\mathbb{R}^{d+1})$ (i.e., smooth and all derivatives decaying faster than any power of the arguments).

(ii) Associated with each interband edge $e \in E_{\text{inter}}$ are a momentum variable $p \in \mathbb{R}^{d,1}$ and a group variable $g \in L$ which runs over the four connected components of the full Lorentz group. These variables are integrated with the measure $d\mu(g, p) = dg dp \theta(p^0) |\varphi(gp)|^2$ involving the right invariant Haar measure dg on L . We notice that the variables g enter only in the combination gp ; thus effectively, the group integrations extend only over the Lorentz boosts and the time inversion in the rest frames of the momenta p . In fact, up to the redundant integral over the compact stabilizer group $O(d)$ of p , the measure is $d\mu(g, p) \propto dp \theta(p^0) dk (k^2)^{\frac{1-d}{2}} |\varphi(k)|^2 \delta(p^2 - k^2)$, where $k = gp \in \mathbb{R}^{d,1}$.

(iii) The combinatorial weights $G_{\text{inter}}!$ and $G_{i,\text{dom}}!$ are given by $\prod_{v,v'} m(v, v')!$ where $m(v, v')$ is the number of edges connecting the vertices v and v' , and the product extends over all pairs of vertices in different bands, and within the band i , respectively.

(iv) The integrand contains a “coupling constant” $r!a_r$ for every internal vertex v of degree r along with the momentum conservation delta function $\delta_v(q, gp)$ for the momentum flow at that vertex involving the domestic momenta q and the Lorentz transformed interband momenta gp .

(v) Finally, the momentum transfer at each band is given by $P = P[p] = \sum_{\text{out}} p - \sum_{\text{in}} p$, where the sums refer to all interband momenta flowing out of the band, resp. into the band. There contributes to the integrand a factor $\hat{f}_i(P_i)$ for every band, where \hat{f}_i are the Fourier transforms of the real test functions $f_i \in \mathcal{S}(\mathbb{R}^{d+1})$. It is worth noting that the momentum transfer of the field operators (involving the interband momenta p) is decoupled from the momentum flow within a graph (involving the transformed momenta gp and the domestic momenta q).

The rapid decay of the integrand along with the momentum conservation delta functions guarantees that every integral in Eq. (2) converges absolutely. Due to the limitations S of the number of vertices per band and R of the degree

of vertices, there are only finitely many different banded graphs, so the functional (2) is well defined. It is evident that it is translation invariant since the arguments of the test functions sum up to zero, $\sum_i P_i = 0$. It is invariant under the orthochronous Lorentz group L^\uparrow since for $\gamma \in L^\uparrow$ the change of integration variables $p \mapsto \gamma p$, $g \mapsto g\gamma^{-1}$ and $q \mapsto q$ takes the functional (2) into its Lorentz transform. We note here that the domestic variables q , being coupled by momentum conservation to the Lorentz invariant variables $k = gp$, may also be considered as Lorentz invariant. This explains why the momentum cutoff does not violate Lorentz invariance: it affects only Lorentz invariant variables.

The spectrum condition is satisfied since for every $j \leq n$, the sum of momentum transfers $\sum_{i \leq j} P_i$ is a sum of interband momenta p restricted to the forward light-cone. The uniqueness of the vacuum will become apparent later when we identify the Hilbert space.

The axiom with the most unprecedented realization in the new models is Locality. We shall limit ourselves here to the simple core of the exact but tedious argument given in [1]. Namely, we shall discuss the commutativity of $\phi(f)$ and $\phi(f')$ when f and f' are delta functions at space-like separated points x and $x' \in \mathbb{R}^{d,1}$. Although these are not a priori admitted as test functions (the convergence argument given in [1] will fail for such functions), we may argue as follows for the validity of the simplified argument: Given the spectrum condition and Poincaré invariance, it is a standard result [2] that at the Jost points where all coordinates are space-like separated, the Wightman distributions are in fact functions. By the Reeh-Schlieder theorem [2], it is then sufficient to test the commutativity of $\phi(x)$ and $\phi(x')$ within Wightman functions at Jost points.

Consider therefore the change in the sum (2) when the field entries $\phi(x)$ at position i and $\phi(x')$ at position $i + 1$ are interchanged. Along with every banded graph contributing to $\mathcal{W}'_n = \mathcal{W}'_n(\dots x, x' \dots)$ there corresponds a banded graph contributing to $\mathcal{W}_n = \mathcal{W}_n(\dots x', x \dots)$ which differs only by the numbering of bands and therefore by the orientation of the interband edges extending from band i to band $i + 1$. Let there be σ such edges in a given graph. The associated integration variables g and p , here and later on collectively indicated by $(g, p)_\sigma$, enter the arguments of the test functions $\hat{f}_i(P) = \exp -iPx$ and $\hat{f}_{i+1}(P) = \exp -iPx'$ through $P = P[p]$, the measure factors $\theta(p^0)|\varphi(gp)|^2$, and the momentum conservation delta functions. Due to Poincaré invariance, we are free to choose the Lorentz frame such that $x^0 = x'^0$, hence $P(x - x') = -\mathbf{P}(\mathbf{x} - \mathbf{x}')$. In this frame, each integral contributing to \mathcal{W}'_n is of the form

$$\int \dots \times \int \left(\prod_1^\sigma dg dp \theta(p^0) |\varphi(gp)|^2 \exp(i\mathbf{p}(\mathbf{x} - \mathbf{x}')) \right) \prod_v \delta_v(\dots, (gp)_\sigma) \times \dots, \tag{3a}$$

while the corresponding integral contributing to \mathcal{W}_n is of the form

$$\int \dots \times \int \left(\prod_1^\sigma dg dp \theta(p^0) |\varphi(gp)|^2 \exp(i\mathbf{p}(\mathbf{x}' - \mathbf{x})) \right) \prod_v \delta_v(\dots, (-gp)_\sigma) \times \dots, \tag{3b}$$

where ... stands for further factors and dependences on other variables common to both contributions to \mathcal{W}'_n resp. \mathcal{W}'_n , and independent of $(g, p)_\sigma$. By the change of integration variables $(g, p)_\sigma \mapsto (gT, Pp)_\sigma$, where T resp. P are the time resp. space inversion in the given Lorentz frame, the integrands are transformed into each other (note that the cutoff function Θ is real, hence $\hat{\Theta}(k) = \hat{\Theta}(-k)$, so $|\varphi(gp)|^2$ is invariant under $gp \mapsto -gp$). This establishes Locality, graph by graph.

Finally, Positivity of the Wightman functional (2) becomes manifest if one views every graph integral as an operator product of integration kernels (= the square brackets in Eq. (2)) in the Fock space $\mathcal{F}(H)$ over the underlying Hilbert space $H = L^2(L \times \mathbb{R}^{d,1}, d\mu)$ with the measure $d\mu(g, p) = dg dp \theta(p^0)|\varphi(gp)|^2$. More precisely, every band subgraph with ν_{out} resp. ν_{in} interband edges oriented out of resp. into it, and μ "bypassing" interband edges linking some lower band to some higher band, corresponds to a kernel interpolating from the $(m = \nu_{\text{out}} + \mu)$ - "particle" subspace $\mathcal{F}_m \equiv H^{\otimes m}$ to the $(n = \nu_{\text{in}} + \mu)$ - "particle" subspace $\mathcal{F}_n \equiv H^{\otimes n}$. As a kernel, it is a function of m pairs of variables (g, p) to be integrated (with the measure $d\mu$) with the variables of a wave function in \mathcal{F}_m ("annihilation part"), and n free pairs of variables (g, p) ("creation part"). It is given by the expression in square brackets in Eq. (2) which is a function of the integration variables $(g, p)_{\text{out}}$ and $(g, p)_{\text{in}}$ associated with the out- and ingoing interband edges, while the bypassing edges contribute as delta function kernels $\delta(g, g')\delta(p - p')/|\varphi(gp)|^2$. (Note that every application of an integral kernel operator in H or $\mathcal{F}(H)$ involves the integral measure $d\mu(g, p) = dg dp \theta(p^0)|\varphi(gp)|^2$, so one has not to worry about denominators.)

A careful analysis of the combinatorics reveals that actually every field operator in Eq. (1) arises as a finite sum of integral kernels, sandwiched between the completely symmetrizing projection operator $\Pi = \bigoplus_n \Pi_n$ onto the symmetric Fock space $\mathcal{F}_+(H) = \Pi \cdot \mathcal{F}(H)$. Namely, let $(k)_{\text{in}}$ resp. $(k)_{\text{out}}$ denote two collections of ν_{in} ingoing resp. ν_{out} outgoing momentum variables, and let the functions $K_{\nu_{\text{out}}}^{\nu_{\text{in}}}((k)_{\text{in}}; (k)_{\text{out}})$ be given by the domestic integrals summed over all band graphs with these variables assigned to the given number of in- and outgoing external lines:

$$K_{\nu_{\text{out}}}^{\nu_{\text{in}}}((k)_{\text{in}}; (k)_{\text{out}}) := \sum_{\substack{\text{band} \\ \text{graphs}}} \frac{1}{G_{\text{dom}}!} \int \left(\prod_{e \in E_{\text{dom}}} dq \theta(q^0) |\varphi(q)|^2 \right) \prod_{v \in V} (r! a_r \delta_v(q, k)), \tag{4}$$

as read off Eq. (2). Since there are no according band graphs, $K_{\nu_{\text{out}}}^{\nu_{\text{in}}}$ vanish for $\nu_{\text{in}} + \nu_{\text{out}} > S(R - 2) + 1$. Let furthermore

$$\begin{aligned} \hat{\Phi}(f)_{\nu_{\text{out}}, \mu}^{\nu_{\text{in}}, \mu}((g, p)_{\text{in}}, (g, p)_\mu; (g, p)_{\text{out}}, (g', p')_\mu) := \\ := \hat{f}(P[p]) \cdot [K_{\nu_{\text{out}}}^{\nu_{\text{in}}}((gp)_{\text{in}}; (gp)_{\text{out}}) \otimes (\text{delta kernels})^{\otimes \mu}] \end{aligned} \tag{5}$$

be integral kernels interpolating between $H^{\otimes \nu_{\text{out}} + \mu}$ and $H^{\otimes \nu_{\text{in}} + \mu}$, where \hat{f} resp. $K_{\nu_{\text{out}}}^{\nu_{\text{in}}}$ are functions of the indicated in- and outgoing interband variables $P[p] = \sum_{\text{out}} p - \sum_{\text{in}} p$ resp. $(gp)_{\text{in}}$ and $(gp)_{\text{out}}$ only, and $(\text{delta kernels})^{\otimes \mu}$ stands for the

delta functions in μ pairs of bypassing interband variables $(g, p)_\mu$ as explained before.

Then $\phi(f)$ is represented by the integral kernels

$$\phi(f) = \sum_{\nu_{in}, \nu_{out}, \mu} \prod_{\nu_{in} + \mu} \left(\frac{\sqrt{(\nu_{in} + \mu)! (\nu_{out} + \mu)!}}{\mu! \nu_{in}! \nu_{out}!} \Phi(f)_{\nu_{out}, \mu}^{\nu_{in}, \mu} \right) \prod_{\nu_{out} + \mu} \cdot \quad (6)$$

We shall refer to the functions (4) as the “reduced kernels.” They are symmetric in both their in- and outgoing sets of variables. They encode the entire dependence of the model on the choice of the constants a_r ($r \leq R$) and the number S (limiting the number of graphs), as well as the previously mentioned restriction on the orientations of domestic edges. Since the remaining input parameters determine the integral measure of the Hilbert space H , the model is now specified by the reduced kernels $K_{\nu_{out}}^{\nu_{in}}$ and the measure $d\mu$.

The assertion that the prescriptions (2) and (4–6) produce the same Wightman functional is the central claim of this section. Since it is essential for the rest of this communication, we formulate it as a lemma.

Lemma. *Let the fields $\phi(f)$ be represented as integral kernels (4–6) on the symmetrized Fock space $\mathcal{F}_+(H)$. Let the vacuum vector Ω be represented by the number $1 \in \mathbb{C} \equiv \mathcal{F}_0 \subset \mathcal{F}_+(H)$. Then the vacuum correlations of $\phi(f)$ coincide with Eq. (2).*

Proof. A vacuum correlation of n operators $\phi(f_i)$ is a finite sum, extending over single band graphs contributing to every reduced kernel (4) and therefore to (5), and over single permutations contributing to the symmetrizing projections in Eq. (6). Every such contribution clearly corresponds to a banded graph contributing to the sum (2), and vice versa. However, the correspondence is not always one to one. It only has to be checked that multiple counting and numerical coefficients according to Eqs. (4–6) together produce the correct combinatorial weights $(G_{inter}! \prod_i G_{i, dom}!)^{-1}$ as in Eq. (2). The domestic factors are explicitly present in Eq. (4) and need not be considered any longer.

At this point, in order to get the global factor $1/G_{inter}!$, it is crucial that the sum (2) extends over all inequivalent labellings of the vertices of the banded graphs, while the sum (6) extends over all inequivalent labellings of the vertices and external lines of the band graphs.

We start with a two-point function and consider a banded graph G contributing to Eq. (2) with its two band subgraphs G_1 and G_2 . We label the vertices of G_1 as v_i , those of G_2 as w_j (for this matter not distinguishing the external vertex of each band graph from its internal vertices). Let m_{ij} edges connect v_i with w_j , thus $\nu_i = \sum_j m_{ij}$ interband edges connect to v_i , and $\kappa_j = \sum_i m_{ij}$ interband edges connect to w_j . Let finally $\nu = \sum_i \nu_i = \sum_j \kappa_j$ denote the total number of interband edges of G . Apart from the domestic combinatorial weights (which are common to Eq. (2) and Eq. (4)), the banded graph enters Eq. (2) with the weight $1/G_{inter}! = \prod_{ij} 1/m_{ij}!$.

Let us compute the coefficient according to the prescription (4–6). Assume for the moment that the symmetry groups of the vertices of G_1 and G_2 considered as abstract subgraphs of G , i.e., ignoring the labelling of vertices and orientation of edges, are trivial. This assumption is equivalent to the assumption that each labelling of the vertices gives rise to a different labelled band graph. Then there are $\nu! / \prod_i \nu_i!$ inequivalent assignments of ν distinguished momenta (k) = (gp) to the external lines of G_1 , and similarly $\nu! / \prod_j \kappa_j!$ assignments of momenta to the external lines of G_2 , each giving rise to one term in the sums (4). Furthermore, each set of ν_i lines extending from v_i can be partitioned in $\nu_i! / \prod_j m_{ij}!$ ways to join the vertices w_j with multiplicities m_{ij} , and there is a similar number of partitions for the vertices w_j . Finally, each of the $m_{ij}!$ contractions of m_{ij} lines between v_i and w_j is counted separately. Thus the total number of contractions of band graphs which give rise to the same banded graph G equals

$$\frac{\nu!}{\prod_i \nu_i!} \frac{\nu!}{\prod_j \kappa_j!} \prod_i \frac{\nu_i!}{\prod_j m_{ij}!} \prod_j \frac{\kappa_j!}{\prod_i m_{ij}!} \prod_{ij} m_{ij}! = \frac{(\nu!)^2}{G_{\text{inter}}!} .$$

Since the symmetrizing projection operator between the two kernels contributes a weight $1/\nu!$ for each permutation, and the explicit numerical coefficients in Eq. (6) contribute another factor $(\sqrt{\nu!}/\nu!)^2 = 1/\nu!$ to the two-point function, the weight of each term is $1/(\nu!)^2$ and the combinatorial weight as in Eq. (2) is reproduced.

Now let the vertices of the abstract band graphs G_i possess some symmetry groups S_i and let $S \subset S_1 \times S_2$ be the symmetry group of the vertices of G . Then one may sum in Eq. (2) over all labellings of the internal vertices of G_1 and over all labellings of the internal vertices of G_2 independently, if one includes a correction factor $1/|S|$ for overcounting of banded graphs. Similarly, one may sum in Eq. (4) over all labellings of internal vertices and over all labellings of external lines independently, if one includes a correction factor $|S_i|$ for each of the two kernels. On the other hand, the counting of inequivalent contractions of two band graphs with labelled vertices and external lines which give rise to the same banded graph with unlabelled interband edges provides an additional multiplicity factor $|S_1 \times S_2/S|$ which cancels the correction factors for overcounting. We conclude that for two-point functions, the prescriptions (2) and (4–6) produce the same combinatorial weights.

Turning now to higher n -point functions, we repeat the previous reasonings with the obvious generalization. The total number of contractions of band graphs G_i ($i = 1, \dots, n$) contributing to Eq. (4) which give rise to the same banded graph G is found to exceed the expected weight $1/G_{\text{inter}}!$ by the factor $\prod_i \nu_{\text{in}}^i! \nu_{\text{out}}^i!$, where ν_{in}^i and ν_{out}^i are the number of in- and outgoing interband edges of G_i . This factor is cancelled by the corresponding factors in the denominators of the numerical coefficients in Eq. (6). The square root numerators of the latter (which arise twice each) are compensated by the weights $1/(\nu + \mu)!$ of each permutation within the projections $\prod_{\nu+\mu}$ between two kernels, while the remaining factors $1/\mu!$ in the denominators of Eq. (6) are cancelled by the number of permutations

of the sets of bypassing variables at each kernel. The discussion of symmetries of the vertices of band graphs also parallels the two-point case. This completes the proof of the lemma. \square

Due to the lemma, Positivity of the Wightman functional (2) is manifest. Let us now complete the list of arguments that Eq. (2) fulfills all Wightman axioms.

The orthochronous Lorentz group L^\uparrow is represented on H , and therefore on the Fock space by second quantization, by the tensor product of the natural action (on $p \in \mathbb{R}^{d,1}$) and the right regular action (on $g \in L$). Since the measure is invariant under this action, the representation is unitary. For the same reason, functions of $k = gp$ represent Lorentz invariant elements of H or $\mathcal{F}_+(H)$, and the reduced kernels (4) are scalar quantities. Consequently, the full kernels (5) and finally the fields (6) transform like scalar fields.

The invariant Wightman domain of the field operators $\phi(f)$ is $\mathcal{D}_0^\phi = \text{span}(\prod \phi(f_i))\Omega \subset \mathcal{F}_+(H)$. It is clear that Ω is the only translation invariant vector in the Hilbert space $\overline{\mathcal{D}_0^\phi}$. The property of the reduced kernels

$$\overline{K_{\nu_{\text{out}}}^{\nu_{\text{in}}}((k)_{\text{in}}; (k)_{\text{out}})} = K_{\nu_{\text{in}}}^{\nu_{\text{out}}}((k)_{\text{out}}; (k)_{\text{in}}) \tag{7}$$

ensures that $\langle \Phi, \phi(f)\Psi \rangle = \langle \phi(f)\Phi, \Psi \rangle$ for $\Phi, \Psi \in \mathcal{D}_0^\phi$, i.e., ϕ is a hermitian field.

Corollary. ([1]) *The n -point distributions given by Eq. (2) define a hermitian scalar local Wightman field.*

3. An Enlarged Class of Local Wightman Fields

We make the following crucial observation. As was remarked before, the fields $\phi(f)$ are completely specified as operators on $\mathcal{F}_+(H)$ by the reduced kernels (and the measure) while it is irrelevant how these kernels were produced by domestic integrals over band graphs. One may indeed *choose* a sequence of reduced kernels as the primary model input.

None of the arguments for Finiteness, Positivity, Translation Invariance, Lorentz Invariance and Spectrum Condition along with the Cluster Property is affected if one replaces the reduced integral kernels $K_{\nu_{\text{out}}}^{\nu_{\text{in}}}((k)_{\text{in}}; (k)_{\text{out}})$ given by Eq. (4) by arbitrary smooth polynomially bounded functions of the respective in- and outgoing variables $k = gp$, and defines $\phi(f)$ by Eqs. (5) and (6). Since the kernels only act between symmetrizing projections, these functions may be chosen symmetric in both of their two sets of variables. Furthermore, the argument for Hermiticity of $\phi(f)$ is unaffected provided the reduced kernels satisfy condition (7) above. These assertions are obvious except, maybe, the one concerning Finiteness, for which we refer to estimate (11) in the lemma below.

Finally, the above simplified argument for Locality remains unaffected when in Eqs. (3) the delta functions due to each band (integrated over the domestic variables) are replaced by the respective reduced kernels, provided

$$K_{\mu}^{\nu+\sigma}((k)_{\nu} \cup (k)_{\sigma}; (k)_{\mu}) = K_{\sigma+\mu}^{\nu}((k)_{\nu}; (-k)_{\sigma} \cup (k)_{\mu}) , \tag{8}$$

where \cup indicates the union of the respective sets of variables. (The reduced kernels (4) have this symmetry.) Namely, when K resp. K' refer to the reduced kernels due to the field entries $\phi(x)$ resp. $\phi(x')$, then in a typical contribution to \mathcal{W}_n products of reduced kernels

$$K_{\sigma+\kappa}^{\lambda}(\dots; (gp)_{\sigma} \cup \dots) K_{\mu}^{\nu+\sigma}(\dots \cup (gp)_{\sigma}; \dots)$$

replace the delta functions in Eq. (3a), while in the corresponding contribution to \mathcal{W}'_n ,

$$K_{\sigma+\mu}^{\nu}(\dots; (gp)_{\sigma} \cup \dots) K_{\kappa}^{\lambda+\sigma}(\dots \cup (gp)_{\sigma}; \dots)$$

replace the delta functions in Eq. (3b). As before in Sect. 2, the dots indicate dependences on other variables which are common to both contributions. With the same change of the integration variables $(g, p)_{\sigma}$ as before, the crossing symmetry (8) ensures $\mathcal{W}_n = \mathcal{W}'_n$.

In view of the two conditions (7) and (8), it is sufficient to specify a terminating (in order not to spoil temperedness of the ensuing distribution) sequence $F \equiv (F_{\nu}(k_1, \dots, k_{\nu}))_{\nu \in \mathbb{N}}$ of smooth polynomially bounded symmetric functions satisfying

$$\overline{F_{\nu}(k_1, \dots, k_{\nu})} = F_{\nu}(-k_1, \dots, -k_{\nu}) \tag{9}$$

(i.e., the Fourier transforms of real functions). We shall call functions satisfying the symmetry (9) “hermitian.” Then the reduced kernels

$$K_{\mu}^{\nu}((k)_{\nu}; (k)_{\mu}) := F_{\nu+\mu}((k)_{\nu} \cup (-k)_{\mu}) \tag{10}$$

satisfy conditions (7) and (8).

The reduced kernels (10) inserted into Eqs. (5) and (6) define a manifestly finite hermitian scalar local Wightman field ϕ_F . Due to the lemma above, this class of fields extends the class constructed in [1].

The preceding arguments apply also without substantial change to mixed Wightman functionals with field entries $\phi_{F^{(i)}}(f_i)$ specified by different sequences $F^{(i)}$ of reduced kernels. The following conclusion is immediate.

Corollary. *Every terminating sequence $F = (F_{\nu}(k_1, \dots, k_{\nu}))_{\nu \in \mathbb{N}}$ of (smooth polynomially bounded symmetric) hermitian functions defines, upon insertion of the reduced kernels (10) into Eqs. (5) and (6), a hermitian scalar local Wightman field ϕ_F on the symmetric Fock space $\mathcal{F}_{\pm}(H)$. The fields ϕ_F associated with different sequences F are defined (as operator-valued tempered distributions) on the joint Wightman domain $\mathcal{D}_0 = \text{span}(\prod \phi_{F^{(i)}}(f_i))\Omega \subset \mathcal{F}_{\pm}(H)$. They are relatively local with respect to each other.*

We shall call a field ϕ_F “of order ν ” if $F_{\nu} \neq 0$ and all other F_{μ} vanish. In the general case, ϕ_F is a finite sum over its components of order ν .

The assignment $F \mapsto \phi_F$ is clearly real linear (in the obvious sense for each component of order ν). It is continuous in the following sense.

Lemma. *The correlations of fields $\phi_{F^{(i)}}$ (of fixed order ν_i) are bounded by*

$$|\langle \Omega, \phi_{F^{(1)}}(f_1) \cdots \phi_{F^{(n)}}(f_n) \Omega \rangle| \leq C_{(\nu_i)} \prod_{i=1}^n \|F_{\nu_i}^{(i)}\|, \tag{11}$$

where the constants $C_{(\nu_i)} < \infty$ depend on the test functions $f_i \in \mathcal{S}(\mathbb{R}^{d+1})$, and

$$\|F_{\nu}\|^2 := \sup_{m_0^2 \leq m_i^2 = p_i^2 \leq m_1^2} \int \left(\prod_{i=1}^{\nu} dg_i |\varphi(g_i p_i)|^2 \right) |F_{\nu}(g_1 p_1, \dots, g_{\nu} p_{\nu})|^2. \tag{12}$$

Thus, for each test function, the assignment $F \mapsto \phi_F(f)$ is weakly continuous on the joint Wightman domain $\mathcal{D}_0 \subset \mathcal{F}_+(H)$.

Proof. Every correlation of field operators in the vacuum state is a finite sum of integrals of the form

$$\int \left(\prod dp \theta(p^0) \right) \prod_{i=1}^n \hat{f}_i(P_i[p]) \int \left(\prod dg |\varphi(gp)|^2 \right) \prod_{i=1}^n K^{(i)}((gp)_{\nu_i - \mu_i}; (gp)_{\mu_i})$$

such that each momentum gp enters precisely one of the reduced kernels $K^{(i)} \equiv K^{(i)\nu_i - \mu_i}_{\mu_i}$ as an ingoing (creation) variable, and another one as an outgoing (annihilation) variable. The g -integrals over products of reduced kernels with the measure $dg |\varphi(gp)|^2$ can be estimated, by repeated use of the Cauchy-Schwarz inequality, by the product of the corresponding finite L^2 -norms of the reduced kernels. The latter are functions of the involved p^2 only, and can in turn be estimated by the supremum over the mass interval $[m_0, m_1]$, i.e., the norms given by Eq. (12). After these crude estimates, which are common to all terms in the sum, the remaining p -integrals of the form $\int (\prod dp \theta(p^0)) \prod_i \hat{f}_i(P_i[p])$ still converge absolutely due to the decay of the test functions [1]. Summing all these integrals yields the finite constants $C_{(\nu_i)}$. □

4. Reduction to Generalized Free Fields

Let us now study some elementary cases, starting with fields of order 1 which we denote by φ_F with $F_1(k) = F(k)$ a hermitian polynomially bounded smooth function. These comprise the fields in [1] when all “coupling constants” vanish, or when $S = 0$, hence every band graph has only its external vertex of degree 1 and $F_1(gp) = 1$. One finds that

$$\langle \Omega, \varphi_{F'}(f') \varphi_F(f) \Omega \rangle = \int dp \theta(p^0) \left(\int dg |\varphi(gp)|^2 \overline{F'(gp)} F(gp) \right) \cdot \hat{f}'(p) \hat{f}(-p), \tag{13}$$

while all truncated higher n -point functions for fields of order 1 vanish. Thus, φ_F are generalized free fields with mass distribution ρ_F supported in the mass interval $m_0^2 \leq p^2 \leq m_1^2$,

$$\rho_F(p^2) = \int dg |\varphi(gp)|^2 |F(gp)|^2 . \tag{14}$$

The generalized free fields for different functions F are in general not independent, i.e., their correlations (13) do not vanish. There are in fact only countably many independent such fields. To see this, it is convenient to view a function $F(k)$ in the two-sheeted region $M = \{k \in \mathbb{R}^{d,1} : m_0^2 \leq k^2 \leq m_1^2\}$ as a family (labelled by the mass) of pairs of functions on velocity space $F_{\pm}(m; v) := F(k)$, where $k = \pm g_v p_m$ with $p_m = (m, \mathbf{0})$ a momentum vector in its rest frame and $g_v \in L^1$ the Lorentz boost by the velocity v . At each mass $p^2 = m^2$, the value of the mass distribution in the mixed two-point function (13) is a scalar product in the space $L^2(\{v \in \mathbb{R}^d : v^2 < 1\}; d\eta_m) \otimes \mathbb{C}^2$,

$$\int dg |\varphi(gp)|^2 \overline{F'(gp)} F(gp) = \frac{1}{2} \sum_{\varepsilon=+,-} \int d\eta_m(v) \overline{F'_\varepsilon(m; v)} F_\varepsilon(m; v) \tag{15}$$

with the measure $d\eta_m(v) = d^d v (1 - v^2)^{-(d+1)/2} |\varphi(g_v p_m)|^2$. The real linear space of hermitian functions corresponds to the +1 eigenspaces of the real linear symmetric involutive operator $(F_+, F_-) \mapsto (\overline{F_-}, \overline{F_+})$. These eigenspaces possess countable orthogonal real bases of the form $(F_+^\alpha(m; v), F_-^\alpha(m; v))$ with F_\pm^α smooth and polynomially bounded in $u \equiv v/\sqrt{1 - v^2}$ and $F_-^\alpha = \overline{F_+^\alpha}$. Since the measure $d\eta_m$ varies smoothly with m , the family of bases can be chosen to vary also smoothly with m . The real span of the functions $(k^2)^n F^\alpha(k) := m^{2n} F_\pm^\alpha(m; v)$ at $k = \pm g_v p_m \in M$ is dense (in the topology (12)) in the space of hermitian polynomially bounded smooth functions on M . Since $\varphi_{k^2 F}(f) = \varphi_F(-\square f)$, it follows from Eq. (11) that the countable family of independent generalized free fields $\varphi^\alpha = \varphi_{F^\alpha}$ has a Wightman domain which is dense in the Hilbert space generated from the vacuum by all order 1 fields φ_F .

The next case is an order ν field with F_ν a constant function. One finds that

$$\phi_F(f) = \frac{F_\nu}{\nu!} : \varphi_1^\nu : (f) \tag{16}$$

is just a Wick power of the generalized free field $\varphi_{F=1}$ of order 1. Similarly, if F_ν is a symmetrized tensor product of ν single variable hermitian functions

$$F_\nu(k_1, \dots, k_\nu) = \sum_{\pi \in S_\nu} \prod_{i=1}^\nu F^{(i)}(k_{\pi(i)}) , \tag{17}$$

then the associated field ϕ_F of order ν is the Wick product

$$\phi_F(f) = : \prod_{i=1}^\nu \varphi_i : (f) \tag{18}$$

of the generalized free fields $\varphi_i = \varphi_{F^{(i)}}$ of order 1. It is not very difficult to prove Eqs. (16) and (18) by verifying that the combinatorics of the integral kernels produces precisely the products of two-point functions required by Wick ordering. Namely, every factor in Eq. (17) arising in a creation kernel will be eventually

integrated with another such factor in an annihilation kernel, yielding a two-point function with mass distribution of the form (15), while the numerical coefficients in Eq. (6) cancel against the combinatorial factors due to symmetrization.

Now, every symmetric hermitian function F_ν in ν variables can be approximated (in the topology (12)) by real linear combinations of symmetrized tensor products of ν hermitian functions in one variable. We conclude:

Corollary. *The fields ϕ_F of order 1 form a countable system of generalized free fields φ . The fields ϕ_F of order ν are approximated (in the sense of the lemma of Sect. 3) by homogeneous real Wick polynomials $:P(\varphi):$ of degree ν in the latter. The vacuum vector is cyclic in the Hilbert space $\mathcal{H} := \overline{\mathcal{D}_0} \subset \mathcal{F}_+(H)$ with respect to the fields of order 1, which consequently act irreducibly in \mathcal{H} .*

5. Conclusion and Discussion

We can now apply the classical results in [5] to conclude that the new fields ϕ_F , and in particular the fields constructed in [1] which are finite sums of fields ϕ_F of order ν , $\nu \leq S(R-2)+1$, belong to the Borchers class of the countable system of generalized free fields defined by their two-point functions (13). (As a reminder: the Borchers class of an irreducible (multi-component) field ϕ_0 consists of all fields on the same Hilbert space which are relatively local with respect to ϕ_0 , and which are therefore automatically relatively local with respect to each other.) For fields with a sharp mass (such that the scattering matrix is defined), coincidence of the Borchers class implies coincidence of the scattering matrix [5].

Let us therefore insert a remark concerning a limit of sharp mass $m_1 \searrow m_0$ which is desirable for a particle interpretation. Note, however, that scattering aspects of fields without a sharp mass were also considered in, e.g., [6]. In the original class of models [1], the naive limit attained by sharpening the bare mass distribution $\rho(k^2)$ is severely obstructed since due to momentum conservation for the flow within a graph, there will always occur powers of several measure factors $|\varphi(q)|^2$ at the same argument in the integrands (e.g., in two-point functions the domestic momenta associated with the edges connected to the external vertices coincide due to momentum conservation but are independently integrated; the problem will be aggravated whenever the coupling $a_2 \neq 0$). In order to keep the highest of such powers regular in the sharp mass limit, all the lower powers must become suppressed, so the limiting Wightman functional will consist of products of two-point functions only and one ends up with a free field. Apart from this obstruction, every contribution from the cubic coupling a_3 will die out exactly as soon as $m_1 < 2m_0$, due to momentum conservation at the triple vertex.

On the other hand, no such obstruction prevents us in the enlarged class of models ϕ_F from sharpening the mass distribution $\rho(k^2)$ independently from the cutoff function Θ and the reduced kernels F_ν . The singularity obstruction is absent since there are no domestic integrations, and the latter effect is absent since there is no momentum conservation at the integration kernels. However, as

we have seen, in the limit $\rho(k^2) \rightarrow \delta(k^2 - m^2)$, the system of generalized free fields becomes a countable family of independent Klein-Gordon fields φ_m^α , and the limiting fields ϕ_F will be Wick polynomials therein.

To summarize the previous remarks, we have found that although the sharp mass limit is much more flexible within the enlarged class of models, the limiting fields will belong to the Borchers class of a countable family of massive free fields, and hence will not describe scattering [5]. Even if it remains to be clarified in which precise sense the corresponding conclusion can be maintained for generalized free fields [6], in view of the results of Sect. 4 we do not share the optimism expressed in ref. [1] that the new fields might describe interaction as long as the mass remains smeared. However, the approach of ref. [1], and in particular the surprising mechanism which restores locality upon integration over the "inner degrees of freedom" associated with the Lorentz group, might well contribute some new and interesting stimulations to constructive quantum field theory.

Acknowledgement. The author is very much indebted to C. Read, who circulated his work prior to publication, and to D. Buchholz and J. Yngvason for valuable comments and pointing out some of the classical literature.

References

- 1 Read, C. J.: Quantum field theories in all dimensions. Univ. Cambridge (UK) preprint (1994), to appear in *Commun. Math. Phys.*
- 2 Streater, R. F., Wightman, A. S.: *PCT, Spin and Statistics, and All That*. New York-Amsterdam: W. A. Benjamin, 1964
For a profound guide to more recent achievements of the theory see also:
Streater, R. F.: *Outline of axiomatic relativistic quantum field theory*. *Rep. Progr. Phys.* **38**, 771–846 (1975), especially Chapter 3
- 3 Greenberg, O. W.: Generalized free fields and models of local field theory. *Ann. Phys. (N.Y.)* **16**, 158–176 (1961)
- 4 Borchers, H.-J.: Algebraic aspects of Wightman field theory. In: R. N. Sen and C. Weil (eds.), *Statistical Mechanics and Field Theory, Haifa Lectures 1971*; New York: Halsted Press, 1972
- 5 Borchers, H.-J.: Über die Mannigfaltigkeit der interpolierenden Felder zu einer kausalen S -Matrix. *Nuovo Cim.* **15**, 784–794 (1960)
- 6 Licht, A. L.: A generalized asymptotic condition. *Ann. Phys. (N.Y.)* **34**, 161–186 (1965)

Communicated by D. Brydges

This article was processed by the author
using the Springer-Verlag \TeX QPMZGHB macro package 1991.

