

TOPOLOGICAL  $K$ -THEORY OF THE INTEGERS AT THE PRIME 2

LUKE HODGKIN

*(communicated by James Stasheff)**Abstract*

Recent results of Voevodsky and others have effectively led to the proof of the Lichtenbaum-Quillen conjectures at the prime 2, and consequently made it possible to determine the 2-homotopy type of the  $K$ -theory spectra for various number rings. The basic case is that of  $BGL(\mathbb{Z})$ ; in this note we use these results to determine the 2-local (topological)  $K$ -theory of the space  $BGL(\mathbb{Z})$ , which can be described as a completed tensor product of two quite simple components; one corresponds to a real ‘image of  $J$ ’ space, the other to  $BBSO$ .

**1. Introduction**

As a result of Voevodsky’s solution of the Milnor conjecture [V] and related work by Bloch, Lichtenbaum, Voevodsky and Suslin [B-L], [S-V], Weibel in [W] calculated the algebraic  $K$ -theory of the integers  $\pi_i(BGL(\mathbb{Z})^+)$  at the prime 2 in terms which essentially confirmed the appropriate version of the Lichtenbaum-Quillen conjectures [L,D-F]. (Much stronger and more general versions of the prime-2 conjectures have since been proved, see in particular [RWK], [R-W].) This result, since it expresses the space  $BGL(\mathbb{Z})^+$  in terms of rather well-known spaces, makes it relatively easy to deduce other invariants. Arlettaz et al. in [A-M-N-Y] have done this for the mod 2 cohomology; in this paper, I shall do the same for the (topological) 2-local  $K$ -theory; the result, which is, perhaps predictably, quite different from that for cohomology is stated in theorem 4.1 and corollary 5.1 below. The use of 2-local, rather than the more usual 2-complete theory requires a little more work, but perhaps can be considered as giving a more interesting result.

While Weibel’s results are more general in character, and could lead to similar calculations for various other rings e.g.  $\mathbb{Z}[\sqrt{-1}]$ , I shall here confine my attention to the integers, partly because of their ‘historical’ interest, and partly because of the link with the stable mapping class group  $B\Gamma = \lim_{\rightarrow} B\Gamma_n$  via the composite  $B\Gamma \rightarrow BSp(\mathbb{Z}) \rightarrow BGL(\mathbb{Z})$ , which arises from the action of surface homeomorphisms on  $H_1$ .

Without attempting a complete survey of recent related work, I should draw attention to the most important:

(i) The corresponding decomposition of spectra for the  $p$ -adics  $BGL(\mathbb{Z}_p^\wedge)$  has been known for some time — for arbitrary  $p$  — through work of Bökstedt, Madsen and Rognes [B-M], [R2]. (Here the Milnor conjecture is not needed.)

(ii) Dwyer and Mitchell, in a sequence of papers, [D-M], [Mi1], [Mi2], have attacked precisely the problem of finding the  $K$ -theory of the spectrum associated with  $BGL(R)_p^\wedge$  when  $R$  is a ring of algebraic integers, which they have (essentially) solved in terms of the ‘Iwasawa module’  $M_\infty$  of  $R$ . The remaining difficulties are those associated with the structure of  $M_\infty$ , and are not trivial.

---

Received 26 May 2000, revised 19 October 2000; published on 13 November 2000.

2000 Mathematics Subject Classification: 19D99, 55N15, 55P15.

Key words and phrases:  $K$ -theory, general linear group of integers, Rothenberg-Steenrod spectral sequence, Bousfield localization.

© 2000, Luke Hodgkin. Permission to copy for private use granted.

(iii) In an important special case, Østvær [Ø] has found the homotopy type of  $BGL(R)_2^\wedge$  when  $R$  is ‘2-regular and non-exceptional’. Examples include cyclotomic rings  $R = \mathbb{Z}[\zeta_{2^r}]$ ,  $r \geq 2$ , (in particular the Gaussian integers); but no rings of integers in totally real fields. The homotopy type is very simple, a product of a  $K(\mathbb{F}_p)$  (finite field) and copies of  $U, SU$ .

Having said this, the result presented here is perhaps principally of interest in the way that it displays the intertwining of the factors which go to make up  $BGL(\mathbb{Z})$ . The justification which Dwyer and Mitchell give for studying the topological  $K$ -theory of algebraic spectra (that it sheds light on the  $K$ -localization of the spectra, and hence can provide evidence about the Lichtenbaum-Quillen conjectures) is not really at issue at the prime 2 where the conjectures are solved; however, the structure has independent interest.

We begin by describing the space which we shall study. Following [Bö], but working in the category of 2-local spaces, we define the ‘étale  $K$ -space’  $JK(\mathbb{Z})$  as the (2-local) homotopy fibre of the composite map:

$$(1) \quad c(\psi^3 - 1) : BO \xrightarrow{\psi^3 - 1} BSpin \xrightarrow{c} BSU$$

(Here  $\psi^3$  is the Adams operation and  $c$  denotes complexification.) This space can be realized through a number of other fibrations, of which we shall note particularly (cf [R2], (2.3))

$$(2) \quad JR_2 \rightarrow JK(\mathbb{Z}) \rightarrow BBSO$$

where  $JR_2$  is the real image of  $J$  space at 2, defined as the fibre of  $\psi^3 - 1 : BO \rightarrow BSpin$ , localized at 2. (See e.g. [Ma].) The 2-completion of  $JK(\mathbb{Z})$  is equivalent to the space which is named  $K^{ét}(\mathbb{Z})$  in [D-F] and elsewhere. Bökstedt defined a map on 2-completions from  $((BGL(\mathbb{Z})^+)_2^\wedge)$  to  $JK(\mathbb{Z})_2^\wedge$ ; it is a consequence of Voevodsky’s theorem and subsequent work that this is a homotopy equivalence. However, it is not obvious that this map is an equivalence — even that it exists — in the localized sense. [I am grateful to the referee for pointing this fact out.] I shall therefore, in sections 2-4, find the  $K$ -theory of  $JK(\mathbb{Z})$  with coefficients in  $\mathbb{Z}_2^\wedge$  and  $\mathbb{Z}_2$ , by a simple application of the Rothenberg-Steenrod spectral sequence. Having done that, in §5 I shall deduce the corresponding results for  $BGL(\mathbb{Z})^+$ ; the completed case is easy, by the above remarks, but the local case requires a special investigation, using Bousfield’s  $K$ -localization functor  $L_K$  [Bou] to identify  $JK(\mathbb{Z})$  with  $L_K(BGL(\mathbb{Z})^+)$ .

Where not otherwise stated, all spaces are supposed localized at 2 in what follows.

## 2. The 2-complete theory

The natural procedure is in any case to begin with the 2-completed theory, and proceed to integrate it with the rational to obtain a 2-local statement. With this in mind, we begin with the following commutative diagram:

$$(3) \quad \begin{array}{ccccc} O & \xrightarrow{c(\psi^3 - 1)} & SU & \xrightarrow{\eta} & JK(\mathbb{Z}) \\ \parallel & & \uparrow c & & \uparrow j \\ O & \xrightarrow{\psi^3 - 1} & Spin & \xrightarrow{\theta} & JR_2 \end{array}$$

Both rows in this diagram are fibrations; the top row derived in the obvious way from the fibration (1), the second similarly from the definition of  $JR_2$ . The two squares are commutative by construction (compare the diagram on p.8 of [R2]); and the right hand square is fibred. The most ‘natural’ approach for such  $K$ -theory computations is usually via the geometric spectral sequence of Rothenberg-Steenrod (see e.g. [A-H]), which gives the  $K$ -theory of a quotient of groups (for example) in terms of those of the group and subgroup; and we can apply this spectral sequence to the fibre square provided that one of the  $Spin$ -actions is free. It will

be convenient to suppose this for the action on  $SU$  (of course nearly free, but not quite...), by the usual device of replacing  $SU$  by  $SU \times ESpin$ . We are accordingly using a homotopy equivalence of  $JK(\mathbb{Z})$  with  $(SU \times ESpin) \times_{Spin} JR_2$ ; and we need the cohomology version of the Rothenberg-Steenrod sequence, which uses derived functors in the category of comodules over a coalgebra — in our case the coalgebra  $K^*(Spin; \mathbb{Z}_2^\wedge)$ .

Much of the argument can be simplified in this case, as we shall see, since we are dealing with a trivial comodule. To clarify the details of the application we want, i.e. to the 2-complete  $K$ -cohomology, one or two technical points should be made. First, there is (as usual in  $K$ -cohomology of large spaces) the question of topology on the coalgebra and comodules  $K^*(\ ; \mathbb{Z}_2^\wedge)$ . Second, the coefficients are not a field, and the modules may not be free or even projective. We need to deal with these objections together so as to obtain a reasonable cotensor product functor. To begin with, the category  $\mathcal{C}_2$  of *profinite* modules over  $\mathbb{Z}_2^\wedge$  is abelian, and the appropriate tensor product is the completed one,  $\hat{\otimes}$ . Because  $\varprojlim$  is exact in the category, the functor  $-\hat{\otimes}A$  is exact if  $A$  is an inverse limit of finitely generated free  $\mathbb{Z}_2^\wedge$ -modules. We shall call such a module ‘flat’, by analogy with the usual case. In particular, this applies to the Hopf algebras  $K^*(Spin; \mathbb{Z}_2^\wedge)$  and  $K^*(SU; \mathbb{Z}_2^\wedge)$ . (This is a consequence of [H], but the detail will be given later.) Hence  $K^*(\ ; \mathbb{Z}_2^\wedge)$  translates products of spaces into completed tensor products of  $\mathbb{Z}_2^\wedge$ -algebras, when one of the spaces is  $SU$  or  $Spin$ .

If  $A$  is a (profinite) flat cocommutative  $\mathbb{Z}_2^\wedge$ -coalgebra, and  $B, C$  are compact comodules over  $A$ , we define the *completed* cotensor product  $B \square_A C$  to make the sequence (cf [M-M])

$$0 \longrightarrow B \hat{\square}_A C \longrightarrow B \hat{\otimes} C \xrightarrow{\Delta \otimes 1 - 1 \otimes \Delta} B \hat{\otimes} A \hat{\otimes} C$$

exact. ( $\Delta$  denotes the structural morphisms for the comodules.) This bifunctor is left exact on sequences of  $A$ -comodules which are split-exact over  $\mathbb{Z}_2^\wedge$ . Its derived functors will be written  $\widehat{Cotor}_p^A(B, C)$ . Recall the spectral sequence — stated here in the appropriate form for our purpose.

**Proposition 2.1.** *Let  $G$  be a group, and let  $X, Y$  be  $G$ -spaces with either  $X$  or  $Y$  free (all in a suitably small category, e.g. 2-local CW-complexes). If  $G, X$  have  $K^*(\ ; \mathbb{Z}_2^\wedge)$  flat in  $\mathcal{C}_2$ , then there is a strongly convergent spectral sequence with*

$$E_2^p = \widehat{Cotor}_{K^*(G; \mathbb{Z}_2^\wedge)}^p(K^*(X; \mathbb{Z}_2^\wedge), K^*(Y; \mathbb{Z}_2^\wedge))$$

$$E_\infty \sim K^*(X \times_G Y; \mathbb{Z}_2^\wedge)$$

Its edge homomorphism is the ‘standard’ map

$$\eta : K^*(X \times_G Y; \mathbb{Z}_2^\wedge) \rightarrow K^*(X; \mathbb{Z}_2^\wedge) \hat{\square}_{K^*(G; \mathbb{Z}_2^\wedge)} K^*(Y; \mathbb{Z}_2^\wedge)$$

(which follows from the definitions).

The proof is the usual geometric one, using the bar resolution. Again because the inverse limit is exact in  $\mathcal{C}_2$  there are no convergence problems.

**Note.** Since we are interested in the 2-local theory, we shall also need a local version of this. Here arguments using profiniteness naturally break down, and alternative methods must be used. The best option is to use the corresponding sequence for  $K$ -homology, which involves the ordinary tensor product and the ordinary  $Tor$  groups over  $K_*(G; \mathbb{Z}_2)$ . Again (since homology theories behave well with respect to direct limits) the sequence is strongly convergent; in this case the proof is clearly simpler. We then need to dualize the results in the appropriate way to derive the  $K$ -cohomology.

We are now ready to state the structure theorem for the map  $c : Spin \rightarrow SU$ ; for maximum generality we shall need the local version.

**Proposition 2.2.** (i) *The Hopf algebras  $K^*(Spin; \mathbb{Z}_2)$  resp.  $K^*(SU; \mathbb{Z}_2)$  are completed exterior algebras on submodules of primitive generators, say  $P_R, P_C$  respectively; and the map*

$c$  induces an epimorphism from  $P_C$  to  $P_R$ , whose kernel  $Q$  is a direct summand. Accordingly, writing  $\hat{E}(\ )$  for the completed exterior algebra on primitive elements, we have:

$$K^*(Spin; \mathbb{Z}_2) = \hat{E}(P_R)$$

$$K^*(SU; \mathbb{Z}_2) = \hat{E}(P_C) \cong \hat{E}(P_R \oplus Q) = \hat{E}(P_R) \hat{\otimes} E(Q)$$

as a tensor product of Hopf algebras.

(ii) The same statements hold for  $K$ -theory with  $\mathbb{Z}_2^\wedge$  coefficients, and  $K^*(Spin; \mathbb{Z}_2^\wedge)$  resp.  $K^*(SU; \mathbb{Z}_2^\wedge)$  is isomorphic to  $K^*(Spin; \mathbb{Z}_2) \otimes_{\mathbb{Z}_2} \mathbb{Z}_2^\wedge$  resp.  $K^*(SU; \mathbb{Z}_2) \otimes_{\mathbb{Z}_2} \mathbb{Z}_2^\wedge$

From this will follow:

**Proposition 2.3.** For any space  $X$  with an action of  $Spin$ , the edge homomorphism of the spectral sequence defines a natural isomorphism

$$\eta : K^*((SU \times ESpin) \times_{Spin} X; \mathbb{Z}_2^\wedge) \rightarrow \hat{E}(Q) \hat{\otimes} K^*(X; \mathbb{Z}_2^\wedge)$$

We postpone the proof of proposition 2.2 to the next section, and show that it implies proposition 2.3.

For this, it is sufficient to identify  $K^*(SU; \mathbb{Z}_2^\wedge) \hat{\square}_{K^*(Spin; \mathbb{Z}_2^\wedge)} K^*(X; \mathbb{Z}_2^\wedge)$ . From the splitting of proposition 2.2, we can deduce that

$$\mu^* : K^*(SU; \mathbb{Z}_2^\wedge) \rightarrow K^*(Spin; \mathbb{Z}_2^\wedge) \hat{\otimes} K^*(SU; \mathbb{Z}_2^\wedge)$$

is identified with

$$\hat{E}(P_R) \hat{\otimes} E(Q) \xrightarrow{\Delta \otimes 1} \hat{E}(P_R) \hat{\otimes} \hat{E}(P_R) \hat{\otimes} \hat{E}(Q)$$

Now we know that  $(\hat{E}(P_R) \hat{\otimes} \hat{E}(Q)) \hat{\square}_{\hat{E}(P_R)} K^*(X; \mathbb{Z}_2^\wedge) \cong \hat{E}(Q) \hat{\otimes} K^*(X; \mathbb{Z}_2^\wedge)$ ; in fact, this is the dual of the well-known analogous formula for the tensor product, and the isomorphism is natural. However,  $\hat{E}(Q) \hat{\otimes} K^*(X; \mathbb{Z}_2^\wedge)$  is exact as a functor of the comodule  $K^*(X; \mathbb{Z}_2^\wedge)$ , since  $\hat{E}(Q)$  is flat, and so its derived functors are trivial:

$$\widehat{Cotor}_{K^*(Spin; \mathbb{Z}_2^\wedge)}^0(K^*(X; \mathbb{Z}_2^\wedge), K^*(SU; \mathbb{Z}_2^\wedge)) = K^*(X; \mathbb{Z}_2^\wedge) \hat{\otimes} \hat{E}(Q)$$

$$\widehat{Cotor}_{K^*(Spin; \mathbb{Z}_2^\wedge)}^p(K^*(X; \mathbb{Z}_2^\wedge), K^*(SU; \mathbb{Z}_2^\wedge)) = 0 \quad (p > 0)$$

Using the edge homomorphism of the spectral sequence, proposition 2.3 follows.

### 3. Structure of $K^*(Spin), K^*(SU)$

We now proceed to the proof of proposition 2.2. Let  $\lambda_r^i$  resp.  $\lambda_c^i$  be the  $i$ th ‘stabilized’ exterior power of the standard representation  $\theta$  from  $Spin(2n+1)$  resp.  $SU(2n+1)$  to  $U$ , considered as an element of the representation ring. That is,  $\lambda_r^i$  is the result of applying the operation  $\lambda^i$  to  $\theta - (2n+1)$ . Then it is obvious that under inclusion maps of  $Spin(2n+1)$ ’s and  $SU(2n+1)$ ’s the  $\lambda^i$ ’s are preserved; and that  $c^*(\lambda_c^i) = c^*(\bar{\lambda}_c^i) = \lambda_r^i$ .

Let now  $\beta$  be the operation (see [H]) which to any representation  $\rho$  of  $G$  assigns its class  $\beta(\rho)$  in  $K^1(G) = [G, U]$  considered as a map from  $G$  to  $U$ . The basic theorem of [H] gives us that  $K^*(SU(2n+1); \mathbb{Z}_2)$  is the exterior algebra

$$E_{\mathbb{Z}_2}(\beta(\lambda_c^1), \dots, \beta(\lambda_c^n), \beta(\bar{\lambda}_c^1), \dots, \beta(\bar{\lambda}_c^n))$$

since these can be seen to be equivalent to the basic representations modulo a little manipulation. (The generators are also, as usual, the primitives for the Hopf algebra structure.) The similar result is not quite true for  $Spin(2n+1)$ , as is well known, the picture being complicated by the Spin representation  $\Delta_n$ , of dimension  $2^n$ . We have:

$$K^*(Spin(2n+1); \mathbb{Z}_2) = E_{\mathbb{Z}_2}(\beta(\lambda_r^1), \dots, \beta(\lambda_r^{n-1}), \beta(\Delta_n))$$

However, there is a relation between  $\lambda_r^n$  and  $\Delta_n$ , since  $(\Delta_n)^2 = \lambda_r^n +$  a sum of terms in  $\lambda_r^1, \dots, \lambda_r^{n-1}$ . Writing  $\Delta_n = 2^n + \tilde{\Delta}_n$ , and applying the usual relations for  $\beta$ , we have that  $\beta(\Delta_n)^2 = 2^{n+1}\beta(\Delta_n)$ . Hence,  $\beta(\lambda_r^n) = 2^{n+1}\beta(\Delta_n) \pmod{\beta(\lambda_r^1), \dots, \beta(\lambda_r^{n-1})}$ .

Write  $M_n$  for the  $\mathbb{Z}_2$ -module which generates the exterior algebra  $K^*(Spin(2n+1); \mathbb{Z}_2)$  and  $N_n$  for the submodule generated by the  $\beta(\lambda_r^i)$ 's. We can deduce a short exact sequence

$$(E_n) \quad 0 \rightarrow N_n \rightarrow M_n \rightarrow \mathbb{Z}/2^{n+1} \cdot \beta(\Delta_n) \rightarrow 0$$

The restrictions from  $E_{n+1}$  to  $E_n$  are straightforward if we take into account that  $\Delta_{n+1}$  restricts to  $2\Delta_n$ . Hence the map from  $\mathbb{Z}/2^{n+2}$  to  $\mathbb{Z}/2^{n+1}$  in the above sequence multiplies the generator by 2. It is easy to deduce that the inverse limit of the  $\mathbb{Z}/2^{n+1}$ 's is zero; and so (since they are finite) is the  $\varprojlim^1$ . Hence the map from  $\varprojlim\{N_n\}$  to  $\varprojlim\{M_n\}$  — the primitives of  $K^*(Spin)$  — is an isomorphism, and we have:

**Proposition 3.1.** *The  $K$ -cohomology rings of  $Spin$ ,  $SU$  are as follows:*

$$\begin{aligned} K^*(Spin; \mathbb{Z}_2) &= \hat{E}_{\mathbb{Z}_2}(\beta(\lambda_r^1), \beta(\lambda_r^2), \dots) \\ K^*(SU; \mathbb{Z}_2) &= \hat{E}_{\mathbb{Z}_2}(\beta(\lambda_c^1), \beta(\lambda_c^2), \dots; \beta(\bar{\lambda}_c^1), \beta(\bar{\lambda}_c^2), \dots) \end{aligned}$$

and the restriction  $c^*$  from  $SU$  to  $Spin$  maps  $\beta(\lambda_c^i), \beta(\bar{\lambda}_c^i)$  to  $\beta(\lambda_r^i)$  ( $i = 1, 2, \dots$ )

From this, proposition 2.2 clearly follows.

We next deduce:

**Proposition 3.2.** *The local  $K$ -theory of the quotient is given by*

$$K^*((SU \times ESpin)/Spin; \mathbb{Z}_2) \cong \hat{E}_{\mathbb{Z}_2}(\beta(\lambda_c^1) - \beta(\bar{\lambda}_c^1), \dots) \cong \hat{E}_{\mathbb{Z}_2}(Q)$$

in the terminology of proposition 2.2.

**Proof.** As stated above, the best way to prove this is as follows. First, dualize proposition 3.1 to give a result on the local  $K$ -homology (the map  $c$  now induces a split monomorphism). Next, apply the Rothenberg-Steenrod sequence in local  $K$ -homology; this is well-behaved, and strong convergence is easily established, as well as flatness (in the usual sense) for the  $K$ -algebras involved. We find a natural isomorphism in  $K$ -homology in a form dual to that of proposition 2.3. In the special case where  $X$  is a point, this can now simply be dualized back to give the required result.

This procedure is of course roundabout, but seems preferable to developing a theory of topological modules which will deal properly with very large algebras over  $\mathbb{Z}_2$  of the kind we are considering here in  $K$ -cohomology.

#### 4. The $K$ -theory of $JK(\mathbb{Z})$

We are now in a position to put the pieces together. The key point is that  $JR_2$  is a 2-adic space, so the local theory and the 2-adic theory coincide for it.

**Theorem 4.1.** *There is a natural isomorphism:*

$$\begin{aligned} K^*(JK(\mathbb{Z}); \mathbb{Z}_2) &\cong \hat{E}_{\mathbb{Z}_2}(Q) \hat{\otimes} K^*(JR_2; \mathbb{Z}_2) \\ &\cong K^*(BBSO; \mathbb{Z}_2) \hat{\otimes} K^*(JR_2; \mathbb{Z}_2) \end{aligned}$$

with an analogous isomorphism for  $\mathbb{Z}_2^\wedge$  coefficients.

**Proof.** We'd like to use a basepoint in  $JR_2$ , but of course can't suppose there is one which is fixed under  $Spin$ . Consider instead the equivariant embedding of  $JR_2$  in the unreduced cone  $C^+JR_2$ . If we can prove the result for  $C^+JR_2$  and for the pair  $P = (C^+JR_2, JR_2)$

separately, then it will follow for  $JR_2$  by the 5-lemma. Now for  $C^+JR_2$  it is already proved (by proposition 3.2). For  $P$ , we consider the commutative diagram:

$$\begin{array}{ccc}
 K^*(SU \times_{Spin} P; \mathbb{Z}_2) & \xrightarrow{\eta} & K^*(SU; \mathbb{Z}_2) \hat{\square}_{K^*(Spin; \mathbb{Z}_2)} K^*(P; \mathbb{Z}_2) \\
 \downarrow \alpha & & \downarrow \beta \\
 K^*(SU \times_{Spin} (P); \mathbb{Z}_2^\wedge) & \xrightarrow{\eta} & K^*(SU; \mathbb{Z}_2^\wedge) \hat{\square}_{K^*(Spin; \mathbb{Z}_2^\wedge)} K^*(P; \mathbb{Z}_2^\wedge)
 \end{array}$$

The arrow  $\eta$  in the lower row is an isomorphism by proposition 2.3. The two vertical arrows are induced by the coefficient homomorphism. Since the reduced homology of  $JR_2$  is finite in every dimension, the same is true for the pair  $SU \times_{Spin} P$ ; so  $K^*(SU \times_{Spin} P; \mathbb{Z}_2)$  is a 2-adic group. Hence the arrow marked  $\alpha$  is an isomorphism. On the other hand, we can embed  $K^*(SU; \mathbb{Z}_2) \hat{\square}_{K^*(Spin; \mathbb{Z}_2)} K^*(P; \mathbb{Z}_2)$  in  $K^*(SU; \mathbb{Z}_2) \hat{\otimes} K^*(P; \mathbb{Z}_2)$  and identify the latter with

$$K^*(SU; \mathbb{Z}_2) \hat{\otimes} (\mathbb{Z}_2^\wedge \hat{\otimes} K^*(P; \mathbb{Z}_2))$$

(again because  $K^*(P; \mathbb{Z}_2)$  is 2-adic). Using this, and the definition of the cotensor product, we find that the right hand vertical arrow  $\beta$  is also an isomorphism. Hence the upper arrow  $\eta$  is one.

Now by the argument used in proposition 2.3, this implies that  $K^*(SU \times_{Spin} P; \mathbb{Z}_2)$  is isomorphic to  $\hat{E}_{\mathbb{Z}_2}(Q) \hat{\otimes} K^*(P; \mathbb{Z}_2)$ . This proves the first line of the theorem. The second results from

**Lemma 4.1.** *The fibre  $SU/Spin$  of  $c : BSpin \rightarrow BSU$  can be identified with the Hopf map  $\eta : BBSO \rightarrow BSpin$ .*

**Proof.** The composite  $c \circ \eta$  is trivial and so lifts to a map  $\tilde{\eta} : BBSO \rightarrow Fib(c)$ . A check on the homotopy sequence shows that this is a homotopy equivalence.  $\square$

A comparison with the fibre sequence (2) shows that the sequence splits from the viewpoint of 2-local  $K$ -theory. Finally, it is worth noting that the  $K$ -theory of  $JR_2$  has been known for a long time, see [H-S]; it is essentially the completed representation ring of the infinite symmetric group  $\Sigma_\infty$ .

### 5. The results for $BGL(\mathbb{Z})^+$

As was remarked in §1, theorem 4.1 immediately gives us the 2-adic  $K$ -theory of  $BGL(\mathbb{Z})^+$ , since  $(BGL(\mathbb{Z})^+)^\wedge$  is homotopy equivalent to  $JK(\mathbb{Z})^\wedge$ . To deal with the localizations, we shall prove the following result:

**Theorem 5.1.** *Let  $L_K$  denote Bousfield's  $K$ -theory localization functor on spaces [Bou]. There is a homotopy equivalence of  $JK(\mathbb{Z})$  with  $L_K(BGL(\mathbb{Z})^+)$ .*

Since  $K^*(L_K(X; \mathbb{Z}_2)) \cong K^*(X; \mathbb{Z}_2)$  for any  $X$ , this shows:

**Corollary 5.1.** *The  $K$ -theory of  $BGL(\mathbb{Z})^+$  with  $\mathbb{Z}_2$  or  $\mathbb{Z}_2^\wedge$  coefficients is computed by theorem 4.1.*

For the proof of theorem 5.1, we shall simplify notation by writing  $K(\mathbb{Z})$  for the ring  $\mathbb{Z} \times BGL(\mathbb{Z})^+$ . We follow the arguments of [Bö] and (§2 of) [R1]. In [Bö], the rational component is considered as well as the 2-adic, but the discussion is essentially concerned with  $\Omega K(\mathbb{Z})$ ; while in [R1], the argument is at the level of spectra, but is purely 2-adic. Our concern is to use  $K$ -localization to circumvent these restrictions.

We first define a map  $s$  as the composite  $BSO \xrightarrow{i} BSG \xrightarrow{\eta} SG$ , where  $i$  is the 'forgetful' map from bundles to spherical fibrations, and  $\eta$  is multiplication by the Hopf map. Both maps

are 2-locally defined, and are infinite loop maps. If  $r : SG \rightarrow K(\mathbb{Z})_1$  is the map into the ‘one-component’ induced by  $QS^0 \rightarrow K(\mathbb{Z})$ , then  $r$  is again a local infinite loop map. In [R1] it is shown that the 2-completion of  $r \circ s$  is nullhomotopic at the level of spectra. However, since  $SG$  has finite homotopy groups,  $r \circ s$  is 2-locally trivial if it is trivial after completion; and as in [R1], the nullhomotopy deloops to give a map  $g : Bfib(s) \rightarrow K(\mathbb{Z})_1$ .

**Lemma 5.1.** *The localizations  $L_K(Bfib(s))$  and  $L_K(JK(\mathbb{Z}))$  are homotopy equivalent.*

**Proof.** This follows directly from diagram 2.8 of [R1], since the fibre sequence  $C_\infty \rightarrow Bfib(s) \rightarrow JK(\mathbb{Z})$  can be constructed 2-locally using  $\rho^3$ . Since  $L_K(C_\infty)$  is a point by [H-S], the fibration becomes an equivalence after  $K$ -localization.  $\square$

Finally, we need:

**Lemma 5.2.** *The  $K$ -theory localization of  $g$ ,  $L_K(g)$ , is a homotopy equivalence from  $L_K(Bfib(s))$  to  $L_K(K(\mathbb{Z})_1)$ .*

**Proof.** Since the homotopy groups in each case are finitely generated, it will be enough to show that  $g$  induces isomorphisms on homotopy (a) when 2-completed and (b) when tensored with  $\mathbb{Q}$ . For the 2-completion, we again use Rognes’ diagram 2.8. The preceding lemma implies that the map there called ‘ $h$ ’ exists after  $K$ -localization; we shall call it ‘ $L_K(h)$ ’, ignoring the question of whether  $h$  exists. And  $L_K(g)$  is an equivalence if and only if  $L_K(h)$  is. But by the subsequent arguments of Rognes,  $L_K(h)$  is a right inverse for the more usual map  $\Phi : L_K(K(\mathbb{Z})_1)_2^\wedge \rightarrow L_K JK(\mathbb{Z})_2^\wedge$ . (As pointed out in [R1], this involves an essentially 2-adic argument.) We now know, however, that  $\Phi$  is a 2-adic equivalence, and so  $L_K(h)$  is. For the rational version we use the argument on pp.31-2 of [Bö]; the homotopy groups of  $fib(s)_2^\wedge$  and  $(\Omega K(\mathbb{Z})_1)_2^\wedge$  are the same after inverting 2, and hence  $\Omega g$  is an isomorphism on rational homotopy. The same result therefore follows for  $g$  and so for  $L_K(g)$ .  $\square$

## References

- [A-H] Anderson, D.M. and Hodgkin, L.H., *On the  $K$ -theory of Eilenberg-MacLane complexes*, Topology **7** (1968), 317-329.
- [A-M-N-Y] Arlettaz, D., Mimura, M., Nakahata, K. and Yagita, K., *The mod 2 cohomology of the linear groups over the ring of integers* Proc. Amer. Math. Soc. **127** (1999), 2199-2212.
- [B-L] Bloch, S. and Lichtenbaum, S., *A spectral sequence for motivic cohomology*,  $K$ -theory preprint archive no. 62 (1994).
- [Bö] Bökstedt, M., *The rational homotopy type of  $\Omega Wh^{Diff}(*),$*  in Algebraic Topology, Aarhus 1982 (Lect. Notes Math. vol. 1051), Berlin Heidelberg New York, Springer, 1984, 25-37.
- [B-M] Bökstedt, M. and Madsen, I., *Topological cyclic homology of the integers* Astérisque **226** (1994), 57-143.
- [Bou] Bousfield, A.K., *The localization of spectra with respect to homology* Topology **18** (1979), 257-281.
- [D-F] Dwyer, W. and Friedlander, E., *Conjectural calculations of general linear group cohomology*, Applications of Algebraic  $K$ -theory to Algebraic Geometry and Number Theory, Boulder 1983 (Contemp. Math. vol. 55 part I), AMS Providence, 1986, 135-147.
- [D-M] Dwyer, W. and Mitchell, S.A., *On the  $K$ -theory spectrum of a ring of algebraic integers*  $K$ -theory **14** (1998), 201-263.
- [H] Hodgkin, L.H.,  *$K$ -theory of Lie groups* Topology **6** (1967), 1-36.

- [H-S] Hodgkin, L.H. and Snaith, V.P. , *The K-theory of some more well-known spaces* Illinois J. Math. **22** (1978), 270-278 .
- [L] Lichtenbaum, S. , *Values of zeta functions, étale cohomology, and algebraic K-theory*, Classical Algebraic K-theory and Connections with Arithmetic. (Lect. Notes in Math. vol. 342), Berlin Heidelberg New York, Springer, 1973, 489-501.
- [Ma] May, J.P.  *$E_\infty$  Ring Spaces and  $E_\infty$  Ring Spectra* (Lect. Notes in Math. vol. 577), Berlin Heidelberg New York, Springer, 1977 .
- [M-M] Milnor, J.W. and Moore, J.C. , *On the Structure of Hopf algebras* Ann. of Math. **81** (1965), 211-264 .
- [Mi1] Mitchell, S.A. , *Topological K-theory of algebraic K-theory spectra*, K-theory preprint archive no.345 (1999) .
- [Mi2] Mitchell, S.A. , *K-theory hypercohomology spectra of number rings at the prime 2*, K-theory preprint archive no. 386 (1999) .
- [Ø] Østvær, P.-A. , *On algebraic K-theory spectra*, preprint .
- [Q] Quillen, D. , *On the cohomology and K-theory of the general linear groups over a finite field* Ann. of Math. **96** (1972), 552-586 .
- [R1] Rognes, J. , *The Hatcher-Waldhausen map is a spectrum map* Math. Ann. **299** (1994), 529-549 .
- [R2] Rognes, J. , *Algebraic K-theory of the two-adic integers* J. Pure Applied Algebra **134** (1999), 287-326 .
- [R-W-K] Rognes, J., Weibel, C., and Kolster, M. , *2-primary algebraic K-theory of rings of integers in number fields* J. Amer. Math. Soc. **13** (2000), 1-54 .
- [R-W] Rognes, J. and Weibel, C. , *Étale descent for 2-primary algebraic K-theory of totally imaginary number fields* K-theory **16** (1999), 101-104 .
- [S] Sullivan, D. , *Genetics of homotopy theory and the Adams conjecture* Ann. of Math. **100** (1974), 1-79 .
- [S-V] Suslin, A. and Voevodsky, V. , *Bloch-Kato conjecture and motivic cohomology with finite coefficients*, K-theory preprint archive no. 83 (1995); see also Friedlander, E., *Motivic complexes of Suslin and Voevodsky*, Astérisque **245** (1997), 355-378 .
- [V] Voevodsky, V. , *The Milnor conjecture*, K-theory preprint archive no. 170 (1996); see also Kahn, B., *The Milnor conjecture after V. Voevodsky*, Astérisque **245** (1997), 379-418 .
- [W] Weibel, C. , *The 2-torsion in the K-theory of the integers*, K-theory preprint archive no.141 (1996).

This article may be accessed via WWW at <http://www.rmi.acnet.ge/hha/> or by anonymous ftp at [ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2000/n9/n9.\(dvi,ps,dvi.gz,ps.gz\)](ftp://ftp.rmi.acnet.ge/pub/hha/volumes/2000/n9/n9.(dvi,ps,dvi.gz,ps.gz))

Luke Hodgkin [luke.hodgkin@kcl.ac.uk](mailto:luke.hodgkin@kcl.ac.uk)

King's College  
London