

DIFFERENTIAL GAMES FOR STOCHASTIC PARTIAL DIFFERENTIAL EQUATIONS

W.H. FLEMING AND M. NISIO

1. Introduction

In this paper we are concerned with zero-sum two-player finite horizon games for stochastic partial differential equations (SPDE in short). The main aim is to formulate the principle of dynamic programming for the upper (or lower) value function and investigate the relationship between upper (or lower) value function and viscosity solution of min-max (or max-min) equation on Hilbert space.

Let us consider SPDE (1.1), a so-called controlled Zakai-equation, arising from zero-sum two-player stochastic differential game with partial information,

$$(1.1) \quad d\xi(x, t) = \left[\sum_{i,j=1}^n \frac{\partial}{\partial x_i} (\sigma^{ij}(x, Y(t), Z(t)) \frac{\partial}{\partial x_i} \xi(x, t) \right. \\ \left. + \sum_{i=1}^n r^i(x, Y(t), Z(t)) \frac{\partial}{\partial x_i} \xi(x, t) - c\xi(x, t) \right] dt \\ + \sum_{j=1}^m f^j(x, Y(t), Z(t)) \xi(x, t) dW_j(t)$$

with initial condition

$$\xi(x, 0) = \eta(x).$$

The pay-off function J is defined by

$$(1.2) \quad J(t, \eta, Y, Z, g) = E \int_0^t h(\xi(s), Y(s), Z(s)) ds + g(\xi(t)), \quad t \leq T,$$

where T is a given positive number and $h(\cdot, y, z)$ and g are functions on Hilbert space. Moreover, $W = (W_1, \dots, W_m)$ is a standard m -dimensional Brownian motion and Y and Z are admissible controls for player I and II respectively, (Definition 2.1).

According to W.H. Fleming and P.E. Souganidis [4], we introduce a strategy (Definition 4.2) and formulate a notion of upper (or lower) value function (see (6.2)). When h and g are tame functions, we can prove the principle of dynamic programming (see (5.40) and Theorem 6.2)). This fact implies that the upper (or lower) value function turns out to be a viscosity solution of the min-max (or max-min) equation corresponding to (1.1) and (1.2).

Here we modify Lions' definition of viscosity solution [8, Part 2], which seems suitable for our problems. If a viscosity solution is unique, then the Isaacs' condition yields the existence of value of game. The finite dimensional case, namely differential game for stochastic differential equations, is investigated in [4] and W.H. Fleming and P.E. Souganidis proved that the upper value function equals the Elliott-Kalton value and turns out to be the unique viscosity solution of the min-max equation. But, in our case, the uniqueness problem is still open and we will consider a simple example and remark on its uniqueness problem in Section 8.

Recently H. Ishii [6] introduced a slightly different notion of viscosity solution and proved the unique existence of solution for nonlinear second order partial differential equations, including the min-max one. But it seems to be still open, whether the upper value function satisfies the min-max equation in Ishii's sense.

In Section 2, we recall some results on SPDE for later use. In particular, the continuity with respect to time will be studied in Section 3. Since the lower value function is investigated in the same way as the upper one, we will only consider the upper one. Applying semi-discretization arguments, we introduce π -admissible strategy (see Definition 4.2) and the upper value function V_π . If the terminal function is tame, then $\{V_\pi, \pi = \text{partition of } [0, T]\}$ is compact (Theorem 4.1) and the principle of dynamic programming will be proved in Section 6. We will show that the upper value function turns out to be a viscosity solution of the min-max equation in Section 7.

2. Preliminaries

Let \mathcal{Y} and \mathcal{Z} be convex compact subsets of R^p and R^q respectively. W denotes a standard m -dimensional Brownian motion, defined on a probability space (Ω, \mathcal{F}, P) . For simplicity, we assume $m = 1$. Put $\mathcal{F}_t = \sigma$ -field generated by $\{W(s), s \leq t\}$.

First we will define control and strategy, according to [4].

DEFINITION 2.1. An admissible control process Y (resp. Z) for player I (resp. II) is an \mathcal{F}_t -progressively measurable process taking values \mathcal{Y} (resp. \mathcal{Z}). The set of all admissible controls for player I (resp. II) is denoted by \mathcal{M} (resp. \mathcal{N}).

We say that Y and Y^* ($\in \mathcal{M}$) are the same on $[t, s]$ and we write $Y = Y^*$ on $[t, s]$, if

$$P(Y = Y^* \text{ a.e. on } [t, s]) = 1.$$

A similar convention is assumed to hold for elements of \mathcal{N} .

DEFINITION 2.2. An admissible strategy α (resp. β) for player I (resp. II) is a mapping $\alpha; \mathcal{N} \rightarrow \mathcal{M}$ (resp. $\beta; \mathcal{M} \rightarrow \mathcal{N}$), such that if $Z = Z^*$ (resp. $Y = Y^*$) on $[0, s]$ then $\alpha(Z) = \alpha(Z^*)$ (resp. $\beta(Y) = \beta(Y^*)$) on $[0, s]$ for any $s \in [0, T]$. The set of all admissible strategies of player I (resp. II) is denoted by Γ (resp. Δ).

Let H^p be a Sobolev space of functions whose generalized derivatives up to the order p belong to $L^2(\mathbb{R}^n)$. Denote its inner product and norm by $(\cdot, \cdot)_p$ and $\|\cdot\|_p$ respectively, for $p = 0$, $H^0 = H$, $(\cdot, \cdot)_0 = (\cdot, \cdot)$ and $\|\cdot\|_0 = \|\cdot\|$, for simplicity.

For $Y \in \mathcal{M}$ and $Z \in \mathcal{N}$, we consider SPDE (1.1) with $m = 1$, namely

$$(2.1) \quad \begin{cases} d\xi(x, t) = (\partial_i(\sigma^{ij}(x, Y(t), Z(t)) \partial_j \xi(x, t)) \\ \quad + \gamma^i(x, Y(t), Z(t)) \partial_i \xi(x, t) - c\xi(x, t)) dt \\ \quad + f(x, Y(t), Z(t)) \xi(x, t) dW(t), \quad x \in \mathbb{R}^n, t > 0, \\ \xi(0) = \eta \in L^2(\mathbb{R}^n), \end{cases}$$

where $\partial_i = \frac{\partial}{\partial x_i}$ and we employ the convention of summation over repeated indices.

DEFINITION 2.3. We say that \mathcal{F}_t -progressively measurable process ξ is a solution of (2.1), if

$$(i) \quad \xi \in L^2(\Omega \times (0, T]; H^1) \cap L^2(\Omega; C([0, T]; H))$$

and

(ii) for $t \in [0, T]$ and $\zeta \in C_0^\infty(\mathbb{R}^n)$ (= infinitely differentiable function with compact support),

$$(\xi(t), \zeta) = (\eta, \zeta) - \int_0^t (\sigma^{ij}(Y(s), Z(s)) \partial_j \xi(s), \partial_i \zeta) ds$$

$$\begin{aligned}
& + \int_0^t (\gamma^i(Y(s), Z(s)) \partial_j \xi(s) - c \xi(s), \zeta) ds \\
& + \int_0^t (f(Y(s), Z(s)) \xi(s), \zeta) dW(s), \text{ with probability 1.}
\end{aligned}$$

Namely we study the Zakai equation in the frame work of [7] and [10]. Since this definition is consistent with a weak solution [1] of partial differential equation, we can split (2.1) into two parts to obtain suitable evaluations (see (3.9)).

Hereafter we always assume the conditions (A1)~(A4) below.

(A1) $\sigma^{ij}(\cdot, y, z), \gamma^i(\cdot, y, z), f(\cdot, y, z) \in C^4(R^n)$ and

$$(2.2) \quad \sup_{(y,z) \in \mathcal{Y} \times \mathcal{Z}} \|g(\cdot, y, z)\|_{C^4(R^n)} \leq b < \infty, \quad \text{for } g = \sigma^{ij}, \gamma^i, f$$

(A2) Lipschitz condition; there is a positive number $l = l_g$ such that

$$\begin{aligned}
& |g(x, y, z) - g(x', y', z')| \\
& \leq l(|x - x'| + |y - y'| + |z - z'|) \text{ for } g = \sigma^{ij}, \gamma^i, f
\end{aligned}$$

where $|\cdot| =$ Euclidian norm.

(A3) uniformly elliptic; $\sigma^{ij} = \sigma^{ji}$ and there is a positive number $\tilde{\lambda}_0$ such that, for any $(x, y, z) \in R^n \times \mathcal{Y} \times \mathcal{Z}$

$$\sum_{ij=1}^n \sigma^{ij}(x, y, z) \theta_i \theta_j \geq \tilde{\lambda}_0 |\theta|^2, \quad \text{for } \theta = (\theta_1, \dots, \theta_n) \in R^n$$

(A4) c is a non-negative constant.

First we recall Theorem 2.1, due to N.V. Krylov and B.L. Rozovskii [5] and E. Pardoux [8], in our convenient form.

THEOREM 2.1. *There exists a unique solution $\xi = \xi(\cdot, \eta, Y, Z)$ of (2.1) and the following properties hold*

$$(2.3) \quad E(\sup_{t \leq T} \|\xi(t, \eta, Y, Z)\|^2) \leq K \|\eta\|^2,$$

$$(2.4) \quad E\left(\int_0^T \|\xi(t, \eta, Y, Z)\|_1^2 dt\right) \leq K \|\eta\|^2$$

$$(2.5) \quad E(\|\xi(t, \eta, Y, Z)\|^{2l}) \leq K \|\eta\|^{2l}, \quad \text{for } t \in [0, T], l = 2, 3, 4,$$

with a constant K depending only on $T, \tilde{\lambda}_0$ and the bound of coefficients.

Since the linearity of equation (2.1) yields, with probability 1,

$$(2.6) \quad \xi(t, \eta, Y, Z) - \xi(t, \tilde{\eta}, Y, Z) = \xi(t, \eta - \tilde{\eta}, Y, Z) \text{ for } t \in [0, T],$$

the solution depends on the initial data continuously.

Let us put

$$\begin{aligned} L(y, z)\zeta &= \partial_i(\sigma^{ij}(\cdot, y, z)\partial_j\zeta) + \gamma^i(\cdot, y, z)\partial_i\zeta, \\ L(t, Y, Z) &= L(Y(t), Z(t)), L(t, Y, \beta) = L(t, Y, \beta(Y)), \\ L(t, \alpha, Z) &= L(t, \alpha(Z), Z), f(t, Y, Z) = f(\cdot, Y(t), Z(t)). \end{aligned}$$

Then (A3) implies the coercive condition, namely there is $\lambda_0 > 0$, such that

$$(2.7) \quad -\langle \zeta, L(y, z)\zeta \rangle + \lambda_0 \|\zeta\|^2 \geq 0, \text{ for } \zeta \in H^1 \text{ and } (y, z) \in \mathcal{Y} \times \mathcal{Z},$$

where $\langle \cdot, \cdot \rangle =$ duality pairing between H^1 and H^{-1} under $H = H^*$ (= dual space of H).

Next we consider SPDE (2.8) below

$$(2.8) \quad \begin{cases} d\zeta(t) = (L(t, Y, Z)\zeta(t) - c\zeta(t))dt + f(t, Y, Z)\zeta(t)dW(t) + F(t)dt \\ \quad + G(t)dW(t), \text{ for } t > 0, \\ \zeta(0) = 0, \end{cases}$$

where F and G are \mathcal{F}_t -progressively measurable and satisfy

$$E\left(\int_0^T \|F(t)\|^2 dt\right) < \infty \quad \text{and} \quad E\left(\int_0^T \|G(t)\|_1^2 dt\right) < \infty.$$

Then, there is a unique solution $\zeta \in L^2(\Omega \times [0, T]; H^1) \cap L^2(\Omega; C([0, T]; H))$ and the following evaluations hold [7],

$$(2.9) \quad E(\sup_{t \leq T} \|\zeta(t; Y, Z)\|^2) \leq \tilde{K} \int_0^T E(\|F(t)\|^2 + \|G(t)\|_1^2) dt$$

$$(2.10) \quad E\left(\int_0^T \|\zeta(t; Y, Z)\|_1^2 dt\right) \leq \tilde{K} \int_0^T E(\|F(t)\|^2 + \|G(t)\|_1^2) dt$$

with a constant \tilde{K} depending only on $T, \tilde{\lambda}_0$ and the bound of coefficients.

Now we will recall the evaluation of negative norm $\|\cdot\|_{-2}$ (= $\|\cdot\|_{H^{-2}}$) due to P.L. Lions [8]. Let $\zeta \in C^\infty(\mathbb{R}^n)$ satisfy the following condition;

$$\zeta = 0 \text{ on } (|x| \leq \frac{1}{2}), \zeta = 1 \text{ on } (|x| \geq 1) \text{ and } 0 \leq \zeta(x) \leq 1 \text{ on } \mathbb{R}^n.$$

Put $\zeta_R(x) = \zeta\left(\frac{x}{R}\right)$. Then we have

THEOREM 2.2 [8]. *There exists a constant K^* , which is independent of Y and Z , such that*

$$(2.11) \quad E \|\xi(t, (1 - \zeta_R)\eta, Y, Z)\|_{-2}^2 \leq K^* \|(1 - \zeta_R)\eta\|_{-2}^2, \quad \text{for } t \in [0, T], R > 0.$$

Moreover, for any $e \in H^2$,

$$(2.12) \quad E |(\xi(t, \eta, Y, Z), e)| \leq K^* \|(1 - \zeta_R)\eta\|_{-2} \|e\|_2 + \left(\varepsilon_R(e) + \frac{K}{R}\right) \|\eta\|$$

where $\varepsilon_R(e) = \int_{|x| > (R/2)} e(x)^2 dx$.

Later, we will see that (2.12) implies weakly sequential continuity of value function.

Remark. Let ξ_0 be a solution of (2.1) with $c = 0$. Then we can easily see

$$(2.13) \quad \xi(t, \eta, Y, Z) = e^{-ct} \xi_0(t, \eta, Y, Z).$$

So, we employ ξ_0 instead of ξ , when we stress $c = 0$.

3. Continuity with respect to time

In this section we study continuous dependency of a solution on time. Fixing Y and Z , we set $L(t, \omega) = L(t, Y(\omega), Z(\omega))$. Suppose $v; R^n \times \Omega \rightarrow R^1$ satisfies the condition,

$$(3.1) \quad v(x, \omega) \exp(-r|x|^2) \in H \quad \text{for all } \omega \in \Omega,$$

with a constant $r \geq 0$.

We consider the Cauchy problem

$$(3.2) \quad \begin{cases} \frac{\partial u}{\partial t} = L(t, \omega)u & \text{for } t > s, \\ u(s) = v(\omega). \end{cases}$$

Since coefficients σ^{ij} , γ^i and f of $L(t, \omega)$ may not be continuous in t , we employ a weak solution [1] of (3.2) and recall the following theorem, due to D.G. Aronson [1].

THEOREM 3.1. *There exists a unique weak fundamental solution for $L(\omega)$, say $\Gamma(x, t, y, s; \omega)$, and (3.3)~(3.6) hold.*

$$(3.3) \quad u(x, t; \omega) = \int_{R^n} \Gamma(x, t, y, s; \omega) v(y, \omega) dy$$

is a unique solution of (3.2)

$$(3.4) \quad \int_{R^n} \Gamma(x, t, y, s; \omega) dy = 1$$

$$(3.5) \quad u(x, t; \omega) = \int_{R^n} \Gamma(x, t, y, \theta; \omega) u(y, \theta; \omega) dy, \quad \text{for } \theta \in [s, t].$$

(3.6) *There exist positive numbers a_1, a_2 and p , independent of ω , such that*

$$p^{-1} g_1(x - y, t - s) \leq \Gamma(x, t, y, s; \omega) \leq p g_2(x - y, t - s)$$

where

$$g_i(x, \theta) = \sqrt{(4\pi\theta a_i)^{-n}} \exp(-|x|^2 / (4\theta a_i)), \quad i = 1, 2.$$

It follows from (3.4) and (3.5) that

$$\begin{aligned} \|u(t, \omega) - u(\theta, \omega)\|^2 &= \int \left[\int \Gamma(x, t, y, \theta; \omega) (u(y, \theta, \omega) - u(x, \theta, \omega)) dy \right]^2 dx \\ &\leq p \int \int g_2(x, t - \theta) (u(y + x, \theta, \omega) - u(y, \theta, \omega))^2 dy dx \\ &= p \int g_2(x, t - \theta) \|u(\cdot + x, \theta, \omega) - u(\cdot, \theta, \omega)\|^2 dx \end{aligned}$$

and, denoting $\Delta_x u(\cdot) = u(\cdot + x) - u(\cdot)$,

$$(3.7) \quad \|u(t, \omega) - u(\theta, \omega)\|^2 \leq p \int g_2(x, t - \theta) \|\Delta_x u(\theta, \omega)\|^2 dx.$$

Let us set $\xi_0(t, \eta, \omega) = \xi_0(t, \eta, Y, Z, \omega)$ and $G(t, \omega) = f(Y(t, \omega), Z(t, \omega)) \xi_0(t, \eta, \omega)$, for simplicity, and define X_1 and $X_2 \in L^2(\Omega; C([s, T]; H))$ as follows,

$$(3.8) \quad \begin{cases} dX_1(t, \omega) = L(t, \omega) X_1(t, \omega) dt, & \text{for } t > s \\ X_1(s, \omega) = \xi_0(s, \eta, \omega) \end{cases}$$

and

$$(3.9) \quad \begin{cases} dX_2(t) = L(t, Y, Z)X_2(t) dt + G(t)dW(t), & \text{for } t > s \\ X_2(s) = 0. \end{cases}$$

Since $\xi_0(s, \eta, \omega) \in H$, (3.3) asserts

$$X_1(x, t, \omega) = \int_{\mathbb{R}^n} \Gamma(x, t, y, s; \omega) \xi_0(y, s, \eta; \omega) dy.$$

Hence (3.7) implies

$$(3.10) \quad \|X_1(t, \omega) - \xi_0(s, \eta, \omega)\|^2 \leq p \int g_2(x, t-s) \|\Delta_x \xi_0(s, \eta, \omega)\|^2 dx.$$

On the other hand, (2.7) yields

$$E \|X_2(t)\|^2 \leq \lambda_0 \int_s^t E \|X_2(\theta)\|^2 d\theta + \int_s^t E \|G(\theta)\|^2 d\theta.$$

Thus, using (2.3), we have

$$(3.11) \quad \begin{aligned} E \|X_2(t)\|^2 &\leq \int_s^t \exp(\lambda_0(t-\theta)) E \|G(\theta)\|^2 d\theta \\ &\leq K |f|_\infty^2 \|\eta\|^2 [\exp(\lambda_0(t-s)) - 1] \lambda_0^{-1}, \end{aligned}$$

and $X(t) = X_1(t) + X_2(t)$ satisfies SPDE (3.12) below,

$$(3.12) \quad \begin{cases} dX(t) = L(t, Y, Z)X(t) dt + f(t, Y, Z)X(t)dW(t) \\ X(s) = \xi_0(s, \eta, Y, Z). \end{cases}$$

Hence, by the uniqueness of solution, X coincides with ξ_0 ,

$$(3.13) \quad X(t) = \xi_0(t, \eta, Y, Z), \quad t \geq s, \quad \text{with probability 1.}$$

Noting $\xi_0(t, \eta, Y, Z) - \xi_0(s, \eta, Y, Z) = X_1(t) - X_1(s) + X_2(t)$, we get

$$(3.14) \quad \begin{aligned} E \|\xi_0(t, \eta, Y, Z) - \xi_0(s, \eta, Y, Z)\|^2 \\ \leq 2p \int g_2(x, t-s) E \|\Delta_x \xi_0(s, \eta, Y, Z)\|^2 dx + 2K |f|_\infty^2 [\exp(\lambda_0(t-s)) - 1] \lambda_0^{-1} \|\eta\|^2. \end{aligned}$$

Hereafter k_i stands for a constant independent of Y and Z . Let us evaluate $\Delta_x \xi_0$. Putting $q_x(t) = \Delta_x \xi_0(t, \eta, Y, Z)$, we can derive the equation (3.15) below,

$$(3.15) \quad \begin{cases} dq_x(t) = L(t, Y, Z)q_x(t)dt + f(t, Y, Z)q_x(t)dW(t) \\ \quad + [\partial_i(\Delta_x\sigma^{ij}(t, Y, Z))\partial_j\tilde{X}(t) + \Delta_x\gamma^i(t, Y, Z)\partial_i\tilde{X}(t)] dt \\ \quad + \Delta_x f(t, Y, Z)\tilde{X}(t)dW(t), \\ q_x(t) = \Delta_x\eta, \end{cases}$$

where $\tilde{X}(t) = \xi_0(\cdot + x, t, \eta, Y, Z)$. Applying the standard arguments, we get

$$(3.16) \quad E \|q_x(t)\|^2 \leq k_1(\|\Delta_x\eta\|^2 + |x|^2 \|\eta\|^2) \quad \text{for } x \in R^n.$$

Combining (3.16) with (3.14), we have

$$(3.17) \quad \begin{aligned} E \| \xi_0(t, \eta, Y, Z) - \xi_0(s, \eta, Y, Z) \|^2 \\ \leq k_2 \left[\int g_2(x, t-s) \|\Delta_x\eta\|^2 dx + (t-s) \|\eta\|^2 \right]. \end{aligned}$$

Since $\|\Delta_x\eta\|$ tends to 0, as $x \rightarrow 0$, the modulus of continuity in t is independent of Y and Z , but depends on the initial data η . Applying the same arguments as [8, part 2], we obtain

$$(3.18) \quad E \| \xi_0(t, \eta, Y, Z) - \xi_0(s, \eta, Y, Z) \|_{-1}^2 \leq k_3 |t-s| \|\eta\|^2.$$

The above evaluation together with (2.13) yields the following proposition,

PROPOSITION 3.1. *There is a constant C_1 such that, for $Y \in \mathcal{M}$, $Z \in \mathcal{N}$ and $\eta \in H$*

$$(3.19) \quad E \| \xi(t, \eta, Y, Z) - \xi(s, \eta, Y, Z) \|^2 \leq C_1(|t-s| \|\eta\|^2 + \int g_2(x, t-s) \|\Delta_x\eta\|^2 dx)$$

$$(3.20) \quad E \| \xi(t, \eta, Y, Z) - \xi(s, \eta, Y, Z) \|_{-1}^2 \leq C_1 |t-s| \|\eta\|^2$$

$$(3.21) \quad E \| \xi(t, \eta, Y, Z) - \xi(s, \eta, Y, Z) \|_{-1}^2 \leq C_1 |t-s| \|\eta\|_1^2 \quad \text{for } \eta \in H^1.$$

For $\Phi \in \text{LUC}(H)$ (= uniformly continuous functions on H with linear growth), we put

$$(3.22) \quad \tilde{\Phi}(t, \eta; Y, Z) = E\Phi(\xi(t, \eta, Y, Z)).$$

Then Proposition 3.1 implies the following one.

PROPOSITION 3.2.

(i) For $\varepsilon > 0$, there is $\tau_0 = \tau_0(\varepsilon, \eta, \Phi) > 0$, such that for $\mathcal{Y} \in \mathcal{M}$ and $Z \in \mathcal{N}$

$$|\tilde{\Phi}(t, \eta, Y, Z) - \tilde{\Phi}(s, \eta, Y, Z)| < \varepsilon, \quad \text{if } |t - s| < \tau_0.$$

(ii) For $\varepsilon > 0$, there is $\delta_0 = \delta_0(\varepsilon, \Phi) > 0$, such that for $Y \in \mathcal{M}$, $Z \in \mathcal{N}$ and $t \in [0, T]$

$$|\tilde{\Phi}(t, \eta, Y, Z) - \tilde{\Phi}(t, \eta', Y, Z)| < \varepsilon, \quad \text{if } \|\eta - \eta'\| < \delta_0.$$

PROPOSITION 3.3. Suppose that, for any bounded set B of H , $\Phi_n \rightarrow \Phi$ uniformly in B and $\sup_n \sup_{\zeta \in B} |\Phi_n(\zeta)| < \infty$. Then, for any bounded set $B \subset H$, $\tilde{\Phi}_n$ tends to $\tilde{\Phi}$ uniformly in $[0, T] \times B \times \mathcal{M} \times \mathcal{N}$.

Proof. Recalling (2.3), we have

$$P(\|\xi(t, \eta, Y, Z)\| > r) \leq K \|\eta\|^2 / r^2.$$

Now taking $r = r(\varepsilon, B) = K(\text{diam } B)^2 \varepsilon^{-2}$ and $N = N(\varepsilon, B)$ such that

$$\sup_{n \geq N} |\Phi_n - \Phi| < \varepsilon \text{ on } \{\eta \in H; \|\eta\| \leq r\}, \text{ we get}$$

$$\begin{aligned} & |\tilde{\Phi}_n(t, \eta, Y, Z) - \tilde{\Phi}(t, \eta, Y, Z)| \\ & < \varepsilon + 2\beta P(\|\xi(t, \eta, Y, Z)\|^2 > r) < \varepsilon + 2\varepsilon\beta \end{aligned}$$

where $\sup |\Phi_n(\zeta)| < \beta$ on B . This completes the proof.

Hereⁿ we call $\varphi: H \rightarrow R^1$, a tame function, if φ is denoted by

$$(3.23) \quad \varphi(\eta) = \hat{\varphi}((e_1, \eta), \dots, (e_j, \eta))$$

with suitable $e_i \in H^2$ and $\hat{\varphi} \in C^2(R^j)$ whose first and second derivatives are BUC (= bounded and uniformly continuous function). So, φ is Lipschitz continuous and linear growth, say

$$(3.24) \quad |\varphi(\eta)| \leq k_1 + k_2 \|\eta\| \quad \text{with } k_i = k_i(\hat{\varphi}).$$

Putting $\bar{\varphi}(t, \eta, Y, Z) = E\varphi(\xi(t, \eta, Y, Z))$, we have

PROPOSITION 3.4. *There is a constant $C_2(\varphi)$ such that*

(3.25)

$$|\bar{\varphi}(t, \eta, Y, Z) - \bar{\varphi}(s, \eta, Y, Z)| \leq C_2(\varphi) \|\eta\| \sqrt{|t - s|}, \quad \text{for } Y \in \mathcal{M} \text{ and } Z \in \mathcal{N}.$$

Proof. $l(\varphi)$ denotes Lipschitz constant of $\hat{\varphi}$ of (3.23).

$$\begin{aligned} & |E\varphi(\xi(t, \eta, Y, Z)) - E\varphi(\xi(s, \eta, Y, Z))| \\ & \leq l(\varphi) \sum_{i=1}^j E |(\xi(t, \eta, Y, Z) - \xi(s, \eta, Y, Z), e_i)| \\ & \leq l(\varphi) \sum_{i=1}^j E \|\xi(t, \eta, Y, Z) - \xi(s, \eta, Y, Z)\|_{-1} \|e_i\|_1. \end{aligned}$$

So, Proposition 3.1 concludes the proof.

4. Tame terminal function

In this section, we deal with pay-off tame terminal function and show some compactness of value functions, employing semi-discretization arguments.

Suppose $h; H \times \mathcal{Y} \times \mathcal{Z} \rightarrow \mathbf{R}^1$ is continuous and satisfies (A5),

(A5) $h(\cdot, y, z)$ is tame, say $h(\eta, y, z) = \mathfrak{h}((\eta, e_1), \dots, (\eta, e_q), y, z)$ and $|\partial_i \mathfrak{h}(x, y, z)|$ and $|\partial_i \partial_j \mathfrak{h}(x, y, z)|$ are bounded in $(x, y, z) \in \mathbf{R}^q \times \mathcal{Y} \times \mathcal{Z}$. Let $\varphi; H \rightarrow \mathbf{R}^1$ be a tame function, say

$$\varphi(\eta) = \hat{\varphi}((\eta, \hat{e}_1), \dots, (\eta, \hat{e}_p)).$$

Since we may assume that $\{\hat{e}_i, i = 1, \dots, p\}$ contains $\{e_i, i = 1, \dots, q\}$, we will drop $\hat{\cdot}$.

Now we define pay-off J and value $V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}$ as follows.

$$(4.1) \quad \begin{cases} J(t, \eta, Y, Z, \varphi) = E \int_0^t h(\xi(\theta, \eta, Y, Z), Y(\theta), Z(\theta)) d\theta \\ \hspace{20em} + \varphi(\xi(t, \eta, Y, Z)) \\ V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}(t, \eta, \varphi) = \inf_{z \in \hat{\mathcal{N}}} \sup_{Y \in \hat{\mathcal{M}}} J(t, \eta, Y, Z, \varphi) \text{ for } \hat{\mathcal{M}} \subset \mathcal{M}, \hat{\mathcal{N}} \subset \mathcal{N}. \end{cases}$$

Since $|h(\eta, y, z)| \leq m_1 + m_2 \|\eta\|$ with m_1 and m_2 , which are independent of y and z , (2.3) derives

$$(4.2) \quad |J(t, \eta, Y, Z, \varphi)| \leq m_1 t + k_1 + K(m_2 + k_2) \|\eta\|$$

whenever $|\varphi(\eta)| \leq k_1 + k_2 \|\eta\|$.

Appealing to Proposition 3.4, Theorem 2.2 and (2.6), we can take a positive numbers $C_3 = C_3(\varphi)$ such that

$$(4.3) \quad \begin{cases} |J(t, \eta, Y, Z, \varphi) - J(s, \zeta, Y, Z, \varphi)| \\ \leq C_3 [(1 + \|\eta\|) \sqrt{|t-s|} + \|\eta - \zeta\|] \text{ for } Y \in \mathcal{M} \text{ and } Z \in \mathcal{N}, \\ |V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}(t, \eta, \varphi) - V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}(s, \zeta, \varphi)| \leq C_3 [(1 + \|\eta\|) \sqrt{|t-s|} + \|\eta - \zeta\|] \\ \text{for } \hat{\mathcal{M}} \subset \mathcal{M} \text{ and } \hat{\mathcal{N}} \subset \mathcal{N}, \end{cases}$$

and, putting $S(R) = \{|x| \geq \frac{R}{2}\}$ and $\varepsilon_R = \sum_{i=1}^p \int_{S(R)} e_i^2(x) dx$, we have

$$(4.4) \quad |V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}(t, \eta, \varphi) - V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}(s, \zeta, \varphi)| \\ \leq C_3 \left[(1 + \|\eta\|) \sqrt{|t-s|} + \|(1 - \zeta_R)(\eta - \zeta)\|_{-2} + \left(\varepsilon_R + \frac{1}{R}\right) \|\eta - \zeta\| \right]$$

with ζ_R of Theorem 2.2. Moreover (4.4) implies

$$(4.5) \quad V_{\hat{\mathcal{M}}\hat{\mathcal{N}}}(\cdot, \varphi) \in C([0, T] \times H_w)$$

where H_w denotes the space H carrying the weak topology.

Let $\pi = \{0 = t_0 < t_1 < \cdots < t_N = T\}$ be a partition on $[0, T]$ and denote $\|\pi\| = \max_i (t_i - t_{i-1})$ its mesh. We put $\mathcal{P}(\pi) = \{t_i, i = 0, \cdots, N\}$.

DEFINITION 4.1. $Y \in \mathcal{M}$ (resp. $Z \in \mathcal{N}$) is called π -admissible for player I (resp. II), if

$$Y(t) = Y(t_j) \text{ (resp. } Z(t) = Z(t_j)), \text{ for } t \in [t_j, t_{j+1}).$$

\mathcal{M}_π (resp. \mathcal{N}_π) denotes the set of all π -admissible controls for player I (resp. II).

DEFINITION 4.2. $\alpha \in \Gamma$ is called a π -admissible strategy for player I, if $\alpha; \mathcal{N} \rightarrow \mathcal{M}_\pi$ such that

- (i) $\alpha(Z)(s) = Z$ -independent for $s < t_1$,
- (ii) $\alpha(Z)(t_j) = \alpha(\hat{Z})(t_j)$ a.e., if $Z = \hat{Z}$ a.e. on $[0, t_j)$.

$\beta \in \Delta$ is called a π -admissible strategy for player II, if $\beta; \mathcal{M} \rightarrow \mathcal{N}_\pi$ such that

- (i) $\beta(Y)(s) = Y$ -independent for $s < t_1$,
- (ii) $\beta(Y)(t_j) = \beta(\hat{Y})(t_j)$ a.e., if $Y = \hat{Y}$ a.e. on $[0, t_j]$.

Γ_π (resp. Δ_π) denotes the set of all π -admissible strategies for player I (resp. II).

Let us set

$$J(t, \eta, Y, \beta, \varphi) = J(t, \eta, Y, \beta(Y), \varphi)$$

and the upper value function V_π for π is defined by (4.6) below,

$$(4.6) \quad V_\pi(t, \eta, \varphi) = \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(t, \eta, Y, \beta, \varphi).$$

From (4.3) and (4.4), we can easily see

PROPOSITION 4.1. $V_\pi(\cdot, \varphi)$ has π -independent continuity.

In particular, (4.4) implies compactness of upper value functions, namely, we have the following theorem.

THEOREM 4.1. Let $\pi_n, n = 1, 2, \dots$, be a sequence of partitions on $[0, T]$. Then, $V_{\pi_n}(\cdot, \varphi), n = 1, 2, \dots$, has a sequence which converges uniformly in any bounded set of $[0, T] \times H$.

For the proof, we will show the following lemma.

LEMMA. Let $\varepsilon > 0$ and $B = \{\eta \in H; \|\eta\| \leq r\}$. Then we can take a finite set $D \subset [0, T] \times B$, such that for $(t, \eta) \in [0, T] \times B$ there is $(s, \zeta) \in D$ satisfying

$$(4.7) \quad |V_\pi(t, \eta, \varphi) - V_\pi(s, \zeta, \varphi)| < \varepsilon, \quad \text{for any partition } \pi.$$

Proof. Let us take a large number $R = R(r, \varepsilon)$, such that

$$(4.8) \quad C_3(\varphi) \left(\varepsilon_R + \frac{1}{R} \right) (2r) < \varepsilon.$$

So we may assume $R(r, \varepsilon) \rightarrow \infty$, as $\varepsilon \rightarrow 0$. Denote $S = \{x \in R^n; |x| \leq R\}$ and $\tilde{\zeta} = \zeta_R$ of Theorem 2.2. Then $\text{supp}(1 - \tilde{\zeta})\eta \subset S$. Since B is weakly compact in H , $\{(1 - \tilde{\zeta})\eta, \eta \in B\}$ is compact in $H^{-2}(S)$. Hence we can take a finite set $E \subset B$, such that $\{(1 - \tilde{\zeta})\eta, \eta \in E\}$ is $\varepsilon(3C_3(\varphi))^{-1}$ -net in $H^{-2}(S)$.

Let Θ be a finite $(3C_3(\varphi)(1+r))^{-2}\varepsilon^2$ -net in $[0, T]$. Then (4.4) asserts that, for $(t, \eta) \in [0, T] \times B$, there is $(s, \zeta) \in \Theta \times E$ such that

$$(4.9) \quad |V_\pi(t, \eta, \varphi) - V_\pi(s, \zeta, \varphi)| < \varepsilon, \quad \text{for any } \pi.$$

This completes the proof of Lemma.

Proof of Theorem. Denote D of Lemma by $D(r, \varepsilon)$ and put $\mathcal{D} = \bigcup_{r, k=1}^{\infty} D\left(r, \frac{1}{k}\right)$.

Since $|V_\pi(\cdot, \varphi)|_\infty \leq m_1 T + k_1(\varphi) + K(m_2 + k_2(\varphi)) \|\eta\|$, some subsequence $V_{\pi_n}(s, \zeta, \varphi)$ converges for any $(s, \zeta) \in \mathcal{D}$. Now

Lemma implies Theorem.

Put $V(\cdot, \varphi) = \lim V_{\pi_n}(\cdot, \varphi)$. Then V also satisfies (4.3) and (4.4).

5. Uniformly continuous terminal function with linear growth

Since an upper value function $V_\pi(\cdot, \varphi)$ may not be tame, we will study pay-off with continuous terminal function, in order to prove the principle of dynamic programming in Section 6.

It is clear that $J(t, \cdot, Y, Z, \Phi) \in \text{LUC}(H)$ whenever $\Phi \in \text{LUC}(H)$. Moreover we can easily see the following proposition.

PROPOSITION 5.1.

(i) For $\varepsilon > 0$, there is $\delta_0 = \delta_0(\varepsilon, \Phi) > 0$, such that

$$(5.1) \quad |J(t, \eta, Y, Z, \Phi) - J(t, \hat{\eta}, Y, Z, \Phi)| < \varepsilon$$

and

$$(5.2) \quad |V_{\hat{\mu}, \hat{\nu}}(t, \eta, \Phi) - V_{\hat{\mu}, \hat{\nu}}(t, \hat{\eta}, \Phi)| < \varepsilon$$

whenever $\|\eta - \hat{\eta}\| < \delta_0$.

(ii) For $\varepsilon > 0$, there is $\hat{\delta}_0 = \hat{\delta}_0(\varepsilon, \phi, \eta) > 0$, such that

$$(5.3) \quad |V_{\hat{\mu}, \hat{\nu}}(t, \eta, \Phi) - V_{\hat{\mu}, \hat{\nu}}(s, \eta, \Phi)| < \varepsilon, \text{ whenever } |t - s| < \hat{\delta}_0.$$

Let us set

$$(5.4) \quad \begin{cases} v(t, \eta, z, \Phi) = \sup_{Y \in \mathcal{M}} J(t, \eta, Y, z, \Phi) \\ v(t, \eta, \Phi) = \inf_{z \in \mathcal{Z}} v(t, \eta, z, \Phi). \end{cases}$$

Then (5.2) implies that $v(t, \cdot, z, \Phi) \in \text{LUC}(H)$ and

$$(5.5) \quad |v(t, \eta, z, \Phi) - v(t, \hat{\eta}, z, \Phi)| < \varepsilon \text{ for any } (t, z) \in [0, T] \times \mathcal{Z},$$

whenever $\|\eta - \hat{\eta}\| < \delta_0$.

First we study the continuity of v with respect to z . Putting

$$\hat{\xi}(t) = \xi(t, \eta, Y, \hat{z}) \text{ and } \zeta(t) = \xi(t, \eta, Y, z) - \hat{\xi}(t)$$

we can see

$$\begin{cases} d\zeta(t) = [L(Y(t), z) - cI]\zeta(t)dt + f(Y(t), z)\zeta(t)dW(t) + F(t)dt + G(t)dW(t) \\ \zeta(0) = 0 \end{cases}$$

where $F(t) = [L(Y(t), z) - L(Y(t), \hat{z})]\hat{\xi}(t)$ and $G(t) = [f(Y(t), z) - f(Y(t), \hat{z})]\hat{\xi}(t)$.

Employing the standard arguments, we obtain the following one.

PROPOSITION 5.2.

(i) There is a constant C_4 independent of Y , z and η such that

$$(5.6) \quad E(\sup_{t \leq T} \|\xi(t, \eta, Y, z) - \xi(t, \eta, Y, \hat{z})\|^2) \leq C_4 \|\eta\|^2 |z - \hat{z}|^2.$$

(ii) For $\varepsilon > 0$ and k , there is $\delta_k = \delta_k(\varepsilon, \Phi) > 0$, such that

$$(5.7) \quad |J(t, \eta, Y, z, \Phi) - J(t, \eta, Y, \hat{z}, \Phi)| < \varepsilon, \\ \text{for } t \in [0, T], Y \in \mathcal{M}, \eta \in B(k) = \{\eta \mid \|\eta\| \leq k\},$$

whenever $|z - \hat{z}| < \delta_k$.

(iii) $v(t, \eta, z, \Phi)$ satisfies (5.7).

Now we will prove the principle of dynamic programming, applying the similar arguments as [4]. Let $\mathfrak{B}_\varepsilon^H (= \mathfrak{B}_{\varepsilon\Phi}^H) = \{\mathcal{E}_j, j = 1, 2, \dots\}$ be a Borel partition of H , with $\text{diam}\mathcal{E}_j < \delta_0(\varepsilon, \Phi)$ (see Proposition 5.1 for δ_0). For simplicity, we may assume $\mathcal{E}_j \subset B(j)$. Since \mathcal{L} is compact, for any fixed $\eta_j \in \mathcal{E}_j$, $v(t, \eta_j, \cdot, \Phi)$ has a minimum point $z_j^* = z^*(t, \eta_j, \Phi)$, namely

$$(5.8) \quad v(t, \eta_j, z_j^*, \Phi) = v(t, \eta_j, \Phi).$$

Taking an approximate maximum point $Y_j^* = Y_j^*(t, \eta_j, \Phi, \varepsilon)$ below,

$$(5.9) \quad v(t, \eta_j, z_j^*, \Phi) - \varepsilon < J(t, \eta_j, Y_j^*, z_j^*, \Phi) \leq v(t, \eta_j, z_j^*, \Phi)$$

we have

$$(5.10) \quad 0 \leq v(t, \eta_j, \Phi) - J(t, \eta_j, Y_j^*, z_j^*, \Phi) < \varepsilon.$$

Hence Proposition 5.2 asserts

$$(5.11) \quad |v(t, \eta, \Phi) - J(t, \eta, Y_j^*, z_j^*, \Phi)| < 3\varepsilon, \quad \text{for } \eta \in \mathcal{E}_j.$$

Appealing to $J(t, \eta_j, Y, z_j^*, \Phi) \leq v(t, \eta_j, z_j^*, \Phi)$ for $Y \in \mathcal{M}$, we get

$$(5.12) \quad J(t, \eta, Y, z_j^*, \Phi) \leq v(t, \eta, \Phi) + 2\varepsilon.$$

For $\pi = \{0 = t_0 < t_1 < \dots < t_N = T\}$ and $\tau \in (t_{p-1}, t_p]$, we will define $v(t) ; \text{LUC}(H) \rightarrow \text{LUC}(H)$ and $\Phi_j (= \Phi_j^\tau)$ as follows

$$(5.13) \quad v(t)\Phi = v(t, \cdot, \Phi) \\ \Phi_p = \Phi, \quad \Phi_{p-1} = v(\tau - t_{p-1})\Phi_p, \quad \Phi_{p-2} = v(s_{p-1})\Phi_{p-1}, \dots, \Phi_0 = v(s_1)\Phi_1$$

where $s_j = t_j - t_{j-1}$. Appealing to (5.8), we will choose an approximately optimal strategy $\beta_\varepsilon \in \Delta_\pi$ in the following way. Fixing $Y \in \mathcal{M}$ and $\eta_{j,k} \in \mathcal{E}_j (\in \mathfrak{P}_{\varepsilon\Phi_k}^H)$ arbitrarily, $\beta_\varepsilon(Y)$ is defined by

$$(5.14) \quad \begin{aligned} \beta_\varepsilon(Y)(\theta) &= I_{[0,t_1]}(\theta) \sum_{(\Phi_1)} I_{\mathcal{E}_1}(\eta) z^*(s_1, \eta_{j,1}, \Phi_1) \\ &\quad + I_{[t_1,t_2]}(\theta) \sum_{(\Phi_2)} I_{\mathcal{E}_2}(\zeta(t_1, \eta)) z^*(s_2, \eta_{j,2}, \Phi_2) + \cdots \\ &\quad + I_{[t_{p-1}, T]}(\theta) \sum_{(\Phi_p)} I_{\mathcal{E}_p}(\zeta(t_{p-1}, \eta)) z^*(\tau - t_{p-1}, \eta_{j,p}, \Phi_p) \end{aligned}$$

where $I_A =$ indicator function of A , $\sum_{(\Phi)} = \sum_{\mathcal{E} \in \mathfrak{P}_{\varepsilon\Phi}^H}$ and $\zeta(t, \eta)$ is a solution of (2.1) for $(Y, \beta_\varepsilon(Y))$ with $\zeta(0) = \eta$, namely,

$$\zeta(t, \eta) = \sum_{(\Phi_1)} I_{\mathcal{E}_1}(\eta) \xi(t, \eta, Y, z^*(s_1, \eta_{j,1}, \Phi_1)) = \xi(t, \eta, Y, \beta_\varepsilon(Y)),$$

for $t \leq t_1$,

Hence $\beta_\varepsilon(Y)$ on $[t_1, t_2]$ is determined by (5.14) and $\zeta(\theta, \eta)$ by $\xi(\theta, \eta, Y, \beta_\varepsilon(Y))$ for $\theta \in [0, t_2]$. Repeating the same arguments, we can determine $\beta_\varepsilon(Y)$ of (5.14) and

$$(5.15) \quad \zeta(t, \eta) = \xi(t, \eta, Y, \beta_\varepsilon(Y)) \text{ for } t \in [0, T].$$

Since $\beta_\varepsilon(Y)(\theta)$ is $\mathcal{F}_{t_{p-1}}$ -measurable and $Y(\theta)$ is a Borel function of $\{W(t), t \leq t_{p-1}\}$ and $\{W(s) - W(t_{p-1}), t_{p-1} \leq s \leq \theta\}$, putting $\xi(t) = \xi(t, \eta, Y, \beta_\varepsilon)$, we have

$$(5.16) \quad \begin{aligned} E \left(\int_{t_{p-1}}^\tau h(\xi(\theta), Y(\theta), \beta_\varepsilon(Y)(\theta)) d\theta + \Phi(\xi(\tau)) / \mathcal{F}_{t_{p-1}} \right) \\ \leq v(\tau - t_{p-1}, \xi(t_{p-1}), \Phi) + 2\varepsilon, \text{ a.s.} \end{aligned}$$

by virtue of (5.12). Thus, it follows that

$$(5.17) \quad \begin{aligned} J(\tau, \eta, Y, \beta_\varepsilon, \Phi) \\ \leq E \left[\int_0^{t_{p-1}} h(\xi(\theta), Y(\theta), \beta_\varepsilon(Y)(\theta)) d\theta + v(\tau - t_{p-1}) \Phi(\xi(t_{p-1})) + 2\varepsilon \right] \\ = E \left[\int_0^{t_{p-1}} h(\xi(\theta), Y(\theta), \beta_\varepsilon(Y)(\theta)) d\theta + \Phi_{p-1}(\xi(t_{p-1})) + 2\varepsilon \right]. \end{aligned}$$

Repeating the same arguments, we obtain

$$(5.18) \quad J(\tau, \eta, Y, \beta_\varepsilon, \Phi) \leq v(s_1)v(s_2) \cdots v(\tau - t_{p-1}) \Phi(\eta) + 2p\varepsilon.$$

Since Y is arbitrary, it follows that

$$(5.19) \quad \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta_\varepsilon, \Phi) \leq \mathbf{v}(s_1)\mathbf{v}(s_2) \cdots \mathbf{v}(\tau - t_{p-1})\Phi(\eta) + 2p\varepsilon.$$

From the fact $\beta_\varepsilon \in \Delta_\pi$, we see, letting $\varepsilon \rightarrow 0$,

$$(5.20) \quad \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi) \leq \mathbf{v}(s_1)\mathbf{v}(s_2) \cdots \mathbf{v}(\tau - t_{p-1})\Phi(\eta).$$

Now we will choose nearly optimal $\alpha_\varepsilon \in \Gamma$ in the following way. Fix $\eta_j \in \mathcal{E}_j \in \mathfrak{B}_{\varepsilon\Phi}^H$ arbitrarily. Noting the compactness of \mathcal{Z} , let us denote $\mathfrak{B}_{\varepsilon\Phi}^{\mathcal{Z}} = \{S_1, \dots, S_m\}$ a finite partition of \mathcal{Z} , with $\text{diam } S_i < \delta_j$, (see (5.7)). For any $z_i \in S_i \in \mathfrak{B}_{\varepsilon\Phi}^{\mathcal{Z}}$, we will take $Y^* = Y^*(t, \eta_j, z_j, \Phi) \in \mathcal{M}$ such that

$$(5.21) \quad J(t, \eta_j, Y^*, z_j, \Phi) > v(t, \eta_j, z_j, \Phi) - \varepsilon.$$

Then it follows from (5.1) and (5.7) that

$$(5.22) \quad J(t, \eta, Y^*, z_j, \Phi) > v(t, \eta, z_j, \Phi) - 3\varepsilon \quad \text{for } \eta \in \mathcal{E}_j$$

and

$$(5.23) \quad J(t, \eta, Y^*, z, \Phi) > v(t, \eta, z, \Phi) - 5\varepsilon \quad \text{for } \eta \in \mathcal{E}_j \text{ and } z \in S_i.$$

When we stress the dependence on Brownian path w , we will denote $Y(\theta)$ by $Y(\theta, w)$. Putting $Y_{p,j,i}^* = Y^*(\tau - t_{p-1}, \eta_j, z_i, \Phi_p)$ and $Y_{k,j,i}^* = Y^*(t_k - t_{k-1}, \eta_j, z_i, \Phi_k)$ for $\eta_j \in \mathcal{E}_j \in \mathfrak{B}_{\varepsilon\Phi_k}^H$ and $z_i \in S_i \in \mathfrak{B}_{\varepsilon\Phi_k}^{\mathcal{Z}}$, $k = p-1, \dots, 0$, we define $\alpha_\varepsilon(Z)$ as follows,

$$(5.24) \quad \begin{aligned} \alpha_\varepsilon(Z)(\theta, w) &= I_{[0, t_1)}(\theta) \sum_{(\Phi_1)}^* I_{\mathcal{E}_j}(\eta) I_{S_i}(Z(0, w)) Y_{1,j,i}^*(\theta, w) \\ &\quad + I_{[t_1, t_2)}(\theta) \sum_{(\Phi_2)}^* I_{\mathcal{E}_j}(\zeta(t_1, \eta)) I_{S_i}(Z(t_1, w)) Y_{2,j,i}^*(\theta - t_1, w_t^+) \\ &\quad + \cdots \\ &\quad + I_{[t_{p-1}, T]}(\theta) \sum_{(\Phi_p)}^* I_{\mathcal{E}_j}(\zeta(t_{p-1}, \eta)) I_{S_i}(Z(t_{p-1}, w)) Y_{p,j,i}^*(\theta - t_{p-1}, w_{t_{p-1}}^+), \end{aligned}$$

where $\sum_{(\Phi)}^* = \sum_{\delta \in \mathfrak{B}_{\varepsilon\Phi}^H} \sum_{S_i \in \mathfrak{B}_{\varepsilon\Phi}^{\mathcal{Z}}}$, w_t^+ is a shifted path, namely $w_t^+(\theta) = w(t + \theta) - w(t)$, and $\zeta(t, \eta) = \xi(t, \eta, \alpha_\varepsilon(Z), Z)$. Applying the same arguments as (5.16)~(5.18), we obtain

$$(5.25) \quad \begin{aligned} J(\tau, \eta, \alpha_\varepsilon, Z, \Phi) &\geq \mathbf{v}(s_1)\mathbf{v}(s_2) \cdots \mathbf{v}(\tau - t_{p-1})\Phi(\eta) - 5p\varepsilon, \\ &\quad \text{for } Z \in \mathcal{N}_\pi. \end{aligned}$$

Thus it follows that

$$(5.26) \quad \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \alpha_\varepsilon, Z, \Phi) \geq \mathbf{v}(s_1)\mathbf{v}(s_2) \cdots \mathbf{v}(\tau - t_{p-1})\Phi(\eta) - 5p\varepsilon.$$

Since $\alpha_\varepsilon \in \Gamma$, letting $\varepsilon \rightarrow 0$, we get

$$(5.27) \quad \sup_{\alpha \in \Gamma} \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \alpha, Z, \Phi) \geq v(s_1)v(s_2) \cdots v(\tau - t_{p-1})\Phi(\eta).$$

Combining with (5.20), we have Proposition 5.3 below,

PROPOSITION 5.3. *Let $\Phi \in \text{LUC}(H)$, $\mathcal{P}_\pi = \{t_j, j = 0, \dots, N\}$ and $\tau \in (t_{p-1}, t_p]$. For $\varepsilon > 0$, there exist $\alpha_\varepsilon \in \Gamma$ and $\beta_\varepsilon \in \Delta_\pi$ such that, for $Y \in \mathcal{M}$ and $Z \in \mathcal{N}_\pi$,*

$$(5.28) \quad \begin{aligned} J(\tau, \eta, Y, \beta_\varepsilon, \Phi) - \varepsilon &< v(t_1)v(t_2 - t_1) \cdots v(\tau - t_{p-1})\Phi(\eta) \\ &< J(\tau, \eta, \alpha_\varepsilon, Z, \Phi) + \varepsilon, \quad \text{for } \eta \in H. \end{aligned}$$

Moreover, according to [4], we obtain

PROPOSITION 5.4.

$$(5.29) \quad \begin{aligned} \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi) &= v(t_1)v(t_2 - t_1) \cdots v(\tau - t_{p-1})\Phi(\eta) \\ &= \sup_{\alpha \in \Gamma} \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \alpha, Z, \Phi). \end{aligned}$$

For the proof, we will recall the following lemma.

LEMMA [4]. *For any $\beta \in \Delta_\pi$ and $\alpha \in \Gamma$, there exist $Y^* \in \mathcal{M}$ and $Z^* \in \mathcal{N}_\pi$ such that*

$$(5.30) \quad J(\tau, \eta, \alpha, Z^*, \Phi) = J(\tau, \eta, Y^*, \beta, \Phi).$$

Outline of proof. For any fixed $z_0 \in \mathcal{Z}$, let us define $Y_k \in \mathcal{M}$ and $z_k \in \mathcal{N}_\pi$, as follows.

$$\begin{aligned} Y_0(s) &= \alpha(z_0)(s), \quad Z_0(s) = \beta(Y_0)(s), \quad Y_1(s) = \alpha(Z_0)(s), \\ Z_1(s) &= \beta(Y_1)(s), \cdots, \quad Y_k(s) = \alpha(Z_{k-1})(s), \quad Z_k(s) = \beta(Y_k)(s), \cdots. \end{aligned}$$

Then we have

$$(5.31) \quad Y_{j+1} = Y_j \text{ on } [0, t_j] \text{ and } Z_{j+1} = Z_j \text{ on } [0, t_j].$$

In fact, $\beta(Y)(s)$, $s < t_1$, is independent of Y , since $\beta \in \Delta_\pi$. Hence it follows that $\beta(Y_0) = \beta(Y_1) = \cdots = \beta(Y_k)$ on $[0, t_1)$, and

$$(5.32) \quad Z_0 = Z_1 = \cdots = Z_k \text{ on } [0, t_1).$$

Now (ii) of Definition (4.2) yields

$$(5.33) \quad \alpha(Z_0)(t_1) = \alpha(Z_1)(t_1) = \cdots = \alpha(Z_k)(t_1)$$

and

$$\alpha(Z_0) = \alpha(Z_1) = \cdots = \alpha(Z_k) \text{ on } [0, t_1].$$

This asserts

$$Y_1 = Y_2 = \cdots = Y_{k+1} \text{ on } [0, t_1].$$

So $\beta(Y_1)(t_1) = \cdots = \beta(Y_{k+1})(t_1)$. Since $\beta \in \Delta_\pi$, we see

$$(5.34) \quad \beta(Y_1) = \cdots = \beta(Y_{k+1}) \text{ on } [t_1, t_2].$$

From (5.32) and (5.34), we can see

$$Z_1 = \cdots = Z_{k+1} \text{ on } [0, t_2].$$

Repeating the same arguments, we can conclude (5.31).

Putting $Y^* = Y_N$ and $Z^* = Z_{N-1}$, we get

$$(5.35) \quad \alpha(Z^*) = Y^* \text{ and } \beta(Y^*) = Z^*.$$

Now the proof is completed.

Proof of Proposition 5.4. From (5.28), we can easily see

$$(5.36) \quad \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi) \leq v(t_1)v(t_2 - t_1) \cdots v(\tau - t_{p-1})\Phi(\eta) \\ \leq \sup_{\alpha \in \Gamma} \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \alpha, Z, \Phi).$$

For $\varepsilon > 0$, we take $\hat{\beta} \in \Delta_\pi$ and $\hat{\alpha} \in \Gamma$ such that

$$(5.37) \quad \begin{cases} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \hat{\beta}, \Phi) - \varepsilon < \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi) \\ \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \hat{\alpha}, Z, \Phi) + \varepsilon > \sup_{\alpha \in \Gamma} \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \alpha, Z, \Phi). \end{cases}$$

Now it follows from Lemma that

$$(5.38) \quad \begin{cases} J(\tau, \eta, Y^*, \hat{\beta}, \Phi) \leq \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi) \\ J(\tau, \eta, \hat{\alpha}, Z^*, \Phi) \geq \inf_{Z \in \mathcal{N}_\pi} J(\tau, \eta, \alpha, Z, \Phi) \\ J(\tau, \eta, Y^*, \hat{\beta}, \Phi) = J(\tau, \eta, \hat{\alpha}, Z^*, \Phi) \end{cases}$$

holds for some $Y^* \in \mathcal{M}$ and $Z^* \in \mathcal{N}_\pi$. Thus, (5.37) and (5.38) complete the proof.

For $\pi = \{0 = t_0 < \dots < t_N = T\}$ and $t \in [t_j, t_{j+1})$, we denote $\pi(t) = \{0 < t_{j+1} - t < t_{j+2} - t < \dots < t_N - t\}$. Putting

$$V_\pi(\tau, \eta, \Phi) = \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi),$$

(5.29) yields the principle of dynamic programming below,

$$(5.39) \quad V_\pi(\tau, \eta, \Phi) = V_\pi(t_j, \eta, V_{\pi(t_j)}(\tau - t_j, \cdot, \Phi)).$$

Namely, for $0 \leq s \leq T - t_j$,

$$(5.40) \quad V_\pi(s + t_j, \eta, \Phi) = V_\pi(t_j, \eta, V_{\pi(t_j)}(s, \cdot, \Phi))$$

holds.

6. Principle of dynamic programming

For partitions π and $\hat{\pi}$, we say $\pi \leq \hat{\pi}$ if $\mathcal{P}_\pi \subset \mathcal{P}_{\hat{\pi}}$, where \mathcal{P}_π denotes the set of division points of π (see Section 4).

PROPOSITION 6.1. Suppose $\pi_n \leq \pi_{n+1}$, $n = 1, 2, \dots$, and put

$$(6.1) \quad V(\tau, \eta, \Phi) = \inf_{\beta \in \bigcup_n \Delta_{\pi_n}} \sup_{Y \in \mathcal{M}} J(\tau, \eta, Y, \beta, \Phi).$$

Then,

$$(6.2) \quad V_{\pi_n}(\cdot, \Phi) \geq V_{\pi_{n+1}}(\cdot, \Phi) \text{ and } V_{\pi_n}(\tau, \eta, \Phi) \rightarrow V(\tau, \eta, \Phi) \text{ for any } \tau, \eta.$$

Proof is easy.

In particular, Theorem 4.1 implies, for a tame function φ

$$(6.3) \quad V_{\pi_n}(\cdot, \varphi) \rightarrow V(\cdot, \varphi) \text{ uniformly in any bounded set of } [0, T] \times H,$$

as $n \rightarrow \infty$.

Moreover we will prove

THEOREM 6.1. Let $\pi_n < \pi_{n+1}$ and $\lim \|\pi_n\| = 0$. Then, $V(\cdot, \Phi)$ of (6.3) does not depend on a sequence $\{\pi_n\}$.

First we will prove the following lemma.

LEMMA. Let $\mathcal{P}_\pi = \{t_j, j = 0, \dots, N\}$, $\mathcal{P}_{\hat{\pi}} = \mathcal{P}_\pi \cup \{\tau\}$ and $t_{p-1} < \tau < t_p$. Then there is a constant $k = k(\varphi)$, independent of π , such that

$$(6.4) \quad |V_\pi(t, \eta, \varphi) - V_{\hat{\pi}}(t, \eta, \varphi)| \leq k\sqrt{t_p - t_{p-1}} (1 + \|\eta\|) \leq k\sqrt{\|\pi\|} (1 + \|\eta\|).$$

Proof of Lemma. For $t \in (t_{j-1}, t_j]$, $j > p$, we put

$$(6.5) \quad \bar{\Phi} = \mathbf{v}(t_{p+1} - t_p)\mathbf{v}(t_{p+2} - t_{p+1}) \cdots \mathbf{v}(t - t_{j-1})\varphi.$$

Since π and $\hat{\pi}$ have the same division points on $[t_p, t]$, (5.29) implies

$$(6.6) \quad \Phi(\eta) = \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(t - t_p, \eta, Y, \beta, \varphi) = \inf_{\beta \in \Delta_{\hat{\pi}}} \sup_{Y \in \mathcal{M}} J(t - t_p, \eta, Y, \beta, \varphi),$$

$$(6.7) \quad \mathbf{v}(t_p - t_{p+1})\Phi(\eta) = \inf_{\beta \in \Delta_\pi} \sup_{Y \in \mathcal{M}} J(t - t_{p-1}, \eta, Y, \beta, \varphi), \quad \text{say } \Phi^*(\eta),$$

and

$$(6.8) \quad \mathbf{v}(t - t_{p+1})\mathbf{v}(t_p - t)\bar{\Phi}(\eta) = \inf_{\beta \in \Delta_{\hat{\pi}}} \sup_{Y \in \mathcal{M}} J(t - t_{p-1}, \eta, Y, \beta, \varphi), \quad \text{say } \hat{\Phi}(\eta).$$

Therefore, we see, from (4.3),

$$(6.9) \quad |\Phi^*(\eta) - \Phi(\eta)| \leq C_3(\varphi) \sqrt{t_p - t_{p-1}} (1 + \|\eta\|)$$

and

$$(6.10) \quad |\hat{\Phi}(\eta) - \Phi(\eta)| \leq C_3(\varphi) \sqrt{t_p - t_{p-1}} (1 + \|\eta\|).$$

Again same arguments yield

$$(6.11) \quad \begin{aligned} & |V_\pi(t, \eta, \varphi) - \mathbf{v}(t_1) \cdots \mathbf{v}(t_{p-1} - t_{p-2})\Phi(\eta)| \\ &= |\mathbf{v}(t_1) \cdots \mathbf{v}(t_{p-1} - t_{p-2})\Phi^*(\eta) - \mathbf{v}(t_1) \cdots \mathbf{v}(t_{p-1} - t_{p-2})\Phi(\eta)| \\ &\leq \sup_{Y \in \mathcal{M}, Z \in \mathcal{N}} E |\Phi^*(\xi(t_{p-1}, \eta, Y, Z)) - \Phi(\xi(t_{p-1}, \eta, Y, Z))| \\ &\leq C_3(\varphi) \sqrt{t_p - t_{p-1}} (1 + \sup_{Y, Z} E \|\xi(t_{p-1}, \eta, Y, Z)\|) \leq \hat{k}(\varphi) \sqrt{t_p - t_{p-1}} (1 + \|\eta\|) \end{aligned}$$

with a constant $\hat{k}(\varphi)$ by (2.3), and similarly for $\hat{\pi}$. This concludes the proof.

More generally, we obtain

$$(6.12) \quad |V_\pi(t, \eta, \varphi) - V_{\hat{\pi}}(t, \eta, \varphi)| \leq k^*(\varphi) \sqrt{\|\pi\|} (1 + \|\eta\|)$$

with a constant $k^*(\varphi)$, if $\mathcal{P}_{\hat{\pi}} = \mathcal{P}_\pi \cup \{\tau_1, \tau_2, \dots, \tau_q\}$.

Proof of Theorem 6.1. Let $\pi \vee \hat{\pi}$ be the partition with division points $\mathcal{P}_\pi \cup \mathcal{P}_{\hat{\pi}}$. Setting $\pi_{n,k} = \pi_\pi \vee \hat{\pi}_k$, $\{\pi_{n,k}, k = 1, 2, \dots\}$ and $\{\pi_{n,k}, n = 1, 2, \dots\}$ again satisfy the condition of Theorem. Put $V = \lim V_{\pi_n}$, $V_k = \lim_{n \rightarrow \infty} V_{\pi_{n,k}}$, $\hat{V} = \lim V_{\hat{\pi}_n}$ and $\hat{V}_n = \lim_{k \rightarrow \infty} V_{\pi_{n,k}}$. Then

$$(6.13) \quad V_k \leq V_{\pi_{n,k}}$$

holds. Moreover, (6.12) implies

$$(6.14) \quad |V_{\pi_n}(t, \eta, \varphi) - V_{\pi_{n,k}}(t, \eta, \varphi)| \leq \hat{k}(\varphi) \sqrt{\|\pi_n\|} (1 + \|\eta\|).$$

Hence, letting n to ∞ , we get

$$(6.15) \quad V = V_k, \quad k = 1, 2, \dots.$$

In the same way, we see

$$(6.16) \quad \hat{V} = \hat{V}_n, \quad n = 1, 2, \dots.$$

Together with (6.13), we have

$$(6.17) \quad V \leq V_{\pi_{n,k}} \quad \text{for any } n, k.$$

As $k \rightarrow \infty$, (6.16) and (6.17) imply

$$(6.18) \quad V \leq \hat{V}_n = \hat{V}.$$

This completes the proof of Theorem 6.1, since we can see the converse inequality in the same way.

Now we will verify the principle of dynamic programming for V with a tame terminal function φ .

THEOREM 6.2. V satisfies the principle of dynamic programming, namely

$$(6.19) \quad V(t + s, \eta, \varphi) = V(t, \eta, V(s, \cdot, \varphi)).$$

Proof. Suppose that $\pi_n < \pi_{n+1}$ and $\|\pi_n\| \rightarrow 0$. Appealing to Theorem 6.1, we may assume $t \in \mathcal{P}_{\pi_n}$, $n = 1, 2, \dots$, for simplicity. Now the principle of dynamic programming (5.29) yields

$$(6.20) \quad V_n(t + s, \eta, \varphi) = V_n(t, \eta, V_n(s, \cdot, \varphi))$$

where $V_n = V_{\pi_n}$.

On the other hand $V_n(\cdot, \varphi)$ is decreasing to $V(\cdot, \varphi)$ uniformly in any bounded set of $[0, T] \times H$. Thus, for any $\varepsilon > 0$ and $r > 0$, there is a large $N = N(\varepsilon, r)$, such that, for $n > N$,

$$(6.21) \quad |V_\pi(t, \eta, V_n(s, \cdot, \varphi)) - V_\pi(t, \eta, V(s, \cdot, \varphi))| \\ \leq \sup_{Y, Z} |J(t, \eta, Y, Z, V_n(s, \cdot, \varphi)) - J(t, \eta, Y, Z, V(s, \cdot, \varphi))|$$

$$\begin{aligned}
&\leq \sup_{Y,Z} E | V_n(s, \xi(t, \eta, Y, Z), \varphi) - V(s, \xi(t, \eta, Y, Z), \varphi) | \\
&\leq \varepsilon + k_3(\varphi) \sup_{Y,Z} E(1 + \|\xi(t, \eta, Y, Z)\|; \|\xi(t, \eta, Y, Z)\| \geq r) \\
&\leq \varepsilon + k_4(\varphi) (1 + \|\dot{\eta}\|) \|\eta\| r^{-1}
\end{aligned}$$

holds. So, as $n \rightarrow \infty$, $V_n(\cdot, \cdot, \varphi)$ is decreasing to $V(\cdot, \cdot, \varphi)$ uniformly in any bounded set of $[0, T] \times H$ and partition π . This fact together with (6.2), implies

$$(6.22) \quad V_n(t, \eta, V_n(s, \cdot, \varphi)) \rightarrow V(t, \eta, V(s, \cdot, \varphi)) \text{ for any } t \text{ and } \eta.$$

Now we can conclude the proof by (6.20).

Remark. We introduce the lower value function in the same way as the upper one, putting

$$W_\pi(t, \eta, \Phi) = \sup_{\alpha \in \Gamma_\pi} \inf_{Z \in \mathcal{N}} J(t, \eta, \alpha, Z, \Phi).$$

Under the same condition of Theorem 6.1, W_{π_n} is increasing to W , defined by

$$W(t, \eta, \Phi) = \sup_{\alpha \in \cup_n \Gamma_{\pi_n}} \inf_{Z \in \mathcal{N}} J(t, \eta, \alpha, Z, \Phi).$$

For a tame terminal function φ , $W(t, \eta, \varphi)$ is independent of a sequence $\{\pi_n\}$ and (6.19) holds.

When V and W coincide, we call it the value function.

7. Viscosity solution of min-max equation

We will study the Cauchy problem of min-max equation (7.1) arising in our stochastic differential game.

$$(7.1) \quad \begin{cases} \frac{\partial}{\partial t} u(t, \eta) + F(D^2 u(t, \eta), Du(t, \eta), \eta) = 0 & \text{in } (0, T) \times H \\ u(0) = \varphi, \text{ (= tame function),} \end{cases}$$

where $D =$ Fréchet derivative in H and

$$(7.2) \quad F(S, p, \eta) = - \min_{z \in \mathcal{Z}} \max_{y \in \mathcal{Y}} \left[\frac{1}{2} (Sf(y, z)\eta, f(y, z)\eta) + \langle p, (L(y, z) - cI)\eta \rangle + h(\eta, y, z) \right]$$

$$= \max_{z \in \mathcal{Z}} \min_{y \in \mathcal{Y}} \left[\frac{1}{2} (-Sf(y, z)\eta, f(y, z)\eta) + \langle \phi, A(y, z)\eta \rangle - h(\eta, y, z) \right]$$

where $\langle \cdot, \cdot \rangle =$ duality pairing between H^2 and H^{-2} under $H = H^*$ and $A(y, z) = -L(y, z) + cI$.

Now we will define a viscosity subsolution (resp. supersolution) of (7.1), according to [2. part 4] and [8. part 2]. Since the min-max equation is more general than Bellman equation, our definition is slightly different from theirs. Let us set

$$\mathcal{D} = \{\phi \in C^{1,2}((0, T) \times H) \cap C([0, T] \times H) ; \text{ weakly lower}$$

semi-continuous (WLSC in short) and the following (7.3)~(7.5) hold,

$$(7.3) \quad \frac{\partial \phi}{\partial t} \text{ is Lipschitz continuous}$$

$$(7.4) \quad D\phi \text{ is Lipschitz continuous and a continuous mapping from } (0, T) \times H \text{ into } H^2 \text{ with } \sup_{t \leq T} \|D\phi(t, \cdot)\|_2 < \infty$$

$$(7.5) \quad (D^2\phi(t, \zeta)\eta, \eta) \text{ is uniformly continuous in any bounded set of } (0, T) \times H \times H\}.$$

DEFINITION 7.1. $u \in UC_{loc}([0, T] \times H)$ is called a viscosity subsolution of (7.1) if

(7.6) u is WUSC (weakly upper semi-continuous) on $(0, T) \times H$ and linear growth,

(7.7) for any $\lambda > 0$ and $\phi \in \mathcal{D}$, the following inequality holds at each global maximum point $(t, \zeta) \in (0, T) \times H$ of $u(\theta, \eta) - \phi(\theta, \eta) - \lambda \|\eta\|^2$,

$$\frac{\partial \phi}{\partial t}(t, \zeta) + F(D^2\phi(t, \zeta) + 2\lambda I, D\phi(t, \zeta), \zeta) \leq -2\lambda(c - \lambda_0) \|\zeta\|^2$$

with λ_0 of (2.7), and

$$(7.8) \quad u(0) = \varphi.$$

DEFINITION 7.2. $u \in UC_{loc}([0, T] \times H)$ is called a supersolution of (7.1), if

(7.9) u is WLSC on $(0, T) \times H$ and linear growth,

(7.10) for any $\lambda > 0$ and $\phi \in \mathcal{D}$, the following inequality holds at each global minimum point $(t, \zeta) \in (0, T) \times H$ of $u(\theta, \eta) + \phi(\theta, \eta) + \lambda \|\eta\|^2$,

$$-\frac{\partial \phi}{\partial t}(t, \zeta) + F(-D^2\phi(t, \zeta) - 2\lambda I, -D\phi(t, \zeta), \zeta) \geq 2\lambda(c - \lambda_0) \|\zeta\|^2$$

with λ_0 of (2.7), and

$$(7.11) \quad u(0) = \varphi.$$

u is a viscosity solution, if it is both a viscosity subsolution and a viscosity supersolution. Hereafter, we suppress viscosity, for simplicity, and call viscosity sub- and super- solutions just sub- and super- solutions, respectively.

Now we state the following theorem.

THEOREM 7.1. *The upper value function $V(\cdot, \varphi)$ is a viscosity solution of min-max equation (7.1).*

Proof. Since $V \in \text{LUC}([0, T] \times H) \cap C([0, T] \times H_w)$, $V(\cdot) - \lambda \|\cdot\|^2$ is weakly USC and bounded above. Suppose that $\phi \in \mathcal{D}$ and $V(\theta, \eta) - \phi(\theta, \eta) - \lambda \|\eta\|^2$ has a global maximum at $(t, \zeta) \in (0, T) \times H$. Now let us assume that

$$(7.12) \quad V(\cdot) = V(\cdot, \varphi) = \lim V_{\pi_n}(\cdot, \varphi),$$

recalling Theorem 6.1. Since V satisfies the principle of dynamic programming, we have

$$(7.13) \quad 0 = \inf_{\beta \in \cup_n \Delta_{\pi_n}} \sup_{Y \in \mathcal{M}} E \int_0^\theta h(\xi(\tau), Y, \beta) d\tau + V(t - \theta, \xi(\theta)) - V(t, \zeta) \\ \leq \inf_{\beta} \sup_Y E \int_0^\theta h(\xi(\tau), Y, \beta) d\tau + \phi(t - \theta, \xi(\theta)) - \phi(t, \zeta) + \lambda (\|\xi(\theta)\|^2 - \|\zeta\|^2)$$

where $\xi(\tau) = \xi(\tau, \zeta, Y, \beta)$. Thus, Itô's formula yields

$$(7.14) \quad 0 \leq \inf_{\beta} \sup_Y E \left[\int_0^\theta h(\xi(\tau), Y, \beta) - \frac{\partial \phi}{\partial t}(t - \tau, \xi(\tau)) \right. \\ \left. - \langle D\phi(t - \tau, \xi(\tau)), A(Y, \beta)\xi(\tau) \rangle \right. \\ \left. + \frac{1}{2} (D^2\phi(t - \tau, \xi(\tau)) (f(Y, \beta)\xi(\tau)), f(Y, \beta)\xi(\tau)) \right. \\ \left. - 2\lambda \langle \xi(\tau), A(Y, \beta)\xi(\tau) \rangle + \|f(Y, \beta)\xi(\tau)\|^2 d\tau \right].$$

Using (7.3)~(7.5) and Proposition 3.1, we evaluate each term of (7.14), namely we have (7.15) ~ (7.22) below, putting $m = \sup_{t \leq T} \|D\phi(t, \cdot)\|_2$, $a = \frac{\partial \phi}{\partial t}(t, \zeta)$, $p = D\phi(t, \zeta)$, $S = D^2\phi(t, \zeta)$ and $M(\tau, \zeta^2) = \tau \|\zeta\|^2 + \int g_2(x, \tau) \|\Delta_x \zeta\|^2 dx$.

$$(7.15) \quad E \left| \frac{\partial \Psi}{\partial t}(t - \tau, \xi(\tau)) - a \right| \leq k_1(|\tau| + M(\tau, \zeta))$$

$$\begin{aligned}
(7.16) \quad & E | \langle D\phi(t - \tau, \xi(\tau)), A(Y, \beta)(\xi(\tau) - \zeta) \rangle | \\
& \leq E \| D\phi(t - \tau, \xi(\tau)) \|_2 \| A(Y, \beta)(\xi(\tau) - \zeta) \|_{-2} \\
& \leq k_2 m E \| \xi(\tau) - \zeta \| \leq k_3 M(\tau, \zeta).
\end{aligned}$$

Since, for any $\varepsilon > 0$, there is $\delta = \delta(\varepsilon, t, \zeta) > 0$ such that $\| D\phi(t - \tau, \eta) - p \|_2 < \varepsilon$, whenever $|\tau| < \delta$ and $\| \eta - \phi \| < \delta$, we get

$$\begin{aligned}
(7.17) \quad & E \| \langle D\phi(t - \tau, \xi(\tau)) - p \|_2 < \varepsilon + 2mP(\| \xi(\tau) - \zeta \| > \delta) \\
& < \varepsilon + 2mM^2(\tau, \zeta) / \delta^2.
\end{aligned}$$

Denoting the Lipschitz constant of $D\phi$ by r , we see

$$(7.18) \quad | (D^2\phi(\theta, \xi)\eta, \eta) | \leq r \| \eta \|^2.$$

Since, for any $\varepsilon > 0$ and $R > 0$, there is $\delta_0 = \delta_0(\varepsilon, R) > 0$ such that

$$| (D^2\phi(\theta, \xi)\eta, \eta) - (D^2\phi(\bar{\theta}, \bar{\xi})\bar{\eta}, \bar{\eta}) | < \varepsilon$$

whenever $|\theta - \bar{\theta}| < \delta_0$, $\| \xi - \bar{\xi} \| < \delta_0$, $\| \eta - \bar{\eta} \| < \delta_0$, and $\| \xi \|, \| \bar{\xi} \|, \| \eta \|, \| \bar{\eta} \| \leq R$, we get

$$\begin{aligned}
(7.19) \quad & E | (D^2\phi(t - \tau, \xi(\tau))(f(Y, \beta)\zeta), f(Y, \beta)\zeta) - (S(f(Y, \beta)\zeta), f(Y, \beta)\zeta) | \\
& < \varepsilon + 2r | f |_\infty^2 \| \zeta \|^2 (P(\| \xi(\tau) \| > R) + P(\| \xi(\tau) - \zeta \| > \delta_0)) \\
& < \varepsilon + 2r | f |_\infty^2 \| \zeta \|^2 ((E \| \xi(\tau) \|^2 / R^2) + (M^2(\tau, \zeta) / \delta_0^2))
\end{aligned}$$

for a large $R > | f |_\infty \| \zeta \|$.

$$\begin{aligned}
(7.20) \quad & E | (D^2\phi(t - \tau, \xi(\tau))(f(Y, \beta)\xi(\tau)), f(Y, \beta)\xi(\tau)) \\
& \quad - (D^2\phi(t - \tau, \xi(\tau))(f(Y, \beta)\zeta), f(Y, \beta)\zeta) | \\
& < r | f |_\infty^2 (E \| \xi(\tau) - \zeta \|^2 + 2 \| \zeta \| E \| \xi(\tau) - \zeta \|) \\
& < r | f |_\infty^2 (M^2(\tau, \zeta) + 2 \| \zeta \| M(\tau, \zeta)).
\end{aligned}$$

$$\begin{aligned}
(7.21) \quad & - 2\lambda E \langle \xi(\tau), A(Y, \beta)\xi(\tau) \rangle \\
& = - 2\lambda E [c \| \xi(\tau) \|^2 - \langle \xi(\tau), L(Y, \beta)\xi(\tau) \rangle] \\
& \leq 2\lambda(\lambda_0 - c) E \| \xi(\tau) \|^2 = 2\lambda(\lambda_0 - c) [\| \zeta \|^2 + E(\| \xi(\tau) \|^2 - \| \zeta \|^2)] \\
& \leq 2\lambda(\lambda_0 - c) \| \zeta \|^2 + k_4 \| \zeta \| M(\tau, \zeta).
\end{aligned}$$

$$(7.22) \quad E | \| f(Y, \beta)\zeta \xi(\tau) \|^2 - \| f(Y, \beta)\zeta \|^2 | \leq k_5 \| \zeta \| M(\tau, \zeta).$$

Inserting above (7.15)~(7.22) into (7.14), we obtain

$$(7.23) \quad 0 \leq \inf_{\beta} \sup_Y E \left[\int_0^{\theta} h(\zeta, Y, \beta) - a - \langle p, A(Y, \beta) \zeta \rangle \right. \\ \left. + \frac{1}{2} ((S + 2\lambda D) f(Y, \beta) \zeta, f(Y, \beta) \zeta) d\tau \right] \\ + 2\lambda(\lambda_0 - c) \|\zeta\|^2 \theta + o(\theta).$$

Next we calculate the main term of (7.23), putting

$$(7.24) \quad K(y, z) = h(\zeta, y, z) - a - \langle p, A(y, z) \zeta \rangle \\ + \frac{1}{2} ((S + 2\lambda D) f(y, z) \zeta, f(y, z) \zeta).$$

Since ζ is fixed, $K(y, z)$ is constant in H . So, it can be regarded as a tame function. Let us consider a differential game, using K instead of h . Putting

$$(7.25) \quad \tilde{V}_{\pi}(\theta, \eta, \varphi) = \inf_{\beta \in \Delta_{\pi}} \sup_{Y \in \mathcal{M}} E \int_0^{\theta} K(Y(s), \beta(Y)(s)) ds + \varphi(\xi(\theta, \eta, Y, \beta))$$

and $\mathcal{P}_{\pi} = \{t_p, p = 0, \dots, N\}$, the principle of dynamic programming (5.39) asserts

$$(7.26) \quad \tilde{V}_{\pi}(\theta, \eta, \varphi) = \tilde{V}_{\pi}(t_p, \eta, \tilde{V}_{\pi(t_p)}(\theta - t_p, \cdot, \varphi)).$$

Setting $\varphi = 0$, we have, for $\theta \in [t_p, t_{p+1})$ and $\tilde{\zeta} \in H$

$$(7.27) \quad \tilde{V}_{\pi(t_p)}(\theta - t_p, \tilde{\zeta}, 0) = \min_{z \in \mathcal{Z}} \sup_{Y \in \mathcal{M}} E \int_0^{\theta - t_p} K(Y(s), z) ds.$$

Noting

$$(7.28) \quad \sup_{Y \in \mathcal{M}} E \int_0^t K(Y(s), z) ds \leq \int_0^t \max_{Y \in \mathcal{Y}} K(Y, z) ds = \max_{y \in \mathcal{Y}} K(y, z) t \\ = \max_{y \in \mathcal{Y}} E \int_0^t K(y, z) ds \leq \sup_{Y \in \mathcal{M}} E \int_0^t K(Y(s), z) ds,$$

we get

$$(7.29) \quad \tilde{V}_{\pi(t_p)}(\theta - t_p, \tilde{\zeta}, 0) = \min_{z \in \mathcal{Z}} \max_{Y \in \mathcal{Y}} K(y, z) (\theta - t_p) = \mu(\theta - t_p)$$

where $\mu = \min_{z \in \mathcal{Z}} \max_{Y \in \mathcal{Y}} K(y, z)$. Again it follows from (5.39) that

$$(7.30) \quad \tilde{V}_{\pi}(t_p, \eta, \mu(\theta - t_p)) = \tilde{V}_{\pi}(t_{p-1}, \eta, \hat{V}_{\pi(t_{p-1})}(t_p - t_{p-1}, \cdot, \mu(\theta - t_p))),$$

and

$$\tilde{V}_\pi(t_{p-1})(t, \tilde{\zeta}, \mu(\theta - t_p)) = \mu t + \mu(\theta - t_p).$$

Thus, we obtain

$$\tilde{V}_\pi(\theta, \eta, 0) = \mu\theta.$$

Therefore

$$(7.31) \quad \inf_{\beta} \sup_Y E \int_0^\theta K(Y(s), \beta(Y)(s)) ds = \lim_{n \rightarrow \infty} \tilde{V}_{\pi_n}(\theta, \eta, 0) = \mu\theta,$$

holds. Now, (7.23) concludes

$$(7.32) \quad -2\lambda(\lambda_0 - c) \|\zeta\|^2 \leq \mu\theta.$$

Recalling the definition of μ , (7.32) asserts that V is a subsolution.

Since we can prove that V is a supersolution in the similar way, the proof is completed.

In the finite dimensional case (see [3], [5]), a subsolution (resp. supersolution) can be defined by super differential (resp. subdifferential) instead of test functions. But, in our case, this equivalence is open. Here we remark on super differential J^+u (resp. subdifferential J^-u) for $u \in \text{UC}_{\text{loc}}((0, T) \times H)$. J^+u is defined by (7.33)

$$(7.33) \quad J^+u(t, \zeta) = \{(a, p, S) \in R^1 \times H^2 \times \mathcal{L}; \\ u(t + \theta, \zeta + \eta) \leq u(t, \zeta) + a\theta + (p, \eta) + \frac{1}{2}(S\eta, \eta) + o(|\theta| + \|\eta\|^2)\} \\ \text{for } (t, \zeta) \in (0, T) \times H,$$

where \mathcal{L} = space of all continuous linear operators from H to $H^*(=H)$ ($J^-(u)$ is defined by (7.33) with the opposite inequality).

Remark. Let $u \in \text{UC}_{\text{loc}}([0, T] \times H)$ be weakly USC and bounded above, say $M = \sup_{\theta, \eta} u(\theta, \eta)$. Suppose

$$(7.34) \quad \sup_{\theta \in [0, T]} u(\theta, \eta) \rightarrow -\infty, \text{ as } \|\eta\| \rightarrow \infty.$$

Let us fix $(t, \zeta) \in (0, T) \times H$ and $\delta > 0$ arbitrarily. Then, for any $\alpha > (M - u(t, \zeta) + 1) / \delta$,

$$(7.35) \quad u(\theta, \eta) - \alpha(\|\eta - \zeta\|_{-1}^2 + |\theta - t|^2) < u(t, \zeta) - 1$$

holds, whenever $\|\eta - \zeta\|_{-1}^2 + |\theta - t|^2 > \delta$. On the other hand, it follows from (7.34) that there is a positive number m , such that

$$(7.36) \quad u(\theta, \eta) < u(t, \zeta) - 1, \quad \text{if } \|\eta\| > m.$$

But, $\|\cdot\|_{-1}$ is WLSC, because $\|\eta\|_{-1} = \sup\{(\eta, e) ; e \in H_1 \text{ and } \|e\| = 1\}$. Hence $u - \alpha(\|\cdot - \zeta\|_{-1}^2 + |\cdot - t|^2)$ has a global maximum point $(\bar{\theta}, \bar{\zeta})$, such that

$$(7.37) \quad \|\bar{\zeta}\| \leq m \text{ and } \|\bar{\zeta} - \zeta\|_{-1}^2 + |\bar{\theta} - t|^2 < \delta.$$

Therefore, setting $B = (I - \Delta)^{-1}$, where $\Delta =$ Laplacian operator,

$$u(\bar{\theta} + \theta, \bar{\zeta} + \eta) \leq u(\bar{\theta}, \bar{\zeta}) + 2\alpha(\bar{\theta} - t)\theta + 2\alpha(B(\bar{\zeta} - \zeta), \eta) + \alpha(B\eta, \eta) + \alpha\theta^2$$

holds. Noting $B(\bar{\zeta} - \zeta) \in H^2$ and $B \in \mathcal{L}$ we get

$$(7.38) \quad (2\alpha(\bar{\theta} - t), 2\alpha B(\bar{\zeta} - \zeta), \alpha B) \in J^+ u(\bar{\theta}, \bar{\zeta}).$$

Since t, ζ and δ are arbitrary, (7.37) and (7.38) yield that $\{(\theta, \eta) \in (0, T) \times H ; J^+ u(\theta, \eta) \neq \emptyset\}$ is dense in $R^1 \times H^{-1}$ -topology.

8. Example

In this section, we will deal with a simple example, where the upper value function is a unique viscosity solution in a class of finite dimensional functions on H .

Setting $\partial_i = \frac{\partial}{\partial x_i}$, let us consider SPDE (8.1) below,

$$(8.1) \quad d\xi(x, t) = \sum_{i=1}^n \sigma^i(Y(t), Z(t)) (\partial_i^2 - (x_i^2/4)) \xi(x, t) dt + f(Y(t), Z(t)) \xi(x, t) dW(t),$$

with initial condition

$$\xi(x, 0) = \eta(x).$$

Although SPDE (8.1) does not satisfy the condition; $c =$ constant, we can easily see the same results, using an Hermite base of H . Let us assume (a1)~(a3) below.

(a1) $\sigma^i(y, z)$ and $f(y, z)$ are Lipschitz continuous in $\mathcal{Y} \times \mathcal{Z}$,

(a2) there is a positive number λ^* such that

$$\sigma^i(y, z) \geq \lambda^* \text{ for any } y \text{ and } z,$$

(a3) $h(\cdot, y, z)$ and an initial function φ are tame functions of Hermite base e_k

and bounded.

Here the Hermite base is defined as follows. For a multi-index $\underline{k} = (k_1, \dots, k_n)$, $k_i \geq 0$.

$$(8.2) \quad \mathbf{e}_{\underline{k}}(\underline{x}) = \prod_{i=1}^n \mathbf{e}_{k_i}(x_i) \quad \text{for } \underline{x} = (x_1, \dots, x_n) \in R^n,$$

where $\mathbf{e}_0(x) = (2\pi)^{-1/4} \exp\left(-\frac{x^2}{4}\right)$ for $x \in R^1$, and

$$\mathbf{e}_m(x) = (-1)^m (\sqrt{2\pi} m!)^{-1/2} \left[\exp\left(\frac{x^2}{4}\right) \right] \left(\frac{d}{dx} \right)^m \exp\left(-\frac{x^2}{2}\right) \text{ for } x \in R^1, m = 1, 2, \dots.$$

Appealing to the following fact,

$$\frac{d}{dx} \mathbf{e}_m(x) = \frac{1}{2} (\sqrt{m+1} \mathbf{e}_{m+1}(x) - \sqrt{m} \mathbf{e}_{m-1}(x)),$$

we have, for $\eta = \sum c_{\underline{k}} \mathbf{e}_{\underline{k}} \in H$,

$$(8.3) \quad \partial_i \eta = \frac{1}{2} (\sum c_{\underline{k}} \sqrt{k_i + 1} \mathbf{e}_{\underline{k} + I_i} - \sum c_{\underline{k}} \mathbf{e}_{\underline{k} - I_i})$$

where $\underline{k} \pm I_i = (k_1, \dots, k_i \pm 1, \dots, k_n)$. Therefore $\partial_i \eta \in H$, if and only if $\sum c_{\underline{k}}^2 k_i < \infty$. Repeating this argument, we see

PROPOSITION 8.1.

If $\sum c_{\underline{k}}^2 |\underline{k}|^p < \infty$, then $\eta \in H^p$, $p = 0, 1, 2, \dots$.

Moreover, \mathbf{e}_m satisfies the following equation

$$(8.4) \quad \mathbf{e}_m'' - (x^2/4) \mathbf{e}_m = -\left(m + \frac{1}{2}\right) \mathbf{e}_m, \quad m = 0, 1, 2, \dots.$$

Hence, employing the formal expansion of $\xi(t)$, say

$$(8.5) \quad \xi(t) = \sum X_{\underline{k}}(t) \mathbf{e}_{\underline{k}}$$

we have SDE

$$(8.6) \quad \begin{cases} dX_{\underline{k}}(t) = -\left(\sum_{i=1}^n \sigma^i(Y(t), Z(t)) \left(k_i + \frac{1}{2}\right)\right) dt + f(Y(t), Z(t)) X_{\underline{k}}(t) dW(t) \\ X_{\underline{k}}(0) = (\eta, \mathbf{e}_{\underline{k}}). \end{cases}$$

(8.6) has a unique solution with continuous path and

$$(8.7) \quad E |X_k(t)|^2 \leq |X_k(0)|^2 \exp\left(-2\lambda^*\left(|k| + \frac{n}{2} + |f|_\infty^2\right)t\right)$$

holds, where $|k| = \sum k_i$ and $|f|_\infty = \sup_{y,z} |f(y, z)|$.

Now for ξ of (8.5), we can easily see that $\xi \in L^2(\Omega \times (0, T); H^2) \cap L^2(\Omega; C([0, T]; H))$ and ξ provides a unique solution of (8.1). Moreover, the evaluation (2.3)~(2.5) and Proposition 3.1 hold. Therefore the upper value function $V(\cdot, \varphi)$ is a viscosity solution of the min-max equation (8.8) below.

$$(8.8) \quad \begin{cases} \frac{\partial u}{\partial t}(t, \eta) - \min_{z \in \mathcal{Z}} \max_{y \in \mathcal{Y}} \left[\frac{1}{2} |f(y, z)|^2 (D^2 u(t, \eta) \eta, \eta) \right. \\ \left. + \langle Du(t, \eta), L(y, z) \eta \rangle + h(\eta, y, z) \right] = 0 \\ u(0, \eta) = \varphi(\eta), \end{cases}$$

where

$$(8.9) \quad L(y, z) = \sum_{i=1}^n \sigma^i(y, z) (\partial_i^2 - (x_i^2/4)).$$

By (a3), h and φ can be written by

$$h(\eta, y, z) = \hat{h}((\eta, e_1), \dots, (\eta, e_N), y, z)$$

and

$$\varphi(\eta) = \hat{\varphi}((\eta, e_1), \dots, (\eta, e_N))$$

where $e_j = e_k$ for $k = k^j = (k_1^j, \dots, k_n^j)$. Let us set $P =$ orthogonal projection onto the linear space generated by (e_1, \dots, e_N) . Then, (8.6) derives that $V(t, \eta, \varphi)$ depends only on $((\eta, e_1), \dots, (\eta, e_N))$, namely

$$(8.10) \quad V(t, \eta, \varphi) = V(t, P\eta, \varphi).$$

Define v by

$$(8.11) \quad v(t, b_1, \dots, b_N) = V(t, b_1 e_1 + \dots + b_N e_N; \varphi).$$

Again (8.6) implies that v is a unique viscosity solution of the following finite dimensional min-max equation (8.12), and bounded by (a3),

$$(8.12) \quad \begin{cases} \frac{\partial v}{\partial t}(t, b) - \min_z \max_y \left[\frac{1}{2} f^2(y, z) \sum_{i,j=1}^N \frac{\partial^2 v}{\partial b_i \partial b_j}(t, b) b_i b_j \right. \\ \left. + \sum_{j=1}^N \sum_{i=1}^n \sigma^i(y, z) \left(k_i^j + \frac{1}{2} \right) b_j \frac{\partial v}{\partial b_j}(t, b) + \hat{h}(b, y, z) \right] = 0 \\ v(0, b) = \hat{\varphi}(b). \end{cases}$$

From (8.10), we see $V(t, \eta, \varphi) = v(t, (\eta, e_1), \dots, (\eta, e_N))$.

THEOREM 8.1. *Suppose that $W \in \text{BUC}([0, T] \times H)$ is a viscosity solution of (8.8) and $W(t, \eta) = W(t, P\eta)$. Then $W = V$.*

Proof. Suppose that $\hat{\psi} \in C^{1,2}((0, T) \times R^N)$ has the two properties below,

(i) $\frac{\partial \hat{\psi}}{\partial t}$ is Lipschitz continuous

and

(ii) $\frac{\partial \hat{\psi}}{\partial x_1}$ and $\frac{\partial^2 \hat{\psi}}{\partial x_i \partial x_j}$ are bounded.

Put $\phi(t, \eta) = \hat{\psi}(t, (\eta, \mathbf{e}_1), \dots, (\eta, \mathbf{e}_N))$. Then $\phi \in \mathcal{D}$ by (i) and (ii).

The function \hat{w} , defined by

$$\hat{w}(t, b_1, \dots, b_N) = W(t, b_1 \mathbf{e}_1 + \dots + b_N \mathbf{e}_N)$$

is continuous, since W is weakly continuous, and

$$(8.13) \quad W(t, \eta) = \hat{w}(t, (\eta, \mathbf{e}_1), \dots, (\eta, \mathbf{e}_N))$$

holds.

Suppose that $\hat{w} - \hat{\psi}$ has a unique maximum at $(\hat{t}, \hat{\mathbf{b}}) \in (0, T) \times R^N$, and $W(t, \eta) - \phi(t, \eta) - \lambda \|\eta\|^2$ has a global maximum at $(t_\lambda, \eta_\lambda)$. Appealing to $\|P\eta\| \leq \|\eta\|$, we have

$$(8.14) \quad W(t, \eta) - \phi(t, \eta) - \lambda \|\eta\|^2 \leq W(t_\lambda, P\eta_\lambda) - \phi(t_\lambda, P\eta_\lambda) - \lambda \|P\eta_\lambda\|^2.$$

So, we may assume $P\eta_\lambda = \eta_\lambda$. Therefore, as $\lambda \rightarrow 0$, t_λ and $(\eta_\lambda, \mathbf{e}_j)$ tend to \hat{t} and $\hat{\mathbf{b}}_j$ respectively and $t_\lambda > 0$ for small λ . Fixing λ arbitrarily and putting $t_\lambda = s$, $(\eta_\lambda, \mathbf{e}_j) = \beta_j$ and $\beta = (\beta_1, \dots, \beta_N)$, we can easily see

$$(8.15) \quad \begin{aligned} \frac{\partial \hat{\psi}}{\partial s}(s, \beta) - \min_z \max_y \left[\sum_{k,j=1}^N \frac{1}{2} f^2(y, z) \frac{\partial^2 \hat{\psi}(s, \beta)}{\partial \beta_k \partial \beta_j} \beta_k \beta_j + \lambda |\beta|^2 \right. \\ \left. + \sum_{j=1}^N \sum_{i=1}^n \sigma^i(y, z) \left(k_i^j + \frac{1}{2} \right) \beta_j \frac{\partial \hat{\psi}(s, \beta)}{\partial \beta_j} + \hat{h}(\beta, y, z) \right] \\ \leq 2\lambda \lambda_0 |\beta|^2. \end{aligned}$$

Tending λ to 0, (8.15) implies that \hat{w} is a subsolution of (8.12).

By the same argument, we can show that \hat{w} is a super solution. Hence we get $\hat{w} = v$. Recalling (8.13), we conclude the proof.

REFERENCES

- [1] D.G. Aronson, Non-negative solution of linear parabolic equation, *Ann. della Scuola Norm. Sup. Pisa*, **22** (1968), 607–694.
- [2] M.G. Crandall and P.L. Lions, Hamilton-Jacobi equations in infinite dimensions, Part 4, *J. Funct. Anal.*, **90** (1990), 273–283, Part 5, **97** (1991), 417–465.
- [3] M.G. Crandall, H. Ishii and P.L. Lions, A user's guide to viscosity solutions, *Bull. Amer. Math. Soc., N. S.*, **27** (1992), 1–67.
- [4] W.H. Fleming and P.E. Souganidis, On the existence of value function of two-player, zero-sum stochastic differential games, *Indiana Univ. Math. J.*, **38**, (1989), 293–314.
- [5] W.H. Fleming and H.M. Soner, *Controlled Markov Processes and Viscosity Solutions*, Springer Verlag, 1992.
- [6] H. Ishii, Viscosity solution of nonlinear second-order partial differential equation in Hilbert spaces, Preprint.
- [7] N.V. Krylov and B.L. Rozovskii. On the Cauchy problem for linear stochastic partial differential equations, *Math. USSR Izv.*, **11** (1977), 1267–1284.
- [8] P.L. Lions, Viscosity solutions and optimal stochastic control in infinite dimensions, Part 1, *Acta Math.*, **161** (1988), 243–278, Part 2 *L. N. M.*, **1390** (1988), 147–170, Part 3, *J. Funct. Anal.*, **86** (1989), 1–18.
- [9] M. Nisio, Optimal control for stochastic differential equations and viscosity solutions of Bellman equations, *Nagoya Math. J.*, **123** (1991), 13–37.
- [10] E. Pardoux, Stochastic partial differential equations and filtering of diffusion processes, *Stochastics*, **3** (1979), 127–167.

W.H. Fleming
Division of Applied Mathematics
Brown University
Providence R. I., 02912, USA

M. Nisio
Department of Mathematics
Kobe University
Rokko, Kobe 657, Japan

Current address:
Department of Mathematics
Osaka Electro-Communication University
Hatsu-cho, Neyagawa 572, Japan

