

**CURVATURE, GEODESICS AND THE BROWNIAN MOTION
 ON A RIEMANNIAN MANIFOLD II
 EXPLOSION PROPERTIES**

KANJI ICHIHARA

§1. Introduction

Let M be an n -dimensional, complete, connected and non compact Riemannian manifold and g be its metric. Δ_M denotes the Laplacian on M .

The Brownian motion on the Riemannian manifold M is defined to be the unique minimal diffusion process $(X_t, \zeta, P_x, x \in M)$ associated with the Laplacian Δ_M where $\zeta(\omega)$ is the explosion time of $X_t(\omega)$ i.e. if $\zeta(\omega) < +\infty$, then $\lim_{t \rightarrow \zeta(\omega)} X_t(\omega) = \infty$.

In the previous paper [3], the author has discussed recurrence and transience of the Brownian motion X on M . This paper may be considered to be a continuation, in which the relation between explosions of the Brownian motion X and geodesics, curvature of the Riemannian manifold M will be investigated. It should be remarked that Yau [7] has given a sufficient condition for no explosion of the Brownian motion in terms of the Ricci curvature.

Let us begin with the Brownian motion $X^0 = (X_t^0, \zeta^0, P_x^0, x \in M_0)$ on a model (M_0, g_0) where the model (M_0, g_0) is defined to be a Riemannian manifold $R^n = [0, +\infty) \times S^{n-1}$ given a metric $dr^2 + g_0(r)^2 d\theta^2$, $(r, \theta) \in (0, +\infty) \times S^{n-1}$. See Ichihara [3] for the precise definition. Then by the same reasoning as in Ichihara [3] Section 1, we obtain from Fellers tests for explosions, McKean [5],

PROPOSITION 1.1. *It holds whether*

$$\begin{array}{ll} P_x^0 \{ \zeta^0 = +\infty \} = 1 & \text{on } M \\ \text{or} & \\ P_x^0 \{ \zeta^0 = +\infty \} = 0 & \text{on } M \end{array}$$

according as

Received July 23, 1980.

$$\int^{+\infty} g_0(r)^{-n+1} dr \int^r g_0(s)^{n-1} ds = +\infty \quad \text{or} \quad < +\infty.$$

§ 2. Tests for explosions of the Brownian motion on a Riemannian manifold M

Let normal, minimal geodesics be defined as in Ichihara [3]. Ric_M and K_M denote the Ricci, and sectional curvatures respectively. $K_0(r)$, $r \geq 0$ is the radial sectional curvature of a model (M_0, g_0) defined in Ichihara [3].

Our main theorems are stated as follows.

THEOREM 2.1. *If for some $p \in M$ there exists a model (M_0, g_0) satisfying the following two conditions (i) and (ii), then no explosion for the Brownian motion X is possible. i.e.*

$$P_x\{\zeta = +\infty\} = 1 \quad \text{on } M.$$

(i) *For every minimal geodesic $m(r) : [0, \ell(m)) \rightarrow M$, $m(0) = p$,*

$$\text{Ric}_M(\dot{m}(r)) \geq (n-1)K_0(r) \quad \text{on } [0, \ell(m)).$$

(ii) $\int^{+\infty} g_0(r)^{-n+1} dr \int^r g_0(s)^{n-1} ds = +\infty.$

THEOREM 2.2. *Let M be simply connected. If for some $p \in M$ there exists a model (M_0, g_0) satisfying the following two conditions (i) and (ii), then explosion for the Brownian motion X is sure. i.e.*

$$P_x\{\zeta < +\infty\} = 1 \quad \text{on } M.$$

(i) *For every normal geodesic $m(r) : [0, +\infty) \rightarrow M$, $m(0) = p$,*

$$K_M(\dot{m}(r), X) \leq K_0(r) \text{ for every unit vector } X \in N(\dot{m}(r)) \text{ on } [0, +\infty)$$

(ii) $\int^{+\infty} g_0(r)^{-n+1} dr \int^r g_0(s)^{n-1} ds < +\infty.$

In order to prove the above theorems, we shall introduce the following notations.

$$\begin{aligned} \sigma_\rho(\omega) &= \inf\{t > 0 \mid d(p, X_t(\omega)) \geq \rho\}, \quad \rho > 0 \\ u_\rho(x) &= E_x\{e^{-\sigma_\rho}\}, \quad \Sigma_\rho = \{x \in M \mid d(p, x) < \rho\} \end{aligned}$$

where $d(x, y)$ is the distance induced by the Riemannian metric. σ_ρ^0 , u_ρ^0 and Σ_ρ^0 denote the corresponding ones of the Brownian motion on a model

(M_0, g_0) centered at $p =$ the origin 0 .

The following proposition will be proved in a similar way to that of Ichihara [2].

PROPOSITION 2.1. *For each $\rho \in (0, +\infty)$, $u_\rho \in C^\infty(\Sigma_\rho)$ and $\Delta_M u_\rho - u_\rho = 0$ in Σ_ρ . Furthermore in case of a model (M_0, g_0)*

$$\lim_{\substack{y \rightarrow x \\ y \in \Sigma_\rho^0}} u_\rho^0(y) = 1$$

for each $x \in \partial(\Sigma_\rho^0)$, the boundary of Σ_ρ^0 .

Proof of Theorem 2.1. Since M_0 is rotationally symmetric about 0 , $u_\rho^0(x)$ is a radial function. i.e.

$$u_\rho^0(x) = u_\rho^0(r) \quad \text{for } x = (r, \theta) \in M_0.$$

Thus $u_\rho^0 \in C^\infty([0, \rho])$ satisfies

$$\frac{d^2 u_\rho^0(r)}{dr^2} + \frac{(n-1)}{g_0(r)} \frac{dg_0(r)}{dr} \frac{du_\rho^0(r)}{dr} = u_\rho^0(r)$$

on $(0, \rho)$. Note that $u_\rho^0(r)$ is, by definition, an increasing function of r . Set $\tilde{u}_\rho(x) = u_\rho^0(d(p, x))$. Therefore following an argument similar to Yau [6], Appendix, we can obtain under the assumption (i) that

$$\Delta_M \tilde{u}_\rho(x) \leq \Delta_{M_0} u_\rho^0(r)$$

for $r = d(p, x) < \rho$, in the distribution sense. Consequently

$$\Delta_M \tilde{u}_\rho - \tilde{u}_\rho \leq \Delta_{M_0} u_\rho^0 - u_\rho^0 = 0 \quad \text{in } \Sigma_\rho.$$

Set

$$\Phi_\rho(x) = u_\rho(x) - \tilde{u}_\rho(x),$$

then it holds that

$$\Delta_M \Phi_\rho - \Phi_\rho = (\Delta_M u_\rho - u_\rho) - (\Delta_M \tilde{u}_\rho - \tilde{u}_\rho) \geq 0.$$

i.e.

$$\Delta_M \Phi_\rho \geq \Phi_\rho \quad \text{in } \Sigma_\rho$$

in the distribution sense.

We shall show that for each $\rho > 0$

$$\Phi_\rho(x) \leq 0 \quad \text{in } \Sigma_\rho.$$

Suppose on the contrary that with some $\rho_0 > 0$

$$\sup_{x \in \Sigma_{\rho_0}} \Phi_{\rho_0}(x) > 0.$$

Since (*) Φ_{ρ_0} is continuous in Σ_{ρ_0} and

$$(**) \quad \overline{\lim}_{\substack{y \rightarrow x \\ y \in \Sigma_{\rho_0}}} \Phi_{\rho_0}(y) \leq 0 \text{ for each } x \in \partial \Sigma_{\rho_0}$$

from Proposition 2.1, there exists a point $x_0 \in \Sigma_{\rho_0}$ such that

$$\Phi_{\rho_0}(x_0) = \sup_{x \in \Sigma_{\rho_0}} \Phi_{\rho_0}(x) > 0.$$

Set

$$C = \{x \in \Sigma_{\rho_0} \mid \Phi_{\rho_0}(x) > 0\}.$$

Denote by C_{x_0} the connected component containing the point x_0 of the set C . Then from the facts (*) and (**),

$$\overline{\lim}_{\substack{y \rightarrow x \\ y \in C_{x_0}}} \Phi_{\rho_0}(y) \leq 0 \quad \text{for each } x \in \partial C_{x_0}.$$

Since Φ_{ρ_0} is weakly A_M -subharmonic in C_{x_0} , applying the strong maximum principle in Littman [4] we obtain

$$\Phi_{\rho_0}(x) = \Phi_{\rho_0}(x_0) \quad \text{for each } x \in C_{x_0},$$

which is a contradiction. Thus we have shown that for each $\rho > 0$,

$$\Phi_{\rho}(x) \leq 0 \quad \text{in } \Sigma_{\rho}.$$

i.e.
$$u_{\rho}(x) \leq \tilde{u}_{\rho}(x) \quad \text{for every } x \in \Sigma_{\rho}.$$

Under the assumption (ii) in Theorem 2.1, the Brownian motion X^0 on the model (M_0, g_0) is conservative. (See Proposition 1.1.)

i.e.
$$P_x^0\{\zeta^0 = +\infty\} = 1 \quad \text{on } M_0.$$

Moreover

$$u_{\rho}^0(r) = u_{\rho}^0(x) = E_x^0\{e^{-\sigma_{\rho}^0}\}$$

converges to

$$E_x^0\{e^{-\zeta^0}\}$$

for each $x = (r, \theta) \in M_0$ because $\sigma_{\rho}^0 \rightarrow \zeta^0$ as $\rho \rightarrow +\infty$. Thus we see that

$$\lim_{\rho \rightarrow +\infty} u_\rho^0(r) = 0 \quad \text{for every } r \geq 0.$$

Hence it follows from the inequality proved above that

$$\lim_{\rho \rightarrow +\infty} u_\rho(x) = 0 \quad \text{for every } x \in M.$$

Since $\sigma_\rho \rightarrow \zeta$ as $\rho \rightarrow +\infty$, we see that

$$0 = \lim_{\rho \rightarrow +\infty} u_\rho(x) = E_x\{e^{-\zeta}\} \quad \text{for every } x \in M.$$

Thus we can conclude

$$P_x\{\zeta = +\infty\} = 1$$

on M .

q.e.d.

Proof of Theorem 2.2. We first note that under the assumptions \exp_p maps $T_p(M)$ diffeomorphically onto M as shown in Ichihara [3]. Thus we have geodesic polar coordinates $(r, \theta) \in (0, +\infty) \times S^{n-1}$ centered at p .

Now define $v = v(r)$, $r \geq 1$ to be the positive increasing solution:

$$v = \sum_{m=0}^{\infty} v_m \quad v_0 = 1$$

$$v_m(r) = \int_1^r g_0(s)^{-n+1} ds \int_0^s g_0(t)^{n-1} v_{m-1}(t) dt, \quad m \geq 1$$

of

$$\frac{1}{g_0(r)^{n-1}} \frac{d}{dr} \left(g_0(r)^{n-1} \frac{dv(r)}{dr} \right) = v(r), \quad r \geq 1.$$

Then it can be easily seen that

$$v(r) \leq \exp\{v_1(r)\}$$

for every $r \geq 1$ and so $v(r)$ is bounded above from the assumption (ii) of Theorem 2.2.

Set $\tilde{v}(x) = v(d(p, x))$. Then with the geodesic polar coordinates (r, θ) and $G(r, \theta) = \sqrt{\det(g_{ij})}(r, \theta)$ where $g = g_{ij} dx_i dx_j$, we have

$$\Delta_M \tilde{v}(x) = \frac{d^2 v(r)}{dr^2} + \frac{1}{G(r, \theta)} \frac{\partial G(r, \theta)}{\partial r} \frac{dv(r)}{dr} \Big|_{r=d(p, x)}.$$

By virtue of Hessian comparison theorem, Greene and Wu [1]

$$\geq \frac{d^2 v(r)}{dr^2} + \frac{(n-1)}{g_0(r)} \frac{dg_0(r)}{dr} \frac{dv(r)}{dr} \Big|_{r=d(p, x)} = v(d(p, x)) = \tilde{v}(x).$$

Now applying Itô's formula to the function $e^{-t}\tilde{v}(x)$, we obtain from the above inequality that

$$v(\rho)E_x\{e^{-\sigma_\rho}, \rho_\rho \leq \tau_1\} + E_x\{e^{-\tau_1}, \sigma_\rho > \tau_1\} \geq \tilde{v}(x)$$

for each $x \in \Sigma_\rho - \bar{\Sigma}_1$ where $\tau_1(\omega) = \inf\{t > 0 | d(p, X_t(\omega)) \leq 1\}$. Letting $\rho \rightarrow +\infty$, we have

$$v(\infty)E_x\{e^{-\zeta}, \zeta < \tau_1\} + E_x\{e^{-\tau_1}, \zeta > \tau_1\} \geq \tilde{v}(x).$$

because $\sigma_\rho \rightarrow \zeta$ as $\rho \rightarrow +\infty$.

We shall show

$$(*) \quad E_x\{e^{-\tau_1}, \tau_1 < \zeta\} \leq P_x\{\tau_1 < \zeta\} \longrightarrow 0 \quad \text{as } d(p, x) \rightarrow +\infty.$$

Set

$$\psi_\rho(r) = \frac{\int_r^\rho g_0(s)^{-n+1} ds}{\int_1^\rho g_0(s)^{-n+1} ds}, \quad \Psi_\rho(x) = \psi_\rho(d(p, x))$$

and

$$\phi_\rho(x) = P_x\{\tau_1 < \sigma_\rho\} \quad \text{for each } \rho > 1.$$

Then it is easy to see that

$$\begin{aligned} \Delta_M \phi_\rho &= 0 && \text{in } \Sigma_\rho - \bar{\Sigma}_1 \\ \phi_\rho(x) &= \begin{cases} 1 & \text{if } d(p, x) = 1 \\ 0 & \text{if } d(p, x) = \rho \end{cases} \end{aligned}$$

and

$$\Psi_\rho(x) = \begin{cases} 1 & \text{if } d(p, x) = 1 \\ 0 & \text{if } d(p, x) = \rho. \end{cases}$$

Furthermore Hessian comparison theorem [1] gives that

$$\Delta_M \Psi_\rho \leq 0 \quad \text{in } \Sigma_\rho - \bar{\Sigma}_1.$$

Consequently we can deduce by virtue of the maximum principle,

$$\phi_\rho(x) \leq \Psi_\rho(x) \quad x \in \Sigma_\rho - \bar{\Sigma}_1.$$

i.e.

$$P_x\{\tau_1 < \sigma_\rho\} \leq \psi_\rho(d(p, x)).$$

Since $\sigma_\rho \rightarrow \zeta$ as $\rho \rightarrow +\infty$, we get

$$P_x\{\tau_1 < \zeta\} \leq \frac{\int_{d(p,x)}^{\infty} g_0(r)^{-n+1} dr}{\int_1^{\infty} g_0(r)^{-n+1} dr} \quad \text{for } d(p, x) > 1,$$

which gives the desired result (*). Thus we obtain from (*)

$$\lim_{x \rightarrow \infty} E_x\{e^{-\zeta}, \zeta < \tau_1\} \geq 1$$

and so

$$\lim_{x \rightarrow \infty} P_x\{\zeta < \infty\} \geq \lim_{x \rightarrow \infty} E_x\{e^{-\zeta}\} \geq 1.$$

By the strong Markov property

$$P_x\{\zeta < +\infty\} = E_x\{P_{X_{\sigma_\rho}}\{\zeta < \infty\}\}$$

for every $\rho > d(p, x)$ and hence

$$= \lim_{\rho \rightarrow +\infty} E_x\{P_{X_{\sigma_\rho}}\{\zeta < +\infty\}\} \geq E_x\{\lim_{\rho \rightarrow +\infty} P_{X_{\sigma_\rho}}\{\zeta < +\infty\}\} \geq 1.$$

This completes the proof.

q.e.d.

§ 3. Some examples

In [7], Yau has shown that no explosion for the Brownian motion is possible if the Ricci curvature of M is bounded from below by a constant. We shall extend this result as follows.

1. If for a fixed $p \in M$ and every minimal geodesic $m(r) : [0, \ell(m)) \rightarrow M$, $m(0) = p$,

$$\text{Ric}_M(\dot{m}(r)) \geq -C_1 r^2 - C_2 \quad \text{on } [0, \ell(m))$$

with positive constants C_i $i = 1, 2$, then no explosion for the Brownian motion X is possible.

Proof. In order to prove this, it is enough to show the existence of a model (M_0, g_0) which satisfies the conditions (i) and (ii) in Theorem 2.1.

Set $K_0(r) = -C_1 r^2 - C_2$, $r \in [0, +\infty)$ and let $g_0(r) \in C([0, +\infty))$ be the unique solution of the following Jacobi equation.

$$\frac{d^2 g_0(r)}{dr^2} = -K_0(r)g_0(r) \quad g_0(0) = 0, \quad \frac{dg_0}{dr}(0) = 1.$$

Then the Sturm comparison theorem asserts that $g_0(r) > r$ for every $r > 0$.

Thus we have obtained a model (M_0, g_0) satisfying (i) in Theorem 2.1.

It remains to verify the condition (ii). In order to do it, we shall introduce the function

$$g_1(r) = \exp\{kr^2\}$$

with a positive constant k . Define

$$K_1(r) = -\frac{1}{g_1(r)} \frac{d^2 g_1(r)}{dr^2} = -4k^2 r^2 - 2k.$$

For a fixed positive number r_0 , it is easily seen that with a sufficiently large k

$$(*) \quad K_1(r) \leq K_0(r) \quad \text{for every } r \geq r_0$$

and

$$(**) \quad \frac{1}{g_1(r_0)} \frac{dg_1}{dr}(r_0) \geq \frac{1}{g_0(r_0)} \frac{dg_0}{dr}(r_0).$$

From the equations $(d^2 g_i(r)/dr^2) = -K_i(r)g_i(r)$, $i = 0, 1$, we have, for every $r \geq r_0$,

$$\begin{aligned} 0 &= g_1(r) \frac{d^2 g_0(r)}{dr^2} - \frac{d^2 g_1(r)}{dr^2} g_0(r) + (K_0(r) - K_1(r))g_1(r)g_0(r) \\ &= \frac{d}{dr} \left(g_1(r) \frac{dg_0(r)}{dr} \right) - \frac{d}{dr} \left(g_0(r) \frac{dg_1(r)}{dr} \right) + (K_0(r) - K_1(r))g_1(r)g_0(r). \end{aligned}$$

Hence we see from (*)

$$\left[g_1(s) \frac{dg_0(s)}{ds} - g_0(s) \frac{dg_1(s)}{ds} \right]_{r_0}^r = \int_{r_0}^r (K_1(s) - K_0(s))g_0(s)g_1(s)ds \leq 0.$$

Therefore it follows from (**) that

$$g_1(r) \frac{dg_0(r)}{dr} - g_0(r) \frac{dg_1(r)}{dr} \leq 0.$$

$$\text{i.e.} \quad \frac{1}{g_1(r)} \frac{dg_1(r)}{dr} \geq \frac{1}{g_0(r)} \frac{dg_0(r)}{dr}$$

for every $r \geq r_0$.

Set

$$G_i(r) = \int_{r_0}^r g_i(u)^{-n+1} du \int_{r_0}^u g_i(v)^{n-1} dv \quad i = 0, 1.$$

Then these functions satisfy

$$\begin{cases} \frac{d^2G_i(r)}{dr^2} + B_i(r)\frac{dG_i(r)}{dr} = 1 & \text{on } [r_0, +\infty) \\ G_i(r_0) = \frac{dG_i}{dr}(r_0) = 0 \end{cases}$$

where
$$B_i(r) = \frac{1}{g_i(r)} \cdot \frac{dg_i(r)}{dr}.$$

Since $B_i(r) \geq B_0(r)$ on $[r_0, +\infty)$ and G_i is an increasing function, we have

$$1 = \frac{d^2G_i(r)}{dr^2} + B_i(r)\frac{dG_i(r)}{dr} \geq \frac{d^2G_0(r)}{dr^2} + B_0(r)\frac{dG_0(r)}{dr}.$$

Solving this differential inequality, we can easily see that

$$G_0(r) \geq G_i(r)$$

for every $r \geq r_0$.

Thus in order to verify the condition (ii), it suffices to show

$$G_i(+\infty) = +\infty.$$

We now compute

$$\begin{aligned} G_i(+\infty) &= \int_{r_0}^{+\infty} dr \int_{r_0}^r \exp\{-(n-1)kr^2 + (n-1)kt^2\} dt \\ &= \int_{r_0}^{+\infty} dr \int_r^{+\infty} \exp\{(n-1)kr^2\} \cdot \exp\{-(n-1)kt^2\} dt. \end{aligned}$$

Using the following inequality

$$\begin{aligned} &\int_r^{+\infty} \exp\{-(n-1)kt^2\} dt \\ &\geq \frac{1}{\sqrt{(n-1)k}} \left(\sqrt{(n-1)k}r + \frac{1}{\sqrt{(n-1)k}r} \right)^{-1} \exp\{-(n-1)kr^2\}, \end{aligned}$$

we have

$$\geq \int_{r_0}^{+\infty} ((n-1)kr + 1)^{-1} dr = +\infty.$$

This completes the proof.

q.e.d.

The next example will be shown in a way similar to the proof of Example 1.

2. Suppose M is simply connected and negatively curved. If for a fixed $p \in M$ and every normal geodesic $m(r) : [0, +\infty) \rightarrow M$, $m(0) = p$

$$K_M(\dot{m}(r), \cdot) \leq -C_1 r^{2+\delta} \quad \text{for every } r \geq C_2$$

with positive constants C_i , $i = 1, 2$ and δ , then explosion for the Brownian motion X on M is sure.

3. Let S_n be an embeded hypersurface in R^{n+1} defined by

$$x_{n+1} = f(x_1, \dots, x_n).$$

Suppose f is a radial function, then the Brownian motion X on S_n is conservative

i.e. $P_x\{\zeta = +\infty\} = 1 \quad \text{on } S_n.$

Proof. Since f is a radial function, using polar coordinates (r, θ) of R^n , we have

$$\begin{aligned} dx_1^2 + \dots + dx_n^2 + dx_{n+1}^2 &= dr^2 + r^2 d\theta^2 + f_r^2 dr^2 \\ &= (1 + f_r^2) dr^2 + r^2 d\theta^2. \end{aligned}$$

As in Example 4 [3], we can obtain the geodesic polar coordinates (s, θ) with

$$dx_1^2 + \dots + dx_n^2 + dx_{n+1}^2 = ds^2 + g_0(s)^2 d\theta^2$$

where

$$\begin{aligned} p(r) &= \int_0^r \sqrt{1 + f_u^2} du, \quad r \geq 0 \\ s &= p(r) \end{aligned}$$

and $g_0(r)$ is the inverse function of p . i.e. $s = p(g_0(s))$.

Notice that

$$B_0(s) = \frac{1}{g_0(s)} \frac{dg_0(s)}{ds} = \frac{1}{r} \frac{1}{\sqrt{1 + f_r^2}}$$

is convergent to zero as $s \rightarrow +\infty$. Set $g_1(s) = e^s$, then we have

$$B_1(s) = \frac{1}{g_1(s)} \frac{dg_1(s)}{ds} = 1.$$

Consequently it holds that for some $r_0 > 0$,

$$B_1(s) \geq B_0(s) \quad \text{on } [r_0, +\infty).$$

Now applying the comparison argument in page 123 we get that

$$(***) \quad \int_{r_0}^r g_0(u)^{-n+1} du \int_{r_0}^u g_0(v)^{n-1} dv \leq \int_{r_0}^r g_1(u)^{-n+1} du \int_{r_0}^u g_1(v)^{n-1} dv.$$

It is easy to see that the right hand of the above inequality (***) is divergent to $+\infty$ when r tends to $+\infty$. Thus we have

$$\int_{r_0}^{+\infty} g_0(r)^{-n+1} dr \int_{r_0}^r g_0(s)^{n-1} ds = +\infty$$

which implies $P_x\{\zeta = +\infty\} = 1$ on S_n .

REFERENCES

- [1] R. E. Greene and H. Wu, Function theory on manifolds which possess a pole, Lecture notes, Springer, no. 699.
- [2] K. Ichihara, Some global properties of symmetric diffusion processes, Publ. R.I.M.S., Kyoto Univ., **14**, no. 2, (1978), 441–486.
- [3] —, Curvature, Geodesics and the Brownian motion on a Riemannian manifold I, Nagoya Math. J., **87** (1982), 101–114.
- [4] W. Littman, A strong maximum principle for weakly L -subharmonic functions, J. Math. and Mech., **8**, no. 5, (1959), 761–770.
- [5] H. P. McKean, Stochastic integrals, Academic press (1969).
- [6] S. T. Yau, Some function-theoretic properties of complete Riemannian manifold and their application to geometry, Indiana Univ. Math. J., **25**, no. 7, (1976), 659–670.
- [7] —, On the heat kernel of a complete Riemannian manifold, J. Math. pures et appl., **57** (1978), 191–201.

*Department of Applied Science
Faculty of Engineering
Kyushu University
Fukuoka, Japan*

*Current address:
Department of Mathematics
Faculty of General Education
Nagoya University
Nagoya, Japan*

