

A NEW VERSION OF \mathfrak{a} -TIGHT CLOSURE

ADELA VRACIU

Abstract. Hara and Yoshida introduced a notion of \mathfrak{a} -tight closure in 2003, and they proved that the test ideals given by this operation correspond to multiplier ideals. However, their operation is not a true closure. The alternative operation introduced here is a true closure. Moreover, we define a joint Hilbert-Kunz multiplicity that can be used to test for membership in this closure. We study the connections between the Hara-Yoshida operation and the one introduced here, primarily from the point of view of test ideals. We also consider variants with positive real exponents.

§1. Introduction

In [HY], Hara and Yoshida introduced a notion of \mathfrak{a} -tight closure, which generalizes the “classical” tight closure of Hochster and Huneke introduced in [HH1]. The main motivation for their work is the connection between the test ideals given by this operation and multiplier ideals, which generalizes previous results of Hara ([H2]) and Smith ([S2]). The advantage of this version of test ideal is that it allows them to study multiplier ideals for pairs, not only the multiplier ideal of a variety.

However, the Hara-Yoshida \mathfrak{a} -tight closure is not a true closure operation, since it gets (potentially) larger when iterated. The version introduced in this paper is a true closure, and it is always contained in the Hara-Yoshida \mathfrak{a} -tight closure. We establish several other connections between these operations. We prove that for a Gorenstein graded algebra of dimension at least 2, the test ideals given by these two operations are the same (Theorem 4.3). The Hara-Yoshida \mathfrak{a} -tight closure of an ideal I is denoted $I^{*\mathfrak{a}}$, while the new version introduced here will be denoted ${}^{\mathfrak{a}}I^*$.

We define a joint Hilbert-Kunz multiplicity associated to two \mathfrak{m} -primary

Received April 5, 2007.

Revised October 11, 2007.

2000 Mathematics Subject Classification: 13A35.

The author thanks N. Epstein, A. Kustin, and K. Yoshida for useful conversations on the topic of this manuscript. The author was partially supported by a grant from the University of South Carolina Research Foundation.

ideals \mathfrak{a} and I , and we prove that this multiplicity can be used to test for membership in our version of \mathfrak{a} -tight closure. This is similar to the way in which the Hilbert-Samuel multiplicity is used to test for membership in the integral closure, and the Hilbert-Kunz multiplicity is used to test for membership in tight closure.

There are versions of both closures, as well as of the joint multiplicity, in which positive real numbers are allowed as exponents. For fixed ideals I and \mathfrak{a} , we study the question of how $I^{*\mathfrak{a}^t}$ and ${}^{\mathfrak{a}^t}I^*$ vary with t . This question is related to the notion of jumping exponents (in characteristic zero), or F-thresholds (in positive characteristic).

In this paper, R will denote a Noetherian ring of positive characteristic $p > 0$ and Krull dimension $d > 0$, and $q = p^e$ will always denote a power of the characteristic. Most of the time, R will be assumed to be either local or graded. R^0 is the set of elements in R that are not in any minimal prime of R . If $I \subset R$ is an ideal, $I^{[q]}$ denotes the ideal generated by all i^q , when $i \in I$.

§2. Definitions and elementary properties

DEFINITION 2.1. ([HY]) Let \mathfrak{a}, I be ideals in R , and $x \in R$. We say that $x \in I^{*\mathfrak{a}}$ if there exists $c \in R^0$ such that $c\mathfrak{a}^q x^q \subseteq I^{[q]}$ for all $q = p^e \gg 0$.

DEFINITION 2.2. Let \mathfrak{a}, I be ideals in R , and $x \in R$. We say that $x \in {}^{\mathfrak{a}}I^*$ if there exists $c \in R^0$ such that $c\mathfrak{a}^q x^q \subseteq \mathfrak{a}^q I^{[q]}$ for all $q = p^e \gg 0$.

In the case when $\mathfrak{a} = R$, both of the above definitions coincide with the definition of the usual tight closure of Hochster and Huneke ([HH1]), which is denoted I^* . Some elementary properties of these operations are summarized below.

OBSERVATION 2.3. (1) For all \mathfrak{a} and I , we have $I^* \subseteq {}^{\mathfrak{a}}I^* \subseteq I^{*\mathfrak{a}}$.

- (2) If $\mathfrak{a} = (f)$ is a principal ideal, then $I^{*\mathfrak{a}} = I^* : f$. In particular, $(I^{*(f)})^{*(f)} \neq I^{*(f)}$ when (R, \mathfrak{m}) is local, I is \mathfrak{m} -primary, and $f \in \mathfrak{m} \setminus I^*$.
- (3) If $\mathfrak{a} = (f)$ is a principal ideal, and f is a non-zerodivisor on R , then ${}^{\mathfrak{a}}I^* = I^*$.
- (4) For all \mathfrak{a} and I , ${}^{\mathfrak{a}}({}^{\mathfrak{a}}I^*)^* = {}^{\mathfrak{a}}I^*$. In other words, ${}^{\mathfrak{a}}I^*$ is a true closure operation.

Proof. (1), (2) and (3) are trivial.

(4) Let $x \in {}^{\mathfrak{a}}({}^{\mathfrak{a}}I^*)^*$. Then there exists $c \in R^o$ such that $c\mathfrak{a}^q x^q \subseteq \mathfrak{a}^q({}^{\mathfrak{a}}I^*)^{[q]}$ for all $q = p^e$. Also, there exists $c' \in R^o$ such that $c'\mathfrak{a}^q({}^{\mathfrak{a}}I^*)^{[q]} \subseteq \mathfrak{a}^q I^{[q]}$. Combining these two inclusions, we get $cc'\mathfrak{a}^q x^q \subseteq \mathfrak{a}^q I^{[q]}$. \square

The notion of *test element for tight closure* was defined in [HH1], and it was proved that test elements (for tight closure) exist in very general classes of rings.

DEFINITION 2.4. An element $c \in R^o$ is called a *test element* for tight closure if we have $cx \in I$ for every ideal I and every $x \in I^*$.

PROPOSITION 2.5. Assume that R has test elements for the usual tight closure. Then for any ideals \mathfrak{a} and I , with I of positive height and $\mathfrak{a} \cap R^o \neq \emptyset$, we have ${}^{\mathfrak{a}}I^* \subseteq \overline{I}$.

Proof. By the usual determinant trick, $cx^q \mathfrak{a}^q \subseteq \mathfrak{a}^q I^{[q]}$ implies $cx^q \in \overline{I^{[q]}} \subseteq \overline{I^q} \subseteq (I^{q-n+1})^*$, where n is the minimal number of generators of I . The last inclusion is by the tight closure version of the Briançon-Skoda theorem ([HH1], Theorem 5.4). Let $d \in R^o$ be a test element, and $f \in I^{n-1} \cap R^o$ a fixed element. Then we have $cdfx^q \in I^q$ for all $q = p^e$, which shows that $x \in \overline{I}$. \square

Versions in which positive real numbers occur as exponents can be defined for both operations:

DEFINITION 2.6. Let $\mathfrak{a}, I \subseteq R$ be ideals, and let $t > 0$ be a real number. Let $x \in R$. For any real number r , $\lceil r \rceil$ denotes the smallest integer greater than or equal to r .

- (1) We say that $x \in I^{*\mathfrak{a}^t}$ if there exists $c \in R^o$ such that $cx^q \mathfrak{a}^{\lceil tq \rceil} \subseteq I^{[q]}$ for all $q = p^e$.
- (2) We say that $x \in \mathfrak{a}^t I^*$ if there exists $c \in R^o$ such that $cx^q \mathfrak{a}^{\lceil tq \rceil} \subseteq \mathfrak{a}^{\lceil tq \rceil} I^{[q]}$ for all $q = p^e$.

Each of these operations gives rise to a test ideal as follows:

DEFINITION 2.7. Let $\mathfrak{a} \subseteq R$ be an ideal, and let $t > 0$ be a real number. We define

$$\tau(\mathfrak{a}^t) := \bigcap (I : I^{*\mathfrak{a}^t}) \quad T_{\mathfrak{a}^t} := \bigcap (I : \mathfrak{a}^t I^*),$$

where each intersection ranges over all the ideals $I \subseteq R$.

Part (1) of the next observation was noted in [HT], where it was used to prove Skoda's theorem for test ideals. Part (2) is an analogue for the new \mathfrak{a} -tight closure.

OBSERVATION 2.8. *Let $\mathfrak{a}, I \subset R$ be ideals, and $t > \nu(\mathfrak{a})$ a real number, where $\nu(\mathfrak{a})$ denotes the minimal number of generators of \mathfrak{a} . Then:*

- (1) $I^{*\mathfrak{a}^t} = (I^{*\mathfrak{a}^{t-1}}) : \mathfrak{a}$.
- (2) $\mathfrak{a}^t I^* = (\mathfrak{a}^{t-1}(\mathfrak{a}I)^*) : \mathfrak{a}$.

Proof. (1) The proof of this statement can be found as part of the proof of Theorem 4.1 in [HT].

(2) Note that we have $\mathfrak{a}^r = \mathfrak{a}^{[q]}\mathfrak{a}^{r-q}$ for all $r > \nu(\mathfrak{a})q$. We have $x \in \mathfrak{a}^t I^* \Leftrightarrow cx^q \mathfrak{a}^{[tq]} \subseteq \mathfrak{a}^{[tq]} I^{[q]} \Leftrightarrow c(\mathfrak{a}x)^{[q]} \mathfrak{a}^{[tq]-q} \subseteq \mathfrak{a}^{[tq]-q}(\mathfrak{a}I)^{[q]}$ and the conclusion follows since $[tq] - q = \lceil (t-1)q \rceil$. \square

We establish two connections between the two versions of \mathfrak{a} -tight closure. The first result, Proposition 2.9, shows that for elements of large enough degree in a graded ring, membership in one of these closures is equivalent to membership in the other. The second result, Proposition 2.12 shows that, under certain assumptions, every element in the Hara-Yoshida \mathfrak{a} -tight closure must satisfy a stronger condition, which bridges the gap between the Hara-Yoshida definition and the one introduced in this paper.

We establish the following notation, which will be in effect throughout this paper when graded rings are considered.

NOTATION. If R is a finitely generated graded algebra over a field, $R = \bigoplus_{n \geq 0} R_n$, we will denote $R_+ := \bigoplus_{n > 0} R_n$ the unique maximal homogeneous ideal of R . We will let y_1, \dots, y_s be algebra generators for R , and let β_1, \dots, β_s be their degrees. Set $\beta = \max(\beta_i)$, $\beta' = \min(\beta_i)$.

We say that R is standard graded if $\beta_i = 1$ for all i .

PROPOSITION 2.9. *Let R be a finitely generated graded algebra over a field and let $\mathfrak{a} \subset R$ be a homogeneous R_+ -primary ideal, so that $R_+^k \subseteq \mathfrak{a} \subseteq R_+^l$ for some integers $l \leq k$. Let $I = (f_1, \dots, f_n)$ be a homogeneous ideal, and $x \in R_N$ with $N \geq \beta k - \beta' l + \max(\deg(f_i))$. Then $x \in I^{*\mathfrak{a}} \Leftrightarrow x \in \mathfrak{a} I^*$.*

Proof. Assume that $x \in I^{*\mathfrak{a}}$ and $\deg(x) \geq \beta k - \beta' l + \max(\deg(f_i))$. For each homogeneous $h \in \mathfrak{a}^q$, we have $\deg(h) \geq \beta' l q$. We can write $cx^q h =$

$\sum_{i=1}^n a_i f_i^q$ with $c \in R^o$, $a_i \in R$ homogeneous elements, so that $\deg(a_i) = \deg(c) + q \deg(x) + \deg(h) - q \deg(f_i) \geq \beta k q$ for each i . Thus, $a_i \in R_{\geq \beta k q} \subseteq R_+^{kq} \subseteq \mathfrak{a}^q$ (the first inclusion follows because any element in $R_{\geq \beta k q}$ can be written as a linear combination of monomials $y_1^{i_1} \cdots y_s^{i_s}$ with $i_1 \beta_1 + \cdots + i_s \beta_s \geq \beta k q$, which implies that $i_1 + \cdots + i_s \geq k q$). \square

OBSERVATION 2.10. *If R is standard graded, so $\beta = \beta'$, $\mathfrak{a} = R_+^r$, and all the generators of I have the same degree, then we have ${}^{\mathfrak{a}}I^* = I^{*\mathfrak{a}} \cap R_{\geq N}$, where N denotes the common degree of the generators of I .*

Proof. Let $x \in {}^{\mathfrak{a}}I^*$, so there exists $c \in R^o$ (which can be assumed homogeneous) such that $cx^q \mathfrak{a}^q \subseteq \mathfrak{a}^q I^{[q]}$. Taking degrees of both sides yields $\deg(c) + q \deg(x) + q r \geq q r + q N$, so that $\deg(x) \geq N$. This shows that ${}^{\mathfrak{a}}I^* \subseteq I^{*\mathfrak{a}} \cap R_{\geq N}$. The other inclusion is contained in Proposition 2.9. \square

This observation might suggest that ${}^{\mathfrak{a}}I^* = I^{*\mathfrak{a}} \cap \bar{I}$ for \mathfrak{m} -primary ideals I . This is in fact not true (however, ${}^{\mathfrak{a}}I^* \subseteq I^{*\mathfrak{a}} \cap \bar{I}$ is always true), as seen in the following example.

EXAMPLE 2.11. Let $R = k[x, y]$, $I = (x^2, y^4)$, $\mathfrak{a} = (x, y)^3$. Then we have $I^{*\mathfrak{a}} = I + (xy^2, y^3) = I : (x, y)^2$, $I^{*\mathfrak{a}} \cap \bar{I} = I + (xy^2)$, and ${}^{\mathfrak{a}}I^* = I + (xy^3) = I : (x, y)$.

Proof. If $i + j \geq 3q$, we have $i \geq q$ or $j \geq 2q$. In either case we have $x^i y^j x^q y^{2q} \in (x^{2q}, y^{4q})$, and thus $xy^2 \in I^{*\mathfrak{a}}$. Similarly, if $i + j \geq 3q$ we have $i \geq 2q$ or $j \geq q$; in either case, $x^i y^j y^{3q} \in (x^{2q}, y^{4q})$, and thus $y^3 \in I^{*\mathfrak{a}}$. Also note $(xy^2)^2 \in I^2$, so $xy^2 \in \bar{I}$. However, $y^3 \notin \bar{I}$ (one can see this from the Newton polygon, for instance).

To see that $xy^2 \notin {}^{\mathfrak{a}}I^*$, we prove the stronger fact that $xy^2 \notin (x, y)^n I^*$ for any $n \geq 3$. This will suffice to prove the last statement, since all the ideals under consideration are monomial. Assume the contrary, so that there exists $c \in R^o$ such that $cx^i y^j x^q y^{2q} \in (x, y)^{nq} I^{[q]}$ for all i, j with $i + j = nq$, for some n .

Choose $i = \lceil \frac{q}{2} \rceil$, $j = (n-1)q + \lfloor \frac{q}{2} \rfloor$. We obtain $cx^{\lceil 3q/2 \rceil} y^{(n-3)q + \lfloor q/2 \rfloor} y^{4q} = ax^{2q} + by^{4q}$ with $a, b \in (x, y)^{nq}$. This is clearly impossible since the degree of $x^{\lceil 3q/2 \rceil} y^{(n-3)q + \lfloor q/2 \rfloor}$ is $(n-1)q$, and the degree of c is a constant. \square

PROPOSITION 2.12. *Let (R, \mathfrak{m}) be an excellent normal domain such that its completion is a domain. Let $I, \mathfrak{a} \subset R$ be ideals, and assume that \mathfrak{a} is not a principal ideal. Then there exists a $Q_0 = p^{e_0}$ and a $c \in R^o$ such that for all $x \in I^{*\mathfrak{a}}$ we have $cx^q \mathfrak{a}^q \subseteq \mathfrak{m}^{q/Q_0} I^{[q]}$ for all $q \gg 0$.*

Note that if \mathfrak{a} is \mathfrak{m} -primary, then we can replace \mathfrak{m}^{q/Q_0} by \mathfrak{a}^{q/Q_0} by choosing a possibly larger Q_0 .

Proof. First note that there is no loss of generality in assuming that I is $*$ -independent, i.e. $I = (f_1, \dots, f_n)$ with $f_i \notin (f_1, \dots, \hat{f}_i, \dots, f_n)^*$ for all i . That is because one can find a $*$ -independent $I_0 \subseteq I$ with $I_0^* = I^*$ (by omitting generators of I that are redundant up to tight closure), and it is easy to see that $I_0^* = I^*$ implies $I_0^{*\mathfrak{a}} = I^{*\mathfrak{a}}$.

Let $\mathfrak{a} = (a_1, \dots, a_s)$, with $s = \nu(\mathfrak{a}) \geq 2$, and $I = (f_1, \dots, f_n)$. The $*$ -independence assumption implies that there exists q_1 such that

$$(f_1^q, \dots, \hat{f}_i^q, \dots, f_n^q) : f_i^q \subseteq \mathfrak{m}^{[q/q_1]}$$

for all q and all i (cf. Proposition 2.4 in [Ab]).

Since R is normal, we have $a_l \notin \overline{(a_k)}$ for any $1 \leq k \neq l \leq s$, and we can choose $q_2 \gg 0$ such that $a_l \notin \overline{(a_k, \mathfrak{m}^{q_2/q_1})}$. Also choose $q_2 \geq s$. In particular, $a_l \notin (a_k, \mathfrak{m}^{q_2/q_1})^*$ and thus we can choose q_0 such that $(a_k^q, \mathfrak{m}^{[qq_2/q_1]}) : a_l^q \subseteq \mathfrak{m}^{q/q_0}$ (using Proposition 2.4 in [Ab] again).

We have $cx^{qq_2} \mathfrak{a}^{qq_2} \subseteq I^{[qq_2]}$ for a fixed $c \in R^\circ$ and all q . Fix an element $a_1^{i_1} \cdots a_s^{i_s} \in \mathfrak{a}^{qq_2}$ and write $2cx^{qq_2} a_1^{i_1} \cdots a_s^{i_s} = b_1 f_1^{qq_2} + \cdots + b_n f_n^{qq_2}$. The choice of q_2 guarantees that $i_k \geq q$ for some k . Choose an index $l \neq k$ and consider the element $a_1^{j_1} \cdots a_s^{j_s} \in \mathfrak{a}^{qq_2}$ with $j_k = i_k - q$, $j_l = i_l + q$, and $j_\tau = i_\tau$ for all other $\tau = 1, \dots, s$. We have $cx^{qq_2} a_1^{j_1} \cdots a_s^{j_s} = b'_1 f_1^{qq_2} + \cdots + b'_n f_n^{qq_2}$. Multiplying the first equation by a_l^q and the second equation by a_k^q yields $(b_i a_l^q - b'_i a_k^q) \in (f_1^{qq_2}, \dots, \hat{f}_i^{qq_2}, \dots, f_n^{qq_2}) : f_i^{qq_2} \subseteq \mathfrak{m}^{[qq_2/q_1]}$, and therefore $b_i \in (a_k^q, \mathfrak{m}^{[qq_2/q_1]}) : a_l^q \subseteq \mathfrak{m}^{q/q_0}$. This holds for all $i = 1, \dots, n$, and for any choice of the multi-index (i_1, \dots, i_s) . We get the desired conclusion by choosing $Q_0 = q_2 q_0$. \square

§3. Joint Hilbert-Kunz multiplicities

The idea of associating a multiplicity to a pair or more ideals (the so-called mixed multiplicity) first appeared in [Bt], and the notion was extensively studied by many other authors, including B. Tossier, D. Rees and I. Swanson. The idea of a multiplicity coming from length functions involving both ordinary and Frobenius powers can be found in work of Hanes ([Hn]). The joint Hilbert-Kunz multiplicity introduced here bares a resemblance to each of these previous multiplicities, but is different from them.

Assume that (R, \mathfrak{m}) is local with Krull dimension $d \geq 1$, let $I, \mathfrak{a} \subset R$ be \mathfrak{m} -primary ideals, M a finitely generated R -module, and $t > 0$ a fixed real number.

We study the function

$$\ell_M(q) := \lambda\left(\frac{M}{\mathfrak{a}^{\lceil qt \rceil} I^{[q]} M}\right),$$

where $q = p^e$. Note that $\mathfrak{a}^{\lceil qt \rceil}$ is an ordinary power where the exponent is obtained by taking the least integer which is greater than or equal to tq , while $I^{[q]}$ is a Frobenius power. We will write $\ell(q)$ for $\ell_R(q)$.

THEOREM 3.1. *Let R, I, \mathfrak{a}, M, t be as above, and let d be the Krull dimension of R . Then there is a $c > 0$ such that*

$$\ell_M(q) = cq^d + \mathcal{O}(q^{d-1}).$$

If $M = R$, we call c the *joint Hilbert-Kunz multiplicity* of the pair (\mathfrak{a}^t, I) and we denote it $e_{HK}(\mathfrak{a}^t, I)$. Note that the usual Hilbert-Samuel and Hilbert-Kunz multiplicities can be recovered as special cases of joint Hilbert-Kunz multiplicity: $e_{HK}(\mathfrak{a}, R) = e(\mathfrak{a})$ and $e_{HK}(R, I) = e_{HK}(I)$.

The proof of the Theorem follows essentially the same steps as in Mon-sky's paper ([Mo]). We will follow closely the outline of his paper.

LEMMA 3.2. *Assume that there is an $h \in R^o$ such that $hM = 0$. Then there exists a $a > 0$ such that $\ell_M(q) \leq aq^{d-1}$.*

Proof. Let n be the number of generators of I . Then we have $I^{[q]} \supseteq I^{nq}$. Also, $\lceil tq \rceil \geq \lceil tq \rceil$, so $\mathfrak{a}^{\lceil tq \rceil} \supseteq \mathfrak{a}^{\lceil tq \rceil} I^{nq}$, and it follows that

$$\ell_M(q) \leq \lambda\left(\frac{M}{(\mathfrak{a}^{\lceil tq \rceil} I^{nq})^q M}\right),$$

which is a Hilbert-Samuel function over the ring R/h , a ring of Krull dimension at most $d - 1$, and thus it is bounded by aq^{d-1} for some $a > 0$. \square

LEMMA 3.3. *Let M, N be finitely generated R -modules such that $M_{p_i} \cong N_{p_i}$ for every minimal prime p_i of R . Then $|\ell_M(q) - \ell_N(q)| \leq \mathcal{O}(q^{d-1})$.*

Proof. Let $S = R \setminus \bigcup p_i$. We have $S^{-1}M \cong S^{-1}N$. Since $S^{-1}\mathrm{Hom}_R(M, N) \cong \mathrm{Hom}_{S^{-1}R}(S^{-1}M, S^{-1}N)$, we have a homomorphism $\phi : M \rightarrow N$ such that $S^{-1}\phi$ is bijective. Unlocalizing, we get an element $h \in S$ such that h annihilates the cokernel C of ϕ . Consider the exact sequence

$$\frac{M}{\mathfrak{a}^{[tq]}I^{[q]}M} \longrightarrow \frac{N}{\mathfrak{a}^{[tq]}I^{[q]}N} \longrightarrow \frac{C}{\mathfrak{a}^{[tq]}I^{[q]}C} \longrightarrow 0$$

Lemma 3.2 gives

$$\ell_N(q) - \ell_M(q) \leq \ell_C(q) \leq aq^{d-1}$$

for some $a > 0$. Now repeat the argument with the roles of M, N reversed in order to get

$$\ell_M(q) - \ell_N(q) \leq bq^{d-1}$$

for some $b > 0$. □

DEFINITION 3.4. Let $M_{(e)}$ be M viewed as an R -module via the Frobenius map $F^e : R \rightarrow R$. Note that (e) is an exact functor, and, if we assume that the residue field of R is perfect, we have

$$\ell_{M_{(e)}}(q) = \lambda\left(\frac{M}{(\mathfrak{a}^{[tq]})^{[p^e]}I^{[qp^e]}M}\right).$$

The following is the one essential ingredient we need in addition to Monsky's ideas:

OBSERVATION 3.5. Let R, \mathfrak{a}, I, t be as above, $e > 0$ a fixed integer. By prime avoidance, we can choose generators f_1, \dots, f_n of \mathfrak{a} that are in R° . Let $f := f_1 \cdots f_n$. Assume that $e \gg 0$ so that $n < p^e$. Then:

a. $\mathfrak{a}^{[tqp^e]} \subseteq (\mathfrak{a}^{[tq]})^{[p^e]} : fp^e$.

b. Assume that the residue field of R is perfect. Then $|\ell_{M_{(e)}}(q) - \ell_M(p^e q)| \leq \mathcal{O}(q^{d-1})$.

Proof. **a.** The generators of $\mathfrak{a}^{[tqp^e]}$ are of the form $F = f_1^{a_1 p^e + i_1} \cdots f_n^{a_n p^e + i_n}$, where $0 \leq i_k \leq p^e - 1$ for all k , and

$$(a_1 + \cdots + a_n)p^e + i_1 + \cdots + i_n = [tqp^e] \geq ([tq] - 1)p^e,$$

with all $a_k, i_k \in \mathbf{Z}$. It follows that $a_1 + \cdots + a_n \geq [tq] - 1 - n/p^e$. Since $a_k \in \mathbf{Z}$ for all k , it must be that $a_1 + \cdots + a_n \geq [tq] - n$ and thus $(a_1 + 1) + \cdots + (a_n + 1) \geq [tq]$, and so $f_1^{a_1 + 1} \cdots f_n^{a_n + 1} F = (f_1^{a_1 + 1} \cdots f_n^{a_n + 1})p^e \in (\mathfrak{a}^{[tq]})^{[p^e]}$.

b. We have

$$\begin{aligned} \ell_{M_{(e)}}(q) - \ell_M(p^e q) &= \lambda\left(\frac{\mathfrak{a}^{[tq^{p^e}]} I^{[qp^e]} M}{(\mathfrak{a}^{[tq]})^{[p^e]} I^{[qp^e]} M}\right) \\ &\leq \lambda\left(\frac{(\mathfrak{a}^{[tq]})^{[p^e]} I^{[qp^e]} M : f^{p^e}}{(\mathfrak{a}^{[tq]})^{[p^e]} I^{[qp^e]} M}\right) = \lambda\left(\frac{M}{(\mathfrak{a}^{[tq]})^{[p^e]} I^{[qp^e]} M + (f^{p^e}) M}\right). \end{aligned}$$

The inequality above follows from part **a**. The second equality follows from the general fact that for any \mathfrak{m} -primary ideal $J \subset R$, and any element $g \in R$, we have

$$\lambda\left(\frac{JM : g}{JM}\right) = \lambda\left(\frac{M}{(J, g)M}\right)$$

applied to $J = (\mathfrak{a}^{[tq]})^{[p^e]} I^{[qp^e]}$ and $g = f^{p^e}$. (Proof of the general fact: consider the short exact sequence

$$0 \longrightarrow \frac{M}{JM : g} \longrightarrow \frac{M}{JM} \longrightarrow \frac{M}{JM + (g)M} \longrightarrow 0$$

where the first map is multiplication by g .) Lemma 3.2 now gives the desired conclusion, since

$$\lambda\left(\frac{M}{(\mathfrak{a}^{[tq]})^{[p^e]} I^{[qp^e]} M + (f^{p^e}) M}\right)$$

is a joint Hilbert-Kunz function over the $d-1$ dimensional ring $R/(f^{p^e})$. \square

LEMMA 3.6. *Assume that the residue field of R is perfect. Let*

$$0 \longrightarrow M' \longrightarrow M \longrightarrow M'' \longrightarrow 0$$

be a short exact sequence of finitely generated R -modules. Then we have

$$\ell_M(q) = \ell_{M'}(q) + \ell_{M''}(q) + \mathcal{O}(q^{d-1}).$$

Proof. **Case 1:** Assume that R is reduced. For each minimal prime p_i of R , R_{p_i} is a field and it follows that $M_{p_i} \cong (M' \oplus M'')_{p_i}$. The conclusion follows from Lemma 3.3.

Case 2: Let \mathfrak{n} denote the nilradical of R , and choose e such that $\mathfrak{n}^{[p^e]} = 0$. Note that $M_{(e)}$ is annihilated by \mathfrak{n} for every module M . We get a short exact sequence of R/\mathfrak{n} modules:

$$0 \longrightarrow M'_{(e)} \longrightarrow M_{(e)} \longrightarrow M''_{(e)} \longrightarrow 0,$$

and now we can apply the result from case 1 in conjunction with Obs. 3.5. \square

LEMMA 3.7. *Assume that R is a domain with perfect residue field. Then there exists $c > 0$ such that*

$$\ell(q) = cq^d + \mathcal{O}(q^{d-1}).$$

Proof. It is known that the rank of $R_{(1)}$ as an R -module is p^d . Apply Lemma 3.3 to the R -modules $R_{(1)}$ and R^{p^d} . We get

$$\left| \lambda \left(\frac{R}{(\mathfrak{a}^{\lceil tq \rceil})^{[p]} I^{[pq]}} \right) - p^d \ell(q) \right| = |\ell_{R_{(1)}}(q) - p^d \ell(q)| \leq a' q^{d-1}$$

for some $a' > 0$, and by Obs. 3.5 it follows that

$$|\ell(pq) - p^d \ell_R(q)| \leq a q^{d-1}$$

for some a . Thus, we have

$$\left| \frac{\ell(pq)}{(pq)^d} - \frac{\ell(q)}{q^d} \right| \leq \frac{a}{p^d q}.$$

It follows that

$$\left| \frac{\ell(q'q)}{(q'q)^d} - \frac{\ell(q)}{q^d} \right| \leq \frac{a}{p^d q} \frac{1 - \frac{1}{q'}}{1 - \frac{1}{p}},$$

thus $\{\ell(q)/q^d\}$ is a Cauchy sequence. Let $c := \lim_{q \rightarrow \infty} \ell(q)/q^d$. If we keep q fixed and let $q' \rightarrow \infty$, we get

$$|\ell(q)/q^d - c| \leq \frac{\alpha'}{q}$$

for some α' and all q , and thus $|\ell(q) - cq^d| \leq \alpha' q^{d-1}$, or in other words $\ell(q) = cq^d + \mathcal{O}(q^{d-1})$. \square

Now we are ready to prove the general case of Theorem 3.1.

Proof. Since every finitely generated module M has a filtration $(0) = M_0 \subset M_1 \subset \cdots \subset M_n = M$ with quotients $M_{i+1}/M_i \cong R/P_i$, with P_i prime ideals, the general case follows from Lemma 3.7 by repeated application of Lemma 3.6.

In order to remove the assumption that the residue field is perfect, note that length is preserved by faithfully flat base change. Thus, we can pass to completion, so that R is a quotient of a formal power series ring $K[[X_1, \dots, X_n]]$, and we can replace R by $R \otimes_K F$, where F is an algebraic closure of K . \square

LEMMA 3.8. *Let $\mathfrak{a} \subset R$ be an \mathfrak{m} -primary ideal and $t > 0$ a real number. Then*

$$\lim_{q \rightarrow \infty} \lambda\left(\frac{R}{\mathfrak{a}^{\lceil tq \rceil}}\right)/q^d = \frac{t^d e(\mathfrak{a})}{d!}.$$

Proof. First note that there exists a sequence of rational numbers $\{k_n/q_n\}$ with denominators of the form $q_n = p^{e_n}$ such that $k_n/q_n \leq t < (k_n + 1)/q_n$, and $q_n < q_{n+1}$, so that $t = \lim_{n \rightarrow \infty} q_n/k_n$. For instance, take $q_n = p^n$, $k_n = \lfloor tp^n \rfloor$.

For n fixed and $q = p^e \gg 0$, we have $k_n \frac{q}{q_n} \leq \lceil tq \rceil \leq (k_n + 1) \frac{q}{q_n}$, and

$$\begin{aligned} \lambda\left(\frac{R}{\mathfrak{a}^{(k_n+1)q/q_n}}\right)/q^d &= e(\mathfrak{a}) \frac{(k_n + 1)^d (q/q_n)^d}{d!} + \mathcal{O}(q^{d-1}), \quad \text{and} \\ \lambda\left(\frac{R}{\mathfrak{a}^{k_n q/q_n}}\right)/q^d &= e(\mathfrak{a}) \frac{(k_n)^d (q/q_n)^d}{d!} + \mathcal{O}(q^{d-1}), \end{aligned}$$

so for all n we have

$$e(\mathfrak{a}) \frac{(k_n)^d}{d! q_n^d} \leq \lim_{q \rightarrow \infty} \lambda\left(\frac{R}{\mathfrak{a}^{\lceil tq \rceil}}\right)/q^d \leq e(\mathfrak{a}) \frac{(k_n + 1)^d}{d! q_n^d}$$

and the desired result follows by taking the limit when $n \rightarrow \infty$. \square

THEOREM 3.9. *If \mathfrak{a}, I are fixed \mathfrak{m} -primary ideals, then $e_{HK}(\mathfrak{a}^t, I)$ is a continuous function of t .*

Proof. Let $t < t'$ be positive real numbers. Then

$$e_{HK}(\mathfrak{a}^{t'}, I) - e_{HK}(\mathfrak{a}^t, I) = \lim_{q \rightarrow \infty} \frac{\lambda(\mathfrak{a}^{\lceil tq \rceil} I^{[q]} / \mathfrak{a}^{\lceil t'q \rceil} I^{[q]})}{q^d}.$$

Let $I = (f_1, \dots, f_n)$. Then we have a composition series

$$\begin{aligned} \mathfrak{a}^{\lceil t'q \rceil} I^{[q]} &\subseteq \mathfrak{a}^{\lceil t'q \rceil} I^{[q]} + f_1^q \mathfrak{a}^{\lceil tq \rceil} \subseteq \dots \subseteq \mathfrak{a}^{\lceil t'q \rceil} I^{[q]} + (f_1^q, \dots, f_i^q) \mathfrak{a}^{\lceil tq \rceil} \\ &\subseteq \dots \subseteq \mathfrak{a}^{\lceil tq \rceil} I^{[q]}. \end{aligned}$$

For $i = 1, \dots, n$, let $K_i = \mathfrak{a}^{\lceil t'q \rceil} I^{[q]} + (f_1^q, \dots, f_{i-1}^q) \mathfrak{a}^{\lceil tq \rceil}$. Then we have

$$\lambda\left(\frac{\mathfrak{a}^{\lceil tq \rceil} I^{[q]}}{\mathfrak{a}^{\lceil t'q \rceil} I^{[q]}}\right) = \sum_{i=1}^n \lambda\left(\frac{K_i + \mathfrak{a}^{\lceil tq \rceil} f_i^q}{K_i}\right).$$

Note that we have a surjective map given by multiplication by f_i^q :

$$\frac{\mathfrak{a}^{[tq]}}{(K_i : f_i^q) \cap \mathfrak{a}^{[tq]}} \rightarrow \frac{\mathfrak{a}^{[tq]} f_i^q}{K_i \cap \mathfrak{a}^{[tq]} f_i^q} \cong \frac{K_i + \mathfrak{a}^{[tq]} f_i^q}{K_i}.$$

It is clear that $\mathfrak{a}^{[t'q]} \subseteq K_i : f_i^q$, so that the length of this term is bounded above by the length of $\mathfrak{a}^{[tq]}/\mathfrak{a}^{[t'q]}$. Thus,

$$\begin{aligned} e_{HK}(\mathfrak{a}^{t'}, I) - e_{HK}(\mathfrak{a}^t, I) &\leq n \lim_{q \rightarrow 0} \frac{\lambda(R/\mathfrak{a}^{[t'q]}) - \lambda(R/\mathfrak{a}^{[tq]})}{q^d} \\ &= ne(\mathfrak{a})(t'^d - t^d) \end{aligned}$$

where the last equality is from Lemma 3.8. \square

We now show how the joint Hilbert-Kunz multiplicity is related to tight closure, integral closure, and \mathfrak{a} -tight closure. The result pertaining to \mathfrak{a} -tight closure, Proposition 3.11 is an analog of testing tight closure via Hilbert-Kunz multiplicities (cf. [HH1], Theorem 8.17), and testing for integral closure via Hilbert-Samuel multiplicities (cf. [NR]).

PROPOSITION 3.10. *If $I \subseteq J \subseteq I^*$ and $\mathfrak{a} \subseteq \mathfrak{b} \subseteq \bar{\mathfrak{a}}$, then $e_{HK}(\mathfrak{a}^t, I) = e_{HK}(\mathfrak{b}^t, J)$ for all $t > 0$.*

Proof. The hypothesis implies that there exists $h \in R^o$, such that $h\mathfrak{b}^{[tq]} J^{[q]} \subseteq \mathfrak{a}^{[tq]} I^{[q]}$ (if $\mathfrak{b} = (b_1, \dots, b_n)$, for each b_i there exists $h_i \in R^o$ such that $h_i b_i^n \in \mathfrak{a}^n$ for all n ; $\mathfrak{b}^{[tq]}$ is generated by $b_1^{i_1} \cdots b_n^{i_n}$ with $i_1 + \cdots + i_n = [tq]$, and choosing $h' = h_1 \cdots h_n$ we have $h' b_1^{i_1} \cdots b_n^{i_n} \in \mathfrak{a}^{i_1} \cdots \mathfrak{a}^{i_n} = \mathfrak{a}^{[tq]}$.) It follows that

$$\lambda\left(\frac{\mathfrak{b}^{[tq]} J^{[q]}}{\mathfrak{a}^{[tq]} I^{[q]}}\right) \leq \lambda\left(\frac{\mathfrak{a}^{[tq]} I^{[q]} : h}{\mathfrak{a}^{[tq]} I^{[q]}}\right) = \lambda\left(\frac{R}{(\mathfrak{a}^{[tq]} I^{[q]}, h)}\right),$$

which is a joint Hilbert-Kunz function over R/h , and thus bounded by $\mathcal{O}(q^{d-1})$. \square

PROPOSITION 3.11. *Assume that R is analytically unramified and formally equidimensional, and has test elements for the usual tight closure. Let \mathfrak{a} , I , J be \mathfrak{m} -primary ideals in R and let $t > 0$. Assume $I \subseteq J$. Then $J \subseteq {}^{\mathfrak{a}^t} I^*$ if and only if $e_{HK}(\mathfrak{a}^t, I) = e_{HK}(\mathfrak{a}^t, J)$.*

Proof. Assume that $J \subseteq \mathfrak{a}^t I^*$, so that $c\mathfrak{a}^{[tq]} J^{[q]} \subseteq \mathfrak{a}^{[tq]} I^{[q]}$ for some $c \in R^o$. Then

$$\lambda\left(\frac{\mathfrak{a}^{[tq]} J^{[q]}}{\mathfrak{a}^{[tq]} I^{[q]}}\right) \leq \lambda\left(\frac{\mathfrak{a}^{[tq]} I^{[q]} : c}{\mathfrak{a}^{[tq]} I^{[q]}}\right) = \lambda\left(\frac{R}{(c, \mathfrak{a}^{[tq]} I^{[q]})}\right),$$

which is a joint Hilbert-Kunz function over the $d-1$ dimensional ring R/c , and therefore it is bounded by $\mathcal{O}(q^{d-1})$. This shows that $e_{HK}(\mathfrak{a}^t, I) = e_{HK}(\mathfrak{a}^t, J)$.

Conversely, assume that $e_{HK}(\mathfrak{a}^t, I) = e_{HK}(\mathfrak{a}^t, J)$. Fix an element $x \in J$. We want to show $x \in \mathfrak{a}^t I^*$.

Fix $q_0 = p^{e_0}$, and fix generators g_1, \dots, g_m for $\mathfrak{a}^{[tq_0]}$. Let f denote the product of a minimal set of generators for \mathfrak{a} , chosen in R^o . Note that $[tq_0q] \leq [tq_0]q$, and thus we have $\mathfrak{a}^{[tq_0]q} \subseteq \mathfrak{a}^{[tq_0q]} \subseteq (\mathfrak{a}^{[tq_0]})^{[q]} : f^q$ (the last inclusion is Observation 3.5 (a.)).

For each g_i , we have

$$\begin{aligned} & \lambda\left(\frac{(\mathfrak{a}^{[tq_0q]} I^{[q_0q]}, \mathfrak{a}^{[tq_0q]} x^{q_0q})}{\mathfrak{a}^{[tq_0q]} I^{[q_0q]}}\right) \\ & \geq \lambda\left(\frac{(\mathfrak{a}^{[tq_0q]} I^{[q_0q]}, g_i^q x^{q_0q})}{\mathfrak{a}^{[tq_0q]} I^{[q_0q]}}\right) = \lambda\left(\frac{R}{\mathfrak{a}^{[tq_0q]} I^{[q_0q]} : g_i^q x^{q_0q}}\right) \\ & \geq \lambda\left(\frac{R}{\mathfrak{a}^{[tq_0]^{[q]}} I^{[q_0q]} : (fg_i x^{q_0})^q}\right) = \lambda\left(\frac{(\mathfrak{a}^{[tq_0]} I^{[q_0]}, fg_i x^{q_0})^{[q]}}{(\mathfrak{a}^{[tq_0]} I^{[q_0]})^{[q]}}\right). \end{aligned}$$

On the other hand, our assumption implies that

$$\lambda\left(\frac{(\mathfrak{a}^{[tq_0q]} I^{[q_0q]}, \mathfrak{a}^{[tq_0q]} x^{q_0q})}{\mathfrak{a}^{[tq_0q]} I^{[q_0q]}}\right) \leq \lambda\left(\frac{\mathfrak{a}^{[tq_0q]} J^{[q_0q]}}{\mathfrak{a}^{[tq_0q]} I^{[q_0q]}}\right)$$

is bounded above by $\mathcal{O}(q^{d-1})$. Since q_0 is fixed, Theorem 8.17 in [HH1] implies $f g_i x^{q_0} \subseteq (\mathfrak{a}^{[tq_0]} I^{[q_0]})^*$.

Since g_i ranges through the generators of $\mathfrak{a}^{[tq_0]}$, we have $f \mathfrak{a}^{[tq_0]} x^{q_0} \subseteq (\mathfrak{a}^{[tq_0]} I^{[q_0]})^*$. But this is true for all q_0 ; if we let $c \in R^o$ be a test element for R , we get

$$c f \mathfrak{a}^{[tq]} x^q \subseteq \mathfrak{a}^{[tq]} I^{[q]}$$

for all $q \gg 0$. Since $f \in R^o$, this gives the desired conclusion. \square

We end this section with some formulas relating the joint Hilbert-Kunz multiplicity to the usual Hilbert-Kunz multiplicity and Hilbert-Samuel multiplicity.

THEOREM 3.12. *Assume that $\dim(R) \geq 1$. Let \mathfrak{a} , $I \subset R$ be \mathfrak{m} -primary ideals.*

a. *For all $t > 0$ we have*

$$(1) \quad e_{HK}(\mathfrak{a}^t, I) \leq e_{HK}(I) + \frac{\ell e(\mathfrak{a})t^d}{d!},$$

where ℓ denotes the $*$ -spread of I , i.e. the minimal number of generators of an ideal J minimal with respect to the condition $J^* \supseteq I$.

b. *If we assume that R is excellent and analytically irreducible then there exists a $t_0 > 0$ such that the inequality in part a. is equality for all $0 \leq t \leq t_0$.*

c.

$$\lim_{t \rightarrow \infty} \frac{e_{HK}(\mathfrak{a}^t, I)}{t^d} = \frac{e(\mathfrak{a})}{d!}.$$

In particular, if $\ell > 1$ then the inequality in part a. is strict for $t \gg 0$.

Proof. First note that we can replace I by any ideal J with $J \subseteq I \subseteq J^*$ without affecting the result. Thus, we may assume that $I = (f_1, \dots, f_\ell)$, where f_1, \dots, f_ℓ are $*$ -independent, i.e. none of them is in the tight closure of the ideal generated by the others. We can also choose all $f_i \in R^o$ by prime avoidance.

We have a filtration

$$\mathfrak{a}^{[tq]} I^{[q]} \subseteq (\mathfrak{a}^{[tq]} I^{[q]}, f_1^q) \subseteq \dots \subseteq (\mathfrak{a}^{[tq]} I^{[q]}, f_1^q, \dots, f_{\ell-1}^q) \subseteq I^{[q]},$$

and therefore we have

$$(2) \quad \lambda\left(\frac{I^{[q]}}{\mathfrak{a}^{[tq]} I^{[q]}}\right) = \sum_{i=1}^{\ell} \lambda\left(\frac{R}{(\mathfrak{a}^{[tq]} I^{[q]}, f_1^q, \dots, f_{i-1}^q) : f_i^q}\right).$$

Since the denominator in each term in the right hand sum contains $\mathfrak{a}^{[tq]}$, the inequality follows by Lemma 3.8.

The second statement follows from Theorem 3.5 (a) in [Vr].

In order to see the last statement, it is enough to restrict to integer exponents t . Note that the denominators appearing in the terms on the right hand side of Equation (2) contain $(\mathfrak{a}^{tq}, f_1^q, \dots, f_{i-1}^q)$, and thus

$$\lambda\left(\frac{I^{[q]}}{\mathfrak{a}^{tq} I^{[q]}}\right) \leq \sum_{i=1}^{\ell} \lambda\left(\frac{R}{(\mathfrak{a}^{tq}, f_1^q, \dots, f_{i-1}^q)}\right)$$

Consider $i > 1$. We have

$$\begin{aligned} \lim_{q \rightarrow \infty} \lambda \left(\frac{R}{(\mathfrak{a}^{tq}, f_1^q, \dots, f_{i-1}^q)} \right) / q^d &\leq e_{HK}((\mathfrak{a}^t, f_1, \dots, f_{i-1})) \\ &\leq \lambda \left(\frac{R}{(\mathfrak{a}^t, f_1, \dots, f_{i-1})} \right) e_{HK}(\mathfrak{m}) \end{aligned}$$

(the last inequality follows by taking a filtration of $R/(\mathfrak{a}^t, f_1, \dots, f_{i-1})$ with quotients equal to R/\mathfrak{m} ; also see Lemma 4.2 in [WY]). As a function of t , $\lambda(R/(\mathfrak{a}^t, f_1, \dots, f_{i-1}))$ is a Hilbert-Samuel function over the ring $R/(f_1, \dots, f_{i-1})$, which has Krull dimension less than d , and therefore dividing by t^d and taking the limit when $t \rightarrow \infty$ yields a limit equal to zero for each of the terms corresponding to $i > 1$ in Equation (2).

Thus we have

$$\begin{aligned} \limsup_{t \rightarrow \infty} \frac{e_{HK}(\mathfrak{a}^t, I) - e_{HK}(I)}{t^d} &= \limsup_{t \rightarrow \infty} \lim_{q \rightarrow \infty} \lambda \left(\frac{R}{\mathfrak{a}^{tq} I^{[q]} : f_1^q} \right) / t^d q^d \\ &\leq \lim_{t, q \rightarrow \infty} \lambda \left(\frac{R}{\mathfrak{a}^{tq}} \right) / t^d q^d = \frac{e(\mathfrak{a})}{d!}. \end{aligned}$$

On the other hand, we have $\mathfrak{a}^{tq} I^{[q]} \subset \mathfrak{a}^{tq}$, and thus $e_{HK}(\mathfrak{a}^t, I) \geq t^d e(\mathfrak{a})/d!$. This proves the equality in part c. \square

The following provides a concrete example where part b. in Theorem 3.12 works with $t_0 = 1$.

EXAMPLE 3.13. Assume (R, \mathfrak{m}) is a Cohen-Macaulay ring, and let $\mathfrak{a} = I = (x_1, \dots, x_d)$ be generated by a regular sequence. If $0 \leq t \leq 1$, then

$$e_{HK}(\mathfrak{a}^t, I) = \frac{t^d e(\mathfrak{a})}{(d-1)!} + e(\mathfrak{a}).$$

More generally, assume that (R, \mathfrak{m}) is excellent equidimensional reduced local ring, and x_1, \dots, x_d is a system of parameters. If $\mathfrak{a} = I = (x_1, \dots, x_d)$, and $0 \leq t \leq 1$, then

$$e_{HK}(\mathfrak{a}^t, I) = \frac{t^d e(\mathfrak{a})}{(d-1)!} + e(\mathfrak{a}).^1$$

¹We are grateful to the referee for pointing out this more general statement.

Proof. For this choice of \mathfrak{a} and I , each term in the sum on the right hand side of Equation (2) for $0 \leq t \leq 1$ is equal to $\lambda(R/\mathfrak{a}^{\lceil tq \rceil})$, and therefore

$$\lambda\left(\frac{R}{\mathfrak{a}^{\lceil tq \rceil} I^{\lceil q \rceil}}\right) = d\lambda\left(\frac{R}{\mathfrak{a}^{\lceil tq \rceil}}\right) + \lambda\left(\frac{R}{I^{\lceil q \rceil}}\right).$$

To prove the general case, note that the i th term in the sum on the right hand side of Equation (2) is

$$\begin{aligned} \mathfrak{b}_{i,q} &:= (\mathfrak{a}^{\lceil tq \rceil}, x_1^q, \dots, x_{i_1}^q) : x_i^q \subseteq \mathfrak{a}^{\lceil tq \rceil} + (x_1^q, \dots, x_{i-1}^q) : x_i^q \\ &\subseteq \mathfrak{a}^{\lceil tq \rceil} + (x_1^q, \dots, x_{i-1}^q)^* \subseteq \mathfrak{a}^{\lceil tq \rceil} + (\mathfrak{a}^q)^* \end{aligned}$$

by the colon capturing property of systems of parameters.

Let c be a test element for R . By the Artin-Rees Lemma, there exists an $s > 0$ such that $c(\mathfrak{a}^q)^* \subseteq (c) \cap \mathfrak{a}^q \subseteq (c)\mathfrak{a}^{q-s}$, and therefore $(\mathfrak{a}^q)^* \subseteq \mathfrak{a}^{q-s}$. We have $\mathfrak{a}^{\lceil tq \rceil} \subseteq \mathfrak{b}_{i,q} \subseteq \mathfrak{a}^{\lceil tq \rceil} + \mathfrak{a}^{q-s}$.

If $t < 1$, then for $q \gg 0$ we have $\mathfrak{a}^{q-s} \subseteq \mathfrak{a}^{\lceil tq \rceil}$, and we get $\mathfrak{b}_{i,q} = \mathfrak{a}^{\lceil tq \rceil}$ for all i . If $t = 1$, then

$$\frac{\lambda(R/\mathfrak{a}^q)}{q^d} \leq \frac{\lambda(R/\mathfrak{b}_{i,q})}{q^d} \leq \frac{\lambda(R/\mathfrak{a}^{q-s})}{q^d}.$$

The limit of each of the outer terms in the inequality above when $q \rightarrow \infty$ is equal to $e(\mathfrak{a})/d!$, and the result follows. \square

§4. Test ideals

The main result of this section, Theorem 4.3 shows that the test ideal for the new version of \mathfrak{a} -tight closure coincides with the test ideal for the Hara-Yoshida \mathfrak{a} -tight closure for R_+ -primary ideals \mathfrak{a} in a graded Gorenstein ring.

For the first lemma below, we recall that a local ring R is approximately Gorenstein if there exists a sequence of \mathfrak{m} -primary irreducible ideals $\{I_t\}$, such that for every k there exists t with $\mathfrak{m}^k \subseteq I_t$. This is a relatively weak condition; for instance normal rings are approximately Gorenstein. Approximately Gorenstein rings were studied in [Ho].

LEMMA 4.1. *Let (R, \mathfrak{m}) be a local approximately Gorenstein ring of characteristic $p > 0$. Let $\{I_t\}$ be a sequence of \mathfrak{m} -primary irreducible ideals, such that for every k there exists t with $\mathfrak{m}^k \subseteq I_t$.*

Then $\tau(\mathfrak{a}) = \bigcap_t (I_t : I_t^{\mathfrak{a}})$ and $T_{\mathfrak{a}} = \bigcap_t (I_t : {}^{\mathfrak{a}}I_t^*)$.*

Proof. We'll prove the second statement (the proof for the first one is slightly easier). The inclusion $T_{\mathfrak{a}} \subseteq \bigcap (I_t : {}^{\mathfrak{a}}I_t^*)$ is clear by definition. Consider $c \in \bigcap (I_t : {}^{\mathfrak{a}}I_t^*)$. First we show that $c({}^{\mathfrak{a}}I^*) \subseteq I$, where I is an arbitrary \mathfrak{m} -primary ideal. The assumption guarantees that there exists t such that $I_t \subseteq I$, and since I_t is irreducible, we can write $I = I_t : K$ for some ideal K . Let $x \in {}^{\mathfrak{a}}I^*$. Then there exists $d \in R^{\circ}$ such that $dx^q \mathfrak{a}^q \subseteq \mathfrak{a}^q I^{[q]}$. Thus, $dx^q K^{[q]} \mathfrak{a}^q \subseteq \mathfrak{a}^q (KI)^{[q]} \subseteq \mathfrak{a}^q I_t^{[q]}$, which shows that $xK \subseteq {}^{\mathfrak{a}}(I_t)^*$. We have $cxK \subseteq I_t$ by the choice of c , and thus $cx \in I_t : K = I$.

Now consider I an arbitrary ideal. We can write $I = \bigcap_n (I + \mathfrak{m}^n)$, an intersection of \mathfrak{m} -primary ideals. Let $x \in {}^{\mathfrak{a}}I^*$. We need to show that $cx \in I$. Note that $x \in {}^{\mathfrak{a}}(I + \mathfrak{m}^n)^*$ for all n , and therefore $cx \in I + \mathfrak{m}^n$ for all n since we have already proved this for \mathfrak{m} -primary ideals. Intersecting over all n yields the desired conclusion. \square

Throughout the rest of this section, R will be assumed to be a Gorenstein positively graded algebra over a field of Krull dimension d and a -invariant a . We let x_1, \dots, x_d be a system of parameters with $\deg(x_i) = \alpha$ for all i , and $I_t := (x_1^t, \dots, x_d^t)$. Let u denote a homogeneous socle generator for (x_1, \dots, x_d) , i.e. $u \in (x_1, \dots, x_d) : R_+ \setminus (x_1, \dots, x_d)$, and let $\delta := \deg(u)$. Note that $\delta = \alpha d + a$, since the a -invariant may be defined as the degree of

$$\left[\frac{u}{x_1 \cdots x_d} \right] \in H_{R_+}^d(R).$$

Note that $I_t^{[q]} = I_{tq}$, and its socle is generated by $(x_1 \cdots x_d)^{tq-1}u$. We will use δ_t to denote the degree of the socle generator for I_t . More precisely, $\delta_t = \deg((x_1 \cdots x_d)^{t-1}u) = (t-1)\alpha d + \delta$. Note that we have $\delta_{tq} = q\delta_t - (q-1)a$.

Fix the notation established before Proposition 2.9.

LEMMA 4.2. *With notations as above, we have*

$$I_t : R_+^N \subseteq I_t + R_{\geq \delta_t - (N-1)\beta}.$$

Moreover, if R is standard graded, then we have equality.

Proof. We prove the claim by induction on N . For the case $N = 1$, $I_t : R_+ = (I_t, (x_1 \cdots x_d)^{t-1}u) \subseteq I_t + R_{\geq \delta_t}$ by the definition of δ_t .

To see that the other inclusion holds in the standard graded case, note that every homogeneous element not in I_t must have a multiple in the socle of I_t , and thus must have degree $\leq \delta_t$.

Assume the claim is true for $N - 1$. Note that $I_t : R_+^N = (I_t : R_+^{N-1}) : R_+$. By the induction hypothesis, we can write $I_t : R_+^{N-1} = (I_t, v_1, \dots, v_r)$ with $\deg(v_i) \geq \delta_t - (N-2)\beta$ for all i . If $v \in (I_t : R_+^N) \setminus I_t : R_+$, then we have $vy_j \equiv a_1v_1 + \dots + a_rv_r \pmod{I_t}$ for some $1 \leq j \leq s$, where $a_i \in R$ can be assumed homogeneous and not all zero. Thus, $\deg(v) + \deg(y_j) \geq \deg(v_i)$ for some i , and the desired inclusion follows.

For the other inclusion in the standard graded case: if $x \in R_{\geq \delta_t - (N-1)}$, then for all $y \in R_+$ we have $xy \in R_{\geq \delta_t - (N-2)} \subseteq I_t : (R_+)^{N-1}$ by the induction hypothesis. \square

THEOREM 4.3. *Let R be a Gorenstein finitely generated graded algebra over a field of positive characteristic. Assume that the Krull dimension d of R is at least 2. Let \mathfrak{a} be a homogeneous ideal which is primary to R_+ . Then $\tau(\mathfrak{a}) = T_{\mathfrak{a}}$.*

NOTE 4.4. The statement of the theorem is not true if the Krull dimension is $d = 1$, since then we can take $\mathfrak{a} = (f)$ to be a principal ideal, and we have ${}^{\mathfrak{a}}I^* = I^*$ and $I^{*\mathfrak{a}} = I^* : f$ for every ideal I . It follows that $T_{\mathfrak{a}} = \tau$, and $\tau(\mathfrak{a}) = \bigcap_I (I : (I^* : f)) = \bigcap_I (I : (I : \tau f)) = \bigcap_I (I, \tau f) = \tau f$, where the intersection is taken over all parameter ideals I (see Lemma 4.1).

Proof. Fix $c \in R^o$ a homogeneous element such that for all ideals $I \subset R$ we have $x \in I^{*\mathfrak{a}} \Rightarrow cx^q \mathfrak{a}^q \subseteq I^{[q]}$. Such a c is called a test element for \mathfrak{a} -tight closure, and the existence of such an element is guaranteed by Theorem 1.7 in [HY]. Fix $k \geq l$ integers such that $R_+^k \subseteq \mathfrak{a} \subseteq R_+^l$.

Due to Lemma 4.1, it is enough to prove that $I_t^{*\mathfrak{a}} = {}^{\mathfrak{a}}I_t^*$ for all $t \gg 0$. Since both $I_t^{*\mathfrak{a}}$ and ${}^{\mathfrak{a}}I_t^*$ are homogeneous ideals, Proposition 2.9 implies that it is enough to show that $x \in I_t^{*\mathfrak{a}} \Rightarrow \deg(x) \geq k\beta - l\beta' + t\alpha$ when $t \gg 0$.

Since $R_+^k \subseteq \mathfrak{a}$, we have $x \in I_t^{*\mathfrak{a}} \Rightarrow cR_+^{kq}x^q \subseteq I_t^{[q]}$. Thus, it follows that $cx^q \in I_{tq} : R_+^{kq}$.

Applying Lemma 4.2, we see that $x \in I_t^{*\mathfrak{a}}$ implies that for all $q = p^e$ we have either $cx^q \in I_t^{[q]}$, or else $\deg(c) + q \deg(x) \geq d(tq - 1)\alpha + \delta - (kq - 1)\beta$. If $cx^q \in I_t^{[q]}$ for all $q = p^e \gg 0$, then $x \in I_t^* \subseteq {}^{\mathfrak{a}}(I_t)^*$. Otherwise, it follows that $\deg(x) \geq dt\alpha - k\beta$. Since $d > 1$, when we choose $t \gg 0$ we have $dt\alpha - k\beta \geq t\alpha + k\beta - l\beta'$, and thus Proposition 2.9 can be applied to show that $x \in {}^{\mathfrak{a}}I_t^*$. \square

We end this section with explicit an computation of test ideals for $\mathfrak{a} = R_+^r$, when R is a Gorenstein graded ring. We note that our result is similar to that in Proposition 5.8 in [HY], but under different assumptions.

PROPOSITION 4.5. *Let R be a standard graded Gorenstein algebra over a field. With notations as above, we have*

$$I_t^{*R^r} = I_t^* + I_t : R_+^{a+1+[r]}$$

for all $r \geq 0$. Thus, $\tau(R_+^r) = \tau(R) \cap R_+^{a+1+[r]}$.

In particular, if R is F-rational, we have $\tau(R) = R$, and thus $\tau(R_+^r) = R_+^{a+1+[r]}$. This also follows from Proposition 5.8 in [HY], where R is not necessarily graded (instead, F-rationality of the Rees ring $R[R_+t]$ is required).

Also, the results of [HS] and [H1] imply that when R is obtained from a characteristic zero ring by reduction to positive characteristic $p \gg 0$, we have $\tau(R) = R_+^{a+1}$, and thus $\tau(R_+^r) = R_+^{a+1+[r]}$ also holds.

Proof. Let $x \in I_t^{*R^r}$ be a homogeneous element, so that $cx^q R_+^{[rq]} \subseteq I_t^{[q]} = I_{tq}$ for some homogeneous $c \in R^o$. Then $cx^q \in I_{tq} : R_+^{[rq]} = I_{tq} + R_{\geq \delta_{tq} - [rq] + 1}$ be Lemma 4.2. Thus we either have $cx^q \in I_t^{[q]}$ for all $q \gg 0$, in which case $x \in I_t^*$, or else we have $\deg(c) + q \deg(x) \geq \delta_{tq} - [rq] + 1 = q\delta_t - (q-1)a - [rq] + 1$ for infinitely many values of $q = p^e$. Dividing each side by q and taking the limits when $q \rightarrow \infty$ yields $\deg(x) \geq \delta_t - a - r$, and since $\deg(x)$ is an integer, this means $\deg(x) \geq \delta_t - a - [r]$. For every homogeneous element $y \in R_{a+1+[r]}$, we have $xy \in R_{\geq \delta_t + 1} \subset I_t$. This proves $I_t^{*R^r} \subseteq I_t^* + I_t : R_+^{a+1+[r]}$.

Conversely, consider $x \in I_t : R_+^{a+1+[r]} = I_t + R_{\geq \delta_t - a - [r]}$. If $x \in I_t$, there is nothing to prove. Otherwise, we have $\deg(x^q) \geq q\delta_t - aq - [r]q \geq q\delta_t - aq - [rq]$. Choosing $c \in R_{\geq a+1}$ yields $\deg(cx^q) \geq \delta_{tq} - [rq] + 1$, so that $cx^q R_+^{[rq]} \subseteq R_{\geq \delta_{tq} + 1} \subset I_t^{[q]}$, and thus $x \in I_t^{*R^r}$. We note that this inclusion can also be obtained as a Corollary of Theorem 2.7 in [HY]. \square

§5. Jumping numbers

The results of this section address the following question:

QUESTION. Given ideals \mathfrak{a} , $I \subset R$, and a fixed $t_0 \geq 0$, does there exist an $\epsilon > 0$ such that $I^{*\mathfrak{a}^t} = I^{*\mathfrak{a}^{t_0}}$, and $\mathfrak{a}^t I^* = \mathfrak{a}^{t_0} I^*$ for all $t \in [t_0, t_0 + \epsilon]$?

This question is somewhat related to the notion of jumping numbers for test ideals. The jumping numbers are defined to be the positive real

numbers c such that $\tau(\mathfrak{a}^c) \neq \tau(\mathfrak{a}^{c-\epsilon})$ for any $\epsilon > 0$ (a similar notion for multiplier ideals has been introduced in [ELSV]). These have been studied extensively in recent research ([MTW], [BMS]). In our context, if for a given t_0 an ϵ can be found that does not depend on the ideal I , it follows that there are no jumping numbers between t_0 and $t_0 + \epsilon$. We give positive answers to our question in several particular cases. A positive answer implies that for a given I , $I^{*\mathfrak{a}^t}$ and $\mathfrak{a}^t I^*$ are constant on intervals of the form $[t_0, t_1)$. We will call t_1 a jumping number for the ideal I if $I^{*\mathfrak{a}^{t_0}} = I^{*\mathfrak{a}^t}$ for all $t \in [t_0, t_1)$ for some $t_0 < t_1$, but $I^{*\mathfrak{a}^{t_0}} \neq I^{*\mathfrak{a}^{t_1}}$ (or $\mathfrak{a}^{t_0} I^* = \mathfrak{a}^t I^*$ but $\mathfrak{a}^{t_0} I^* \neq \mathfrak{a}^{t_1} I^*$).

The following observation shows that it will be enough to check only one inclusion in order to answer the above question in the affirmative.

OBSERVATION 5.1. *Let $I, \mathfrak{a} \subset R$ be fixed ideals, and $0 \leq t < t'$ real numbers. Then $I^{*\mathfrak{a}^t} \subseteq I^{*\mathfrak{a}^{t'}}$, and $\mathfrak{a}^t I^* \subseteq \mathfrak{a}^{t'} I^*$.*

However, note that it is not always true that $\mathfrak{a} \subseteq \mathfrak{b} \Rightarrow {}^b I^ \subseteq {}^a I^*$, while the corresponding statement is trivially true for the Hara-Yoshida version.*

Proof. The statement for the Hara-Yoshida version is trivial, since $\mathfrak{a}^{[t'q]} \subseteq \mathfrak{a}^{[tq]}$.

Consider $x \in \mathfrak{a}^t I^*$, so that $cx^q \mathfrak{a}^{[tq]} \subseteq \mathfrak{a}^{[tq]} I^{[q]}$. Multiplying each side by arbitrary elements in $\mathfrak{a}^{[t'q]-[tq]}$ yields the desired conclusion.

For the claim in the last paragraph, take for example $\mathfrak{a} = (f)$, with $f \in (\mathfrak{b}) := (x, y)^2$, $I = (x^2, y^2)$, in the ring $R = k[x, y]$. Then ${}^b I^* = (x^2, y^2, xy)$, while ${}^a I^* = I$. \square

PROPOSITION 5.2. *Assume that (R, \mathfrak{m}) is an excellent reduced equidimensional local ring, and \mathfrak{a}, I are \mathfrak{m} -primary ideals. Then for every $t_0 \geq 0$, there exists $\epsilon > 0$ such that $\mathfrak{a}^t I^* = \mathfrak{a}^{t_0} I^*$ for all $t \in [t_0, t_0 + \epsilon]$.*

Proof. First note that for each $x \notin \mathfrak{a}^{t_0} I^*$, there exists $\epsilon > 0$ such that $x \notin \mathfrak{a}^{t_0+\epsilon} I^*$. This follows from Theorem 3.9, and Proposition 3.11, applied to the ideals I and $J = (I, x)$.

Construct a sequence $t_1 > t_2 > \dots > t_n > \dots > t_0$ recursively as follows: Choose $x_1 \notin \mathfrak{a}^{t_0} I^*$, and let $t_1 > t_0$ such that $x_1 \notin \mathfrak{a}^{t_1} I^*$ (the existence of such a t_1 is guaranteed by the previous claim). If t_1, \dots, t_k have been constructed, then we either have $\mathfrak{a}^{t_0} I^* = \mathfrak{a}^{t_k} I^*$, in which case the proof is complete (take $\epsilon = t_k - t_0$), or else we can choose an $x_{k+1} \in \mathfrak{a}^{t_k} I^* \setminus \mathfrak{a}^{t_0} I^*$, and, by the previous claim, there exists $t'_{k+1} > t_0$ such that $x_{k+1} \notin \mathfrak{a}^{t'_{k+1}} I^*$

(note that we must have $t_k > t_{k+1}$ by Observation 5.1). Thus, we have a chain of ideals $\mathfrak{a}^{t_0} I^* \subseteq \dots \subseteq \mathfrak{a}^{t_n} I^* \subseteq \dots \subseteq \mathfrak{a}^{t_2} I^* \subseteq \mathfrak{a}^{t_1} I^*$. Note that the construction of t_k shows that the inclusions are strict unless the recursive process stops. This contradicts the fact that $\mathfrak{a}^{t_0} I^*$ is \mathfrak{m} -primary. \square

The next result deals with the case when $\mathfrak{a} = (f)$ is a principal ideal, with $f \in R^o$. Note that in this case we only need to consider the Hara-Yoshida version, since $\mathfrak{a}^t I^* = I^*$ for all t . It turns out that a positive answer to the question considered here is related to the existence of test exponents. We review the definition.

DEFINITION 5.3. Let $I \subset R$ be an ideal, and $c \in R^o$ a test element for the usual tight closure. We say that $q_0 = p^{e_0}$ is a test exponent for the pair (I, c) if $cx^q \in I^{[q]}$ for any one choice of $q \geq q_0$ implies $x \in I^*$.

Test exponents were introduced in [HH2], where it is shown that their existence is closely related to the localization problem for tight closure.

LEMMA 5.4. Assume $\mathfrak{a} = (f)$ is a principal ideal, with $f \in R^o$. Assume that R has test elements for the usual tight closure. Then $x \in I^{*\mathfrak{a}^t} \Leftrightarrow x^q f^{[tq]} \in (I^{[q]})^*$ for all q .

In particular, if there exists q_1 such that $tq_1 \in \mathbf{Z}$, then $x \in I^{*\mathfrak{a}^t} \Leftrightarrow x^{q_1} f^{tq_1} \in (I^{[q_1]})^*$.

Proof. Note that the following inequalities hold for all q :

$$[tq] - 1 < tq \leq [tq] < tq + 1.$$

It follows that

$$q_1 [tq_2] - q_1 < [tq_1 q_2] < q_1 [tq_2] + 1.$$

Assume that $x \in I^{*\mathfrak{a}^t}$, and let $q = q_1 q_2$. Then there exists $c \in R^o$ such that $cx^{q_1 q_2} f^{[tq_1 q_2]} \in I^{[q_1 q_2]}$, which implies $cfx^{q_1 q_2} f^{[tq_2]q_1} \in I^{[q_1 q_2]}$. Since $cf \in R^o$, this shows that $x^{q_2} f^{[tq_2]} \in (I^{[q_2]})^*$ for all q_2 .

Conversely, assume that $x^{q_2} f^{[tq_2]} \in (I^{[q_2]})^*$ for some q_2 , and let $c \in R^o$ be a test element for the usual tight closure. Then $cx^{q_1 q_2} f^{[tq_2]q_1} \in I^{[q_1 q_2]}$, which implies $cx^{q_1 q_2} f^{[tq_1 q_2] + q_1} \in I^{[q_1 q_2]}$ for all q_1 . Fix q_1 and allow q_2 to vary $cf^{q_1} x^q f^{[tq]} \in I^{[q]}$ for all $q \gg 0$. Since $cf^{q_1} \in R^o$, this shows that $x \in I^{*\mathfrak{a}^t}$. \square

PROPOSITION 5.5. *Let $\mathfrak{a} = (f)$ with $f \in R^\circ$, $I \subset R$ an arbitrary ideal, and $t_0 \geq 0$. Assume that there exists q_1 such that $t_0 q_1$ is an integer, and assume that $c \in R^\circ$ is a test element for the closure such that there exists a test exponent q_0 for the ideal $I^{[q_1]}$ and the test element cf^{q_1} .*

Then we have $I^{\mathfrak{a}^t} = I^{*\mathfrak{a}^{t_0}}$ for all $t \in [t_0, t_0 + 1/q_0]$.*

Proof. Let $t = t_0 + 1/q_0$. Let $q = q_1 q_0$ so that $tq = t_0 q + q_1$ is an integer. Assume that $x \in I^{*\mathfrak{a}^t}$; by Lemma 5.4, this implies that $x^q f^{t_0 q} f^{q_1} \in (I^{[q]})^*$. Since c is a test element, we have $cf^{q_1} (x^{q_1} f^{t_0 q_1})^{q_0} \in (I^{[q_1]})^{[q_0]}$. Since q_0 is a test exponent, this implies $x^{q_1} f^{t_0 q_1} \in (I^{[q_1]})^*$, and thus $x \in I^{*\mathfrak{a}^{t_0}}$ by Lemma 5.4. \square

COROLLARY 5.6. *If (R, \mathfrak{m}) is a regular local ring, $\mathfrak{a} = (f)$ is a principal ideal, and $t_0 \geq 0$ is such that $t_0 q_1 \in \mathbf{Z}$ for some $q_1 = p^{e_1}$, and q_0 is such that $f^{q_1} \notin \mathfrak{m}^{[q_0]}$, then there are no jumping numbers for the test ideals $\tau(\mathfrak{a}^t)$ in the interval $[t_0, t_0 + 1/q_0]$.*

Proof. Since $I^* = I$ for every ideal I , we can take $c = 1$, and note that a q_0 with the property that $f^{q_1} \notin \mathfrak{m}^{[q_0]}$ is a test exponent for $(I^{[q_1]}, f^{q_1})$ for any ideal I . Indeed, if $x \notin I^{[q_1]}$ and $f^{q_1} x^{q_0} \in I^{[q_1 q_0]} \Rightarrow f^{q_1} \in I^{[q_1 q_0]} : x^{q_0} = (I^{[q_1]} : x)^{[q_0]} \subseteq \mathfrak{m}^{[q_0]}$, contradicting the choice of q_0 . \square

Note that if $t_0 = 0$, then the converse of Proposition 5.5 holds, in the sense that a positive answer to the question discussed here implies existence of test exponents for the usual tight closure. Recall that $I^{*\mathfrak{a}^0} = I^{*R} = I^*$ is the usual tight closure.

PROPOSITION 5.7. *Let $\mathfrak{a} = (f)$ with $f \in R^\circ$ a test element for tight closure, $I \subset R$ an arbitrary ideal.*

Assume that $q_0 = p^{e_0}$ is such that $I^{\mathfrak{a}^{1/q_0}} = I^*$. Then q_0 is a test exponent for the pair (I, f) .*

Proof. Let $x \in R$ be such that $fx^{q_1} \in I^{[q_1]}$ for some $q_1 \geq q_0$. Then for all $q \geq q_1$ we have $f^{q/q_1} x^q \in I^{[q]}$, and therefore $f^{q/q_0} x^q \in I^{[q]}$, which shows that $x \in I^{*\mathfrak{a}^{1/q_0}} = I^*$ by assumption. \square

In the next result, I and \mathfrak{a} are arbitrary ideals, but we restrict attention to $t_0 = 0$.

PROPOSITION 5.8. *Assume that (R, \mathfrak{m}) is an excellent analytically irreducible local domain. Let $I, \mathfrak{a} \subset R$ be ideals, $\mathfrak{a} \neq 0$. Let $x \notin I^*$. Then there exists $q_0 = p^{e_0}$ such that $x \notin I^{*\mathfrak{a}^{1/q_0}}$.*

Note that our result is not quite sufficient to give an affirmative answer to the question raised in the beginning of the section for this case, since q_0 is allowed to depend on x .

Proof. By Proposition 2.4 in [Ab], there exists q_1 such that $I^{[q]} : x^q \subseteq \mathfrak{m}^{[q/q_1]}$ for all $q \geq q_1$. Assume by contradiction that $x \in I^{*\mathfrak{a}^{1/q_0}}$ for every q_0 . This means that $cx^q\mathfrak{a}^{q/q_0} \subseteq I^{[q]}$ for all $q \gg 0$, so that $c\mathfrak{a}^{q/q_0} \subseteq I^{[q]} : x^q \subseteq \mathfrak{m}^{[q/q_1]}$. Let $q = q_0q_1Q$. Then we obtain $c\mathfrak{a}^{q_1Q} \subseteq \mathfrak{m}^{[q_0Q]}$, which implies $\mathfrak{a}^{q_1} \subseteq (\mathfrak{m}^{[q_0]})^*$. Since q_1 is fixed, this is false for $q_0 \gg 0$.

At the other end of the spectrum, we ask the following question.

QUESTION. If I, \mathfrak{a} are fixed ideals, and N is such that $\mathfrak{a}^N I^* = \mathfrak{a}^{N'} I^*$ for all $N' \geq N$ (note that such an N exists by the Noetherian property) describe $\mathfrak{a}^N I^*$.

We will use $\mathfrak{a}^\infty I^*$ to denote $\mathfrak{a}^N I^*$ when N is as above. Note that a similar definition for the Hara-Yoshida version of \mathfrak{a} -tight closure would yield the whole ring when I is an \mathfrak{m} -primary ideal, since for $N \gg 0$ we have $\mathfrak{a}^N \subseteq I$, and $\mathfrak{a}^{[Nq/k]} \subseteq I^{[q]}$, where k is the number of generators of \mathfrak{a} . When $\mathfrak{a} = (f)$ is a principal ideal with f a non-zerodivisor, we have $\mathfrak{a}^\infty I^* = I^*$ for every ideal I .

We always have $\mathfrak{a}^\infty I^* \subseteq \overline{I}$. Observation 2.10 implies that when R is standard graded, $\mathfrak{a} = R_+^r$ for some $r > 0$, and I is homogeneous with all generators of the same degree, we have ${}^*\mathfrak{a}^\infty I^* = \overline{I}$. However, Example 2.11 shows that for $R = k[x, y]$, $I = (x^2, y^4)$, and $\mathfrak{a} = (x, y)$ we have $\mathfrak{a}^\infty I^* \neq \overline{I}$. In fact in this example it is easy to check that $\mathfrak{a}^\infty I^* = (x^2, y^4, xy^3)$. More generally, we note the following:

PROPOSITION 5.9. *Let R be a two-dimensional standard graded normal domain, let $I = (f_1, f_2)$ be a homogeneous parameter ideal. Let $d = \max(\deg(f_1), \deg(f_2))$. Then ${}^{R^\infty}_+ I^* = I + R_{\geq d}$.*

Proof. Say that $d = \deg(f_1)$.

Let $x \in {}^{R_+^\infty}I^*$, so $x \in {}^{R_+^n}I^*$ for some n . Assume that $\deg(x) < d$. For some homogeneous $c \in R^o$, and for all y among a minimal set of generators of R_+^{nq} we have $cx^qy = a_1f_1^q + a_2f_2^q$ with $a_1, a_2 \in R_+^{nq}$. If $\deg(x) < d$, it follows by comparing degrees that $cx^qy = a_2f_2^q$, so that $cR_+^{nq}x^q \subseteq R_+^{nq}f_2^q$. But this implies that $x \in \overline{(f_2)} = (f_2)$.

Conversely, assume that $\deg(x) \geq d$. For $n \gg 0$ we have $x \in I^{*R_+^n}$, so that there exists $c \in R^o$ such that for all $y \in R_+^{nq}$, $cx^qy = a_1f_1^q + a_2f_2^q$ with $a_1, a_2 \in R$. Comparing degrees, we see that $a_1, a_2 \in R_+^{nq}$. \square

REFERENCES

- [Ab] I. Aberbach, *Extensions of weakly and strongly F-regular rings by flat maps*, J. Algebra, **241** (2001), 799–807.
- [Bt] P. B. Bhattacharya, *The Hilbert function of two ideals*, Proc. Cambridge Philos. Soc., **53** (1957), 568–575.
- [BMS] M. Blickle, M. Mustata, and K. E. Smith, *Discreteness and rationality of F-thresholds*, preprint.
- [ELSV] L. Ein, R. Lazarsfeld, K. E. Smith and D. Varolin, *Jumping coefficients for multiplier ideals*, Duke Math. J., **123** (2004), 469–506.
- [Ep] N. Epstein, *A tight closure analogue of analytic spread*, Math. Proc. Cambridge Philos. Soc., **139** (2005), no. 2, 371–383.
- [Hn] D. Hanes, *Notes on the Hilbert-Kunz function*, J. Algebra, **265** (2003), no. 2, 619–630.
- [H1] N. Hara, *Characterization of rational singularities in terms of the injectivity of Frobenius*, Amer. J. Math., **120** (1998), no. 5, 981–996.
- [H2] N. Hara, *Geometric interpretation of test ideals*, Trans. Amer. Math. Soc., **353** (2001), no. 5, 1885–1906.
- [HT] N. Hara and S. Tagaki, *On a generalization of test ideals*, Nagoya Math. J., **175** (2004), 59–74.
- [HY] N. Hara and K. Yoshida, *A generalization of tight closure and multiplier ideals*, Trans. Amer. Math. Soc., **355** (2003), no. 8, 3143–3174.
- [Ho] M. Hochster, *Cyclic purity versus purity in excellent Noetherian rings*, Trans. Amer. Math. Soc., **231** (1997), no. 2, 463–488.
- [HH1] M. Hochster and C. Huneke, *Tight closure, invariant theory, and the Briançon-Skoda theorem*, J. Amer. Math. Soc., **3** (1990), 31–116.
- [HH2] M. Hochster and C. Huneke, *Localization and test exponents for tight closure*, Michigan Math. J., **48** (2000), 305–329.
- [HS] C. Huneke and K. Smith, *Tight closure and the Kodaira vanishing theorem*, J. Reine Angew. Math., **484** (1997), 127–152.
- [Mo] P. Monsky, *The Hilbert-Kunz function*, Math. Ann., **263** (1983), 43–49.
- [MTW] M. Mustata, S. Tagaki, and K.-i. Watanabe, *F-thresholds and Bernstein-Sato*

polynomials, European Congress of Mathematics, Eur. Math. Soc., Zrich, 2005, pp. 341–364.

- [NR] D. G. Northcott, and D. Rees, *Reductions of ideals in local rings*, Proc. Cambridge Philos. Soc., **50** (1954), 145–158.
- [S1] K. E. Smith, *Tight closure in graded rings*, J. Math. Kyoto Univ., **37** (1997), no. 1, 35–53.
- [S2] K. E. Smith, *The multiplier ideal is a universal test ideal*, Comm. Algebra, **28** (2000), 5915–5929.
- [Vr] A. Vraciu, **-Independence and special tight closure*, J. Algebra, **249** (2002), 544–565.
- [WY] K. Watanabe and K. Yoshida, *Hilbert-Kunz multiplicity and an inequality between multiplicity and colength*, J. Algebra, **230** (2000), 295–317.

Department of Mathematics
University of South Carolina
Columbia SC 29205
U.S.A.

`vraciu@math.sc.edu`