

EXTENDIBILITY AND STABLE EXTENDIBILITY OF NORMAL BUNDLES ASSOCIATED TO IMMERSIONS OF REAL PROJECTIVE SPACES

Dedicated to the Memory of Professor Katsuo Kawakubo

TEIICHI KOBAYASHI, HARUO MAKI and TOSHIO YOSHIDA

(Received July 28, 2000)

1. Introduction

The extension problem is one of the fundamental problems in topology. We consider the problem for vector bundles over real projective spaces.

Let F be the real field \mathbf{R} , the complex field \mathbf{C} or the quaternion field \mathbf{H} . Let X be a space and A be a subspace. A t -dimensional F -vector bundle ζ over A is called *extendible* (respectively *stably extendible*) to X , if there is a t -dimensional F -vector bundle over X whose restriction to A is equivalent (respectively stably equivalent) to ζ as F -vector bundles, that is, if ζ is equivalent (respectively stably equivalent) to $i^*\alpha$ for a t -dimensional F -vector bundle α over X , where $i: A \rightarrow X$ is the inclusion (cf. [13] and [5]).

As is seen in [7, Theorem 6.4] and [11, Theorem 2.2], the extendibility (or the stable extendibility) is closely related to the span, i.e., the maximum number of linearly independent cross-sections of an F -vector bundle, and one can see in the proof of Theorem C of this paper how the stable extendibility is related to the immersion problem.

Let \mathbf{R}^n be the n -dimensional Euclidean space and $F\mathbf{P}^n$ be the n -dimensional F -projective space. Concerning stably extendible F -vector bundles for $F = \mathbf{R}$ and \mathbf{C} , R.L.E. Schwarzenberger obtained the following results (cf. [2], [3], [7], [12] and [13]).

Theorem (Schwarzenberger). *Let $F = \mathbf{R}$ or \mathbf{C} . If a k -dimensional F -vector bundle ζ over $F\mathbf{P}^n$ is stably extendible to $F\mathbf{P}^m$ for every $m > n$, then ζ is stably equivalent to a sum of k F -line bundles.*

In the original results of Schwarzenberger, the F -vector bundles are assumed to be extendible, but his results are also valid for the stably extendible F -vector bundles.

Recently, M. Imaoka and K. Kuwana have proved in [5] that if a k -dimensional

H -vector bundle ζ over HP^n is stably extendible to HP^m for every $m > n$ and its top non-zero Pontrjagin class is not zero mod 2, then ζ is stably equivalent to a sum of k H -line bundles provided $k \leq n$.

We study the question: Determine the necessary and sufficient condition that a R -vector bundle over RP^n is stably extendible to RP^m for every $m > n$. We have obtained the results for the tangent bundle $\tau = \tau(RP^n)$ of RP^n (cf. [7] and [9]), for the normal bundle ν associated to an immersion of RP^n in R^{2n+1} (cf. [10]) and for the complexification $c\nu$ of ν (cf. [10]) as follows:

- 1) τ is stably extendible to RP^m for every $m > n$ if and only if $n = 1, 3$ or 7 .
- 2) ν is stably extendible to RP^m for every $m > n$ if and only if $1 \leq n \leq 8$.
- 3) $c\nu$ is stably extendible to RP^m for every $m > n$ if and only if $1 \leq n \leq 9$.

The purpose of this paper is to improve 2) and 3) for the normal bundle ν associated to an immersion of RP^n in R^{n+k} where k is any positive integer and for the complexification $c\nu$ of ν .

Let $\phi(n)$ be the number of integers s such that $0 < s \leq n$ and $s \equiv 0, 1, 2$ or $4 \pmod{8}$. Then we have

Theorem A. *Let ν be the normal bundle associated to an immersion of RP^n in R^{n+k} , where $k > 0$. Then ν is stably extendible to RP^m for every $m > n$ if and only if $k \geq 2^{\phi(n)} - n - 1$.*

Theorem B. *Let ν be the normal bundle associated to an immersion of RP^n in R^{n+k} , and let $n+1 \leq k \leq n+12$. Then the following three conditions are equivalent:*

- (1) ν is extendible to RP^m for every $m > n$.
- (2) ν is stably extendible to RP^m for every $m > n$.
- (3) $1 \leq n \leq 8$.

These are improvements of Theorem A in [10].

Let $[x]$ denote the integral part of a real number x . Then for the complexification of the normal bundle, we have

Theorem C. *Let $c\nu$ be the complexification of the normal bundle ν associated to an immersion of RP^n in R^{n+k} , where $k > 0$. Then the following hold.*

- (i) *For $n \geq 6$, $c\nu$ is stably extendible to RP^m for every $m > n$ if and only if $k \geq 2^{\lfloor n/2 \rfloor} - n - 1$.*
- (ii) *For $1 \leq n \leq 5$, $c\nu$ is stably extendible to RP^m for every $m > n$.*

The following is an improvement of Theorem 4.4 in [10].

Theorem D. *Let $c\nu$ be the complexification of the normal bundle ν associated to an immersion of RP^n in R^{n+k} , and let $n \leq k \leq n+8$. Then the following three*

conditions are equivalent:

- (1) $c\nu$ is extendible to \mathbf{RP}^m for every $m > n$.
- (2) $c\nu$ is stably extendible to \mathbf{RP}^m for every $m > n$.
- (3) $1 \leq n \leq 9$.

This note is arranged as follows. In Section 2 we study relations between extendibility and stable extendibility. In Section 3 we prove Theorem A. We prove Theorem B and give some examples in Section 4. In Section 5 we prove Theorem C. We prove Theorem D and give some examples in Section 6.

2. Extendibility and stable extendibility

In the following, we use the same letter for a vector bundle and its equivalence class, and use an integer k for a k -dimensional trivial bundle.

Let d denote $\dim_{\mathbf{R}} F$, where $F = \mathbf{R}, \mathbf{C}$ or \mathbf{H} . The following fact is known (cf. [4, Theorem 1.5, p.100]).

(2.1). If α and β are two t -dimensional F -vector bundles over an n -dimensional CW-complex X such that $\langle (n+2)/d - 1 \rangle \leq t$ and $\alpha \oplus k = \beta \oplus k$ for some k -dimensional trivial F -bundle k over X , then $\alpha = \beta$, where \oplus denotes the Whitney sum and $\langle x \rangle$ denotes the smallest integer m with $x \leq m$.

Theorem 2.2. *Let X be a subcomplex of a finite dimensional CW-complex Y and let ζ be an \mathbf{R} -vector bundle over X such that $\dim \zeta > \dim X$. Then ζ is extendible to Y if and only if ζ is stably extendible to Y .*

In case $\dim \zeta = \dim X$, this does not hold in general.

Proof. The “only if” part is clear. Suppose that ζ is stably equivalent to $i^*(\alpha)$ for some \mathbf{R} -vector bundle α over Y , where $i: X \rightarrow Y$ is the inclusion. In case $\dim \zeta > \dim X$, ζ is equivalent to $i^*(\alpha)$ by (2.1).

A counter example is given by the n -sphere S^n in the $(n + 1)$ -sphere S^{n+1} and the tangent bundle $\tau = \tau(S^n)$ of S^n for $n \neq 1, 3, 7$. In fact, $\tau \oplus 1$ is the $(n + 1)$ -dimensional trivial bundle over S^n and so $\tau \oplus 1 = i^*(n) \oplus 1$, where $i: S^n \rightarrow S^{n+1}$ is the inclusion and n denotes the n -dimensional trivial \mathbf{R} -vector bundle over S^{n+1} . Hence τ is stably extendible to S^{n+1} . On the other hand, if there is an n -dimensional \mathbf{R} -vector bundle α over S^{n+1} such that $\tau = i^*(\alpha)$, τ is trivial, since $i: S^n \rightarrow S^{n+1}$ is homotopic to a constant map. Hence $n = 1, 3$ or 7 . So τ is not extendible to S^{n+1} for $n \neq 1, 3, 7$. □

The following is proved in the way similar to the former part of the proof of Theorem 2.2.

Theorem 2.3. *Let X be a subcomplex of a finite dimensional CW-complex Y and let ζ be a \mathbf{C} -vector bundle over X such that $\dim \zeta \geq \langle (\dim X)/2 \rangle$. Then ζ is extendible to Y if and only if ζ is stably extendible to Y .*

Corollary 2.4. *Let M be a submanifold of a finite dimensional differentiable manifold N and $c\tau(M)$ be the complexification of the tangent bundle $\tau(M)$ of M . Then $c\tau(M)$ is extendible to N if and only if $\tau(M)$ is stably extendible to N .*

3. Proof of Theorem A

Let ξ_n be the canonical line bundle over $\mathbf{R}P^n$.

Lemma 3.1. *Let ν be the normal bundle associated to an immersion of $\mathbf{R}P^n$ in \mathbf{R}^{n+k} , where $k > 0$. Then the equality*

$$\nu = (a2^{\phi(n)} - n - 1)\xi_n + n + k + 1 - a2^{\phi(n)}$$

holds in $KO(\mathbf{R}P^n)$, where a is any integer.

Proof. Let $\tau = \tau(\mathbf{R}P^n)$ be the tangent bundle of $\mathbf{R}P^n$. Then we have $\tau \oplus \nu = n + k$ and $\tau \oplus 1 = (n + 1)\xi_n$. Hence

$$\nu = n + k + 1 - (n + 1)\xi_n = (a2^{\phi(n)} - n - 1)\xi_n + n + k + 1 - a2^{\phi(n)}$$

in $KO(\mathbf{R}P^n)$ for any integer a , since $\xi_n - 1$ is of order $2^{\phi(n)}$ (cf. [1, Theorem 7.4]). \square

Theorem 3.2. *Let ν be the normal bundle associated to an immersion of $\mathbf{R}P^n$ in \mathbf{R}^{n+k} , where $k > 0$. Then ν is stably extendible to $\mathbf{R}P^m$ for every $m > n$ if $k \geq 2^{\phi(n)} - n - 1$, and if $k > n$, in addition, ν is extendible to $\mathbf{R}P^m$ for every $m > n$.*

Proof. By Lemma 3.1, we have $\nu = A\xi_n + B$, where $A = 2^{\phi(n)} - n - 1$ and $B = n + k + 1 - 2^{\phi(n)}$. Clearly $A \geq 0$, and $B \geq 0$ by the assumption. For $m > n$, $i^*(A\xi_m \oplus B) = A\xi_n \oplus B$, where $i: \mathbf{R}P^n \rightarrow \mathbf{R}P^m$ is the standard inclusion. Hence ν is stably extendible to $\mathbf{R}P^m$ for every $m > n$, since ν is stably equivalent to $A\xi_n \oplus B$. If $k > n$, in addition, $\dim \mathbf{R}P^n = n < k = \dim \nu = A + B$, and so we obtain $\nu = A\xi_n \oplus B$ by (2.1). Thus ν is extendible to $\mathbf{R}P^m$ for every $m > n$. \square

The following result ([9, Theorem 4.1]) is the ‘‘stably extendible version’’ of Theorem 6.2 in [7].

(3.3). Let ζ be a t -dimensional \mathbf{R} -vector bundle over $\mathbf{R}P^n$. Assume that there is a positive integer l such that ζ is stably equivalent to $(t + l)\xi_n$ and $t + l < 2^{\phi(n)}$. Then

$n < t + l$ and ζ is not stably extendible to \mathbf{RP}^{t+l} .

Using (3.3), we have obtained the following in [10, Theorem 2.4] (cf. [11, Proposition 6.4(iii)(b)]).

(3.4). The normal bundle associated to an immersion of \mathbf{RP}^n in \mathbf{R}^{n+k} is not stably extendible to \mathbf{RP}^{n+k+1} , if $0 < k < 2^{\phi(n)} - n - 1$.

Theorem 3.5. *Let ν be the normal bundle associated to an immersion of \mathbf{RP}^n in \mathbf{R}^{n+k} . Then ν is not stably extendible to \mathbf{RP}^m for $m = \min\{2^{\phi(n)} - n - 1, n + k + 1\}$, if $0 < k < 2^{\phi(n)} - n - 1$.*

Proof. Put $\zeta = \nu$, $t = k$ and $l = 2^{\phi(n)} - n - k - 1$ in (3.3). Then clearly $t + l < 2^{\phi(n)}$, and $l > 0$ by the assumption. So ν is not stably extendible to \mathbf{RP}^m for $m = 2^{\phi(n)} - n - 1$. By (3.4), ν is not stably extendible to \mathbf{RP}^m for $m = n + k + 1$. \square

Putting $n = 9$ in Theorem 3.5, we have

Corollary 3.6. *If $1 \leq k \leq 21$, the normal bundle associated to an immersion of \mathbf{RP}^9 in \mathbf{R}^{9+k} is not stably extendible to \mathbf{RP}^m for $m = \min\{22, k + 10\}$.*

Proof of Theorem A. The “if” part follows from Theorem 3.2 and the “only if” part follows from Theorem 3.5. \square

4. Proof of Theorem B

Let ξ_n be the canonical line bundle over \mathbf{RP}^n .

Theorem 4.1. *Let $\nu = \nu(f_n)$ be the normal bundle associated to an immersion $f_n: \mathbf{RP}^n \rightarrow \mathbf{R}^{n+k}$, where $k > 0$. Then, for $1 \leq n \leq 10$, we have the equalities*

$$\begin{aligned} \nu(f_1) &= k, & \nu(f_2) &= \xi_2 + k - 1, & \nu(f_3) &= k, \\ \nu(f_4) &= 3\xi_4 + k - 3, & \nu(f_5) &= 2\xi_5 + k - 2, & \nu(f_6) &= \xi_6 + k - 1, \\ \nu(f_7) &= k, & \nu(f_8) &= 7\xi_8 + k - 7, & \nu(f_9) &= 22\xi_9 + k - 22 \\ \text{and } \nu(f_{10}) &= 53\xi_{10} + k - 53 \end{aligned}$$

in $KO(\mathbf{RP}^n)$.

If $1 \leq n \leq 8$ and $k > n$ or if $n \geq 9$ and $k \geq 2^{\phi(n)} - n - 1$, the equalities hold in the set of equivalence classes of \mathbf{R} -vector bundles over \mathbf{RP}^n .

Proof. By Lemma 3.1, we have

$$\nu = n + k + 1 - (n + 1)\xi_n = (a2^{\phi(n)} - n - 1)\xi_n + n + k + 1 - a2^{\phi(n)}$$

in $KO(\mathbf{RP}^n)$ for any integer a . So we have the former part by putting $a = 1$.

The latter part is a consequence of the former part by (2.1), since $\nu = A\xi_n + B$ for non-negative integers A and B such that $\dim \mathbf{RP}^n = n < k = \dim \nu = A + B$, if $1 \leq n \leq 8$ and $k > n$ or if $n \geq 9$ and $k \geq 2^{\phi(n)} - n - 1$. \square

Corollary 4.2. *If $1 \leq n \leq 8$ and $k > n$ or if $n \geq 9$ and $k \geq 2^{\phi(n)} - n - 1$, $\nu(f_n)$ is extendible to \mathbf{RP}^m for every $m > n$.*

Proof. Since ξ_n and the trivial \mathbf{R} -bundles over \mathbf{RP}^n are extendible to \mathbf{RP}^m for every $m > n$, the result follows from the latter part of Theorem 4.1. \square

Theorem B is a consequence of the following

Theorem 4.3. *Let ν be the normal bundle associated to an immersion of \mathbf{RP}^n in \mathbf{R}^{n+k} . Then we have*

- (i) ν is stably extendible to \mathbf{RP}^m for every $m > n$ if $1 \leq n \leq 8$ and $k \geq n$, and ν is extendible to \mathbf{RP}^m for every $m > n$ if $1 \leq n \leq 8$ and $k > n$.
- (ii) ν is not stably extendible to \mathbf{RP}^{n+k+1} if $n \geq 9$ and $1 \leq k \leq n + 12$.

Proof. The former part of Theorem 4.1 implies the former part of (i). In fact, if $k \geq n$, the \mathbf{R} -vector bundles k , $\xi_2 \oplus (k-1)$, k , $3\xi_4 \oplus (k-3)$, $2\xi_5 \oplus (k-2)$, $\xi_6 \oplus (k-1)$, k and $7\xi_8 \oplus (k-7)$ over \mathbf{RP}^n , where $1 \leq n \leq 8$ respectively, are extendible to \mathbf{RP}^m for every $m > n$, and they are stably equivalent to $\nu(f_n)$ respectively.

The latter part of (i) follows from the former part of (i) by Theorem 2.2.

(ii) is a consequence of (3.4), because $0 < k < 2^{\phi(n)} - n - 1$ if $n \geq 9$ and $1 \leq k \leq n + 12$. \square

In [6, Theorem 1], the following (4.4) is proved (cf. [11, Corollary 2.3 (2)]).

(4.4). Let ζ be a t -dimensional \mathbf{R} -vector bundle over \mathbf{RP}^n . If $n < t$, ζ is extendible to \mathbf{RP}^m for every m with $n < m \leq t$.

The next example is due to (4.4) and Corollary 3.6.

EXAMPLE 4.5. The normal bundle associated to an immersion of \mathbf{RP}^9 in \mathbf{R}^{30} is extendible to \mathbf{RP}^{21} , but is not stably extendible to \mathbf{RP}^{22} .

5. Proof of Theorem C

Lemma 5.1. *Let $c\nu$ be the complexification of the normal bundle ν associated to an immersion of \mathbf{RP}^n in \mathbf{R}^{n+k} , where $k > 0$. Then the equality*

$$c\nu = (b2^{\lfloor n/2 \rfloor} - n - 1)c\xi_n + n + k + 1 - b2^{\lfloor n/2 \rfloor}$$

holds in $K(\mathbf{RP}^n)$, where b is any integer.

Proof. Complexifying the equality in Lemma 3.1 and considering that $c\xi_n - 1$ is of order $2^{\lfloor n/2 \rfloor}$, we have the equality above, since $\lfloor n/2 \rfloor \leq \phi(n)$. \square

Theorem 5.2. *Let $c\nu$ be the complexification of the normal bundle ν associated to an immersion of \mathbf{RP}^n in \mathbf{R}^{n+k} , where $k > 0$. Then $c\nu$ is stably extendible to \mathbf{RP}^m for every $m > n$ if $k \geq 2^{\phi(n)} - n - 1$, or if $k \geq 2^{\lfloor n/2 \rfloor} - n - 1 \geq 0$. And if $2k \geq n$, in addition, $c\nu$ is extendible to \mathbf{RP}^m for every $m > n$.*

Proof. To prove the first part, by Lemma 5.1, we have $c\nu = Ac\xi_n + B$, where $A = 2^{\phi(n)} - n - 1$ and $B = n + k + 1 - 2^{\phi(n)}$, since we may take $b = 1$ if $n \equiv 6, 7$ or $0 \pmod 8$ and $b = 2$ otherwise. Clearly $A \geq 0$, and $B \geq 0$ by the assumption. For $m > n$, $i^*(Ac\xi_m \oplus B) = Ac\xi_n \oplus B$, where $i: \mathbf{RP}^m \rightarrow \mathbf{RP}^n$ is the standard inclusion. Hence $c\nu$ is stably extendible to \mathbf{RP}^m for every $m > n$, since $c\nu$ is stably equivalent to $Ac\xi_n \oplus B$.

To prove the second part, taking $b = 1$ in Lemma 5.1, we have $c\nu = Ac\xi_n + B$, where $A = 2^{\lfloor n/2 \rfloor} - n - 1$ and $B = n + k + 1 - 2^{\lfloor n/2 \rfloor}$. By the assumption $A \geq 0$ and $B \geq 0$. So $c\nu$ is stably extendible to \mathbf{RP}^m for every $m > n$, in the way similar to the proof above.

If $2k \geq n$, in addition, $\langle (\dim \mathbf{RP}^n)/2 \rangle = \langle n/2 \rangle \leq k = \dim c\nu = A + B$, and so we obtain $c\nu = Ac\xi_n \oplus B$ by (2.1). Thus $c\nu$ is extendible to \mathbf{RP}^m for every $m > n$. \square

We recall the following result ([9, Theorem 2.1]) which is the ‘‘stably extendible version’’ of Theorem 4.2 for $d = 1$ in [8].

(5.3). Let ζ be a t -dimensional \mathbf{C} -vector bundle over \mathbf{RP}^n . Assume that there is a positive integer l such that ζ is stably equivalent to $(t+l)c\xi_n$ and $t+l < 2^{\lfloor n/2 \rfloor}$. Then $\lfloor n/2 \rfloor < t+l$ and ζ is not stably extendible to \mathbf{RP}^m for every m with $t+l \leq \lfloor m/2 \rfloor$.

Theorem 5.4. *Let $c\nu$ be the complexification of the normal bundle ν associated to an immersion of \mathbf{RP}^n in \mathbf{R}^{n+k} , where $k > 0$. Then $c\nu$ is not stably extendible to \mathbf{RP}^m for every m with $2^{\lfloor n/2 \rfloor + 1} - 2n - 2 \leq m$, if $k < 2^{\lfloor n/2 \rfloor} - n - 1$.*

Proof. Put $\zeta = c\nu$, $t = k$ and $l = 2^{\lfloor n/2 \rfloor} - n - k - 1$ in (5.3). Then clearly $t+l < 2^{\lfloor n/2 \rfloor}$, and $l > 0$ by the assumption. So $c\nu$ is not stably extendible to \mathbf{RP}^m for every m with $2^{\lfloor n/2 \rfloor} - n - 1 \leq \lfloor m/2 \rfloor$. \square

Proof of Theorem C. (i) For $n \geq 6$, the ‘‘only if’’ part follows from Theorem 5.4, and the ‘‘if’’ part follows from Theorem 5.2, since $2^{\lfloor n/2 \rfloor} - n - 1 \geq 0$ if $n \geq 6$.

(ii) As is well-known, $\mathbf{RP}^1 \subseteq \mathbf{R}^2$, $\mathbf{RP}^2 \subseteq \mathbf{R}^3$, $\mathbf{RP}^3 \subseteq \mathbf{R}^4$, $\mathbf{RP}^4 \subseteq \mathbf{R}^7$ and $\mathbf{RP}^5 \subseteq \mathbf{R}^7$, where we denote by $\mathbf{RP}^n \subseteq \mathbf{R}^N$ the existence of an immersion of \mathbf{RP}^n in \mathbf{R}^N , and these immersions are best possible, that is, there do not exist immersions of \mathbf{RP}^n in

\mathbf{R}^{N-1} . Hence we have $k \geq 2^{\phi(n)} - n - 1$ for $1 \leq n \leq 5$. So the result follows also from Theorem 5.2. \square

6. Proof of Theorem D

Theorem 6.1. *Let $c\nu = c\nu(f_n)$ be the complexification of the normal bundle $\nu = \nu(f_n)$ associated to an immersion $f_n: \mathbf{R}P^n \rightarrow \mathbf{R}^{n+k}$, where $k \geq n$. Then we have the Whitney sum decompositions as follows:*

$$\begin{aligned} c\nu(f_1) &= k, & c\nu(f_2) &= c\xi_2 \oplus (k - 1), & c\nu(f_3) &= k, \\ c\nu(f_4) &= 3c\xi_4 \oplus (k - 3), & c\nu(f_5) &= 2c\xi_5 \oplus (k - 2), & c\nu(f_6) &= c\xi_6 \oplus (k - 1), \\ c\nu(f_7) &= k, & c\nu(f_8) &= 7c\xi_8 \oplus (k - 7) \text{ and } c\nu(f_9) &= 6c\xi_9 \oplus (k - 6). \end{aligned}$$

Proof. Complexifying the equalities in the former part of Theorem 4.1, we have the equalities above for $1 \leq n \leq 8$ using (2.1). So it suffices to prove the equality for $n = 9$. By the former part of Theorem 4.1, $\nu(f_9) = 22\xi_9 + k - 22$, and so $c\nu(f_9) = 22c\xi_9 + k - 22$. According to [1, Theorem 7.3], $c\xi_9 - 1$ is of order 16. Hence $16c\xi_9 - 16 = 0$ in $K(\mathbf{R}P^9)$, and so $c\nu(f_9) = 6c\xi_9 + k - 6$. Therefore, $c\nu(f_9) = 6c\xi_9 \oplus (k - 6)$ by (2.1). \square

Corollary 6.2. *If $1 \leq k \leq 20$, the complexification $c\nu(f_{10})$ of the normal bundle $\nu(f_{10})$ associated to an immersion $f_{10}: \mathbf{R}P^{10} \rightarrow \mathbf{R}^{10+k}$ is not stably extendible to $\mathbf{R}P^{42}$.*

Proof. By the former part of Theorem 4.1, $\nu(f_{10}) = 53\xi_{10} + k - 53$, and so $c\nu(f_{10}) = 53c\xi_{10} + k - 53 = 21c\xi_{10} + k - 21$, since $c\xi_{10} - 1$ is of order 32. Hence we have the result from (5.3) by putting $n = 10$, $\zeta = c\nu(f_{10})$, $t = k$ and $l = 21 - k$, since $l > 0$ for $1 \leq k \leq 20$ and since $t + l = 21 < 2^{\lceil 10/2 \rceil} = 32$. \square

Define $l(n) = 2^{\lceil n/2 \rceil} - n - k - 1$. Then we have

Lemma 6.3. *$l(n) > 0$ for any k and n such that $10 \leq n \leq k \leq n + 8$, and $k + l(n) < 2^{\lceil n/2 \rceil}$, for any k and n .*

Proof. For $10 \leq n \leq 17$, the inequalities hold clearly. For $n \geq 18$, we prove the inequalities by induction. \square

Theorem D is a consequence of the following

Theorem 6.4. *Let $c\nu$ be the complexification of the normal bundle ν associated to an immersion of $\mathbf{R}P^n$ in \mathbf{R}^{n+k} . Then we have*

- (i) *$c\nu$ is extendible to $\mathbf{R}P^m$ for every $m > n$ if $1 \leq n \leq 9$ and $k \geq n$.*

(ii) $c\nu$ is not stably extendible to \mathbf{RP}^m for every m with $2^{\lfloor n/2 \rfloor + 1} - 2n - 2 \leq m$ if $n \geq 10$ and $n \leq k \leq n + 8$.

Proof. Since $c\xi_n$ and the trivial C -vector bundles over \mathbf{RP}^n are extendible to \mathbf{RP}^m for every $m > n$, Theorem 6.1 implies (i).

By Lemma 5.1, we have

$$c\nu = \{b2^{\lfloor n/2 \rfloor} - (n+1)\}c\xi_n + n + k + 1 - b2^{\lfloor n/2 \rfloor}$$

for any integer b . (ii) follows from (5.3), Lemma 6.3 and the equality above by putting $\zeta = c\nu$, $t = k$ and $l = 2^{\lfloor n/2 \rfloor} - n - k - 1$. \square

In [6, Theorem 2], the following (6.5) is proved (cf. [11, Corollary 2.3 (2)]).

(6.5). Let ζ be a t -dimensional C -vector bundle over \mathbf{RP}^n . If $n < 2t + 1$, ζ is extendible to \mathbf{RP}^m for every m with $n < m \leq 2t + 1$.

The next example is due to (6.5) and Corollary 6.2 for $k = 20$.

EXAMPLE 6.6. The complexification of the normal bundle associated to an immersion of \mathbf{RP}^{10} in \mathbf{R}^{30} is extendible to \mathbf{RP}^{41} , but is not stably extendible to \mathbf{RP}^{42} .

References

- [1] J.F. Adams, *Vector fields on spheres*, Ann. of Math. **75** (1962), 603–632.
- [2] J.F. Adams and Z. Mahmud, *Maps between classifying spaces*, Invent. Math. **35** (1976), 1–41.
- [3] F. Hirzebruch, *Topological Methods in Algebraic Geometry*, 3rd ed., Appendix I by R.L.E. Schwarzenberger, Springer-Verlag, 1978.
- [4] D. Husemoller, *Fibre Bundles*, Second Edition, Graduate Texts in Mathematics 20, Springer-Verlag, 1975.
- [5] M. Imaoka and K. Kuwana, *Stably extendible vector bundles over the quaternionic projective spaces*, Hiroshima Math. J. **29** (1999), 273–279.
- [6] T. Kobayashi and K. Komatsu, *Extendibility and stable extendibility of vector bundles over real projective spaces*, Hiroshima Math. J. **31** (2001), 99–106.
- [7] T. Kobayashi, H. Maki and T. Yoshida, *Remarks on extendible vector bundles over lens spaces and real projective spaces*, Hiroshima Math. J. **5** (1975), 487–497.
- [8] T. Kobayashi, H. Maki and T. Yoshida, *Extendibility with degree d of the complex vector bundles over lens spaces and projective spaces*, Mem. Fac. Sci. Kochi Univ. Ser. A Math. **1** (1980), 23–33.
- [9] T. Kobayashi, H. Maki and T. Yoshida, *Stably extendible vector bundles over the real projective spaces and the lens spaces*, Hiroshima Math. J. **29** (1999), 631–638.
- [10] T. Kobayashi, H. Maki and T. Yoshida, *Stable extendibility of normal bundles associated to immersions of real projective spaces and lens spaces*, Mem. Fac. Sci. Kochi Univ. Ser. A Math. **20** (2000), 31–38.

- [11] H. Maki, *On the extendibility of vector bundles over the lens spaces and the projective spaces*, Hiroshima Math. J. **13** (1983), 1–28.
- [12] E. Rees, *On submanifolds of projective space*, J. London Math. Soc. **19** (1979), 159–162.
- [13] R.L.E. Schwarzenberger, *Extendible vector bundles over real projective space*, Quart. J. Math. Oxford **17** (1966), 19–21.

T. Kobayashi
Kochi Asakura-ki 292-21
Kochi 780-8066, Japan

H. Maki
Department of Mathematics
Faculty of Culture and Education
Saga University
Saga 840-8502, Japan

T. Yoshida
Department of Mathematics
Faculty of Integrated Arts and Sciences
Hiroshima University
Higashi-Hiroshima 739-8521, Japan