

A KIRILLOV MODEL OF A PRINCIPAL SERIES REPRESENTATION OF $\mathrm{GL}_2(\mathcal{D})$

YOSHIHIDE NAKAMURA

(Received April 30, 2003)

0. Introduction

Let F be a non-Archimedean local field of arbitrary characteristic and \mathcal{D} a central finite dimensional division algebra over F . Godement [1] constructed a model of an irreducible admissible representation (π, V) of $\mathrm{GL}_2(F)$, which is called the Kirillov model of (π, V) and is denoted by $\mathcal{K}(\pi)$. $\mathcal{K}(\pi)$ is realized as a certain space consisting of locally constant functions on F^* that vanish outside some compact subset of F . On $\mathcal{K}(\pi)$, upper triangular matrices act as

$$\pi \left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \right) f(x) = \psi_F(d^{-1}xb)\omega_\pi(d)f(d^{-1}xa),$$

where ω_π is the central character of π and ψ_F is a non-trivial additive character of F . Godement obtained an irreducibility criterion of principal series representations by using the theory of Kirillov models, and then classified principal series representations of $\mathrm{GL}_2(F)$.

Prasad and Raghuram [2] developed the theory of Kirillov models for admissible representations of $\mathrm{GL}_2(\mathcal{D})$. Let (π, V) be an admissible representation of $\mathrm{GL}_2(\mathcal{D})$ and $V_{N,\psi}$ the twisted Jacquet module of (π, V) with respect to a non-trivial additive character Ψ of \mathcal{D} . The Kirillov model of (π, V) is defined to be a certain space consisting of $V_{N,\psi}$ -valued locally constant functions on \mathcal{D}^* . If f is an element of the Kirillov model of (π, V) , f vanishes outside some compact subset of \mathcal{D} and upper triangular matrices act as

$$\pi \left(\begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \right) f(X) = \Psi(D^{-1}XB)\pi_{N,\psi} \left(\begin{pmatrix} D & 0 \\ 0 & D \end{pmatrix} \right) f(D^{-1}XA).$$

In this paper we study a Kirillov model of a principal series representation $V(\pi_1, \pi_2)$ of $\mathrm{GL}_2(\mathcal{D})$ induced from an irreducible representation $(\pi_1 \otimes \pi_2, V_1 \otimes V_2)$ of $\mathcal{D}^* \times \mathcal{D}^*$. Any element of $V(\pi_1, \pi_2)$ is a $V_1 \otimes V_2$ -valued locally constant function on $\mathrm{GL}_2(\mathcal{D})$ and $\mathrm{GL}_2(\mathcal{D})$ acts on $V(\pi_1, \pi_2)$ by right translations. Even if $V(\pi_1, \pi_2)$ is not irreducible, we construct its Kirillov model as follows. The element ξ_φ of the Kirillov model of $V(\pi_1, \pi_2)$ corresponding to $\varphi \in V(\pi_1, \pi_2)$ is given as a distri-

bution on $C_c^\infty(\mathcal{D})$ by the form

$$\xi_\varphi(X) = |X|^{1/2} 1 \otimes \pi_2(X) \sum_{n \in \mathbb{Z}} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)}_\varphi \left(\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \right) \left(\begin{pmatrix} 1 & Y \\ 0 & 1 \end{pmatrix} \right) dY,$$

where \mathfrak{v} denotes an additive valuation on \mathcal{D} . Raghuram [3] proved that the defining infinite series of ξ_φ converges. We give a proof of this fact by a different way from Raghuram in Lemma 2.2. As a consequence of the convergence of the series, we know that the Kirillov model is realized as a certain space of functions on \mathcal{D}^* . The asymptotic behavior of ξ_φ around 0 characterizes a principal series representation $V(\pi_1, \pi_2)$. Although Raghuram studied a behavior of $\hat{\phi}$ around 0, our statement in Theorem 2.3 is more precise than Raghuram’s one.

Moreover, we give a condition under when the map $\phi \mapsto \hat{\phi}$ is injective in Proposition 2.4 and Theorem 2.6. From this theorem we get a sufficient condition for irreducibility of the principal series representations in Corollary 2.7. If the characteristic of F is 0, an irreducibility criterion of the principal series representations of $GL_n(\mathcal{D})$ was given by Tadić [4] by using the theories of the Langlands classification and Hopf algebras. If we apply the results of Tadić to $GL_2(\mathcal{D})$ case, the principal series representation $V(\pi_1, \pi_2)$ is reducible if and only if $\pi_2(X) = |X|^{\pm 1} \pi_1(X)$ for all $X \in \mathcal{D}^*$ when the characteristic of F is 0. As a consequence of this fact and Theorem 2.6 we know that if $\dim_F \mathcal{D} \neq 1$ and the characteristic of F is 0, there exists a reducible principal series representation $V(\pi_1, \pi_2)$ such that the maps from $V(\pi_1, \pi_2)$ to its Kirillov model and from $V(\pi_1, \pi_2)^\vee$ to its Kirillov model are injective. If $\dim_F \mathcal{D} = 1$, such representations do not exist.

1. Preliminaries

1.1. Notations. In this paper \mathbb{Z} denotes the ring of integers and \mathbb{C} the field of complex numbers as usual. Let F be a non-Archimedean local field of arbitrary characteristic, \mathfrak{O}_F the integer ring of F , \mathfrak{P}_F the unique maximal ideal of \mathfrak{O}_F , q the cardinality of $\mathfrak{O}_F/\mathfrak{P}_F$, and ϖ_F the prime element of F . The additive valuation \mathfrak{v}_F and the multiplicative valuation $|\cdot|_F$ on F are normalized so that $|\varpi_F|_F = q^{-\mathfrak{v}_F(\varpi_F)} = q^{-1}$. We fix a nontrivial additive character ψ_F of F so chosen that the maximal fractional ideal in F on which ψ_F is trivial is \mathfrak{O}_F . Let \mathcal{D} denote a central division algebra of dimension d^2 over F , \mathfrak{O} the maximal order of \mathcal{D} , and \mathfrak{P} the unique maximal ideal of \mathfrak{O} . Notice that the cardinality of $\mathfrak{O}/\mathfrak{P}$ is equal to q^d . There is a generator ϖ of \mathfrak{P} as $\varpi^d = \varpi_F$. The additive valuation and the multiplicative valuation $|\cdot|$ on \mathcal{D} are normalized so that $|\varpi| = q^{-\mathfrak{v}(\varpi)} = q^{-d}$. Let $T_{\mathcal{D}/F}$ be the reduced trace from \mathcal{D} to F . Let Ψ be the additive character of \mathcal{D} obtained by composing $T_{\mathcal{D}/F}$ and the character ψ_F . Let dX be the Haar measure on \mathcal{D} normalized so that the volume of \mathfrak{O}^* is $(1 - q^{-d})^{-1}$.

Let $M_2(\mathcal{D})$ be the matrix algebra of 2×2 matrices with entries in \mathcal{D} , $G =$

$GL_2(\mathcal{D}) = M_2(\mathcal{D})^*$ the unit group of $M_2(\mathcal{D})$, P the subgroup of upper triangular matrices in G and N the unipotent radical of P consisting of matrices with 1's on diagonal. The Shalika subgroup S is defined to be the subgroup of G consisting of the matrices of the form $\begin{pmatrix} A & B \\ 0 & A \end{pmatrix}$ for $A \in \mathcal{D}^*$ and $B \in \mathcal{D}$. The subgroup of S consisting of the matrices of the form $\begin{pmatrix} A & 0 \\ 0 & A \end{pmatrix}$ for all $A \in \mathcal{D}^*$ is denoted by $\Delta(\mathcal{D}^*)$.

For a totally disconnected locally compact topological space X and an arbitrary vector space V , let $C^\infty(X, V)$ be the space consisting of V -valued locally constant functions on X and $C_c^\infty(X, V)$ be the subspace of $C^\infty(X, V)$ consisting of compactly supported functions. If V is one dimensional, we write simply $C^\infty(X)$ and $C_c^\infty(X)$ for $C^\infty(X, V)$ and $C_c^\infty(X, V)$, respectively.

Proposition 1.1. *Let $w = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. Then G is decomposed into the disjoint union of P and $PwP = PwN = NwP$.*

The subset PwP is called the big cell.

Proposition 1.2. *The additive character Ψ of \mathcal{D} is a constant on \mathfrak{P}^{1-d} .*

For the proof, refer to [5, Chapter 10].

1.2. Admissible representations and Kirillov models. Let (π, V) be a representation of G . In this paper, the representation space V is always a vector space over \mathbb{C} . (π, V) is called admissible if the stabilizer subgroup of v in G is open for all $v \in V$ and the subspace which consists of all elements that are invariant under G' is finite dimensional for all open subgroup G' of G .

Let (π_1, V_1) and (π_2, V_2) be two irreducible representations of \mathcal{D}^* . We extend π_1, π_2 to a representation of P on which N acts trivially. Let $V(\pi_1, \pi_2)$ denote the representation of G induced from $\pi_1 \otimes \pi_2$ of P . Namely,

$$V(\pi_1, \pi_2) = \left\{ \varphi \in C^\infty(G, V_1 \otimes V_2) \left| \begin{array}{l} \varphi \left(\begin{pmatrix} A & B \\ 0 & D \end{pmatrix} g \right) = |AD^{-1}|^{1/2} \times \pi_1(A) \otimes \pi_2(D) \varphi(g) \\ \left(\text{for all } \begin{pmatrix} A & B \\ 0 & D \end{pmatrix} \in P \text{ and } g \in G \right) \end{array} \right. \right\}$$

and G acts on $V(\pi_1, \pi_2)$ by right translations. Then we obtain an admissible representation. Such a representation is called a principal series representation.

The following lemma is proved in the same way as [1, Theorem 5].

Lemma 1.3. *The contragredient representation of $V(\pi_1, \pi_2)$ is isomorphic to $V(\pi_1^\vee, \pi_2^\vee)$, where π_i^\vee denote the contragredient representation of π_i .*

We study the Kirillov model in order to investigate when a principal series repre-

sentation is irreducible. Let (π, V) be an admissible representation of G . Let $V(N, \Psi)$ be the subspace of V spanned by $\pi\left(\begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix}\right)v - \Psi(X)v$ for all v in V and X in \mathcal{D} . The twisted Jacquet module $V_{N,\Psi}$ of V is defined as $V/V(N, \Psi)$. $V_{N,\Psi}$ is an S -module and the maximal quotient of V on which N acts via Ψ . It is known that if (π, V) is irreducible, $V_{N,\Psi}$ is finite dimensional. The next lemma was proved by Prasad and Raghuram in [2, Theorem 2.1].

Lemma 1.4. *The twisted Jacquet module $V(\pi_1, \pi_2)_{N,\Psi}$ of a principal series representation $V(\pi_1, \pi_2)$ is isomorphic with $V_1 \otimes V_2$ as $\Delta(\mathcal{D}^*)$ -modules.*

DEFINITION 1.1. For any infinite dimensional admissible representation (π, V) of G , let L be the natural projection from V to $V_{N,\Psi}$. Let ξ_v be the function on \mathcal{D}^* defined by $\xi_v(X) = L\left(\pi\left(\begin{pmatrix} X & 0 \\ 0 & 1 \end{pmatrix}\right)v\right)$. Let $\mathcal{K}(\pi)$ denote the space consisting of functions ξ_v for all v in V . $\mathcal{K}(\pi)$ is called the Kirillov model of π .

The action of any element $\begin{pmatrix} A & B \\ 0 & D \end{pmatrix}$ of P on $\mathcal{K}(\pi)$ is easy to describe, which is

$$\pi\left(\begin{pmatrix} A & B \\ 0 & D \end{pmatrix}\right)\xi(X) = \Psi(D^{-1}XB)\pi_{N,\Psi}\left(\begin{pmatrix} D & 0 \\ 0 & D \end{pmatrix}\right)\xi(D^{-1}XA)$$

for all ξ in $\mathcal{K}(\pi)$ and X in \mathcal{D}^* . From this formula, each $V_{N,\Psi}$ -valued function ξ of $\mathcal{K}(\pi)$ is locally constant on \mathcal{D}^* and vanishes outside some compact subset of \mathcal{D} because the stabilizer subgroup of ξ is open. The G -intertwining operator $v \mapsto \xi_v$ is injective if (π, V) is irreducible. Prasad and Raghuram proved the following lemma [2, Theorem 3.1].

Lemma 1.5. *For an admissible representation π , the Kirillov model $\mathcal{K}(\pi)$ contains the space $C_c^\infty(\mathcal{D}^*, V_{N,\Psi})$. Moreover, if π is a principal series representation, $C_c^\infty(\mathcal{D}^*, V_{N,\Psi})$ is a proper subspace of $\mathcal{K}(\pi)$.*

2. Main results

2.1. Asymptotic behavior of an element of a Kirillov model. In this section, we study the Kirillov model of a principal series representation of $GL_2(\mathcal{D})$. Since \mathcal{D}^* is not always commutative, the irreducible representation of \mathcal{D}^* is not one-dimensional. However since \mathcal{D}^* is compact modulo the center F^* , the irreducible representation is finite-dimensional. Let $(\pi_1, V_1), (\pi_2, V_2)$ be two irreducible representations of \mathcal{D}^* .

The element ξ_φ in the Kirillov model of $V(\pi_1, \pi_2)$ corresponding to φ is defined as

$$\xi_\varphi(X) = |X|^{1/2} \otimes \pi_2(X) \sum_{n \in \mathbb{Z}} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)} \varphi\left(w^{-1} \begin{pmatrix} 1 & Y \\ 0 & 1 \end{pmatrix}\right) dY.$$

This map $\varphi \mapsto \xi_\varphi$ is a G -intertwining operator, but not always injective.

We introduce the functions ϕ on \mathcal{D} such that $\phi(X) = \varphi(w^{-1} \begin{pmatrix} 1 & X \\ 0 & 1 \end{pmatrix})$. Let $\mathcal{F}(\pi_1, \pi_2)$ denote the space of such functions on \mathcal{D} . All functions ϕ of $\mathcal{F}(\pi_1, \pi_2)$ are locally constant on \mathcal{D} and $|X|\pi_1(X) \otimes \pi_2(X^{-1})\phi(X)$ are constant vectors for $|X|$ large. We define $\hat{\phi}$ of ϕ as

$$(1) \quad \hat{\phi}(X) = \sum_{n \in \mathbb{Z}} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)}\phi(Y) dY.$$

$\hat{\phi}$ makes sense if this is regarded as a Fourier transform of ϕ in the sense of distribution on $C_c^\infty(\mathcal{D}^*)$.

Lemma 2.1. *The map $\varphi \mapsto \xi_\varphi$ is injective if and only if the map $\phi \mapsto \hat{\phi}$ is injective.*

Proof. The map $\varphi \mapsto \xi_\varphi$ is a composition of the maps $\varphi \mapsto \phi$, $\phi \mapsto \hat{\phi}$ and $\hat{\phi} \mapsto \xi_\varphi$. The map $\hat{\phi} \mapsto \xi_\varphi$ is obviously isomorphic.

Since the big cell is dense in G , φ is completely determined on G by the corresponding ϕ . Hence the map $\varphi \mapsto \phi$ is an isomorphism from $V(\pi_1, \pi_2)$ to $\mathcal{F}(\pi_1, \pi_2)$. □

As a consequence of this lemma, it is important to consider the map $\phi \mapsto \hat{\phi}$. We start to consider of the convergence of the series of (1).

Lemma 2.2. *The series of (1) converges and the function vanishes outside some compact subset of \mathcal{D} .*

Proof. It is clear that $\mathcal{F}(\pi_1, \pi_2)$ is the direct sum of $C_c^\infty(\mathcal{D}, V_1 \otimes V_2)$ and the subspace spanned by the functions

$$\phi_v(X) = \begin{cases} |X|^{-1}\pi_1(X^{-1}) \otimes \pi_2(X)v & \text{if } |X| \geq 1 \\ 0 & \text{if } |X| < 1 \end{cases}$$

for all $v \in V_1 \otimes V_2$. If $\phi \in C_c^\infty(\mathcal{D}, V_1 \otimes V_2)$, $\phi \mapsto \hat{\phi}$ is a usual Fourier transform and therefore the series converges on every compact subset of \mathcal{D}^* .

Before considering ϕ_v , we give a filtration to $V_1 \otimes V_2$. We denote by f the minimal number such that $\pi_1(X) \otimes \pi_2(Y)v = v$ for all v in $V_1 \otimes V_2$ and X, Y in $1 + \mathfrak{P}^f$. Let

$$\begin{aligned} W'_f &= V_1 \otimes V_2, \\ W'_{i-1} &= \{v \in W'_i \mid \pi_1(X) \otimes \pi_2(Y)v = v \text{ (for all } X, Y \in 1 + \mathfrak{P}^{i-1})\} \quad \text{for } 2 \leq i \leq f, \\ W'_0 &= \{v \in W'_1 \mid \pi_1(X) \otimes \pi_2(Y)v = v \text{ (for all } X, Y \in \mathfrak{O}^*)\}. \end{aligned}$$

There exists an $\mathfrak{D}^* \times \mathfrak{D}^*$ -invariant scalar product $\langle \cdot, \cdot \rangle$ on $V_1 \otimes V_2$. Indeed, if we fix a scalar product (\cdot, \cdot) on $V_1 \otimes V_2$, then $\langle \cdot, \cdot \rangle$ may be given by

$$\langle v, w \rangle = \int_{\mathfrak{D}^*} \int_{\mathfrak{D}^*} (\pi_1(X) \otimes \pi_2(Y)v, \pi_1(X) \otimes \pi_2(Y)w) d^*Y d^*X.$$

Let

$$W_i = \{v \in W'_i \mid \langle v, v' \rangle = 0 \text{ (for all } v' \in W'_{i-1})\},$$

for $1 \leq i \leq f$ and $W_0 = W'_0$. Then $V_1 \otimes V_2 = \bigoplus_{i=0}^f W_i$ and if $i \neq j$, $\langle v_i, v_j \rangle = 0$ for all $v_i \in W_i$ and $v_j \in W_j$. Notice that if W_0 is not $\{0\}$, $V_1 \otimes V_2$ is one-dimensional because all $\pi_1(X) \otimes \pi_2(Y)$, $X, Y \in \mathfrak{D}^*$, are commutative with each other on W_0 . If v_i is an element of W_i , then

$$\phi_{v_i}(X) = \begin{cases} |X|^{-1} \pi_1(X^{-1}) \otimes \pi_2(X)v_i & \text{if } |X| \geq 1 \\ 0 & \text{if } |X| < 1, \end{cases}$$

and $\hat{\phi}_{v_i}$ is equal to

$$\sum_{n \leq 0} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y.$$

If $i = 0$, then

$$\begin{aligned} & \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_0 d^*Y \\ &= \int_{\mathfrak{D}^*} \overline{\Psi(X\varpi^n Y)} \pi_1(Y^{-1}\varpi^{-n}) \otimes \pi_2(\varpi^n Y)v_0 d^*Y \\ &= \pi_1(\varpi^{-n}) \otimes \pi_2(\varpi^n)v_0 \int_{\mathfrak{D}^*} \overline{\Psi(X\varpi^n Y)} d^*Y \\ &= \pi_1(\varpi^{-n}) \otimes \pi_2(\varpi^n)v_0 \int_{\mathfrak{D}} (\overline{\Psi(X\varpi^n Y)} - |\varpi| \overline{\Psi(X\varpi^{n+1} Y)}) dY. \end{aligned}$$

Since Ψ is trivial on \mathfrak{P}^{1-d} , $\int_{\mathfrak{D}} (\overline{\Psi(X\varpi^n Y)} - |\varpi| \overline{\Psi(X\varpi^{n+1} Y)}) dY \neq 0$ is equivalent to $X\varpi^{n+1} \in \mathfrak{P}^{1-d}$. Hence $\hat{\phi}_{v_0}$ vanishes outside some compact subset of \mathfrak{D} and the series turns out to be a finite sum whenever $\mathfrak{v}(X)$ is fixed.

Let $i \neq 0$. Since $v_i \in W_i$,

$$\begin{aligned} & \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y \\ &= \int_{\mathfrak{D}^*/1+\mathfrak{P}^i} \int_{1+\mathfrak{P}^i} \overline{\Psi(X\varpi^n AB)} \pi_1(B^{-1}A^{-1}\varpi^{-n}) \otimes \pi_2(\varpi^n AB)v_i d^*B d^*A \end{aligned}$$

$$\begin{aligned} &= \int_{\mathcal{D}^*/1+\mathfrak{P}^i} \pi_1(A^{-1}\varpi^{-n}) \otimes \pi_2(\varpi^n A) \int_{1+\mathfrak{P}^i} \overline{\Psi(X\varpi^n AB)} v_i d^*B d^*A \\ &= \int_{\mathcal{D}^*/1+\mathfrak{P}^i} \overline{\Psi(X\varpi^n A)} \pi_1(A^{-1}\varpi^{-n}) \otimes \pi_2(\varpi^n A) v_i d^*A \int_{\mathfrak{P}^i} \overline{\Psi(X\varpi^n AB)} dB. \end{aligned}$$

Since Ψ is trivial on \mathfrak{P}^{1-d} , $\int_{\mathfrak{P}^i} \overline{\Psi(X\varpi^n AB)} dB \neq 0$ is equivalent to $X\varpi^n A \in \mathfrak{P}^{1-d}$. Hence $\hat{\phi}_{v_i}$ vanishes outside some compact subset of \mathcal{D} and the series turn out to be a finite sum whenever $\mathfrak{v}(X)$ is fixed.

This completes the proof since any function in $\mathcal{F}(\pi_1, \pi_2)$ can be written as a finite sum of the above functions. □

By this lemma the Kirillov model is realized as a certain space consisting of locally constant functions on \mathcal{D}^* .

REMARK 2.1. Raghuram also considered the convergence of the series (1) in [3] as follows. For $\mathfrak{v}(X)$ large, let

$$A(X) = \sum_{n \leq \mathfrak{v}(X)} \int_{\mathfrak{v}(T)=n} \overline{\Psi(T)} (\pi_1(T^{-1}) \otimes \pi_2(T)) d^*T.$$

$A(X)$ is an element of $\text{End}(V_1 \otimes V_2)$. Then

$$\hat{\phi}_{\mathfrak{v}}(X) = (1 \otimes \pi_2(X)^{-1}) \cdot A(X) \cdot (\pi_1(X) \otimes 1)v$$

where the notations are the same as Lemma 2.2. He analyzed $A(X)$ and proved that the defining series of $A(X)$ is a finite sum.

Raghuram also calculated the asymptotic behavior of $\hat{\phi}$ around 0 and obtained

$$\hat{\phi}(X) = (1 \otimes \pi_2(X^{-1})) \cdot A(X) \cdot (\pi_1(X) \otimes 1)v_1 + v_2$$

for $|X|$ enough small. By the proof of Lemma (2.2), we can calculate $A(X)$ more precisely.

Let ω_i be the central characters of π_i for $i = 1, 2$ and $\omega = \omega_1 \cdot \omega_2^{-1}$.

Theorem 2.3. *For each $\phi \in \mathcal{F}(\pi_1, \pi_2)$, there exist four vectors $v_\alpha, v_\beta, v_\gamma, v_\delta$ in $V_1 \otimes V_2$ such that*

$$(2) \quad \hat{\phi}(X) = \left((1 \otimes \pi_2(X^{-1})) \cdot A_1 \cdot (\pi_1(X) \otimes 1) + \sum_{t=0}^{[m/d]} \omega(\varpi^{td}) A_2 + A_3(m) \right) v_\alpha + \pi_1(X) \otimes \pi_2(X^{-1})v_\beta + mv_\gamma + v_\delta$$

for $X \in \mathfrak{P}^m$, $X \notin \mathfrak{P}^{m+1}$ with m large. Here

$$\begin{aligned} A_1 &= \sum_{1-d-f \leq n \leq 1-d} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(Y)} \pi_1(Y) \otimes \pi_2(Y^{-1}) d^*Y, \\ A_2 &= \sum_{1-d \leq n \leq 0} \int_{\mathfrak{v}(Y)=n} \pi_1(Y^{-1}) \otimes \pi_2(Y) d^*Y, \\ A_3(m) &= \sum_{1-d-m \leq n \leq -d-[m/d]d} \int_{\mathfrak{v}(Y)=n} \pi_1(Y^{-1}) \otimes \pi_2(Y) d^*Y, \end{aligned}$$

considered as elements of $\text{End}(V_1 \otimes V_2)$.

Proof. Similarly as in previous lemma, we start from the case ϕ is in $C_c^\infty(\mathcal{D}, V_1 \otimes V_2)$. Since $\phi \mapsto \hat{\phi}$ is Fourier transform, in some neighborhood of 0, $\hat{\phi}(X)$ is a constant vector $\int_{\mathcal{D}} \phi(Y) dY$.

Let $m = \mathfrak{v}(X)$ be enough large. From the proof of the previous lemma, we have

$$\hat{\phi}_v(X) = \sum_{-d-f-m \leq n \leq 0} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)} \pi_1(Y^{-1}) \otimes \pi_2(Y) v d^*Y$$

for v in $V_1 \otimes V_2$. If v_0 is a non-zero element of W_0 , π_1 and π_2 are characters. Then,

$$\begin{aligned} \hat{\phi}_{v_0}(X) &= \sum_{-d-m \leq n \leq 0} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)} \pi_1(Y^{-1}) \pi_2(Y) v_0 d^*Y \\ &= \sum_{-d-m \leq n \leq 0} \pi_1(\varpi^{-n}) \pi_2(\varpi^n) v_0 \int_{\mathcal{D}^*} \overline{\Psi(X\varpi^n Y)} d^*Y \\ &= \sum_{-d-m \leq n \leq 0} \pi_1(\varpi^{-n}) \pi_2(\varpi^n) v_0 \int_{\mathcal{D}} (\overline{\Psi(X\varpi^n Y)} - |\varpi| \overline{\Psi(X\varpi^{n+1} Y)}) dY. \end{aligned}$$

If we assume $\pi_1(\varpi) \pi_2(\varpi^{-1}) \neq 1$, since Ψ is trivial on \mathfrak{P}^{1-d} ,

$$\begin{aligned} \hat{\phi}_{v_0}(X) &= -|\varpi| \pi_1(\varpi^{d+m}) \pi_2(\varpi^{-d-m}) v_0 + (1 - |\varpi|) \sum_{1-d-m \leq n \leq 0} \pi_1(\varpi^{-n}) \pi_2(\varpi^n) v_0 \\ &= -\pi_1(X) \otimes \pi_2(X^{-1}) \\ &\quad \times \left((1 - |\varpi|) \frac{\pi_1(\varpi^d) \otimes \pi_2(\varpi^{-d})}{1 - \pi_1(\varpi) \otimes \pi_2(\varpi^{-1})} + |\varpi| \pi_1(\varpi^d) \otimes \pi_2(\varpi^{-d}) \right) v_0 \\ &\quad + \frac{1}{1 - \pi_1(\varpi) \otimes \pi_2(\varpi^{-1})} v_0. \end{aligned}$$

The last is the behavior of $\hat{\phi}_{v_0}$ around 0 in this case.

If we assume $\pi_1(\varpi)\pi_2(\varpi^{-1}) = 1$,

$$\begin{aligned} \hat{\phi}_{v_0}(X) &= -|\varpi|\pi_1(\varpi^{d+m})\pi_2(\varpi^{-d-m})v_0 + (1 - |\varpi|) \sum_{1-d-m \leq n \leq 0} \pi_1(\varpi^{-n})\pi_2(\varpi^n)v_0 \\ &= -|\varpi|v_0 + (1 - |\varpi|)(d + m)v_0 \\ &= m(1 - |\varpi|)v_0 + ((1 - |\varpi|)d - |\varpi|)v_0. \end{aligned}$$

The last is the behavior of $\hat{\phi}_{v_0}$ around 0 in this case.

Next, we assume v_i is an element of W_i for $i \neq 0$. Since Ψ is trivial on \mathfrak{P}^{1-d} ,

$$\begin{aligned} \hat{\phi}_{v_i}(X) &= \sum_{1-d-f-m \leq n \leq -d-m} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(XY)}\pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y \\ &+ \sum_{1-d-m \leq n \leq 0} \int_{\mathfrak{v}(Y)=n} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y \\ &= (1 \otimes \pi_2(X^{-1})) \\ &\times \left(\sum_{1-d-f \leq n \leq -d} \int_{\mathfrak{v}(Y)=n} \overline{\Psi(Y)}\pi_1(Y^{-1}) \otimes \pi_2(Y)d^*Y \right) (\pi_1(X) \otimes 1)v_i d^*Y \\ &+ \sum_{1-d-[m/d]d \leq n \leq 0} \int_{\mathfrak{v}(Y)=n} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y \\ &+ \sum_{1-d-m \leq n \leq -d-[m/d]d} \int_{\mathfrak{v}(Y)=n} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y \\ &= (1 \otimes \pi_2(X^{-1})) \cdot A_1 \cdot (\pi_1(X) \otimes 1)v_i \\ &+ \sum_{t=0}^{[m/d]} \omega(\varpi^{td}) \left(\sum_{1-d \leq n \leq 0} \int_{\mathfrak{v}(Y)=n} \pi_1(Y^{-1}) \otimes \pi_2(Y)v_i d^*Y \right) + A_3(m)v_i. \end{aligned}$$

Then the asymptotic behavior around 0 is

$$\hat{\phi}_{v_i}(X) = (1 \otimes \pi_2(X^{-1})) \cdot A_1 \cdot (\pi_1(X) \otimes 1)v_i + \sum_{t=0}^{[m/d]} \omega(\varpi^{td})A_2v_i + A_3(m)v_i$$

in this case.

Any function in $\mathcal{F}(\pi_1, \pi_2)$ is a finite sum of above functions. Hence (2) is obtained. \square

2.2. Injectivity of the map to a Kirillov model. Here we study the condition under when the map from $V(\pi_1, \pi_2)$ to its Kirillov model is injective. Since this map is G -intertwining, $V(\pi_1, \pi_2)$ is reducible if the map has non-zero kernel.

Proposition 2.4. *The mapping $\phi \mapsto \hat{\phi}$ is injective unless there exists a non-zero subspace of $V_1 \otimes V_2$ on which $\pi_1(X) \otimes \pi_2(X^{-1})$ acts as $|X|^{-1}$, in which case its kernel is the set of constant vector-valued functions in $\mathcal{F}(\pi_1, \pi_2)$.*

Proof. We fix a basis of n -dimensional vector space $V_1 \otimes V_2$. Then, $\hat{\phi}(X)$ is written as $(\hat{\phi}_1(X), \dots, \hat{\phi}_n(X))$ and also $\phi(X)$ is $(\phi_1(X), \dots, \phi_n(X))$, where each $\hat{\phi}_i$ is the Fourier transform of ϕ_i . If $\hat{\phi}_i = 0$ on \mathcal{D}^* , the measure $\hat{\phi}_i(X)dX$ is proportional to Dirac measure, which means ϕ_i is a constant on \mathcal{D} . Hence ϕ is a constant vector on \mathcal{D} . This happens if and only if there exists a non-zero subspace in $V_1 \otimes V_2$ on which $\pi_1(X) \otimes \pi_2(X^{-1})$ acts as $|X|^{-1}$. \square

Proposition 2.5. *Let H be an arbitrary group, (π_1, V_1) and (π_2, V_2) finite dimensional irreducible representations of H , and χ a one dimensional representation of H . There exists a non-zero element v of $V_1 \otimes V_2$ such that $\pi_1(X) \otimes \pi_2(X^{-1})v = \chi(X)v$ for all $X \in H$ if and only if $\pi_1 = \chi \cdot \pi_2$ and $\dim V_1 = \dim V_2 = 1$.*

Proof. We assume there exists a non-zero element v of $V_1 \otimes V_2$ such that $\pi_1(X) \otimes \pi_2(X^{-1})v = \chi(X)v$ for all $X \in H$ and (π_1, V_1) and (π_2, V_2) are finite dimensional and irreducible. Notice that

$$\pi_1(X) \otimes 1v = \chi(X)(1 \otimes \pi_2(X))v.$$

Any element of $V_1 \otimes V_2$ is written as

$$\sum_i a_i(\pi_1(Y_i) \otimes 1)v,$$

where the sum is finite, $a_i \in \mathbb{C}^*$, and $Y_i \in H$. For any element X of H , one has

$$\begin{aligned} & \pi_1(X) \otimes \pi_2(X^{-1}) \left(\sum_i a_i(\pi_1(Y_i) \otimes 1)v \right) \\ &= \sum_i a_i(1 \otimes \pi_2(X^{-1}))(\pi_1(XY_i) \otimes 1)v \\ &= \sum_i a_i(1 \otimes \pi_2(Y_i))(\pi_1(XY_i) \otimes \pi_2((XY_i)^{-1}))v \\ &= \sum_i a_i\chi(XY_i)(1 \otimes \pi_2(Y_i))v \\ &= \chi(X) \sum_i a_i(\pi_1(Y_i) \otimes 1)v. \end{aligned}$$

Hence $\pi_1(X) \otimes \pi_2(X^{-1})$ acts on $V_1 \otimes V_2$ as $\chi(X)$. Next we consider the action

of $\pi_1(XY) \otimes 1$ on $V_1 \otimes V_2$ for all $X, Y \in H$. If w is any element of $V_1 \otimes V_2$,

$$\begin{aligned} (\pi_1(XY) \otimes 1)w &= \chi(Y)(\pi_1(X) \otimes \pi_2(Y))w \\ &= \chi(Y)(1 \otimes \pi_2(Y))(\pi_1(X) \otimes 1)w \\ &= (\pi_1(YX) \otimes 1)w. \end{aligned}$$

By Schur’s lemma, $\dim V_1 = 1$. Similarly, $\dim V_2 = 1$.

The converse is obvious. □

These two propositions yield immediately the next theorem.

Theorem 2.6. *The map from an induced representation $V(\pi_1, \pi_2)$ to its Kirillov model is injective unless $\pi_1 = | \cdot |^{-1} \cdot \pi_2$ and $\dim V_1 = \dim V_2 = 1$.*

By this theorem we obtain a sufficient condition for the reducibility of a principal series representation.

Corollary 2.7. *If $\dim V_1 = \dim V_2 = 1$ and $\pi_1 = | \cdot |^{\pm 1} \cdot \pi_2$, $V(\pi_1, \pi_2)$ is reducible.*

Proof. Since the map $V(\pi_1, \pi_2) \ni \varphi \mapsto \xi_\varphi \in \mathcal{K}(\pi)$ is a G -intertwining operator, if this map is not injective, $V(\pi_1, \pi_2)$ is reducible. By Lemma 1.3, the map from $V(\pi_1, \pi_2)^\vee$ to its Kirillov model is not injective if $\pi_1 = | \cdot | \cdot \pi_2$ and $\dim V_1 = \dim V_2 = 1$. □

Tadić obtained the irreducibility criterion of principal series representations of $GL_n(\mathcal{D})$ when the characteristic of F is 0 by using theories of Langlands classification and Hopf algebras [4, Lemma 2.5 and 4.2]. The following theorem is a $GL_2(\mathcal{D})$ case of the results of Tadić.

Theorem 2.8 (Tadić). *When the characteristic of F is 0, the representation $V(\pi_1, \pi_2)$ is reducible if and only if $\pi_1 = | \cdot |^{\pm 1} \pi_2$.*

As a consequence of Corollary 2.7 and Theorem 2.8, if $d \geq 2$ and the characteristic of F is 0, there exists a reducible principal series representation $V(\pi_1, \pi_2)$ such that the maps from $V(\pi_1, \pi_2)$ to $\mathcal{K}(\pi)$ and from $V(\pi_1, \pi_2)^\vee$ to $\mathcal{K}(\pi)^\vee$ are injective. If $d = 1$, i.e. \mathcal{D} is a commutative field, such representation $V(\pi_1, \pi_2)$ does not exist [1, Theorem 6].

ACKNOWLEDGEMENTS. The author would like to thank Professor Takao Watanabe for many helpful advice. The author also thank Department of Mathematics of Osaka University for very pleasant studying environments.

References

- [1] R. Godement: Notes on Jacquet-Langlands theory, Institute for Advanced Study, Princeton, 1970.
- [2] D. Prasad and A. Raghuram: *Kirillov theory for $GL_2(\mathcal{D})$ where \mathcal{D} is a division algebra over a non-Archimedean local field*, *Duke Math. J.* **104** (2000), 19–44.
- [3] A. Raghuram: *On representations of p -adic $GL_2(\mathcal{D})$* , *Pacific J. Math.* **206** (2002), 451–464.
- [4] M. Tadić: *Induced representations of $GL(n, A)$ for p -adic division algebras A* , *J. Reine Angew. Math.* **405** (1990), 48–77.
- [5] A. Weil: *Basic number theory*, Springer-Verlag, New York, 1967.

Department of Mathematics
Graduate School of Science
Osaka University
Toyonaka 560-0043, Japan
e-mail: nakamura@gaia.math.wani.osaka-u.ac.jp