

## ON THE CHARACTERISTIC CAUCHY PROBLEM FOR PARTIAL DIFFERENTIAL EQUATIONS

HITOSHI KUMANO-GO and KUSUO ISÉ

(Received March 16, 1965)

We shall consider the differential operator

$$(0.1) \quad L = \sum_{j+\alpha_0 k \leq m} a_{j,k}(t) \frac{\partial^{j+k}}{\partial t^j \partial x^k} \quad (0 < \alpha_0 < 1, a_{m,0}(t) \equiv 1)$$

where  $m$  is a positive integer and  $a_{j,k}(t)$  are complex-valued continuous functions on  $0 \leq t \leq T$ . If the plane  $t=0$  is characteristic with respect to  $L$ , then we can suitably choose  $\alpha_0$  so that  $L$  contains at least one coefficient  $a_{j,k}(0) \neq 0$  for  $j \neq m$  and  $j + \alpha_0 k = m$ . In this case, when the coefficients of  $L$  are constant, L. Hörmander [7] constructed non-trivial  $C^\infty$ -functions  $u = u(t, x)$  which satisfy  $Lu = 0$  in the whole  $(t, x)$ -space and vanish for  $t \leq 0$ . On the other hand, I. M. Gelfand and G. E. Shilov [3] proved the uniqueness of the characteristic Cauchy problem for  $Lu = 0$  under some restrictions on the behavior of  $u$  as  $|x| \rightarrow \infty$ , and their results were improved by K. I. Babenko [1], B. L. Gurevich [5], [6] and A. Friedman [2].

In this note we shall show the existence of non-trivial functions of  $G(\alpha_0, e^{-\gamma|x|^{1/(1-\alpha_0)}})$  (see Definition 1 and Lemma 1) and apply this result to prove a uniqueness theorem (Theorem 2) of the Cauchy problem without aid of generalized functions. To prove an existence theorem (Theorem 1) which is important for the proof of Theorem 2, we shall use the method of E. de Giorgi [4] which was also used by G. Talenti [9] and Tsutsumi [11]. For the case that  $t=0$  is characteristic with respect to  $L$  with constant coefficients  $a_{j,k}$ , we shall show in Theorem 3 that the uniqueness in Theorem 1 and Theorem 2 does not hold for any  $\varepsilon_0 > 0$  if we replace the factor  $(n!)^{\alpha_0} g(x)$  in (2.3) by  $(n!)^{\alpha_0 + \varepsilon} g(x)$  and the factor  $e^{\gamma|x|^{1/(1-\alpha_0)}}$  in (2.11) by  $e^{\gamma|x|^{1/(1-\alpha_0) + \varepsilon_0}}$  respectively.

The authors wish to thank Professor M. Nagumo for stimulating discussions.

1. We define the following class of functions for any positive continuous function  $g(x)$  on  $-\infty < x < +\infty$  and any  $\alpha$  such that  $0 < \alpha < 1$ .

**Definition 1.** We call  $\varphi(x)$  a function of  $G(\alpha, g(x))$  if  $\varphi(x)$  is a  $C^\infty$ -function on  $-\infty < x < +\infty$  satisfying

$$(1.1) \quad |\varphi^{(n)}(x)| \leq C^{n+1}(n!)^\alpha g(x) \quad (n = 0, 1, 2, \dots; -\infty < x < +\infty)$$

for some constant  $C$ .

If  $\varphi(x) (\not\equiv 0)$  is a function of  $G(\alpha, g(x))$  (which is contained in the Gevray class of  $\alpha < 1$ ), it can be extended to be an entire function of order  $1/(1-\alpha)^2$ , so that for any  $\varepsilon > 0$  there exists a sequence  $\{x_n\}_{n=1}^\infty$  such that  $x_n \rightarrow +\infty$  and  $|\varphi(x_n)| > e^{-x_n^{1/(1-\alpha)+\varepsilon}}$  ( $n=1, 2, \dots$ )<sup>2)</sup>. Hence we know that for any  $\gamma > 0$   $G(\alpha, e^{-\gamma|x|^\beta})$  contains only one trivial function ( $\equiv 0$ ) unless  $\beta \leq 1/(1-\alpha)$ . Now we show the following lemma.

**Lemma 1.** Let  $\alpha$  and  $\gamma$  be any positive number and  $0 < \alpha < 1$ . Then we can construct a non-trivial entire function  $F_{\alpha, \gamma}(z)$  of order  $1/(1-\alpha)$  which satisfies

$$(1.2) \quad |F_{\alpha, \gamma}^{(n)}(x)| \leq C^{n+1}(n!)^\alpha e^{-\gamma|x|^{1/(1-\alpha)}} \quad (n = 0, 1, 2, \dots; -\infty < x < +\infty)$$

for some constant  $C$  depending only on  $\alpha$  and  $\gamma$ .

*Proof.* For any positive number  $\rho > 1$  we consider a function

$$(1.3) \quad f_\rho(x) = \int_{-\infty+i}^{+\infty+i} e^{i\alpha_\rho x w} \exp\{e^{i\theta_\rho w^\rho}\} dw \quad (w = u + iv)$$

where  $w^\rho$  is defined by  $e^{\rho \log w}$  such as  $\log 1 = 0$  by setting the slit on the nonpositive imaginary axis, and  $\theta_\rho$  is defined by

$$(1.4) \quad \theta_\rho = (m+1-\rho/2)\pi \quad \text{for } 2m-1 < \rho \leq 2m+1 \quad (m=1, 2, \dots)$$

( $\alpha_\rho > 0$  will be defined later).

Then, for the case  $\rho = 2m+1$  we have  $\Re(e^{i\theta_\rho w^\rho})^{3)} = -\Im w^\rho = -(2m+1)u^{2m}v(1+o(1))$  as  $|u| \rightarrow \infty$  for any fixed  $v > 0$ .

For  $2m-1 < \rho < 2m+1$  we have

$$(1.5) \quad \begin{cases} \pi - \frac{1}{2}\pi < \theta_\rho < \pi + \frac{1}{2}\pi \\ (2m+1)\pi - \frac{1}{2}\pi < \theta_\rho + \rho\pi < (2m+1)\pi + \frac{1}{2}\pi, \end{cases}$$

so that if we set

$$(1.6) \quad \beta_\rho = \frac{\pi}{2} \text{Min} \left\{ \frac{(2m+1)-\rho}{2}, \frac{\rho-(2m-1)}{2} \right\}$$

1) [8], pp. 262-263. 2) [10], p. 273.

3) For a complex number  $w$ ,  $\Re w$  means the real part of  $w$  and  $\Im w$  the imaginary part.

then, we have for  $0 \leq \theta \leq \beta_\rho$ ,

$$(1.7) \quad \begin{cases} \frac{\pi}{2} + \beta_\rho < \theta_\rho + \rho\theta < \frac{3}{2}\pi - \beta_\rho \\ \left(2m + \frac{1}{2}\right)\pi + \beta_\rho < \theta_\rho + \rho(\pi - \theta) < \left(2m + \frac{3}{2}\right)\pi - \beta_\rho \end{cases}$$

and consequently we have for  $v > 0$

$$(1.8) \quad \Re(e^{i\theta_\rho w^\rho}) \leq -|w|^\rho \sin \beta_\rho \quad \text{for } |w| \geq v \cot \theta.$$

These facts mean that, for any  $\rho > 1$ , we can extend  $f_\rho(x)$  to an entire function  $f_\rho(z) = f_\rho(x + iy)$  and we have

$$(1.9) \quad f_\rho^{(n)}(x) = \int_{-\infty+i}^{+\infty+i} (ia_\rho w)^n e^{ia_\rho xw} \exp\{e^{i\theta_\rho w^\rho}\} dw.$$

Now we estimate the values of  $f_\rho^{(n)}(x)$ . First we do this for  $\rho = 2m + 1$  which is fairly complicated. If we set

$$(1.10) \quad \theta_0 = \pi/(2\rho) \quad \text{and} \quad a_\rho = 5 \operatorname{cosec} \theta_0$$

we have, for  $w = |w|e^{i\theta}$  and  $w = |w|e^{i(\pi-\theta)}$  ( $0 \leq \theta \leq \theta_0$ ),

$$(1.11) \quad \Re(e^{i\theta_\rho w^\rho}) = -\Im w^\rho = -|w|^\rho \sin(\rho\theta) \quad (0 \leq \rho\theta \leq \rho\theta_0 = \pi/2).$$

In the case  $x > 0$ , we deform the path of the integral (1.9) to  $\Gamma = C_1 \cup C_2 \cup C_3$  where  $C_1, C_2$  and  $C_3$  are the segments (of straight lines) from  $+\infty e^{i(\pi-\theta_0)}$  to  $x^\delta e^{i(\pi-\theta_0)}$ , from  $x^\delta e^{i(\pi-\theta_0)}$  to  $x^\delta e^{i\theta_0}$ , and from  $x^\delta e^{i\theta_0}$  to  $+\infty e^{i\theta_0}$  respectively by setting  $\delta = 1/(\rho - 1)$ . Let the values of the above integral on  $C_j$  be denoted by  $I_j(x)$  ( $j = 1, 2, 3$ ). Then, by (1.10) and (1.11) we have

$$|I_1(x)| \leq e^{-5x^{1+\delta}} \int_{x^\delta}^{+\infty} \{(a_\rho r)^n e^{-r^{\rho/2}}\} e^{-r^{\rho/2}} dr.$$

Now for positive  $\delta$  and  $\rho$  we use the inequality

$$(1.12) \quad r^n e^{-\delta r^\rho} \leq C_{\delta,\rho}^n (n!)^{1/\rho} \quad (r \geq 0, n = 0, 1, 2, \dots)$$

which will be often used in later discussions. Then, remarking  $1 + \delta = \rho'$  where  $1/\rho + 1/\rho' = 1$ , we have

$$(1.13) \quad |I_1(x)| \leq C_1^{n+1} (n!)^{1/\rho} e^{-5x^{\rho'}}.$$

For  $I_2(x)$ , remarking  $\delta\rho = \rho'$  we have

---

4) In what follows in this section,  $C_i$  are constants depending only on  $\rho$ .

$$\begin{aligned}
 |I_2(x)| &\leq e^{-5x^{\rho'}} (a_\rho x^\delta)^n \int_{-x^\delta \cos \theta_0}^{x^\delta \cos \theta_0} e^{(x^\delta)^\rho} du \\
 &= e^{-2x^{\rho'}} a_\rho^n \{(x^\delta)^n e^{-(x^\delta)^\rho}\} \{2x^\delta \cos \theta_0 \cdot e^{-x^{\rho'}}\},
 \end{aligned}$$

so that by (1.12) we have

$$(1.14) \quad |I_2(x)| \leq C_2^{n+1} (n!)^{1/\rho} e^{-2x^{\rho'}}.$$

Since  $I_3(x)$  can be estimated in the same way as  $I_1(x)$ , we get by (1.13) and (1.14)

$$(1.15) \quad |f_\rho^{(n)}(x)| \leq C_3^{n+1} (n!)^{1/\rho} e^{-2x^{\rho'}} \quad \text{for } x > 0.$$

In the case  $x < 0$ , we deform the path of the integral (1.9) to  $\Gamma' = C'_1 \cup C'_2 \cup C'_3 \cup C'_4 \cup C'_5$  where  $C'_1, C'_2, C'_3, C'_4, C'_5$  are the segments from  $+\infty e^{i(\pi-\theta_1)}$  to  $b|x|^\delta e^{i(\pi-\theta_1)} \operatorname{cosec} \theta_1$ , from  $b|x|^\delta e^{i(\pi-\theta_1)} \operatorname{cosec} \theta_1$  to  $b|x|^\delta e^{i(\pi-\theta_0)} \operatorname{cosec} \theta_0$ , from  $b|x|^\delta e^{i(\pi-\theta_0)} \operatorname{cosec} \theta_0$  to  $b|x|^\delta e^{i\theta_0} \operatorname{cosec} \theta_0$ , from  $b|x|^\delta e^{i\theta_0} \operatorname{cosec} \theta_0$  to  $b|x|^\delta e^{i\theta_1} \operatorname{cosec} \theta_1$ , from  $b|x|^\delta e^{i\theta_1} \operatorname{cosec} \theta_1$  to  $+\infty e^{i\theta_1}$  respectively. ( $0 < b$  and  $0 < \theta_1 < \theta_0$  will be determined later.) Let the values of the integral on  $C'_j$  be denoted by  $I'_j(x)$  ( $j=1, 2, 3, 4, 5$ ). For  $I'_3(x)$  we have by (1.12)

$$(1.16) \quad \begin{aligned} |I'_3(x)| &\leq 2b|x|^\delta \cot \theta_0 (a_\rho b|x|^\delta \operatorname{cosec} \theta_0)^n e^{a_\rho b|x|^{1+\delta} + (b|x|^\delta \operatorname{cosec} \theta_0)^\rho} \\ &\leq C_4^{n+1} (n!)^{1/\rho} e^{b(3a_\rho + b^{\rho-1} \operatorname{cosec}^\rho \theta_0)|x|^{\rho'}}, \end{aligned}$$

for  $I'_2(x)$ , remarking  $\Re(e^{i\theta_\rho} w^\rho) < 0$  by (1.11),

$$(1.17) \quad \begin{aligned} |I'_2(x)| &\leq 2b|x|^\delta \cot \theta_1 (a_\rho b|x|^\delta \operatorname{cosec} \theta_1)^n e^{a_\rho b|x|^{1+\delta}} \\ &\leq C_5^{n+1} (n!)^{1/\rho} e^{3a_\rho b|x|^{\rho'}}, \end{aligned}$$

and for  $I'_1(x)$  we have by (1.11)

$$|I'_1(x)| \leq a_\rho^n \int_{b|x|^\delta \operatorname{cosec} \theta_1}^\infty r^n e^{a_\rho |x| r \sin \theta_1 - r^\rho \sin(\rho \theta_1)} dr.$$

If we determine  $\theta_1$  such as  $a_\rho \leq b^{\rho-1} \operatorname{cosec}^\rho \theta_1 \sin(\rho \theta_1)/2$  for fixed  $b > 0$  which is possible because of  $\operatorname{cosec}^\rho \theta_1 \sin(\rho \theta_1) = \rho \theta_1^{-(\rho-1)}(1+o(1))$  as  $\theta_1 \downarrow 0$ , then we have  $a_\rho |x| r \sin \theta_1 - r^\rho \sin(\rho \theta_1) \leq -r^\rho \sin(\rho \theta_1)/2$  for  $r \geq b|x|^\delta \operatorname{cosec} \theta_1$ . Hence we have by (1.12)

$$(1.18) \quad \begin{aligned} |I'_1(x)| &\leq a_\rho^n \int_0^\infty r^n e^{-\sin(\rho \theta_1) r^{\rho/2}} dr \\ &\leq \left\{ \int_0^\infty e^{-\sin(\rho \theta_1) r^{\rho/4}} dr \right\} C_{\rho, \theta_1}^{n+1} (n!)^{1/\rho}. \end{aligned}$$

For  $I'_4(x)$  and  $I'_5(x)$  we can get the same estimates as  $I'_2(x)$  and  $I'_1(x)$  respectively. Hence if we determine  $b$  such as  $b(3a_\rho + b^{\rho-1} \operatorname{cosec}^\rho \theta_0) \leq 1$  and  $\theta_1 > 0$  such as  $a_\rho \leq b^{\rho-1} \operatorname{cosec}^\rho \theta_1 \sin(\rho \theta_1)/2$ , we have

$$(1.19) \quad |f_\rho^{(n)}(x)| \leq C_6^{n+1}(n!)^{1/\rho} e^{|x|^{1/\rho}} \quad \text{for } x < 0$$

by (1.16), (1.17), (1.18). Consequently, if we set

$$F_{\alpha, \gamma}(z) = f_{1/\alpha}(\gamma^\alpha z) \cdot f_{1/\alpha}(-\gamma^\alpha z)$$

then, using Leibniz' formula and

$$\sum_{j=0}^n \binom{n}{j} (j!)^\alpha ((n-j)!)^\alpha \leq (n!)^\alpha \sum_{j=0}^n \binom{n}{j} = (n!)^\alpha 2^n,$$

we get (1.2) by (1.15) and (1.19).

For  $2m-1 < \rho < 2m+1$ , we set  $a_\rho = 5 \operatorname{cosec}^\rho \beta_\rho$  with  $\beta_\rho$  of (1.6). For  $x > 0$ , in order to estimate (1.9), we deform the path of the integral (1.9) to  $\Gamma'' = C_1'' \cup C_2'' \cup C_3''$  where  $C_1''$ ,  $C_2''$ , and  $C_3''$  are the segments from  $-\infty + ix^\delta$  to  $-x^\delta \cot \beta_\rho + ix^\delta$ , from  $-x^\delta \cot \beta_\rho + ix^\delta$  to  $x^\delta \cot \beta_\rho + ix^\delta$ , and from  $x^\delta \cot \beta_\rho + ix^\delta$  to  $+\infty + ix^\delta$  respectively. For  $x < 0$  we need not deform the path and we evaluate the integral on the intervals  $(-\infty + i, -\cot \beta_\rho + i)$ ,  $(-\cot \beta_\rho + i, \cot \beta_\rho + i)$ , and  $(\cot \beta_\rho + i, +\infty + i)$ . Then, using (1.8), we get the same result as the case of  $\rho = 2m+1$ .

**Lemma 2.** *Let  $\varphi(x)$  be a function of  $G(\alpha, g(x))$  satisfying (1.1). Then, for any real  $y$ ,  $\psi(x) \equiv e^{ixy}\varphi(x)$  is also of  $G(\alpha, g(x))$  and satisfies*

$$|\psi^{(n)}(x)| \leq (C + |y|)^{n+1} (n!)^\alpha g(x) \quad (n = 0, 1, 2, \dots).$$

*Proof.* Omitted, as it is so easy.

2. For any function  $f(t, x)$  such that  $(\partial/\partial x)^{(m/\alpha_0)} f(t, x)^5$  is continuous on  $\{0 \leq t \leq T\} \times \{-\infty < x < +\infty\}$ , we define  $(Hf)(t, x)$  as follows.

$$(2.1) \quad Hf = \sum_{j=1}^m \sum_{k=0}^{(j/\alpha_0)} (-1)^{j+k+1} H_{j,k} f$$

where

$$(2.2) \quad (H_{j,k} f)(t, x) = \int_0^t \frac{(t-\tau)^{j-1}}{(j-1)!} a_{m-j,k}(\tau) \frac{\partial^k}{\partial x^k} f(\tau, x) d\tau.$$

**Theorem 1.** *Let  $g(x)$  be any positive function on  $-\infty < x < +\infty$  and  $f(t, x)$  be a complex-valued function whose derivatives  $\partial^n f/\partial x^n$  ( $n=0, 1, \dots$ ) are continuous on  $\{0 \leq t \leq T\} \times \{-\infty < x < +\infty\}$  and satisfy*

$$(2.3) \quad \left| \frac{\partial^n}{\partial x^n} f(t, x) \right| \leq C^{n+1} (n!)^{\alpha_0} g(x) \quad (n = 0, 1, \dots; 0 \leq t \leq T, -\infty < x < +\infty)$$

---

5) For a real number  $a$ ,  $[a]$  means the maximum of the integers which do not exceed  $a$ .

for some constant  $C$ . Then there exists only one function  $v(t, x)$  that satisfies

$$(2.4) \quad v - Hv = f \quad (0 \leq t < T_0, \quad -\infty < x < +\infty),$$

whose derivatives  $\partial^n v / \partial x^n$  ( $n=0, 1, 2, \dots$ ) are continuous on  $\{0 \leq t < T_0\} \times \{-\infty < x < +\infty\}$  and satisfy for some constant  $K$

$$(2.5) \quad \left| \frac{\partial^n}{\partial x^n} v(t, x) \right| \leq K^{n+1} (n!)^{\alpha_0} g(x) \quad (n=0, 1, \dots; \quad 0 \leq t < T_0, \quad -\infty < x < +\infty)$$

where  $T_0 > 0$  depends only on  $\alpha_0, C$  and

$$(2.6) \quad A \equiv \text{Max} \{ |a_{j,k}(t)|; \quad j + \alpha_0 k \leq m, \quad 0 \leq t \leq T \}.$$

Proof. Without loss of generality we assume  $C > 1$  and estimate  $|H^n f|$ . By (2.1), (2.2), (2.3), (2.6), we get easily

$$(2.7) \quad |(Hf)(t, x)| \leq CA(m/\alpha_0 + 1)g(x) \sum_{j=1}^m C^{j/\alpha_0} ([j/\alpha_0]!)^{\alpha_0} t^j / j!.$$

By Stirling's formula we have

$$(2.8) \quad \frac{([j/\alpha_0]!)^{\alpha_0}}{j!} \leq K_0 \frac{\{(j/\alpha_0)^{j/\alpha_0} \sqrt{2\pi j/\alpha_0} e^{-j/\alpha_0}\}^{\alpha_0}}{j^j \sqrt{2\pi j} e^{-j}} \leq K_1 \alpha_0^{-j} \quad (j = 1, 2, \dots)$$

for some constants  $K_0$  and  $K_1$ . Hence, replacing  $([j/\alpha_0]!)^{\alpha_0} / j!$  of (2.7) by  $K_1 \alpha_0^{-j}$ , we get

$$\begin{aligned} |Hf(t, x)| &\leq CAK_1(m/\alpha_0 + 1)g(x) \sum_{j=1}^m (C^{1/\alpha_0} t / \alpha_0)^j \\ &\leq CAK_1 m(m/\alpha_0 + 1)g(x) C^{1/\alpha_0} t / \alpha_0 \quad \text{for } 0 \leq C^{1/\alpha_0} t / \alpha_0 < 1, \end{aligned}$$

so that

$$|Hf(t, x)| \leq CK_1 g(x) t / (2T_0) \quad \text{for } 0 \leq t < 2T_0$$

where

$$(2.9) \quad 1/(2T_0) = \text{Max} \{ m(m/\alpha_0 + 1)(A+1)C^{1/\alpha_0} / \alpha_0, 1/T \}.$$

Next we estimate  $|H^2 f|$ . Since by (2.2)

$$\begin{aligned} (H_{h,l} \cdot H_{j,k} f)(t, x) \\ = \int_0^t \frac{(t-\sigma)^{h-1}}{(h-1)!} a_{m-h,l}(\sigma) \int_0^\sigma \frac{(\sigma-\tau)^{j-1}}{(j-1)!} a_{m-j,k}(\tau) \frac{\partial^{l+k}}{\partial x^{l+k}} f(\tau, x) d\tau d\sigma \end{aligned}$$

we get, remarking  $l+k \leq [(h+j)/\alpha_0]$ ,

$$\begin{aligned} |H_{h,l} \cdot H_{j,k} f(t, x)| &\leq A^2 \int_0^t \frac{(t-\tau)^{h+j-1}}{(h+j-1)!} \left| \frac{\partial^{l+k}}{\partial x^{l+k}} f(\tau, x) \right| d\tau \\ &\leq CA^2 C^{(h+j)/\alpha_0} ([ (h+j)/\alpha_0 ]!)^{\alpha_0} g(x) t^{h+j} / (h+j)!. \end{aligned}$$

If we replace  $([(h+j)/\alpha_0]!)^{\alpha_0}/(h+j)!$  by  $K_1\alpha_0^{-(h+j)}$  by (2.8), then we get

$$|H^2f(t, x)| \leq CK_1g(x)(t/(2T_0))^2 \quad \text{for } 0 \leq t < 2T_0.$$

Similarly we get for  $n=3, 4, \dots$

$$|H^n f(t, x)| \leq CK_1g(x)(t/(2T_0))^n \quad \text{for } 0 \leq t < 2T_0.$$

Hence we can set

$$v(t, x) = \sum_{n=0}^{\infty} (H^n f)(t, x) \quad \text{for } 0 \leq t < 2T_0,$$

so that  $v(t, x)$  is the unique solution of (2.4).

Next we estimate the value of  $(\partial/\partial x)^n Hf = H(\partial/\partial x)^n f$  ( $n=0, 1, 2, \dots$ ).  
 Remarking  $2^{n+l} \geq (n+l)!/(n!l!)$ , we get in the same way as (2.7)

$$\begin{aligned} \left| \frac{\partial^n}{\partial x^n} Hf(t, x) \right| &\leq CA(m/\alpha_0 + 1)g(x) \sum_{j=1}^m C^{j/\alpha_0 + n} \{(n + [j/\alpha_0])!\}^{\alpha_0} t^j / j! \\ &\leq CA(m/\alpha_0 + 1)g(x) \sum_{j=1}^m (2C)^{j/\alpha_0 + n} (n!)^{\alpha_0} ([j/\alpha_0]!)^{\alpha_0} t^j / j! \\ &\leq (2C)^{n+1} (n!)^{\alpha_0} K_1g(x)t/T_0 \quad \text{for } 0 \leq t < T_0. \end{aligned}$$

Similarly we get for  $l=1, 2, \dots$

$$\left| \frac{\partial^n}{\partial x^n} H^l f(t, x) \right| \leq (2C)^{n+1} (n!)^{\alpha_0} K_1g(x)(t/T_0)^l \quad (0 \leq t < T_0).$$

Hence we get

$$\left| \frac{\partial^n}{\partial x^n} v(t, x) \right| = \left| \sum_{l=0}^{\infty} \left( \frac{\partial}{\partial x} \right)^n H^l f(t, x) \right| \leq (2C)^{n+1} (n!)^{\alpha_0} K_1g(x)T_0/(T_0 - t)$$

for  $0 \leq t < T_0$  so that (2.5) holds.

**Theorem 2.** *Let  $u(t, x)$  be a solution of the Cauchy problem*

$$(2.10) \quad \begin{aligned} (Lu)(t, x) &= 0 \quad (0 \leq t \leq T, \quad -\infty < x < +\infty) \\ \frac{\partial^j u}{\partial t^j}(0, x) &= 0 \quad (j = 0, \dots, m-1; \quad -\infty < x < +\infty) \end{aligned}$$

whose derivatives contained in  $L$  are continuous and satisfy

$$(2.11) \quad \left| \frac{\partial^{j+k}}{\partial t^j \partial x^k} u(t, x) \right| \leq C e^{\gamma|x|^{1/(1-\alpha_0)}} \quad (j + \alpha_0 k \leq m, \quad 0 \leq t \leq T, \quad -\infty < x < +\infty)$$

for some constants  $C$  and  $\gamma > 0$ . Then  $u(t, x)$  vanishes identically for  $0 \leq t \leq T, \quad -\infty < x < +\infty$ .

**Proof.** We write  $\beta_0 = 1/(1-\alpha_0)$ . By Lemma 1 we can take a non-trivial function  $\varphi(x)$  of  $G(\alpha_0, e^{-2\gamma|x|^{\beta_0}})$  satisfying

$$(2.12) \quad |\varphi^{(n)}(x)| \leq C_1^{n+1}(n!)^{\alpha_0} e^{-2\gamma|x|\beta_0} \quad (n = 0, 1, 2, \dots)$$

for some constant  $C_1$  depending only on  $\alpha_0$  and  $\gamma$ . We set

$$(2.13) \quad 1/(2T_0) = \text{Max} \{m(m\alpha_0 + 1)(A + 1)(1 + C_1)^{1/\alpha_0}/\alpha_0, 1/T\}$$

where  $A = \text{Max} \{|a_{j,k}(t)|; 0 \leq t \leq T, j + \alpha_0 k \leq m\}$  (cf. (2.9)).

If the assertion of the theorem is not true, then, for some  $u(t, x)$  satisfying (2.10) and (2.11), there exists  $t_0$  such that  $t_0 = \inf \{0 \leq t \leq T; u(t, x) \neq 0 \text{ for some } x\} < T$ . Taking  $\tau$  such as  $0 < \tau < T' \equiv \text{Min}(T_0, T - t_0)$ , we set  $(-1)^j b_{j,k}(t) = a_{j,k}(t_0 + \tau - t)$  ( $j + \alpha_0 k \leq m$ ) and  $w(t, x) = u(t_0 + \tau - t, x)$  ( $0 \leq t \leq \tau, -\infty < x < +\infty$ ).

Let  $L', H'$  and  $H'_{j,k}$  be the operators given by (0.1), (2.1) and (2.2) replacing  $a_{j,k}(t)$  by  $b_{j,k}(t)$  respectively. Then  $w(t, x)$  is a solution of the Cauchy problem,  $L'w = 0$  ( $0 \leq t \leq \tau, -\infty < x < +\infty$ ),  $(\partial/\partial t)^j w(\tau, x) = 0$  ( $j = 0, \dots, m-1; -\infty < x < +\infty$ ), satisfying the same condition as (2.11). By Lemma 2, (2.13) and the proof of Theorem 1, we obtain for  $-1 \leq y \leq 1$  the solution  $v(t, x; y)$  of  $v - H'v = e^{ixy}\varphi(x)$  on  $\{0 \leq t \leq \tau\} \times \{-\infty < x < +\infty\}$  where  $\tau$  is independent of  $y$ . Furthermore

$$\left| \frac{\partial^n}{\partial x^n} v(t, x; y) \right| \leq C_2^{n+1}(n!)^{\alpha_0} e^{-2\gamma|x|\beta_0} \quad (n = 0, 1, 2, \dots)$$

for some constant  $C_2$ . Integrating by parts, we get easily

$$(-1)^{j+k} \int_{-\infty}^{+\infty} \int_0^\tau (H'_{j,k}v) \frac{\partial^m w}{\partial t^m} dt dx = \int_{-\infty}^{+\infty} \int_0^\tau v b_{m-j,k} \left(\frac{\partial}{\partial t}\right)^{m-j} \left(\frac{\partial}{\partial x}\right)^k w dt dx,$$

and hence

$$\begin{aligned} 0 &= \int_{-\infty}^{+\infty} \int_0^\tau v(L'w) dt dx = \int_{-\infty}^{+\infty} \int_0^\tau (v - H'v) \frac{\partial^m w}{\partial t^m} dt dx \\ &= \int_{-\infty}^{+\infty} e^{ixy}\varphi(x) \frac{\partial^{m-1} w}{\partial t^{m-1}}(0, x) dx \quad \text{for } |y| \leq 1. \end{aligned}$$

Since  $\int_{-\infty}^{+\infty} e^{ixy}\varphi(x) \frac{\partial^{m-1} w}{\partial t^{m-1}}(0, x) dx$  is an entire function of  $y$  by (2.11) and (2.12), this vanishes identically on  $-\infty < y < +\infty$ . Hence the integrable function  $\varphi(x)(\partial^{m-1} w/\partial t^{m-1})(0, x)$  also vanishes identically on  $-\infty < x < +\infty$ . Since  $\varphi(x)$  is a non-trivial entire function, we get

$$(-1)^{m-1} \frac{\partial^{m-1} w}{\partial t^{m-1}}(0, x) = \frac{\partial^{m-1} u}{\partial t^{m-1}}(t_0 + \tau, x) = 0$$

for  $0 \leq \tau < T', -\infty < x < +\infty$ .

By  $(\partial^j u/\partial t^j)(t_0, x) = 0$  ( $j = 0, \dots, m-1; -\infty < x < +\infty$ )

we get  $u(t, x) \equiv 0$  for  $t_0 \leq t < t_0 + T', -\infty < x < +\infty$ ,

which is contradiction.



3. In this section we shall consider a differential operator with constant coefficients

$$(3.1) \quad L\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x}\right) = \sum_{j+\alpha_0 k \leq m} a_{j,k} \frac{\partial^{j+k}}{\partial t^j \partial x^k} \quad (0 < \alpha_0 < 1, a_{m,0} = 1)$$

containing at least one coefficient  $a_{j_0, k_0} \neq 0$  with  $j_0 \neq m$  and  $j_0 + \alpha_0 k_0 = m$ .

**Lemma 3.** *Let  $L(-i\lambda, -i\eta) = \sum_{j+\alpha_0 k \leq m} a_{j,k} (-i\lambda)^j (-i\eta)^k$  be the differential polynomial corresponding to (3.1). Then we have an analytic function  $\eta = \eta(\lambda)$  defined on  $\Im m \lambda \geq K$  for some constant  $K > 0$  such that  $L(-i\lambda, -i\eta(\lambda)) = 0$  and  $|\eta(\lambda)| \leq A |\lambda|^{\alpha_0}$  for some constant  $A$ .*

*Proof.* We write

$$\begin{aligned} L(-i\lambda, -i\eta) &= (-i)^m \prod_{j=1}^m (\lambda - \lambda_j(\eta)) \\ &= (-i)^m (\lambda^m + Q_1(\eta)\lambda^{m-1} + \dots + Q_{m-j_0}(\eta)\lambda^{j_0} + \dots + Q_m(\eta)). \end{aligned}$$

Then  $\lambda_j(\eta)$  have Puiseux series expansions at infinity<sup>6)</sup>

$$\lambda_j(\eta) = \sum_{n=-\infty}^{l_j} \alpha_{j,n} (\eta^{1/p_j})^n \quad (\alpha_{j,l_j} \neq 0, j = 1, \dots, m)$$

on  $|\eta| \geq K'$  for some constant  $K' > 0$ . Hence  $\lambda_j(\eta) = \alpha_{j,l_j} \eta^{l_j/p_j} (1 + o(1))$  as  $\eta \rightarrow \infty$ . If we would assume  $l_j/p_j < 1/\alpha_0$  for all  $j = 1, \dots, m$  then on  $|\eta| \geq K'$

$$(3.2) \quad |Q_{m-j_0}(\eta)| = \left| \sum_{j_1, \dots, j_{m-j_0}} \lambda_{j_1}(\eta) \dots \lambda_{j_{m-j_0}}(\eta) \right| \leq C |\eta|^{(m-j_0)/\alpha_0 - \varepsilon}$$

for some  $C, \varepsilon > 0$ . On the other hand

$$Q_{m-j_0}(\eta) = a\eta^{(m-j_0)/\alpha_0} + \dots \quad (a = i^{m-j_0-k_0} a_{j_0, k_0} \neq 0),$$

so that  $Q_{m-j_0}(\eta) = a\eta^{(m-j_0)/\alpha_0} (1 + o(1))$  as  $\eta \rightarrow \infty$ . This contradicts to (3.2). Consequently we have an analytic function

$$\lambda(\eta) = \sum_{n=-\infty}^l a_n (\eta^{1/p})^n \quad (a_l \neq 0, l/p \geq 1/\alpha_0)$$

on  $|\eta| \geq K'$  such that  $L(-i\lambda(\eta), -i\eta) = 0$ . Since  $\lambda(\eta) = \alpha_l \eta^{l/p} (1 + o(1))$  as  $\eta \rightarrow \infty$ , we have an analytic function  $\eta = \eta(\lambda)$  defined on  $\Im m \lambda \geq K$  for some constant  $K \geq 1$  such that  $L(-i\lambda, -i\eta(\lambda)) = 0$  and  $|\eta(\lambda)| \leq A |\lambda|^{p/l} \leq A |\lambda|^{\alpha_0}$  for some constant  $A$ .

**Theorem 3.** *Let  $L\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x}\right)$  be a differential operator of the form*

6) [12], pp. 50-55.

(3.1) containing at least one coefficient  $a_{j_0 k_0} \neq 0$  with  $j_0 \neq m$  and  $j_0 + \alpha_0 k_0 = m$ . Then for any  $\varepsilon_0 > 0$ , we can construct a  $C^\infty$ -solution  $U(t, x)$  of

$$L\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x}\right)U(t, x) = 0 \quad (-\infty < t < +\infty, -\infty < x < +\infty)$$

such that

$$(3.3) \quad \begin{cases} U(t, x) \neq 0 & \text{for } (-\infty < t < +\infty, -\infty < x < +\infty), \\ U(t, x) = 0 & \text{for } (-\infty < t \leq 0, -\infty < x < +\infty), \end{cases}$$

and

$$(3.4) \quad \left| \frac{\partial^{j+k}}{\partial t^j \partial x^k} U(t, x) \right| \leq C_1^{1+j+k} (j!)^{1+\varepsilon_0} (k!)^{\alpha_0+\varepsilon_0} e^{C_2(|x|^{1/(1-\alpha_0)^{2+\varepsilon_0}} + 1)t}$$

$$(-\infty < t < +\infty, -\infty < x < +\infty)$$

for some positive constants  $C_1$  and  $C_2$  depending only on  $\varepsilon_0$ .

Proof. Consider the equation  $L(-i\lambda, -i\eta) = 0$ . Then by Lemma 3, we have an analytic root  $\eta(\lambda)$  defined on  $\Im m \lambda \geq K$  for some constant  $K > 0$  which satisfies

$$(3.5) \quad |\eta(\lambda)| \leq A |\lambda|^{\alpha_0} \quad (\Im m \lambda \geq K)$$

for some constant  $A$ . Take  $\rho$  such that  $0 < \alpha_0 < \rho < 1$  and consider the function

$$(3.6) \quad U_\rho(t, x) = \int_{-\infty+i\tau}^{+\infty+i\tau} e^{-i(t\lambda+x\eta(\lambda))-(\lambda/i)^\rho} d\lambda^\tau \quad (\tau \geq K)$$

where  $(\lambda/i)^\rho$  is defined such that  $(\lambda/i)^\rho > 0$  on the positive imaginary axis. Since  $\Re(\lambda/i)^\rho \geq |\lambda|^\rho \cos(\rho\pi/2)$ , by (3.5) we can see that  $U_\rho(t, x) \in C^\infty(-\infty < t < +\infty, -\infty < x < +\infty)$ ,  $L\left(\frac{\partial}{\partial t}, \frac{\partial}{\partial x}\right)U_\rho(t, x) = 0$ , and that  $U_\rho$  does not depend on  $\tau (\geq K)$ . For any fixed  $x$ , if we set  $C_\rho = \{2A \cos(\rho\pi/2)\}^{1/(\rho-\alpha_0)}$ , then for  $\Im m \lambda = \tau \geq (C_\rho |x|^{1/(\rho-\alpha_0)} + K)$  we have

$$(3.7) \quad \begin{aligned} |x\eta(\lambda)| - |\lambda|^\rho \cos(\rho\pi/2) &\leq A |\lambda|^{\alpha_0} |x| - |\lambda|^\rho \cos(\rho\pi/2). \\ &\leq -\frac{1}{2} |\lambda|^\rho \cos(\rho\pi/2). \end{aligned}$$

Taking  $t < 0$  and  $x$  arbitrarily, we have

$$\begin{aligned} |U_\rho(t, x)| &\leq e^{t\tau} \int_{-\infty+i\tau}^{+\infty+i\tau} e^{-|\lambda|^\rho \cos(\rho\pi/2)/2} d|\lambda| \\ &\leq A e^{t\tau} \int_{-\infty}^{+\infty} e^{-|y|^\rho \cos(\rho\pi/2)/2} dy \rightarrow 0 \quad (\tau \rightarrow \infty) \end{aligned}$$

7) [7], p. 122.

for  $\tau \geq C_\rho |x|^{1/(\rho-\alpha_0)} + K$ . Hence  $U_\rho(t, x) = 0$  for  $-\infty < t < 0$  and  $-\infty < x < +\infty$ . Making  $x$  fixed, we can write

$$U_\rho(t, x) = e^{t\tau} \int_{-\infty}^{+\infty} e^{-it y} e^{-ix\eta(x+i\tau) - (-iy+\tau)^\rho} dy,$$

so that we have  $U_\rho(t, x) \equiv 0$  as a function of  $t$  on  $(-\infty, \infty)$ . We get (3.3) by these arguments.

Next, for the derivatives of  $U_\rho(t, x)$ , we have by (3.5)

$$(3.8) \quad \left| \frac{\partial^{j+k}}{\partial t^j \partial x^k} U_\rho(t, x) \right| \leq A^k e^{\tau t} \int_{-\infty+i\tau}^{+\infty+i\tau} |\lambda|^j |\lambda|^{\alpha_0 k} e^{-|\lambda|^\rho \cos(\rho\pi/2)} d|\lambda|$$

where  $\tau = C_\rho |x|^{1/(\rho-\alpha_0)} + K$ . If we write the integrand of the right hand side of (3.8) as

$$(|\lambda|^j e^{-|\lambda|^\rho \cos(\rho\pi/2)}) (|\lambda|^{\alpha_0 k} e^{-|\lambda|^{\alpha_0 \rho / \alpha_0 \cos(\rho\pi/2)}}) (e^{-|\lambda|^\rho \cos(\rho\pi/2)})$$

then by (1.12) we have

$$\begin{aligned} \left| \frac{\partial^{j+k}}{\partial t^j \partial x^k} U_\rho(t, x) \right| &\leq A^k e^{\tau t} C_\rho^j (j!)^{1/\rho} C_{\rho, \alpha_0}^k (k!)^{\alpha_0/\rho} \int_{-\infty+i\tau}^{+\infty+i\tau} e^{-|\lambda|^\rho \cos(\rho\pi/2)} d|\lambda| \\ &\leq C^{1+j+k} e^{(C_\rho |x|^{1/(\rho-\alpha_0)} + K)t} (j!)^{1/\rho} (k!)^{\alpha_0/\rho} \int_{-\infty}^{+\infty} e^{-|y|^\rho \cos(\rho\pi/2)} dy \end{aligned}$$

where  $C$  is a constant depending on  $A, \rho$  and  $\alpha_0$ . We fix  $\rho$  ( $\alpha_0 < \rho < 1$ ) such as  $1/\rho \leq 1 + \varepsilon_0$ ,  $\alpha_0/\rho \leq \alpha_0 + \varepsilon_0$ , and  $1/(\rho - \alpha_0) \leq 1/(1 - \alpha_0) + \varepsilon_0$ . Hence we get (3.4).

OSAKA UNIVERSITY  
UNIVERSITY OF OSAKA PREFECTURE

References

[1] K. I. Babenko: *On a new problem of quasi-analyticity and on Fourier transforms of entire functions*, Trudy Moskov. Mat. Obsč. 5 (1956), 523-542.  
 [2] A. Friedman: *Generalized functions and partial differential equations*, Englewood Cliffs N. J., 1963.  
 [3] I. M. Gelfand and G. E. Shilov: *Fourier transforms of rapidly increasing functions and questions of uniqueness for the solution of the Cauchy problem*, Uspehi Mat. Nauk SSSR 8 No. 6 (1953), 3-54.  
 [4] E. de Giorgi: *Un teorema di unicit  per il problema di Cauchy, relativo ad equazioni differenziali lineari a derivate parziali di tipo parabolico*, Ann. Mat. Pura Appl. Ser. IV, 40 (1955), 371-377.

- [ 5 ] B. L. Gurevich: *New types of spaces of fundamental and generalized functions and Cauchy's problem for systems of finite difference equations*, Doklady Akad. Nauk SSSR (N.S.) **99** (1954), 893–895.
- [ 6 ] B. L. Gurevich: *New types of fundamental and generalized spaces and Cauchy's problem for systems of difference equations involving differential operations*, Doklady Akad. Nauk SSSR (N.S.) **108** (1956), 1001–1003.
- [ 7 ] L. Hörmander: *Linear partial differential operators*, Berlin, 1963.
- [ 8 ] B. Le Roy: *Valeurs asymptotiques de certaines séries procédant suivant les puissances entières et positives d'une variable réelle*, Bull. Sci. Math. (2) **24** (1900), 245–268.
- [ 9 ] G. Talenti: *Un problema di Cauchy*, Ann. Scuola Norm. Sup. Pisa (3) **18** (1964), 165–186.
- [10] E. C. Titchmarsh: *The theory of functions*, Oxford University Press, 1939.
- [11] A. Tsutsumi: *A remark on the uniqueness of the noncharacteristic Cauchy problem for equations of parabolic type*, Proc. Jap. Acad. **41** (1965), 65–70.
- [12] B. L. van der Waerden: *Einführung in die algebraische Geometrie*, Berlin, 1939.