

## ON STRONGLY SEPARABLE ALGEBRAS

AKIRA HATTORI

(Received July 27, 1965)

Let  $A$  be a separable algebra over a commutative ring  $R$  [1] with center  $C$ . There exists an element  $\sum u_i \otimes v_i^0 \in A \otimes A^0$  such that

- i)  $\sum xu_i \otimes v_i^0 = \sum u_i \otimes (v_i x)^0 \quad (x \in A),$
- ii)  $\sum u_i v_i = 1$

The mapping  $x \rightarrow \sum u_i x v_i$  is a projection  $A \rightarrow C$ , and  $C$  is a  $C$ -direct summand of  $A$ . While, the mapping  $\lambda : x \rightarrow \sum v_i x u_i$  yields a  $C$ -direct decomposition  $A = \lambda(A) \oplus [A, A]$ , where  $[A, A]$  is the  $C$ -linear subspace spanned by all  $[x, y] = xy - yx$  [2]. In many important cases  $\lambda(A)$  coincides with  $C$ , and we have  $A = C \oplus [A, A]$ . Giving the name strongly separable algebras, T. Kanzaki studied such a class of algebras. He first studied strongly separable algebras over a field [3], and then the general case was studied in utilizing the former results [4]. The objective of this note is to re-establish his results more simply, and to generalize in some respects.

**Theorem 1.** *The following three statements are equivalent for an algebra  $A$  over  $R$ .*

- 1) *There exists an element  $s = \sum v_i \otimes u_i^0 \in A \otimes A^0$  such that*
  - i)  $\sum v_i x \otimes u_i^0 = \sum v_i \otimes (x u_i)^0 \quad (x \in A),$
  - ii)  $\sum v_i u_i = 1.$
- 2)  *$A$  is separable and  $A = C \oplus [A, A]$*
- 3)  *$A$  is separable and there exists a symmetric  $C$ -linear map  $\lambda : A \rightarrow C$  such that  $\lambda(1) = 1.$*

REMARK.  $A$  is called *strongly separable* if  $A$  satisfies these conditions. The original definition by Kanzaki was given by the condition 1), formulated in another way, and he proved the equivalence of 1) and 2) [4, Th. 1].

Proof. 1)  $\Rightarrow$  2). Put  $\lambda(x) = \sum v_i x u_i$ .  $\lambda$  is a  $C$ -linear map  $A \rightarrow A$ , is identity on  $C$ , and vanishes on  $[A, A]$ . ( $\lambda(xy) = \sum (v_i x) y u_i = \sum v_i y (x u_i) = \lambda(yx)$ .) It follows  $C \cap [A, A] = 0$ . For any  $x \in A$ , we have  $x = \sum v_i u_i x$

$= \sum u_i x v_i + \sum [v_i, u_i x] \in C + [A, A]$ . Hence we have  $A = C \oplus [A, A]$ .  $1$  is represented as

$$1 = \sum v_i u_i = \sum u_i v_i + \sum [v_i, u_i] = 1 + 0,$$

so that we have  $\sum u_i v_i = 1$ . Since we have moreover  $\sum x u_i \otimes v_i^0 = \sum u_i \otimes (v_i x)^0$ ,  $A$  is separable.

2)  $\Rightarrow$  3) Take the projection  $\lambda : A \rightarrow C$ .

3)  $\Rightarrow$  1) As  $A$  is central separable over  $C$ , the mapping  $\eta : A \otimes_C A^0 \rightarrow$

$\text{Hom}_C(A, A)$  defined by  $\eta(a \otimes b^0)(x) = axb$  is an isomorphism [1]. Let, in particular,  $\lambda = \eta(\sum v_i \otimes u_i^0)$ . Then we have

ii)  $\sum v_i u_i = \lambda(1) = 1.$

Further

$$\begin{aligned} \eta(\sum v_i y \otimes u_i^0)(x) &= \sum v_i y x u_i = \lambda(yx) \\ &= \lambda(xy) = \sum v_i x y u_i = \eta(\sum v_i \otimes (y u_i)^0). \end{aligned}$$

This means

i)  $\sum v_i y \otimes u_i^0 = \sum v_i \otimes (y u_i)^0.$

Hence  $s_C = \sum v_i \otimes_C u_i^0$  is the desired element in  $A \otimes_C A^0$ .

Now,  $C$  is separable over  $R$ , and there exists  $\sum p_j \otimes q_j \in C \otimes C$  such that  $\sum c p_j \otimes q_j = \sum p_j \otimes q_j c$  ( $c \in C$ ) and  $\sum p_j q_j = 1$ . The mapping  $a \otimes_C b^0 \rightarrow \sum a p_j \otimes_R (b q_j)^0$  is a well-defined ring homomorphism  $\sigma : A \otimes_C A^0 \rightarrow A \otimes_R A^0$  such that  $\tau \sigma = \text{identity}$ , where  $\tau$  denotes the natural epimorphism  $A \otimes_R A^0 \rightarrow A \otimes_C A^0$ . Put  $s = \sigma(s_C) = \sum v_i p_j \otimes (u_i q_j)^0$ . We have

i)  $\sum v_i p_j u_i q_j = \sum v_i u_i \sum p_j q_j = 1.$

Since  $s_C(x \otimes_C 1^0 - 1 \otimes_C x^0) = 0$ , we have, applying  $\sigma$ ,

ii)  $s(x \otimes_R 1^0 - 1 \otimes_R x^0) = 0.$

Hence  $s$  has the desired properties. q.e.d.

**Corollary** [4, Lemma 1]. *A is strongly separable if and only if A is strongly separable over C and C is separable.*

We now study a strongly separable algebra  $A$  over its center  $C$ . Let  $A^*$  be the dual space of  $A : A^* = \text{Hom}_C(A, C)$ , considered as a two-sided  $A$ -module by  $a\xi(x) = \xi(xa)$ ,  $\xi a(x) = \xi(ax)$ . Then the mapping  $x \otimes \xi \rightarrow x\xi$  yields an isomorphism  $A \otimes_C (A^*)^A = A^*$  [1]. But  $(A^*)^A = \{\xi \in A^* \mid$

$a\xi = \xi a, a \in A$  consists of symmetric linear forms. Since  $A = C \oplus [A, A]$ , this set consists of scalar multiples of  $\lambda$ . It follows that the mapping  $\tilde{\lambda} : x \rightarrow x\lambda$  gives an isomorphism of two-sided  $A$ -modules  $A \cong A^*$  (i.e.  $A$  is a symmetric algebra).

In [2], we defined the rank element of a projective module. Since  $A$  is  $C$ -finite projective, we can speak of the rank element  $r_C(A)$  which is an element of  $C$ . Then, as a natural generalization of [3, Theorem], we have

**Theorem 2.** *A central separable algebra  $A$  over  $C$  is strongly separable if and only if  $r_C(A)$  is a unit of  $C$ .*

*Proof.* Assume that  $A$  is strongly separable. Since  $A$  is  $C$ -finite projective, we have an isomorphism  $\theta : A^* \otimes_C A^0 \cong \text{Hom}_C(A, A)$  by  $\theta(\xi \otimes a^0)(x) = \xi(x)a$ . Composing this with  $\tilde{\lambda} \otimes 1 : A \otimes_C A^0 \cong A^* \otimes_C A^0$ , we have an isomorphism  $f : A \otimes_C A^0 \cong \text{Hom}_C(A, A)$ , which is defined by  $f(a \otimes b^0)(x) = \lambda(ax)b$ . Let  $\theta(\sum \xi_i \otimes u_i) = \text{identity of } A$  ( $u_i$  as above), and  $\xi_i = w_i \lambda$  ( $w_i \in A$ ). Then we have  $x = \sum \lambda(xw_i)u_i$ . Since

$$\begin{aligned} f(\sum xw_i \otimes u_i^0)(y) &= \sum (xw_i y)u_i = yx \\ &= \sum (yw_i)u_i x = f(\sum w_i \otimes (u_i x)^0), \end{aligned}$$

we have

$$\sum xw_i \otimes u_i^0 = \sum w_i \otimes (u_i x)^0 \quad (x \in A).$$

It follows that  $\sum w_i u_i$  is in the center  $C$ , and we have

$$r_C(A) = \sum \xi_i(u_i) = \lambda(\sum w_i u_i) = \sum w_i u_i.$$

But

$$\begin{aligned} 1 &= \sum v_j u_j = \sum \lambda(u_j w_j) v_j u_j = \sum \lambda(u_j) (v_j w_j) u_j \\ &= (\sum \lambda(u_j) v_j) (\sum w_j u_j). \end{aligned}$$

This shows that  $r_C(A) = \sum w_i u_i$  is a unit of  $C$ .

Conversely assume that  $r_C(A) = c$  is a unit of  $C$ . Consider the trace :  $\text{Tr}(x) = \sum \xi_i(xu_i)$ , where  $\theta(\sum \xi_i \otimes u_i) = 1$  as above. We see that this is a symmetric linear form  $A \rightarrow C$  and  $r_C(A) = \text{Tr}(1)$  [2]. Put  $\lambda = c^{-1} \text{Tr}$ . Then  $\lambda$  is also a symmetric linear form  $A \rightarrow C$ , and satisfies  $\lambda(1) = c^{-1} \text{Tr}(1) = 1$ . Hence  $A$  is strongly separable. q.e.d.

**References**

- [ 1 ] M. Auslander and O. Goldman, *The Brauer group of a commutative ring*, Trans. Amer. Math. Soc. **97** (1960), 367-409.
- [ 2 ] A. Hattori, *Rank element of a projective module*, Nagoya Math. J. **25** (1965), 113-120.
- [ 3 ] T. Kanzaki, *A type of separable algebras*, J. Math. Osaka City Univ. **13** (1962), 39-43.
- [ 4 ] T. Kanzaki, *Special type of separable algebras over a commutative ring*, Proc. Japan Acad. **40** (1964), 781-786.