

ON HOMOGENEOUS KÄHLER MANIFOLDS WITH NON-DEGENERATE CANONICAL HERMITIAN FORM OF SIGNATURE $(2, 2(n-1))$

HIROHIKO SHIMA

(Received October 19, 1972)

We denote by M a connected homogeneous Kähler manifold of complex dimension n on which a connected Lie group G acts effectively as a group of holomorphic isometries, and by K an isotropy subgroup of G at a point o of M . Let v be the G -invariant volume element corresponding to the Kähler metric. In a local coordinate system $\{z_1, \dots, z_n\}$, v has an expression $v = i^n F dz_1 \wedge \dots \wedge dz_n \wedge d\bar{z}_1 \wedge \dots \wedge d\bar{z}_n$. The G -invariant hermitian form $h = \sum_{i,j} \frac{\partial^2 \log F}{\partial z_i \partial \bar{z}_j} dz_i d\bar{z}_j$ is called the canonical hermitian form of $M=G/K$. It is known that the Ricci tensor of the Kähler manifold M is equal to $-h$. The purpose of this paper is to prove the following:

Theorem 1. *Let $M=G/K$ be a simply connected homogeneous Kähler manifold with non-degenerate canonical hermitian form h of signature $(2, 2(n-1))$. Then, if either G is semi-simple or G contains a one parameter normal subgroup, $M=G/K$ is a holomorphic fibre bundle whose base space is the unit disk $\{z \in \mathbb{C}; |z| < 1\}$, and whose fibre is a homogeneous Kähler manifold of a compact semi-simple Lie group.*

In the case of $\dim_{\mathbb{C}} G/K=2$, the assumption of Theorem 1 is fulfilled and we have

Theorem 2. *Let $M=G/K$ be a complex two dimensional homogeneous Kähler manifold with non-degenerate canonical hermitian form h of signature $(2, 2)$. Then G is semi-simple or G contains a one parameter normal subgroup.*

As an application of these Theorems, we obtain a classification of complex two dimensional homogeneous Kähler manifolds with non-degenerate canonical hermitian form.

1. Let (I, g) be the G -invariant Kähler structure on M , i.e., I is the G -invariant complex structure tensor on M and g is the G -invariant Kähler metric on M . Let \mathfrak{g} be the Lie algebra of all left invariant vector fields on G and let \mathfrak{k} be the subalgebra of \mathfrak{g} corresponding to K . We denote by π the canonical projection

from G onto $M=G/K$ and denote by π_e the differential of π at the identity e of G . Let X_e, I_o and g_o be the values of X, I and g at e and $\pi(e)=o$ respectively. Then there exist a linear endomorphism J of \mathfrak{g} and a skew symmetric bilinear form ρ on \mathfrak{g} such that

$$\pi_e(JX)_e = I_o(\pi_e X_e), \quad \rho(X, Y) = g_o(\pi_e X_e, \pi_e Y_e),$$

for $X, Y \in \mathfrak{g}$. Then $(\mathfrak{g}, \mathfrak{k}, J, \rho)$ satisfies the following properties [2], [3].

- (1.1) $J\mathfrak{k} \subset \mathfrak{k}, \quad J^2 X \equiv -X \pmod{\mathfrak{k}},$
- (1.2) $[W, JX] \equiv J[W, X] \pmod{\mathfrak{k}},$
- (1.3) $[JX, JY] \equiv J[JX, Y] + J[X, JY] + [X, Y] \pmod{\mathfrak{k}},$
- (1.4) $\rho(W, X) = 0,$
- (1.5) $\rho(JX, JY) = \rho(X, Y),$
- (1.6) $\rho(JX, X) > 0, \quad X \notin \mathfrak{k},$
- (1.7) $\rho([X, Y], Z) + \rho([Y, Z], X) + \rho([Z, X], Y) = 0,$

where $X, Y, Z \in \mathfrak{g}, W \in \mathfrak{k}$.

Then $(\mathfrak{g}, \mathfrak{k}, J, \rho)$ will be called the Kähler algebra of $M=G/K$.

Koszul proved that the canonical hermitian form h of a homogeneous Kähler manifold G/K has the following expression [3]. Put

$$\begin{aligned} \eta(X, Y) &= h_o(\pi_e X_e, \pi_e Y_e), \quad \text{and} \\ (1.8) \quad \psi(X) &= \text{Tr}_{\mathfrak{g}/\mathfrak{k}}(\text{ad}(JX) - J \text{ad}(X)), \end{aligned}$$

it follows then

$$(1.9) \quad \eta(X, Y) = \frac{1}{2} \psi([JX, Y]),$$

for $X, Y \in \mathfrak{g}$. The form ψ satisfies the following properties:

- (1.10) $\psi([W, X]) = 0,$
- (1.11) $\psi([JX, JY]) = \psi([X, Y]), \quad \text{for } X, Y \in \mathfrak{g}, W \in \mathfrak{k}.$

Since G acts effectively on $G/K, \mathfrak{k}$ contains no non-zero ideal of \mathfrak{g} and there exists an $\text{ad}(\mathfrak{k})$ -invariant inner product $(,)$ on \mathfrak{g} . Henceforth, we assume that the canonical hermitian form h of G/K is non-degenerate, which is equivalent to the following condition:

Let $X \in \mathfrak{g}$. If $\eta(X, Y) = 0$ for all $Y \in \mathfrak{g}$, then $X \in \mathfrak{k}$.

2. We shall now prepare a few lemmas for later use. The following lemma is due to [2].

Lemma 2.1. For $E, X, Y \in \mathfrak{g}$,

$$\begin{aligned} & \frac{d}{dt} \rho(\exp t \operatorname{ad}(JE)X, \exp t \operatorname{ad}(JE)Y) \\ &= \rho(JE, \exp t \operatorname{ad}(JE)[X, Y]). \end{aligned}$$

Lemma 2.2. The adjoint representation of \mathfrak{g} is faithful.

Proof. Put $\mathfrak{a} = \{X \in \mathfrak{g}; \operatorname{ad}(X) = 0\}$, then \mathfrak{a} is an ideal of \mathfrak{g} . We have for $X \in \mathfrak{a}$

$$\begin{aligned} 2\eta(X, Y) &= \psi([JX, Y]) \\ &= -\psi([X, JY]) = 0, \quad \text{for all } Y \in \mathfrak{g}. \end{aligned}$$

Since h is non-degenerate, we have $X \in \mathfrak{k}$, and hence $\mathfrak{a} \subset \mathfrak{k}$. By the effectiveness, we have $\mathfrak{a} = \{0\}$. Q.E.D.

Lemma 2.3. Let \mathfrak{r} be a commutative ideal of \mathfrak{g} . Then, $\mathfrak{k} \cap \mathfrak{r} = \{0\}$, $\mathfrak{k} \cap J\mathfrak{r} = \{0\}$.

Proof. Let $A \in \mathfrak{k} \cap \mathfrak{r}$. Since \mathfrak{r} is a commutative ideal, we have $\operatorname{ad}(A)^2 = 0$. By the effectiveness, it follows that

$$\begin{aligned} (\operatorname{ad}(A)^2 X, X) + (\operatorname{ad}(A)X, \operatorname{ad}(A)X) &= 0, \\ (\operatorname{ad}(A)X, \operatorname{ad}(A)X) &= 0, \end{aligned}$$

for $X \in \mathfrak{g}$, with respect to the $\operatorname{ad}(\mathfrak{k})$ -invariant inner product (\cdot, \cdot) on \mathfrak{g} . Hence $\operatorname{ad}(A)X = 0$ for all $X \in \mathfrak{g}$, and $A = 0$ by Lemma 2.2, which proves $\mathfrak{k} \cap \mathfrak{r} = \{0\}$. $\mathfrak{k} \cap J\mathfrak{r} = \{0\}$ follows from $\mathfrak{k} \cap \mathfrak{r} = \{0\}$. Q.E.D.

Lemma 2.4. Let \mathfrak{r} be a non-zero commutative ideal of \mathfrak{g} . Then $\psi \neq 0$ on \mathfrak{r} .

Proof. Assume $\psi = 0$ on \mathfrak{r} . For $X \in \mathfrak{r}$, we have $2\eta(X, Y) = -\psi([JY, X]) = 0$ for all $Y \in \mathfrak{g}$. Since h is non-degenerate, we have $X \in \mathfrak{k}$ and hence $\mathfrak{r} \subset \mathfrak{k}$, which contradicts to Lemma 2.3. Q.E.D.

Lemma 2.5. $\operatorname{Tr}_{\mathfrak{g}/\mathfrak{k}} \operatorname{ad}(W) = 0$, for $W \in \mathfrak{k}$.

Proof. Using (1.4), (1.7), we have

$$\begin{aligned} \rho(W, [X, Y]) + \rho(X, [Y, W]) + \rho(Y, [W, X]) &= 0, \\ \rho(X, [Y, W]) + \rho(Y, [W, X]) &= 0, \end{aligned}$$

for $X, Y \in \mathfrak{g}$. Hence it follows that

$$\begin{aligned} \rho(JX, [W, Y]) + \rho([W, JX], Y) &= 0, \\ \rho(JX, [W, Y]) + \rho(J[W, X], Y) &= 0, \end{aligned}$$

for $X, Y \in \mathfrak{g}$. This implies that the endomorphism of $\mathfrak{g}/\mathfrak{k}$ which is induced by

$\text{ad}(W)$ is skew symmetric with respect to the inner product which is defined by $\rho(JX, Y)$. Therefore $\text{Tr}_{\mathfrak{g}/\mathfrak{k}} \text{ad}(W) = 0$. Q.E.D.

Lemma 2.6. *Let $\{E\}$ be a one dimensional ideal of \mathfrak{g} . Then $[E, W] = 0$, for $W \in \mathfrak{k}$. Moreover, there exists an endomorphism \tilde{J} of \mathfrak{g} such that $\tilde{J}X \equiv JX \pmod{\mathfrak{k}}$, $[\tilde{J}E, W] = 0$, for $X \in \mathfrak{g}$, $W \in \mathfrak{k}$.*

Proof. Put $[E, W] = \lambda E$, $\lambda \in \mathbf{R}$. Using (1.10), we have $0 = \psi([E, W]) = \lambda \psi(E)$. Since $\psi(E) \neq 0$ by Lemma 2.4, it follows $\lambda = 0$ and hence $[E, W] = 0$. Put $\mathfrak{h} = \mathfrak{k} + \{JE\}$, then $[\mathfrak{k}, \mathfrak{h}] \subset \mathfrak{h}$. Let $\{L\}$ be the orthogonal complement of \mathfrak{k} in \mathfrak{h} with respect to the $\text{ad}(\mathfrak{k})$ -invariant inner product on \mathfrak{g} . Then $[\mathfrak{k}, \{L\}] \subset \{L\}$. We may assume that $L = W_0 + JE$ where $W_0 \in \mathfrak{k}$. Therefore we can choose a linear endomorphism \tilde{J} on \mathfrak{g} such that $\tilde{J}E = L$, $\tilde{J}X \equiv JX \pmod{\mathfrak{k}}$ for $X \in \mathfrak{g}$. Then it follows

$$\begin{aligned} [\tilde{J}E, \mathfrak{k}] &= [L, \mathfrak{k}] \subset \{L\}, \\ [\tilde{J}E, \mathfrak{k}] &\subset [\mathfrak{h}, \mathfrak{k}] \subset \mathfrak{k}. \end{aligned}$$

This implies $[\tilde{J}E, W] = 0$ for $W \in \mathfrak{k}$.

Q.E.D

Therefore, for any one dimensional ideal $\{E\}$ of \mathfrak{g} , we may assume that $[JE, \mathfrak{k}] = \{0\}$.

3. We shall prove the following theorem.

Theorem 1'. *Let $(\mathfrak{g}, \mathfrak{k}, J, \rho)$ be the Kähler algebra of a homogeneous Kähler manifold G/K with non-degenerate canonical hermitian form h of signature $(2, 2(n-1))$. If there exists a one dimensional ideal \mathfrak{r} of \mathfrak{g} , then we have the following.*

- 1) *With suitable choice of $E \neq 0 \in \mathfrak{r}$, we have $[JE, E] = E$.*
- 2) *Put $\mathfrak{p} = \{P \in \mathfrak{g}; [P, E] = [JP, E] = 0\}$. Then we have the decomposition $\mathfrak{g} = \{JE\} + \{E\} + \mathfrak{p}$ of \mathfrak{g} into the direct sum of vector spaces. We know also that \mathfrak{p} is a compact semi-simple J -invariant ideal of \mathfrak{g} and that the real parts of the eigenvalues of $\text{ad}(JE)$ on \mathfrak{p} are equal to 0.*

The first part of the proof of Theorem 1' is nearly the same as the previous one [6]. But, for the sake of completeness we carry out the proof.

Lemma 3.1. *Let $\{E\}$ be a one dimensional ideal of \mathfrak{g} and put $\mathfrak{p} = \{P \in \mathfrak{g}; [P, E] = [JP, E] = 0\}$. Then we have*

- 1) $\mathfrak{k} \subset \mathfrak{p}$,
- 2) $J\mathfrak{p} \subset \mathfrak{p}$, $\text{ad}(JE)\mathfrak{p} \subset \mathfrak{p}$,
- 3) $\text{ad}(JE)J \equiv J\text{ad}(JE) \pmod{\mathfrak{k}}$ on \mathfrak{p} .

Proof. 1) follows from Lemma 2.6. For $P \in \mathfrak{p}$, we have

$$\begin{aligned} [JE, JP] &= J[JE, P] + J[E, JP] + [E, P] + W_0 \\ &= J[JE, P] + W_0 \end{aligned}$$

for some $W_0 \in \mathfrak{k}$, and hence 3) is proved. For $P \in \mathfrak{p}$, it follows that

$$\begin{aligned} [[JE, P], E] &= [[JE, E], P] + [JE, [P, E]] = 0, \\ [J[JE, P], E] &= [[JE, JP] - W_0, E] = 0, \end{aligned}$$

where $W_0 \in \mathfrak{k}$. Therefore $\text{ad}(JE)P \in \mathfrak{p}$ for all $P \in \mathfrak{p}$, which proves 2).

Lemma 3.2. *Let $\{E\}$ be a one dimensional ideal of \mathfrak{g} . Then $[JE, E] \neq 0$, therefore with suitable choice of $E \neq 0$, we have $[JE, E] = E$.*

Proof. Assume that $[JE, E] = 0$. For $X \in \mathfrak{g}$, we have $J[JE, X] = [JE, JX] - J[E, JX] - [E, X] + W_0 = [JE, JX] - \lambda JE - \mu E + W_0$, where $\lambda, \mu \in \mathbf{R}$, $W_0 \in \mathfrak{k}$. We have

$$\begin{aligned} [[JE, X], E] &= [[JE, E], X] + [JE, [X, E]] = 0, \\ [J[JE, X], E] &= [[JE, JX], E] - \lambda [JE, E] - \mu [E, E] \\ &\quad + [W_0, E] = 0, \end{aligned}$$

which implies that $[JE, X] \in \mathfrak{p}$, and hence we have

$$(3.1) \quad \text{ad}(JE)\mathfrak{g} \subset \mathfrak{p}.$$

Let $P \in \mathfrak{p}$. We have

$$\begin{aligned} \rho(JE, [JE, P]) &= \rho(-E, J[JE, P]) \\ &= -\rho(E, [JE, JP]) \\ &= \rho(JE, [JP, E]) + \rho(JP, [E, JE]) \\ &= 0, \end{aligned}$$

and it follows that for $X \in \mathfrak{g}$

$$(3.2) \quad \rho(JE, \text{ad}(JE)^2 X) = 0.$$

Applying Lemma 2.1, (3.2), we have for $X, Y \in \mathfrak{g}$

$$\begin{aligned} &\frac{d^3}{dt^3} \rho(\exp t \text{ad}(JE)X, \exp t \text{ad}(JE)Y) \\ &= \frac{d^2}{dt^2} \rho(JE, \exp t \text{ad}(JE)[X, Y]) \\ &= \rho(JE, \text{ad}(JE)^2 \exp t \text{ad}(JE)[X, Y]) \\ &= 0. \end{aligned}$$

Hence we may put

$$(3.3) \quad \begin{aligned} & \rho(\exp t \operatorname{ad}(JE)X, \exp t \operatorname{ad}(JE)Y) \\ &= at^2 + bt + c \end{aligned}$$

where a, b and c are real numbers not depending on t . Since $\operatorname{ad}(JE)\mathfrak{p} \subset \mathfrak{p}$, $\operatorname{ad}(JE)\mathfrak{k} = \{0\}$ by Lemma 2.6, (3.1), $\operatorname{ad}(JE)$ induces a linear endomorphism $\widetilde{\operatorname{ad}}(JE)$ on $\mathfrak{p}/\mathfrak{k}$. Let $\alpha + i\beta$ ($\alpha, \beta \in \mathbf{R}$) be an eigenvalue of $\widetilde{\operatorname{ad}}(JE)$. As $\operatorname{ad}(JE)J \equiv J \operatorname{ad}(JE) \pmod{\mathfrak{k}}$ on \mathfrak{p} , there exists an element $P \in \mathfrak{p}$, $P \notin \mathfrak{k}$ such that $[JE, P] \equiv (\alpha + \beta J)P \pmod{\mathfrak{k}}$, and hence $\exp t \operatorname{ad}(JE)P \equiv \exp t(\alpha + \beta J)P \pmod{\mathfrak{k}}$. Therefore we have by Lemma 3.1,

$$\begin{aligned} & \rho(\exp t \operatorname{ad}(JE)JP, \exp t \operatorname{ad}(JE)P) \\ &= \rho(J \exp t \operatorname{ad}(JE)P, \exp t \operatorname{ad}(JE)P) \\ &= \rho(J \exp t(\alpha + \beta J)P, \exp t(\alpha + \beta J)P) \\ &= \rho(\exp t(\alpha + \beta J)JP, \exp t(\alpha + \beta J)P) \\ &= e^{(\alpha + i\beta)t} \overline{e^{(\alpha + i\beta)t}} \rho(JP, P) \\ &= e^{2\alpha t} \rho(JP, P). \end{aligned}$$

From this and (3.3), we have

$$e^{2\alpha t} \rho(JP, P) = at^2 + bt + c.$$

Since $P \notin \mathfrak{k}$, $\rho(JP, P) > 0$ and hence $\alpha = 0$. This fact and $\operatorname{ad}(JE)\mathfrak{k} = \{0\}$ show that the real parts of the eigenvalues of $\operatorname{ad}(JE)$ on \mathfrak{p} are equal to 0. Therefore we have

$$\begin{aligned} \psi(E) &= \operatorname{Tr}_{\mathfrak{g}/\mathfrak{k}}(\operatorname{ad}(JE) - J \operatorname{ad}(E)) \\ &= \operatorname{Tr}_{\mathfrak{g}}(\operatorname{ad}(JE) - J \operatorname{ad}(E)) - \operatorname{Tr}_{\mathfrak{k}}(\operatorname{ad}(JE) - J \operatorname{ad}(E)) \\ &= \operatorname{Tr}_{\mathfrak{g}} \operatorname{ad}(JE) - \operatorname{Tr}_{\mathfrak{g}} J \operatorname{ad}(E) \\ &= \operatorname{Tr}_{\mathfrak{p}} \operatorname{ad}(JE) - \operatorname{Tr}_{(JE)} J \operatorname{ad}(E) \\ &= 0. \end{aligned}$$

However this contradicts to $\psi \neq 0$ on $\{E\}$ by Lemma 2.4.

Q.E.D.

Lemma 3.3 *Let $\{E\}$ be a one dimensional ideal of \mathfrak{g} . Then we get the decomposition*

$$\mathfrak{g} = \{JE\} + \{E\} + \mathfrak{p}$$

of \mathfrak{g} into the direct sum of vector spaces with the following properties:

- 1) $[JE, E] = E$.
- 2) The factors of the decomposition are mutually orthogonal with respect to the form η , and η is positive definite on $\{JE\} + \{E\}$.
- 3) The real parts of the eigenvalues of $\operatorname{ad}(JE)$ on \mathfrak{p} are equal to 0 or $1/2$.
- 4) $\rho(JE, P) = 0$ for $P \in \mathfrak{p}$.

Proof. By lemma 3.2, we may assume that E satisfies the condition $[JE, E]=E$. Since $\{E\}$ is a one dimensional ideal of \mathfrak{g} , we get $[X, E]=\alpha(X)E$, $[JX, E]=\beta(X)E$, for $X \in \mathfrak{g}$, where α, β are linear functions on \mathfrak{g} . It is easily seen that $P=X-\alpha(X)JE-\beta(X)E$ belongs to \mathfrak{p} for any $X \in \mathfrak{g}$. Therefore we have the decomposition $\mathfrak{g}=\{JE\}+\{E\}+\mathfrak{p}$. Now, by Lemma 2.1, we have for $P \in \mathfrak{p}$,

$$\begin{aligned} & \frac{d}{dt} \rho(\exp t \operatorname{ad}(JE)E, \exp t \operatorname{ad}(JE)P) \\ &= \rho(JE, \exp t \operatorname{ad}(JE)[E, P]) \\ &= 0. \end{aligned}$$

Since $\exp t \operatorname{ad}(JE)E=e^t E$, we have

$$\rho(E, \exp t \operatorname{ad}(JE)P) = a'e^{-t}$$

where a' is a constant determined by P and independent of t . We have then

$$\begin{aligned} \rho(JE, \exp t \operatorname{ad}(JE)P) &= -\rho(E, J \exp t \operatorname{ad}(JE)P) \\ &= -\rho(E, \exp t \operatorname{ad}(JE)JP) \\ &= ae^{-t} \end{aligned}$$

where a is a constant determined by JP . Let $X=\lambda JE+\mu E+P \in \mathfrak{g}$, where $\lambda, \mu \in \mathbf{R}, P \in \mathfrak{p}$. Then we have

$$\begin{aligned} \rho(JE, \exp t \operatorname{ad}(JE)X) &= \rho(JE, \lambda JE+\mu e^t E+\exp t \operatorname{ad}(JE)P) \\ &= \mu \rho(JE, E)e^t + \rho(JE, \exp t \operatorname{ad}(JE)P) \\ &= ae^{-t} + be^t \end{aligned}$$

where a, b are constants independent of t . This fact and Lemma 2.1 show that for $X, Y \in \mathfrak{g}$

$$\begin{aligned} & \frac{d}{dt} \rho(\exp t \operatorname{ad}(JE)X, \exp t \operatorname{ad}(JE)Y) \\ &= \rho(JE, \exp t \operatorname{ad}(JE)[X, Y]) \\ &= ae^{-t} + be^t. \end{aligned}$$

Hence we obtain

$$\begin{aligned} & \rho(\exp t \operatorname{ad}(JE)X, \exp t \operatorname{ad}(JE)Y) \\ &= ae^{-t} + be^t + c, \end{aligned}$$

where a, b and c are constants independent of t . Let $\alpha + i\beta$ be an eigenvalue of $\widetilde{\operatorname{ad}}(JE)$ on $\mathfrak{p}/\mathfrak{k}$. As $\operatorname{ad}(JE)J \equiv J \operatorname{ad}(JE) \pmod{\mathfrak{k}}$ on \mathfrak{p} , there exists an element $P \in \mathfrak{p}, P \notin \mathfrak{k}$ such that $\operatorname{ad}(JE)P \equiv (\alpha + \beta J)P \pmod{\mathfrak{k}}$. Hence we have

$$\begin{aligned}
 & \rho(\exp t \operatorname{ad}(JE)JP, \exp t \operatorname{ad}(JE)P) \\
 &= \rho(J \exp t \operatorname{ad}(JE)P, \exp t \operatorname{ad}(JE)P) \\
 &= \rho(J \exp t(\alpha + \beta J)P, \exp t(\alpha + \beta J)P) \\
 &= \rho(\exp t(\alpha + \beta J)JP, \exp t(\alpha + \beta J)P) \\
 &= e^{(\alpha + i\beta)t} \overline{e^{(\alpha + i\beta)t}} \rho(JP, P) \\
 &= e^{2\alpha t} \rho(JP, P).
 \end{aligned}$$

Therefore

$$(3.4) \quad e^{2\alpha t} \rho(JP, P) = ae^{-t} + be^t + c.$$

Since $P \notin \mathfrak{k}$ and $\rho(JP, P) > 0$, we have $\alpha = 0$ or $1/2$ or $-1/2$. Let \tilde{J} be the linear endomorphism of $\tilde{\mathfrak{p}} = \mathfrak{p}/\mathfrak{k}$ which is induced by J and put for $\alpha, \beta \in \mathbf{R}$;

$$\begin{aligned}
 \tilde{\mathfrak{p}}_{(\alpha + i\beta)} &= \{ \tilde{P} \in \tilde{\mathfrak{p}}; (\tilde{\operatorname{ad}}(JE) - (\alpha + \beta \tilde{J}))^m \tilde{P} = 0 \}, \\
 \tilde{\mathfrak{p}}_{\alpha} &= \sum_{\beta} \tilde{\mathfrak{p}}_{(\alpha + i\beta)}.
 \end{aligned}$$

Then we have

$$\tilde{\mathfrak{p}} = \sum_{\alpha + i\beta} \tilde{\mathfrak{p}}_{(\alpha + i\beta)}$$

where $\alpha = 0$ or $1/2$ or $-1/2$.

Let $\tilde{P} \neq 0 \in \tilde{\mathfrak{p}}_{(\alpha + i\beta)}$ and let $P \in \mathfrak{p}$ be a representation of \tilde{P} . Then there exists a positive integer m such that $(\tilde{\operatorname{ad}}(JE) - (\alpha + \beta \tilde{J}))^m \tilde{P} = 0$. Therefore we have

$$\begin{aligned}
 \exp t \tilde{\operatorname{ad}}(JE)\tilde{P} &= \exp t(\alpha + \beta \tilde{J}) \sum_{l=0}^{m-1} \frac{t^l}{l!} (\tilde{\operatorname{ad}}(JE) - (\alpha + \beta \tilde{J}))^l \tilde{P} \\
 &= e^{\alpha t} \left\{ \cos \beta t \sum_{l=0}^{m-1} \frac{t^l}{l!} (\tilde{\operatorname{ad}}(JE) - (\alpha + \beta \tilde{J}))^l \tilde{P} \right. \\
 &\quad \left. + \sin \beta t \sum_{l=0}^{m-1} \frac{t^l}{l!} (\tilde{\operatorname{ad}}(JE) - (\alpha + \beta \tilde{J}))^l \tilde{J}\tilde{P} \right\}.
 \end{aligned}$$

This shows that

$$\begin{aligned}
 \exp t \operatorname{ad}(JE)P &\equiv e^{\alpha t} \left\{ \cos \beta t \sum_{l=0}^{m-1} \frac{t^l}{l!} (\operatorname{ad}(JE) - (\alpha + \beta J))^l P \right. \\
 &\quad \left. + \sin \beta t \sum_{l=0}^{m-1} \frac{t^l}{l!} (\operatorname{ad}(JE) - (\alpha + \beta J))^l JP \right\} \pmod{\mathfrak{k}}.
 \end{aligned}$$

Hence we have

$$\begin{aligned}
 & \rho(JE, \exp t \operatorname{ad}(JE)P) \\
 &= e^{\alpha t} \left\{ \cos \beta t \sum_{l=0}^{m-1} \frac{1}{l!} \rho(JE, (\operatorname{ad}(JE) - (\alpha + \beta J))^l P) t^l \right. \\
 &\quad \left. + \sin \beta t \sum_{l=0}^{m-1} \frac{1}{l!} \rho(JE, (\operatorname{ad}(JE) - (\alpha + \beta J))^l JP) t^l \right\}.
 \end{aligned}$$

Put

$$h(t) = \sum_{l=0}^{m-1} \frac{1}{l!} \rho(JE, (\text{ad}(JE) - (\alpha + \beta J))^l P) t^l,$$

$$k(t) = \sum_{l=0}^{m-1} \frac{1}{l!} \rho(JE, (\text{ad}(JE) - (\alpha + \beta J))^l JP) t^l.$$

Then $h(t)$ and $k(t)$ are polynomials of degree $\leq m-1$. We have then

$$h(t) \cos \beta t + k(t) \sin \beta t = a e^{-(1+\alpha)t},$$

$$\left| \frac{h(t)}{t^m} \cos \beta t + \frac{k(t)}{t^m} \sin \beta t \right| = \left| a \frac{t^{-(1+\alpha)t}}{t^m} \right|.$$

Assume that $a \neq 0$. Since $1 + \alpha > 0$ and since $h(t)$ and $k(t)$ are polynomials of degree $\leq m-1$, the left side of the above formula approaches to 0 and the right side to ∞ , when $t \rightarrow -\infty$. This is a contradiction, and we get $a=0$, which implies that

$$\rho(JE, \exp t \text{ad}(JE) P) = 0$$

where P is a representative of $\tilde{P} \in \tilde{\mathfrak{p}}_{(\alpha + i\beta)}$. Thus we have

$$\rho(JE, \exp t \text{ad}(JE) P) = 0, \quad \text{for all } P \in \mathfrak{p},$$

and hence

$$\rho(JE, P) = 0, \quad \text{for all } P \in \mathfrak{p}.$$

Therefore 4) is proved. Moreover the formula (3.4) is reduced to

$$(3.4)' \quad e^{2\alpha t} \rho(JP, P) = b e^t + c.$$

This implies that $\alpha=0$ or $1/2$. Therefore we know that the real parts of the eigenvalues of $\text{ad}(JE)$ on \mathfrak{p} are equal to 0 or $1/2$. Thus the assertion 3) is proved. Now we shall show 2). The assertion that the decomposition $\mathfrak{g} = \{JE\} + \{E\} + \mathfrak{p}$ is an orthogonal decomposition is clear. Put $f = \text{ad}(JE) - J\text{ad}(E)$. Then we have $f(W)=0$ for $W \in \mathfrak{k}$, $f(JE)=JE$, $f(E)=E$ and $f(P)=[JE, P]$ for $P \in \mathfrak{p}$. Hence it follows that

$$\begin{aligned} \psi(E) &= \text{Tr}_{\mathfrak{g}/\mathfrak{k}}(\text{ad}(JE) - J\text{ad}(E)) \\ &= \text{Tr}_{\mathfrak{g}}(\text{ad}(JE) - J\text{ad}(E)) \\ &= 2 + \text{Tr}_{\mathfrak{p}} \text{ad}(JE) \\ &> 0. \end{aligned}$$

Therefore $2\eta(JE, JE) = 2\eta(E, E) = \psi(E) > 0$.

Q.E.D.

For $\alpha, \beta \in \mathbf{R}$, put

$$\begin{aligned} \mathfrak{p}_{(\alpha+i\beta)} &= \{P \in \mathfrak{p}; (\text{ad}(JE) - (\alpha+i\beta))^m P = 0\}, \\ \mathfrak{p}_\alpha &= \sum_{\beta} \mathfrak{p}_{(\alpha+i\beta)}, \end{aligned}$$

and let π' be the canonical projection from \mathfrak{g} onto $\mathfrak{g}/\mathfrak{k}$. Then we have

$$\begin{aligned} \bar{\mathfrak{p}}_{(\alpha+i\beta)} &= \pi'(\mathfrak{p}_{(\alpha+i\beta)}), \quad \bar{\mathfrak{p}}_\alpha = \pi'(\mathfrak{p}_\alpha), \\ \mathfrak{p} &= \mathfrak{p}_0 + \mathfrak{p}_\frac{1}{2}, \\ J\mathfrak{p}_\alpha &\subset \mathfrak{k} + \mathfrak{p}_\alpha, \\ \text{ad}(JE)\mathfrak{p}_\alpha &\subset \mathfrak{p}_\alpha. \end{aligned}$$

Lemma 3.4. *The form η is positive definite on $\mathfrak{p}_\frac{1}{2}$.*

Proof. We shall first prove that the decomposition $\mathfrak{p}_\frac{1}{2} = \sum_{\beta} \mathfrak{p}_{(\frac{1}{2}+i\beta)}$ is an orthogonal decomposition with respect to η . Let $P \neq 0 \in \mathfrak{p}_{(\frac{1}{2}+i\beta)}$, $Q \neq 0 \in \mathfrak{p}_{(\frac{1}{2}+i\beta')}$, and assume $\beta \neq \beta'$. Then we have

$$\begin{aligned} \exp t \text{ad}(JE)P &\equiv \exp t(1/2 + \beta J) \sum_{l=0}^{r-1} \frac{t^l}{l!} (\text{ad}(JE) - (1/2 + \beta J))^l P \pmod{\mathfrak{k}}, \\ \exp t \text{ad}(JE)Q &\equiv \exp (1/2 + \beta' J) \sum_{l=0}^{s-1} \frac{t^l}{l!} (\text{ad}(JE) - (1/2 + \beta' J))^l Q \pmod{\mathfrak{k}}. \end{aligned}$$

By Lemma 2.1, we have

$$\begin{aligned} (3.5) \quad & \frac{d}{dt} \rho(\exp t \text{ad}(JE)JP, \exp t \text{ad}(JE)Q) \\ &= \rho(JE, \exp t \text{ad}(JE)[JP, Q]). \end{aligned}$$

The left side of this equation is equal to

$$\begin{aligned} & \frac{d}{dt} \rho(J \exp t \text{ad}(JE)P, \exp t \text{ad}(JE)Q) \\ &= \frac{d}{dt} \rho(J \exp t(1/2 + \beta J) \sum_{l=0}^{r-1} \frac{t^l}{l!} (\text{ad}(JE) - (1/2 + \beta J))^l P, \\ & \quad \exp t(1/2 + \beta' J) \sum_{l=0}^{s-1} \frac{t^l}{l!} (\text{ad}(JE) - (1/2 + \beta' J))^l Q) \\ &= \frac{d}{dt} e^t \rho(\exp \beta t J \sum_{l=0}^{r-1} \frac{t^l}{l!} (\text{ad}(JE) - (1/2 + \beta J))^l JP, \\ & \quad \exp \beta' t J \sum_{l=0}^{s-1} \frac{t^l}{l!} (\text{ad}(JE) - (1/2 + \beta' J))^l Q) \\ &= \frac{d}{dt} e^t \rho(\{\cos \beta t + (\sin \beta t)J\}u(t), \{\cos \beta' t + (\sin \beta' t)J\}v(t)) \end{aligned}$$

$$\begin{aligned}
 &= \frac{d}{dt} e^t \{(\cos \beta t \cos \beta' t + \sin \beta t \sin \beta' t) \rho(u(t), v(t)) \\
 &\quad + (\sin \beta t \cos \beta' t - \cos \beta t \sin \beta' t) \rho(Ju(t), v(t))\} \\
 &= \frac{d}{dt} e^t \{h(t) \cos(\beta - \beta')t + k(t) \sin(\beta - \beta')t\} \\
 &= e^t \{a(t) \cos(\beta - \beta')t + b(t) \sin(\beta - \beta')t\},
 \end{aligned}$$

where

$$\begin{aligned}
 u(t) &= \sum_{i=0}^{r-1} \frac{t^i}{i!} (\text{ad}(JE) - (1/2 + \beta J)^i) JP, \\
 v(t) &= \sum_{i=0}^{s-1} \frac{t^i}{i!} (\text{ad}(JE) - (1/2 + \beta' J)^i) Q, \\
 h(t) &= \rho(u(t), v(t)), \quad k(t) = \rho(Ju(t), v(t)), \\
 a(t) &= h(t) + h'(t) + (\beta - \beta')k(t), \\
 b(t) &= k(t) + k'(t) - (\beta - \beta')h(t).
 \end{aligned}$$

Hence $a(t)$ and $b(t)$ are polynomials. Since $[JP, Q] \in [\mathfrak{k} + \mathfrak{p}_1, \mathfrak{p}_2] \subset \{E\} + \mathfrak{p}_2$, we put $[JP, Q] = \lambda E + P'$, where $\lambda \in \mathbf{R}, P' \in \mathfrak{p}_2$. Using Lemma 3.3; 4), the right side of the equation (3.5) is equal to

$$\begin{aligned}
 &\rho(JE, \exp t \text{ad}(JE)[JP, Q]) \\
 &= \rho(JE, \exp t \text{ad}(JE)(\lambda E + P')) \\
 &= e^t \lambda \rho(JE, E) + \rho(JE, \exp t \text{ad}(JE)P') \\
 &= e^t \lambda \rho(JE, E).
 \end{aligned}$$

Therefore we have

$$a(t) \cos(\beta - \beta')t + b(t) \sin(\beta - \beta')t = \lambda \rho(JE, E).$$

Since $a(t) - \lambda \rho(JE, E)$ is a polynomial and since $a(t_n) - \lambda \rho(JE, E) = 0$ for $t_n = \frac{2n\pi}{\beta - \beta'}$, where n integer, it follows that $a(t)$ is a constant a . Similarly $b(t)$ is a constant b . Hence we have

$$a \cos(\beta - \beta')t + b \sin(\beta - \beta')t = \lambda \rho(JE, E).$$

By this formula, we have $(\beta - \beta')^2 \lambda \rho(JE, E) = 0$. Since $\beta - \beta' \neq 0$ and $\rho(JE, E) > 0$, we get $\lambda = 0$. Moreover $\text{ad}(JE)$ is non-singular on \mathfrak{p}_2 and so there exists an element $P'' \in \mathfrak{p}_2$ such that $P' = [JE, P'']$. Thus we have

$$\begin{aligned}
 2\eta(P, Q) &= \psi([JP, Q]) \\
 &= \psi(P') \\
 &= \psi([JE, P'']) \\
 &= -\psi([E, JP'']) \\
 &= 0.
 \end{aligned}$$

This shows that $\mathfrak{p}_{(1/2+i\beta)}$ and $\mathfrak{p}_{(1/2+i\beta')}$ are mutually orthogonal with respect to η . Now, let $P \neq 0 \in \mathfrak{p}_{(1/2+i\beta)}$. Then we have

$$\exp t \operatorname{ad}(JE)P \equiv \exp t(1/2 + \beta J)u(t) \pmod{\mathfrak{k}},$$

where $u(t) = \sum_{i=0}^{m-1} \frac{t^i}{i!} (\operatorname{ad}(JE) - (1/2 + \beta J))^i P$. By Lemma 2.1, it follows that

$$\begin{aligned}
 (3.6) \quad & \frac{d}{dt} \rho(\exp t \operatorname{ad}(JE)JP, \exp t \operatorname{ad}(JE)P) \\
 &= \rho(JE, \exp t \operatorname{ad}(JE)[JP, P])
 \end{aligned}$$

The left side of the equation (3.6) is equal to

$$\begin{aligned}
 & \frac{d}{dt} \rho(J \exp t \operatorname{ad}(JE)P, \exp t \operatorname{ad}(JE)P) \\
 &= \frac{d}{dt} \rho(J \exp t(1/2 + \beta J)u(t), \exp t(1/2 + \beta J)u(t)) \\
 &= \frac{d}{dt} \rho(\exp t(1/2 + \beta J)Ju(t), \exp t(1/2 + \beta J)u(t)) \\
 &= \frac{d}{dt} e^{c(1/2+i\beta)t} e^{c(1/2+i\beta')t} \rho(Ju(t), u(t)) \\
 &= \frac{d}{dt} e^t \rho(Ju(t), u(t)) \\
 &= e^t (h'(t) + h(t))
 \end{aligned}$$

where $h(t) = \rho(Ju(t), u(t))$, and $h(t)$ is a polynomial of degree $\leq 2m - 2$. Since $[JP, P] = \lambda E + P'$, where $\lambda \in \mathbf{R}$, $P' \in \mathfrak{p}_2$, the right side of the equation (3.6) is equal to

$$\begin{aligned}
 & \rho(JE, \exp t \operatorname{ad}(JE)(\lambda E + P')) \\
 &= e^t \lambda \rho(JE, E) + \rho(JE, \exp t \operatorname{ad}(JE)P') \\
 &= e^t \lambda \rho(JE, E).
 \end{aligned}$$

Hence we have

$$h'(t) + h(t) = \lambda \rho(JE, E).$$

The solution of this equation is $h(t) = ce^{-t} + \lambda \rho(JE, E)$, where c is an arbitrary

constant. However, $h(t)$ is a polynomial and so $c=0$. Hence we have

$$h(t) = \lambda\rho(JE, E),$$

and hence it follows that

$$\lambda = \frac{h(t)}{\rho(JE, E)} = \frac{h(0)}{\rho(JE, E)} = \frac{\rho(JP, P)}{\rho(JE, E)} > 0.$$

Therefore we have

$$\begin{aligned} 2\eta(P, P) &= \psi([JP, P]) \\ &= \lambda\psi(E) + \psi(P') \\ &= \lambda\psi(E) > 0. \end{aligned}$$

This shows that η is positive definite on $\mathfrak{p}_{(c\frac{1}{2}+i\beta)}$ and hence on $\mathfrak{p}_i = \sum_{\beta} \mathfrak{p}_{(c\frac{1}{2}+i\beta)}$.

Q.E.D.

Proof of Theorem 1'. Since η is positive definite on $\{JE\} + \{E\} + \mathfrak{p}_i$ and since the signature of h is $(2, 2(n-1))$, we have $\mathfrak{p}_i = \{0\}$, and hence $\mathfrak{p} = \mathfrak{p}_0$. Let $P, Q \in \mathfrak{p}$. Since $[P, Q] \in [\mathfrak{g}_0, \mathfrak{g}_0] \subset \mathfrak{g}_0$, where $\mathfrak{g}_0 = \{JE\} + \mathfrak{p}$, we put $[P, Q] = \lambda JE + P'$, where $\lambda \in \mathbf{R}, P' \in \mathfrak{p}$. It follows that $[E, [P, Q]] = [E, \lambda JE + P'] = -\lambda E$ and $[E, [P, Q]] = [[E, P], Q] + [P, [E, Q]] = 0$. This implies that $\lambda = 0$ and $[P, Q] \in \mathfrak{p}$. Therefore \mathfrak{p} is a subalgebra of \mathfrak{g} and also an ideal of \mathfrak{g} . Moreover we see easily that $(\mathfrak{p}, \mathfrak{k}, J, \rho)$ is an effective Kähler algebra. Since the decomposition $\mathfrak{g} = \{JE\} + \{E\} + \mathfrak{p}$ is orthogonal with respect to η and η is positive definite on $\{JE\} + \{E\}$ and since the signature of h is $(2, 2(n-1))$, we know that $\eta(P, P) < 0$, for $P \in \mathfrak{p}, P \notin \mathfrak{k}$. Now, for $P, Q \in \mathfrak{p}$, put

$$\begin{aligned} \psi'(P) &= \text{Tr}_{\mathfrak{p}/\mathfrak{k}}(\text{ad}(JP) - J\text{ad}(P)), \\ 2\eta'(P, Q) &= \psi'([JP, Q]). \end{aligned}$$

For $P \in \mathfrak{p}, P \notin \mathfrak{k}$, we have $(\text{ad}(JP) - J\text{ad}(P))E = 0, (\text{ad}(JP) - J\text{ad}(P))JE \equiv 0 \pmod{\mathfrak{k}}$ and hence $\psi'(P) = \psi'(P)$. This implies that

$$\begin{aligned} 2\eta'(P, P) &= \psi'([JP, P]) \\ &= \psi([JP, P]) \\ &= 2\eta(P, P) < 0, \end{aligned}$$

which proves that the canonical hermitian form of $(\mathfrak{p}, \mathfrak{k}, J, \rho)$ is negative definite. Therefore we know that \mathfrak{p} is a compact semi-simple subalgebra of \mathfrak{g} [5].

Q.E.D.

Proof of Theorem 1. When G is a semi-simple Lie group, our assertion follows from the results of Borel [1] and Koszul [3]. We shall show the case where G contains a one parameter normal subgroup of G . Let $\{E\}$ be the ideal

of \mathfrak{g} corresponding to the one parameter subgroup. With appropriate choice of J , we may assume that $J^2E = -E$. Put $\mathfrak{g}' = \{JE\} + \{E\}$. Then (\mathfrak{g}', J, ρ) is a Kähler algebra of the unit disk $\{z \in \mathbb{C}; |z| < 1\}$. Now, for $X, Y \in \mathfrak{g}$, we define

$$\bar{\rho}(X, Y) = \rho(X', Y')$$

where X', Y' are the \mathfrak{g}' -components of X, Y with respect to the decomposition $\mathfrak{g} = \mathfrak{g}' + \mathfrak{p}$ respectively. Then $(\mathfrak{g}, \mathfrak{p}, J, \bar{\rho})$ is a Kähler algebra. We denote by G' (resp. P) the connected subgroup of G corresponding to \mathfrak{g}' (resp. \mathfrak{p}). Since $(\mathfrak{g}, \mathfrak{p}, J, \bar{\rho})$ is a Kähler algebra, G/P admits an invariant Kähler structure and is holomorphically isomorphic to the G' -orbit passing through the origin o . We know by Theorem 1' that G/K is a holomorphic fibre bundle whose base space is $G/P \cong \{z \in \mathbb{C}; |z| < 1\}$, and whose fibre is P/K . Q.E.D.

4. Proof of Theorem 2

Let $(\mathfrak{g}, \mathfrak{k}, J, \rho)$ be the Kähler algebra of G/K . We show that, if \mathfrak{g} is not semi-simple, then there exists a one dimensional ideal of \mathfrak{g} . Assume that \mathfrak{g} is not semi-simple. Then there exists a non-zero commutative ideal \mathfrak{r} of \mathfrak{g} . Consider a J -invariant subalgebra $\mathfrak{g}' = \mathfrak{k} + J\mathfrak{r} + \mathfrak{r}$. Then we have

Lemma 4.1. $\dim_{\mathbb{C}} \mathfrak{g}'/\mathfrak{k} = 1$.

Since $\mathfrak{k} \cap \mathfrak{r} = \{0\}$ by Lemma 2.3 and $\dim_{\mathbb{C}} \mathfrak{g}/\mathfrak{k} = 2, \dim_{\mathbb{C}} \mathfrak{g}'/\mathfrak{k} = 1$ or 2 . Suppose that $\dim_{\mathbb{C}} \mathfrak{g}'/\mathfrak{k} = 2$. Then $\dim_{\mathbb{C}} \mathfrak{g}/\mathfrak{k} = \dim_{\mathbb{C}} \mathfrak{g}'/\mathfrak{k}, \dim \mathfrak{g} = \dim \mathfrak{g}'$ and hence $\mathfrak{g} = \mathfrak{k} + J\mathfrak{r} + \mathfrak{r}$. Since $\dim \mathfrak{g}'/\mathfrak{k} = 4$ and $\mathfrak{k} \cap \mathfrak{r} = \{0\}$, we have $2 \leq \dim \mathfrak{r} \leq 4$. Let π' be the projection from \mathfrak{g} onto $\mathfrak{g}/\mathfrak{k}$. Then it follows that

$$\begin{aligned} \dim \pi'(J\mathfrak{r}) \cap \pi'(\mathfrak{r}) &= \dim \pi'(J\mathfrak{r}) + \dim \pi'(\mathfrak{r}) - \dim \pi'(\mathfrak{g}) \\ &= 2 \dim \mathfrak{r} - 4. \end{aligned}$$

First, we shall show $\dim \mathfrak{r} \neq 3, 4$. Suppose $\dim \mathfrak{r} = 3$ or 4 . Then $\dim \pi'(J\mathfrak{r}) \cap \pi'(\mathfrak{r}) > 0$, and so there exist $A \neq 0, B \neq 0 \in \mathfrak{r}$ and $W \in \mathfrak{k}$ such that $JA = B + W$. Therefore we have $2\eta(A, C) = \psi([JA, C]) = \psi([B + W, C]) = 0$ and $2\eta(A, JC) = \psi([JA, JC]) = \psi([A, C]) = 0$ for all $C \in \mathfrak{r}$. Since $\mathfrak{g} = \mathfrak{k} + J\mathfrak{r} + \mathfrak{r}$, we know $\eta(A, X) = 0$ for all $X \in \mathfrak{g}$. This implies $A \in \mathfrak{k}$, which is a contradiction to Lemma 2.3. Next, we shall prove $\dim \mathfrak{r} \neq 2$. Suppose $\dim \mathfrak{r} = 2$. Then $\dim \pi'(J\mathfrak{r}) \cap \pi'(\mathfrak{r}) = 0$, and hence $\mathfrak{g} = \mathfrak{k} + J\mathfrak{r} + \mathfrak{r}$ is a direct sum as vector spaces. Let A be an element in \mathfrak{r} such that $\eta(A, B) = 0$ for all $B \in \mathfrak{r}$. Since $\mathfrak{g} = \mathfrak{k} + J\mathfrak{r} + \mathfrak{r}$ and $2\eta(A, JB) = \psi([JA, JB]) = \psi([A, B]) = 0$, we have $\eta(A, X) = 0$ for any $X \in \mathfrak{g}$, which implies $A \in \mathfrak{k}$, and hence $A = 0$ by Lemma 2.3. This shows that η is non-degenerate on \mathfrak{r} . Therefore there exists a unique non-zero element $E \in \mathfrak{r}$ such that $2\eta(E, A) = \psi(A)$ for all $A \in \mathfrak{r}$. We have then

$$(4.1) \quad \begin{aligned} [JE, E] &= E, \\ \psi(E) &\neq 0. \end{aligned}$$

Indeed, for $A \in \mathfrak{r}$ we have

$$\begin{aligned} 2\eta([JE, E], A) &= \psi([J[JE, E], A]) \\ &= -\psi([[JE, E], JA]) \\ &= \psi([[E, JA], JE]) + \psi([[JA, JE], E]) \\ &= -\psi([E, JA]) + \psi([J[JA, E] + J[A, JE], E]) \\ &= \psi(A) + \psi([JA, E]) + \psi([A, JE]) \\ &= \psi(A). \end{aligned}$$

This shows that $[JE, E] = E$. Let F be an element in \mathfrak{r} independent of E . Put $[JE, F] = \lambda E + \mu F$, where $\lambda, \mu \in \mathbf{R}$. Then $\psi(E) = \text{Tr}_{\mathfrak{g}/\mathfrak{t}}(\text{ad}(JE) - J\text{ad}(E)) = 2(1 + \mu)$. We shall show $\psi(E) \neq 0$. Suppose $\psi(E) = 0$. Then $\mu = -1$ and $\psi(F) = 2\eta(E, F) = \psi([JE, F]) = \lambda\psi(E) - \psi(F) = -\psi(F)$. Therefore $\psi(F) = 0$, and hence $\psi = 0$ on \mathfrak{r} , which is a contradiction to Lemma 2.4.

(4.2) There exists an element F in \mathfrak{r} independent of E such that

$$\begin{aligned} [JE, E] &= E, & [JE, F] &= \alpha F, \\ [JF, E] &= \beta F, & [JF, F] &= -E, \\ \psi(F) &= 0, \end{aligned}$$

where $\alpha, \beta \in \mathbf{R}$.

Proof. By $2\eta(E, E) = \psi(E) \neq 0$, there exists $\tilde{F} \neq 0$ such that $2\eta(E, \tilde{F}) = \psi(\tilde{F}) = 0$. Since η is non-degenerate on \mathfrak{r} and the signature is $(1, 1)$, we have $\eta(E, E)\eta(\tilde{F}, \tilde{F}) < 0$. Put $[JE, \tilde{F}] = \alpha\tilde{F} + \alpha'E$, where $\alpha, \alpha' \in \mathbf{R}$. We have then $0 = \psi(\tilde{F}) = 2\eta(E, \tilde{F}) = \psi([JE, \tilde{F}]) = \alpha\psi(\tilde{F}) + \alpha'\psi(E) = \alpha'\psi(E)$, and hence $\alpha' = 0$, and $[JE, \tilde{F}] = \alpha\tilde{F}$. Similarly we have $[J\tilde{F}, E] = \beta\tilde{F}$. Now, we put $[J\tilde{F}, \tilde{F}] = \gamma E + \delta\tilde{F}$, where $\gamma, \delta \in \mathbf{R}$. Then we have $0 = \psi(\tilde{F}) = \text{Tr}_{\mathfrak{g}/\mathfrak{t}}(\text{ad}(J\tilde{F}) - J\text{ad}(\tilde{F})) = 2\delta$ and so $[J\tilde{F}, \tilde{F}] = \gamma E$. Since $2\eta(\tilde{F}, \tilde{F}) = \psi([J\tilde{F}, \tilde{F}]) = \gamma\psi(E) = 2\gamma\eta(E, E)$, it follows $\gamma = \frac{\eta(\tilde{F}, \tilde{F})}{\eta(E, E)} < 0$. Putting $F = \frac{1}{\sqrt{-\gamma}}\tilde{F}$, we have $[JF, F] = -E$.

Q.E.D.

$$(4.3) \quad \mathfrak{k} = \{0\}.$$

Proof. For $W \in \mathfrak{k}$, put $[W, E] = \lambda E + \mu F$. Since $0 = \psi([W, E]) = \lambda\psi(E) + \mu\psi(F) = \lambda\psi(E)$, $\lambda = 0$ and hence $[W, E] = \mu F$. We have $\psi([JF, [W, E]]) = -\mu\psi(E)$ and $\psi([JF, [W, E]]) = \psi([[JF, W], E]) + \psi([W, [JF, E]]) = \psi([J[F, W], E]) = \psi([JE, [F, W]]) = \psi([F, W]) = 0$. Thus $\mu = 0$ and $[W, E] = 0$. Now, put $[W, F] = \lambda E + \mu F$. By $0 = \psi([W, F]) = \lambda\psi(E) + \mu\psi(F) = \lambda\psi(E)$,

we have $\lambda=0$ and $[W, F]=\mu F$. Hence it follows $\psi([JF, [W, F]])=-\mu\psi(E)$. On the other hand we have $\psi([JF, [W, F]])=\psi([[JF, W], F])+\psi([W, [JF, F]])=\psi([J[F, W], F])-\mu\psi([JF, F])=\mu\psi(E)$. Therefore $2\mu\psi(E)=0$, and hence $\mu=0, [W, F]=0$. Thus $[\mathfrak{k}, \mathfrak{r}]=0$. Since $[\mathfrak{k}, J\mathfrak{r}]\subset \mathfrak{k}, [\mathfrak{k}, \mathfrak{k}]\subset \mathfrak{k}$ and $\mathfrak{g}=\mathfrak{k}+J\mathfrak{r}+\mathfrak{r}$, we know that \mathfrak{k} is an ideal of \mathfrak{g} . By the effectiveness, we have $\mathfrak{k}=\{0\}$.

Q.E.D.

$$(4.4) \quad 2\alpha = \beta + 1 .$$

Proof. Using Jacobi identity and (1.3), we have

$$\begin{aligned} 0 &= [[JE, JF], F] + [[JF, F], JE] + [[F, JE], JF] \\ &= [[J[JE, F] + J[E, JF], F] + [[JF, F], JE] + [[F, JE], JE] \\ &= (\alpha - \beta)[JF, F] - [E, JE] - \alpha[F, JF] \\ &= (-2\alpha + \beta + 1)E . \end{aligned}$$

Hence it follows $2\alpha = \beta + 1$.

Q.E.D.

By (1.7), (4.2) and (4.4), we have

$$\begin{aligned} 0 &= \rho([JE, F], JF) + \rho([F, JF], JE) + \rho([JF, JE], F) \\ &= \alpha\rho(F, JF) + \rho(E, JE) - (\alpha - \beta)\rho(JF, F) \\ &= (-2\alpha + \beta)\rho(JF, F) - \rho(JE, E) \\ &= -\rho(JF, F) - \rho(JE, E) . \end{aligned}$$

This contradicts to $\rho(JE, E) > 0, \rho(JF, F) > 0$. Therefore $\dim \mathfrak{r} \neq 2$ and hence $\dim_{\mathbb{C}} \mathfrak{g}'/\mathfrak{k} \neq 2$. Thus we have proved $\dim_{\mathbb{C}} \mathfrak{g}'/\mathfrak{k} = 1$, this completes the proof of Lemma 4.1.

Q.E.D.

Let $\mathfrak{r} \neq \{0\}$ be a commutative ideal of \mathfrak{g} . Since $\dim \mathfrak{g}'/\mathfrak{k} = 2$ by Lemma 4.1 and $\mathfrak{k} \cap \mathfrak{r} = \{0\}$, it follows that $\dim \mathfrak{r} = 1$ or 2 . Assume $\dim \mathfrak{r} = 2$. Then we have

$$\begin{aligned} \dim \pi'(J\mathfrak{r}) \cap \pi'(\mathfrak{r}) &= \dim \pi'(J\mathfrak{r}) + \dim \pi'(\mathfrak{r}) - \dim (\pi'(J\mathfrak{r}) + \pi'(\mathfrak{r})) \\ &= 2 \dim \mathfrak{r} - 2 \\ &= 2 . \end{aligned}$$

This implies $\pi'(J\mathfrak{r}) = \pi'(\mathfrak{r})$ and hence $J\mathfrak{r} \subset \mathfrak{k} + \mathfrak{r}$. For any $A \in \mathfrak{r}$, we have $JA = A' + W$, where $A' \in \mathfrak{r}, W \in \mathfrak{k}$. It follows then

$$\begin{aligned} \psi(A) &= Tr_{\mathfrak{g}/\mathfrak{k}}(\text{ad}(JA) - J\text{ad}(A)) \\ &= Tr_{\mathfrak{g}/\mathfrak{k}}(\text{ad}(A') - J\text{ad}(A)) + Tr_{\mathfrak{g}/\mathfrak{k}}\text{ad}(W) \\ &= Tr_{\mathfrak{r}+\mathfrak{k}/\mathfrak{k}}(\text{ad}(A') - J\text{ad}(A)) \\ &= 0 . \end{aligned}$$

Hence $\psi=0$ on \mathfrak{r} , which is a contradiction to Lemma 2.4. Thus \mathfrak{r} is a one dimensional ideal of \mathfrak{g} . Therefore Theorem 2 is proved. Q.E.D.

5. We shall classify two dimensional connected simply connected homogeneous Kähler manifolds with non-degenerate canonical hermitian form h . The signature of h is $(4, 0)$ or $(2, 2)$ or $(0, 4)$.

(i) The case $(4, 0)$. Since h is positive definite, G/K is isomorphic to a homogeneous bounded domain. Hence G/K is either $\{z \in \mathbf{C}; |z| < 1\} \times \{z \in \mathbf{C}; |z| < 1\}$ or $\{(z_1, z_2) \in \mathbf{C}^2; |z_1|^2 + |z_2|^2 < 1\}$.

(ii) The case $(0, 4)$. Since h is negative definite, G is a compact semi-simple Lie group by [5]. By a theorem in [4], G/K is a hermitian symmetric space. Hence G/K is either $P_1(\mathbf{C}) \times P_1(\mathbf{C})$ or $P_2(\mathbf{C})$, where $P_n(\mathbf{C})$ is a complex n -dimensional projective space.

(iii) The case $(2, 2)$. Applying Theorem 1 and 2, we obtain that G/K is a holomorphic fibre bundle whose base space is the unit disk $\{z \in \mathbf{C}; |z| < 1\}$, and whose fibre is $P_1(\mathbf{C})$.

YAMAGUCHI UNIVERSITY

References

- [1] A. Borel: *Kählerian coset spaces of semi-simple Lie groups*, Proc. Nat. Acad. Sci. USA **40** (1954), 1147–1151.
- [2] S.G. Gindikin, I.I. Pjateckii-Sapiro, E.B. Vinberg: *Homogeneous Kähler manifolds*, "Geometry of Homogeneous Bounded Domains", Centro Int. Mat. Estivo, 3° Ciclo, Urbino, 1967.
- [3] J.L. Koszul: *Sur la forme hermitienne canonique des espaces homogènes complexes*, Canad. J. Math. **7** (1955), 562–576.
- [4] A. Lichnerowicz: *Espaces homogènes Kählériens*, Colloque Géométrie Différentielle, Strasbourg, 1953, 171–201.
- [5] H. Shima: *On homogeneous complex manifolds with negative definite canonical hermitian form*, Proc. Japan Acad. **46** (1970), 209–211.
- [6] H. Shima: *On homogeneous Kähler manifolds of solvable Lie groups*, J. Math. Soc. Japan **25** (1973), 422–445.

